



Universiteit
Leiden
The Netherlands

Control of sporulation-specific cell division in *Streptomyces coelicolor*

Noens, E.

Citation

Noens, E. (2007, September 25). *Control of sporulation-specific cell division in Streptomyces coelicolor*. Department Microbial Development (LIC) Department Electron Microscopy (LUMC/MCB), Leiden University. Retrieved from <https://hdl.handle.net/1887/12351>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12351>

Note: To cite this publication please use the final published version (if applicable).

REFERENCES

- Addinall, S.G., Cao, C., and Lutkenhaus, J. (1997) FtsN, a late recruit to the septum in *Escherichia coli*. *Mol Microbiol* **25**: 303-309.
- Ainsa, J.A., Parry, H.D., and Chater, K.F. (1999) A response regulator-like protein that functions at an intermediate stage of sporulation in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **34**: 607-619.
- Ainsa, J.A., Ryding, N.J., Hartley, N., Findlay, K.C., Bruton, C.J., and Chater, K.F. (2000) WhiA, a protein of unknown function conserved among gram-positive bacteria, is essential for sporulation in *Streptomyces coelicolor* A3(2). *J Bacteriol* **182**: 5470-5478.
- Aretz W, Meiwes J, Seibert G, Vobis G, Wink J. (2000) Friulimicins: novel lipopeptide antibiotics with peptidoglycan synthesis inhibiting activity from Actinoplanes friuliensis sp. nov. I. Taxonomic studies of the producing microorganism and fermentation. *J Antibiot (Tokyo)* **53**:807-15.
- Ausmees, N., Kuhn, J.R., and Jacobs-Wagner, C. (2003) The bacterial cytoskeleton: an intermediate filament-like function in cell shape. *Cell* **115**: 705-713.
- Autret, S., and Errington, J. (2001) Dynamic proteins in bacteria. *Dev Cell* **1**: 10-11.
- Barilla, D., Rosenberg, M.F., Nobmann, U., and Hayes, F. (2005) Bacterial DNA segregation dynamics mediated by the polymerizing protein ParF. *Embo J* **24**: 1453-1464.
- Bath, J., Wu, L.J., Errington, J., and Wang, J.C. (2000) Role of *Bacillus subtilis* SpoIIIE in DNA transport across the mother cell-prespore division septum. *Science* **290**: 995-997.
- Bennett, J.A., and McCormick, J.R. (2001) Two new loci affecting cell division identified as suppressors of an *ftsQ*-null mutation in *Streptomyces coelicolor* A3(2). *FEMS Microbiol Lett* **202**: 251-256.
- Bentley, S.D., Chater, K.F., Cerdeno-Tarraga, A.M., Challis, G.L., Thomson, N.R., James, K.D., Harris, D.E., Quail, M.A., Kieser, H., Harper, D., Bateman, A., Brown, S., Chandra, G., Chen, C.W., Collins, M., Cronin, A., Fraser, A., Goble, A., Hidalgo, J., Hornsby, T., Howarth, S., Huang, C.H., Kieser, T., Larke, L., Murphy, L., Oliver, K., O'Neil, S., Rabinowitsch, E., Rajandream, M.A., Rutherford, K., Rutter, S., Seeger, K., Saunders, D., Sharp, S., Squares, R., Squares, S., Taylor, K., Warren, T., Wietzorek, A., Woodward, J., Barrell, B.G., Parkhill, J., and Hopwood, D.A. (2002) Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* **417**: 141-147.
- Bernhardt, T.G., and de Boer, P.A. (2003) The *Escherichia coli* amidase AmiC is a periplasmic septal ring component exported via the twin-arginine transport pathway. *Mol Microbiol* **48**: 1171-1182.
- Bernhardt, T.G., and de Boer, P.A. (2004) Screening for synthetic lethal mutants in *Escherichia coli* and identification of EnvC (YibP) as a periplasmic septal ring factor with murein hydrolase activity. *Mol Microbiol* **52**: 1255-1269.
- Bernhardt, T.G., and de Boer, P.A. (2005) SlmA, a nucleoid-associated, FtsZ binding protein required for blocking septal ring assembly over chromosomes in *E. coli*. *Mol Cell* **18**: 555-564.
- Bibb, M. (1996) 1995 Colworth Prize Lecture. The regulation of antibiotic production in *Streptomyces coelicolor* A3(2). *Microbiology* **142**: 1335-1344.
- Bibb, M.J., Molle, V., and Buttner, M.J. (2000) sigma(BldN), an extracytoplasmic function RNA polymerase sigma factor required for aerial mycelium formation in *Streptomyces coelicolor* A3(2). *J. Bacteriol.* **182**: 4606-4616.
- Bierman, M., Logan, R., O'Brien, K., Seno, E.T., Rao, R.N., and Schoner, B.E. (1992) Plasmid cloning vectors for the conjugal transfer of DNA from *Escherichia coli* to *Streptomyces* spp. *Gene* **116**: 43-49.
- Bignell, D.R., Warawa, J.L., Strap, J.L., Chater, K.F., and Leskiw, B.K. (2000) Study of the bldG locus suggests that an anti-anti-sigma factor and an anti-sigma factor may be involved in *Streptomyces coelicolor* antibiotic production and sporulation. *Microbiology* **146**: 2161-2173.
- Bigot, S., Corre, J., Louarn, J.M., Cornet, F., and Barre, F.X. (2004) FtsK activities in Xer recombination, DNA mobilization and cell division involve overlapping and separate domains of the protein. *Mol Microbiol* **54**: 876-886.
- Bishop, A., Fielding, S., Dyson, P., and Herron, P. (2004) Systematic insertional mutagenesis of a streptomycete genome: a link between osmoadaptation and antibiotic production. *Genome Res* **14**: 893-900.
- Blondelet-Rouault, M.H., Weiser, J., Lebrhi, A., Branny, P., and Pernodet, J.L. (1997) Antibiotic resistance gene cassettes derived from the omega interposon for use in *E. coli* and *Streptomyces*. *Gene* **190**: 315-317.
- Bolhuis, A., Broekhuizen, C.P., Sorokin, A., van Roosmalen, M.L., Venema, G., Bron, S., Quax, W.J., and van Dijl, J.M. (1998) SecDF of *Bacillus subtilis*, a molecular Siamese twin required for the efficient secretion of proteins. *J Biol Chem* **273**: 21217-21224.
- Bork, P., Sander, C., and Valencia, A. (1992) An ATPase domain common to prokaryotic cell cycle proteins, sugar kinases, actin, and hsp70 heat shock proteins. *Proc Natl Acad Sci U S A* **89**: 7290-7294.
- Botta, G.A., and Park, J.T. (1981) Evidence for involvement of penicillin-binding protein 3 in murein synthesis during septation but not during cell elongation. *J Bacteriol* **145**: 333-340.
- Bramhill, D., and Thompson, C.M. (1994) GTP-dependent polymerization of *Escherichia coli* FtsZ protein to form tubules. *Proc Natl Acad Sci U S A* **91**: 5813-5817.
- Bramhill, D. (1997) Bacterial cell division. *Annu Rev Cell Dev Biol* **13**: 395-424.
- Braud, S., Lavire, C., Bellier, A., and Mazodier, P. (2006) Effect of SsrA (tmRNA) tagging system on translational regulation in *Streptomyces*. *Arch Microbiol* **184**: 343-352.
- Bucca, G., Brassington, A.M., Hotchkiss, G., Mersinias, V., and Smith, C.P. (2003) Negative feedback regulation of *dnaK*, *clpB* and *lon* expression by the DnaK chaperone machine in *Streptomyces coelicolor*, identified by transcriptome and in vivo DnaK-depletion analysis. *Mol Microbiol* **50**: 153-166.
- Buddelmeijer, N., and Beckwith, J. (2002) Assembly of cell division proteins at the *E. coli* cell center. *Curr Opin Microbiol* **5**: 553-557.
- Buddelmeijer, N., and Beckwith, J. (2004) A complex of the *Escherichia coli* cell division proteins FtsL, FtsB and FtsQ forms independently of its localization to the septal region. *Mol Microbiol* **52**: 1315-1327.
- Burger, A., Brandt, B., Susstrunk, U., Thompson, C.J., and Wohlleben, W. (1998) Analysis of a *Streptomyces coelicolor* A3(2) locus containing the nucleoside diphosphate kinase (*ndk*) and folylpolyglutamate synthetase (*folC*) genes. *FEMS Microbiol Lett* **159**: 283-291.
- Burger, A., Sichler, K., Kelemen, G., Buttner, M., and Wohlleben, W. (2000) Identification and characterization of the *mre* gene region of *Streptomyces coelicolor* A3(2). *Mol Gen Genet* **263**: 1053-1060.
- Bushell, M.E. (1988) Growth, product formation and fermentation technology. In *Actinomycetes in biotechnology*. London, UK: Academic press, pp. 185-217.
- Cabean, M.T., and Jacobs-Wagner, C. (2005) Bacterial cell shape. *Nat Rev Microbiol* **3**: 601-610.
- Calcutt, M.J. (1994) Gene organization in the *dnaA-gyrA* region of the *Streptomyces coelicolor* chromosome. *Gene* **151**: 23-28.

- Carballido-Lopez, R., Formstone, A., Li, Y., Ehrlich, S.D., Noiroit, P., and Errington, J. (2006) Actin homolog MreBH governs cell morphogenesis by localization of the cell wall hydrolase LytE. *Dev Cell* **11**: 399-409.
- Chater, K.F. (1972) A morphological and genetic mapping study of white colony mutants of *Streptomyces coelicolor*. *J Gen Microbiol* **72**: 9-28.
- Chater, K.F., Bruton, C.J., Plaskitt, K.A., Buttner, M.J., Mendez, C., and Helmann, J.D. (1989) The developmental fate of *S. coelicolor* hyphae depends upon a gene product homologous with the motility sigma factor of *B. subtilis*. *Cell* **59**: 133-143.
- Chater, K.F., and Losick, R. (1997) Mycelial life style of *Streptomyces coelicolor* A3(2) and its relatives. In *Bacteria as multicellular organisms*. Shapiro, J.A. and Dworkin, M. (eds). New York: Oxford University Press, pp. 149-182.
- Chater, K.F. (1998) taking a genetic scalpel to the *Streptomyces* colony. *Microbiology* **144**: 1465-1478.
- Chater, K.F. (2001) Regulation of sporulation in *Streptomyces coelicolor* A3(2): a checkpoint multiplex? *Curr Opin Microbiol* **4**: 667-673.
- Chater, K.F., and Chandra, G. (2006) The evolution of development in *Streptomyces* analysed by genome comparisons. *FEMS Microbiol Rev* **30**: 651-672.
- Claessen, D., Wosten, H.A., van Keulen, G., Faber, O.G., Alves, A.M., Meijer, W.G., and Dijkhuizen, L. (2002) Two novel homologous proteins of *Streptomyces coelicolor* and *Streptomyces lividans* are involved in the formation of the rodlet layer and mediate attachment to a hydrophobic surface. *Mol Microbiol* **44**: 1483-1492.
- Claessen, D., Rink, R., de Jong, W., Siebring, J., de Vreugd, P., Boersma, F.G., Dijkhuizen, L., and Wosten, H.A. (2003) A novel class of secreted hydrophobic proteins is involved in aerial hyphae formation in *Streptomyces coelicolor* by forming amyloid-like fibrils. *Genes Dev* **17**: 1714-1726.
- Claessen, D., Stokroos, I., Deelstra, H.J., Penninga, N.A., Bormann, C., Salas, J.A., Dijkhuizen, L., and Wosten, H.A. (2004) The formation of the rodlet layer of streptomycetes is the result of the interplay between rodins and chaplins. *Mol Microbiol* **53**: 433-443.
- Corbin, B.D., Geissler, B., Sadashivam, M., and Margolin, W. (2004) Z-ring-independent interaction between a subdomain of FtsA and late septation proteins as revealed by a polar recruitment assay. *J Bacteriol* **186**: 7736-7744.
- Covarrubias, L., Cervantes, L., Covarrubias, A., Soberon, X., Vichido, I., Blanco, A., Kupersztch-Portnoy, Y.M., and Bolivar, F. (1981) Construction and characterization of new cloning vehicles. V. Mobilization and coding properties of pBR322 and several deletion derivatives including pBR327 and pBR328. *Gene* **13**: 25-35.
- Dalton, K., Thibessard, A., Hunter, J.I.B., and Kelemen, G.H. (2007) A novel compartment, the "sub-apical stem" of the aerial hyphae, is the location of a SigN dependent, developmentally distinct transcription in *Streptomyces coelicolor*. *Mol Microbiol* **In press**.
- Daniel, R.A., Drake, S., Buchanan, C.E., Scholle, R., and Errington, J. (1994) The *Bacillus subtilis* spoVD gene encodes a mother-cell-specific penicillin-binding protein required for spore morphogenesis. *J Mol Biol* **235**: 209-220.
- Daniel, R.A., and Errington, J. (2003) Control of cell morphogenesis in bacteria: two distinct ways to make a rod-shaped cell. *Cell* **113**: 767-776.
- Daniel, R.A., Noiroit-Gros, M.F., Noiroit, P., and Errington, J. (2006) Multiple interactions between the transmembrane division proteins of *Bacillus subtilis* and the role of FtsL instability in divisome assembly. *J Bacteriol* **188**: 7396-7404.
- Datsenko, K.A., and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**: 6640-6645.
- Davis, N.K., and Chater, K.F. (1992) The *Streptomyces coelicolor* whiB gene encodes a small transcription factor-like protein dispensable for growth but essential for sporulation. *Mol Gen Genet* **232**: 351-358.
- de Leeuw, E., Graham, B., Phillips, G.J., ten Hagen-Jongman, C.M., Oudega, B., and Luijink, J. (1999) Molecular characterization of *Escherichia coli* FtsE and FtsX. *Mol Microbiol* **31**: 983-993.
- Den Blaauwen, T., Aarsman, M.E., Vischer, N.O., and Nanninga, N. (2003) Penicillin-binding protein PBP2 of *Escherichia coli* localizes preferentially in the lateral wall and at mid-cell in comparison with the old cell pole. *Mol Microbiol* **47**: 539-547.
- Derouaux, A., Halici, S., Nothaft, H., Neutelings, T., Moutzouris, G., Dusart, J., Titgemeyer, F., and Rigali, S. (2004) Deletion of a cyclic AMP receptor protein homologue diminishes germination and affects morphological development of *Streptomyces coelicolor*. *J Bacteriol* **186**: 1893-1897.
- Du, Y., and Arvidson, C.G. (2003) Identification of ZipA, a Signal Recognition Particle-Dependent Protein from *Neisseria gonorrhoeae*. *J Bacteriol* **185**: 2122-2130.
- Dye, N.A., Pincus, Z., Theriot, J.A., Shapiro, L., and Gitai, Z. (2005) Two independent spiral structures control cell shape in *Caulobacter*. *Proc Natl Acad Sci U S A* **102**: 18608-18613.
- Eccleston, M., Ali, R.A., Seyler, R., Westpheling, J., and Nodwell, J. (2002) Structural and genetic analysis of the BldB protein of *Streptomyces coelicolor*. *J Bacteriol* **184**: 4270-4276.
- Edwards, D.H., and Errington, J. (1997) The *Bacillus subtilis* DivIVA protein targets to the division septum and controls the site specificity of cell division. *Mol Microbiol* **24**: 905-915.
- Elliot, M., Damji, F., Passantino, R., Chater, K., and Leskiw, B. (1998) The bldD gene of *Streptomyces coelicolor* A3(2): a regulatory gene involved in morphogenesis and antibiotic production. *J Bacteriol* **180**: 1549-1555.
- Elliot, M.A., and Leskiw, B.K. (1999) The BldD protein from *Streptomyces coelicolor* is a DNA-binding protein. *J. Bacteriol.* **181**: 6832-6835.
- Elliot, M.A., Bibb, M.J., Buttner, M.J., and Leskiw, B.K. (2001) BldD is a direct regulator of key developmental genes in *Streptomyces coelicolor* A3(2). *Mol. Microbiol.* **40**: 257-269.
- Elliot, M.A., Karoomuthaisiri, N., Huang, J., Bibb, M.J., Cohen, S.N., Kao, C.M., and Buttner, M.J. (2003) The chaplins: a family of hydrophobic cell-surface proteins involved in aerial mycelium formation in *Streptomyces coelicolor*. *Genes Dev* **17**: 1727-1740.
- Ensign, J.C. (1978) Formation, properties, and germination of actinomycete spores. *Annu Rev Microbiol* **32**: 185-219.
- Erickson, H.P., Taylor, D.W., Taylor, K.A., and Bramhill, D. (1996) Bacterial cell division protein FtsZ assembles into protofilament sheets and minirings, structural homologs of tubulin polymers. *Proc Natl Acad Sci U S A* **93**: 519-523.
- Errington, J. (2001) Septation and chromosome segregation during sporulation in *Bacillus subtilis*. *Curr Opin Microbiol* **4**: 660-666.
- Errington, J., Bath, J., and Wu, L.J. (2001) DNA transport in bacteria. *Nature Rev. Mol. Cell Biol.* **2**: 538-544.
- Errington, J. (2003) Regulation of endospore formation in *Bacillus subtilis*. *Nat Rev Microbiol* **1**: 117-126.
- Errington, J., Daniel, R.A., and Scheffers, D.J. (2003) Cytokinesis in bacteria. *Microbiol Mol Biol Rev* **67**: 52-65.
- Fernandez-Moreno, M.A., Caballero, J.L., Hopwood, D.A., and Malpartida, F. (1991) The act cluster contains regulatory and antibiotic export genes, direct targets for translational control by the bldA tRNA gene of *Streptomyces*. *Cell* **66**: 769-780.
- Figge, R.M., Divakaruni, A.V., and Gober, J.W. (2004) MreB, the cell shape-determining bacterial actin homologue, co-ordinates cell wall morphogenesis in *Caulobacter crescentus*. *Mol Microbiol* **51**: 1321-1332.
- Flärdh, K., Findlay, K.C., and Chater, K.F. (1999) Association of early sporulation genes with suggested developmental decision points in *Streptomyces coelicolor* A3(2). *Microbiology* **145**: 2229-2243.

- Flärdh, K., Leibovitz, E., Buttner, M.J., and Chater, K.F. (2000) Generation of a non-sporulating strain of *Streptomyces coelicolor* A3(2) by the manipulation of a developmentally controlled *ftsZ* promoter. *Mol. Microbiol.* **38**: 737-749.
- Flärdh, K. (2003a) Growth polarity and cell division in *Streptomyces*. *Curr Opin Microbiol* **6**: 564-571.
- Flärdh, K. (2003b) Essential role of DivIVA in polar growth and morphogenesis in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **49**: 1523-1536.
- Flärdh, K., and van Wezel, G.P. (2003) Cell division during growth and development of *Streptomyces*. In *Recent developments in bacteriology*. Pandalai, S.G. (ed). Trivandrum, India: Transworld research network.
- Formstone, A., and Errington, J. (2005) A magnesium-dependent mreB null mutant: implications for the role of mreB in *Bacillus subtilis*. *Mol Microbiol* **55**: 1646-1657.
- Fu, X., Shih, Y.L., Zhang, Y., and Rothfield, L.I. (2001) The MinE ring required for proper placement of the division site is a mobile structure that changes its cellular location during the *Escherichia coli* division cycle. *Proc Natl Acad Sci U S A* **98**: 980-985.
- Gill, D.R., and Salmond, G.P. (1990) The identification of the *Escherichia coli* *ftsY* gene product: an unusual protein. *Mol Microbiol* **4**: 575-583.
- Gitai, Z., Dye, N., and Shapiro, L. (2004) An actin-like gene can determine cell polarity in bacteria. *Proc Natl Acad Sci U S A* **101**: 8643-8648.
- Gitai, Z., Dye, N.A., Reisenauer, A., Wachi, M., and Shapiro, L. (2005) MreB actin-mediated segregation of a specific region of a bacterial chromosome. *Cell* **120**: 329-341.
- Glauert, A.M., and Hopwood, D.A. (1961) The fine structure of *Streptomyces violaceoruber* (*S. coelicolor*). III. The walls of the mycelium and spores. *J Biophys Biochem Cytol* **10**: 505-516.
- Goehring, N.W., Gueiros-Filho, F., and Beckwith, J. (2005) Premature targeting of a cell division protein to midcell allows dissection of divisome assembly in *Escherichia coli*. *Genes Dev* **19**: 127-137.
- Goehring, N.W., Gonzalez, M.D., and Beckwith, J. (2006) Premature targeting of cell division proteins to midcell reveals hierarchies of protein interactions involved in divisome assembly. *Mol Microbiol* **61**: 33-45.
- Goodell, E.W. (1985) Recycling of murein by *Escherichia coli*. *J Bacteriol* **163**: 305-310.
- Grantcharova, N., Ubbaysekera, W., Mowbray, S.L., McCormick, J.R., and Flärdh, K. (2003) A missense mutation in *ftsZ* differentially affects vegetative and developmentally controlled cell division in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **47**: 645-656.
- Grantcharova, N., Lustig, U., and Flärdh, K. (2005) Dynamics of FtsZ assembly during sporulation in *Streptomyces coelicolor* A3(2). *J Bacteriol* **187**: 3227-3237.
- Gray, D.I., Gooday, G.W., and Prosser, J.I. (1990) Apical hyphal extension in *Streptomyces coelicolor* A3(2). *J Gen Microbiol* **136**: 1077-1084.
- Grossman, A.D. (1995) Genetic networks controlling the initiation of sporulation and the development of genetic competence in *Bacillus subtilis*. *Annu Rev Genet* **29**: 477-508.
- Gueiros-Filho, F.J., and Losick, R. (2002) A widely conserved bacterial cell division protein that promotes assembly of the tubulin-like protein FtsZ. *Genes Dev* **16**: 2544-2556.
- Gust, B., Challis, G.L., Fowler, K., Kieser, T., and Chater, K.F. (2003) PCR-targeted *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci U S A* **100**: 1541-1546.
- Harris, S.D., Hofmann, A.F., Tedford, H.W., and Lee, M.P. (1999) Identification and characterization of genes required for hyphal morphogenesis in the filamentous fungus *Aspergillus nidulans*. *Genetics* **151**: 1015-1025.
- Henriques, A.O., Glaser, P., Piggot, P.J., and Moran, C.P., Jr. (1998) Control of cell shape and elongation by the *rodA* gene in *Bacillus subtilis*. *Mol Microbiol* **28**: 235-247.
- Higgins, C.F. (1992) ABC transporters: from microorganisms to man. *Annu Rev Cell Biol* **8**: 67-113.
- Higgins, D.G., Thompson, J.D., and Gibson, T.J. (1996) Using CLUSTAL for multiple sequence alignments. *Methods Enzymol* **266**: 383-402.
- Holtje, J.V. (1998) Growth of the stress-bearing and shape-maintaining murein sacculus of *Escherichia coli*. *Microbiol Mol Biol Rev* **62**: 181-203.
- Hu, Z., and Lutkenhaus, J. (1999) Topological regulation of cell division in *Escherichia coli* involves rapid pole to pole oscillation of the division inhibitor MinC under the control of MinD and MinE. *Mol Microbiol* **34**: 82-90.
- Hunt, A.C., Servin-Gonzalez, L., Kelemen, G.H., and Buttner, M.J. (2005) The *bldC* developmental locus of *Streptomyces coelicolor* encodes a member of a family of small DNA-binding proteins related to the DNA-binding domains of the MerR family. *J Bacteriol* **187**: 716-728.
- Ikeda, H., Ishikawa, J., Hanamoto, A., Shinose, M., Kikuchi, H., Shiba, T., Sakaki, Y., Hattori, M., and Omura, S. (2003) Complete genome sequence and comparative analysis of the industrial microorganism *Streptomyces avermitilis*. *Nat Biotechnol* **14**: 14.
- Jakimowicz, D., Chater, K., and Zakrzewska-Czerwinska, J. (2002) The ParB protein of *Streptomyces coelicolor* A3(2) recognizes a cluster of *parS* sequences within the origin-proximal region of the linear chromosome. *Mol Microbiol* **45**: 1365-1377.
- Jakimowicz, D., Gust, B., Zakrzewska-Czerwinska, J., and Chater, K.F. (2005) Developmental-stage-specific assembly of ParB complexes in *Streptomyces coelicolor* hyphae. *J Bacteriol* **187**: 3572-3580.
- Jakimowicz, D., Mouz, S., Zakrzewska-Czerwinska, J., and Chater, K.F. (2006) Developmental control of a *parAB* promoter leads to formation of sporulation-associated ParB complexes in *Streptomyces coelicolor*. *J Bacteriol* **188**: 1710-1720.
- Janssen, G.R., and Bibb, M.J. (1993) Derivatives of pUC18 that have *Bgl*II sites flanking a modified multiple cloning site and that retain the ability to identify recombinant clones by visual screening of *Escherichia coli* colonies. *Gene* **124**: 133-134.
- Jiang, H., and Kendrick, K.E. (2000) Characterization of *ssfR* and *ssgA*, two genes involved in sporulation of *Streptomyces griseus*. *J Bacteriol* **182**: 5521-5529.
- Jones, L.J., Carballido-Lopez, R., and Errington, J. (2001) Control of cell shape in bacteria: helical, actin-like filaments in *Bacillus subtilis*. *Cell* **104**: 913-922.
- Justice, S.S., Garcia-Lara, J., and Rothfield, L.I. (2000) Cell division inhibitors SulA and MinC/MinD block septum formation at different steps in the assembly of the *Escherichia coli* division machinery. *Mol Microbiol* **37**: 410-423.
- Kato, J.Y., Suzuki, A., Yamazaki, H., Ohnishi, Y., and Horinouchi, S. (2002) Control by A-factor of a metalloendopeptidase gene involved in aerial mycelium formation in *Streptomyces griseus*. *J Bacteriol* **184**: 6016-6025.
- Kawamoto, S., and Ensign, J.C. (1995) Cloning and characterization of a gene involved in regulation of sporulation and cell division in *Streptomyces griseus*. *Actinomycetologica* **9**: 136-151.
- Kawamoto, S., Watanabe, H., Hesketh, A., Ensign, J.C., and Ochi, K. (1997) Expression analysis of the *ssgA* gene product, associated with sporulation and cell division in *Streptomyces griseus*. *Microbiology* **143**: 1077-1086.

- Keijser, B.J., van Wezel, G.P., Canters, G.W., and Vijgenboom, E. (2002) Developmental regulation of the *Streptomyces lividans ram* genes: involvement of RamR in regulation of the *ramCSAB* operon. *J Bacteriol* **184**: 4420-4429.
- Keijser, B.J., Noens, E.E., Kraal, B., Koerten, H.K., and van Wezel, G.P. (2003) The *Streptomyces coelicolor ssgB* gene is required for early stages of sporulation. *FEMS Microbiol Lett* **225**: 59-67.
- Kelemen, G.H., Brown, G.L., Kormanec, J., Potuckova, L., Chater, K.F., and Buttner, M.J. (1996) The positions of the sigma-factor genes, *whiG* and *sigF*, in the hierarchy controlling the development of spore chains in the aerial hyphae of *Streptomyces coelicolor* A3(2). *Mol Microbiol* **21**: 593-603.
- Kelemen, G.H., Brian, P., Flärdh, K., Chamberlin, L., Chater, K.F., and Buttner, M.J. (1998) Developmental regulation of transcription of *whiE*, a locus specifying the polyketide spore pigment in *Streptomyces coelicolor* A3 (2). *J Bacteriol* **180**: 2515-2521.
- Kelemen, G.H., Viollier, P.H., Tenor, J., Marri, L., Buttner, M.J., and Thompson, C.J. (2001) A connection between stress and development in the multicellular prokaryote *Streptomyces coelicolor* A3(2). *Mol Microbiol* **40**: 804-814.
- Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F., and Hopwood, D.A. (2000) *Practical Streptomyces genetics*. Norwich, U.K.: John Innes Foundation.
- Kim, H.J., Calcutt, M.J., Schmidt, F.J., and Chater, K.F. (2000) Partitioning of the linear chromosome during sporulation of *Streptomyces coelicolor* A3(2) involves an *oriC*-linked *parAB* locus. *J Bacteriol* **182**: 1313-1320.
- Koonin, E.V. (1993) A superfamily of ATPases with diverse functions containing either classical or deviant ATP-binding motif. *J Mol Biol* **229**: 1165-1174.
- Kormanec, J., and Sevcikova, B. (2002) The stress-response sigma factor sigma(H) controls the expression of *ssgB*, a homologue of the sporulation-specific cell division gene *ssgA*, in *Streptomyces coelicolor* A3(2). *Mol Genet Genomics* **267**: 536-543.
- Kruse, T., Moller-Jensen, J., Lobner-Olesen, A., and Gerdes, K. (2003) Dysfunctional MreB inhibits chromosome segregation in *Escherichia coli*. *Embo J* **22**: 5283-5292.
- Kruse, T., Bork-Jensen, J., and Gerdes, K. (2005) The morphogenetic MreBCD proteins of *Escherichia coli* form an essential membrane-bound complex. *Mol Microbiol* **55**: 78-89.
- Kruse, T., and Gerdes, K. (2005) Bacterial DNA segregation by the actin-like MreB protein. *Trends Cell Biol* **15**: 343-345.
- Kwak, J., Dharmatilake, A.J., Jiang, H., and Kendrick, K.E. (2001) Differential regulation of *ftsZ* transcription during septation of *Streptomyces griseus*. *J Bacteriol* **183**: 5092-5101.
- Larson, J.L., and Hershberger, C.L. (1986) The minimal replicon of a streptomycete plasmid produces an ultrahigh level of plasmid DNA. *Plasmid* **15**: 199-209.
- Lawlor, E.J., Baylis, H.A., and Chater, K.F. (1987) Pleiotropic morphological and antibiotic deficiencies result from mutations in a gene encoding a tRNA-like product in *Streptomyces coelicolor* A3(2). *Genes Dev* **1**: 1305-1310.
- Leaver, M., and Errington, J. (2005) Roles for MreC and MreD proteins in helical growth of the cylindrical cell wall in *Bacillus subtilis*. *Mol Microbiol* **57**: 1196-1209.
- Lee, J.C., and Stewart, G.C. (2003) Essential nature of the mreC determinant of *Bacillus subtilis*. *J Bacteriol* **185**: 4490-4498.
- Lee, K., and Cohen, S.N. (2003) A *Streptomyces coelicolor* functional orthologue of *Escherichia coli* RNase E shows shuffling of catalytic and PNPase-binding domains. *Mol Microbiol* **48**: 349-360.
- Leskiw, B.K., Bibb, M.J., and Chater, K.F. (1991a) The use of a rare codon specifically during development? *Mol Microbiol* **5**: 2861-2867.
- Leskiw, B.K., Lawlor, E.J., Fernandez-Abalos, J.M., and Chater, K.F. (1991b) TTA codons in some genes prevent their expression in a class of developmental, antibiotic-negative, *Streptomyces* mutants. *Proc Natl Acad Sci U S A* **88**: 2461-2465.
- Leskiw, B.K., and Mah, R. (1995) The bldA-encoded tRNA is poorly expressed in the *bldI* mutant of *Streptomyces coelicolor* A3(2). *Microbiology* **141** (Pt 8): 1921-1926.
- Levin, P.A., Kurtser, I.G., and Grossman, A.D. (1999) Identification and characterization of a negative regulator of FtsZ ring formation in *Bacillus subtilis*. *Proc Natl Acad Sci U S A* **96**: 9642-9647.
- Liu, G., Draper, G.C., and Donachie, W.D. (1998) FtsK is a bifunctional protein involved in cell division and chromosome localization in *Escherichia coli*. *Mol Microbiol* **29**: 893-903.
- Locci, R. (1980) Response of developing branched bacteria to adverse environments. II. Micromorphological effects of lysozyme on some aerobic actinomycetes. *Zentralbl Bakteriol A* **247**: 374-382.
- Locher, K.P. (2004) Structure and mechanism of ABC transporters. *Curr Opin Struct Biol* **14**: 426-431.
- Longtime, M.S., DeMarini, D.J., Valencik, M.L., Al-Awar, O.S., Fares, H., De Virgilio, C., and Pringle, J.R. (1996) The septins: roles in cytokinesis and other processes. *Curr Opin Cell Biol* **8**: 106-119.
- Lowe, J., van den Ent, F., and Amos, L.A. (2004) Molecules of the bacterial cytoskeleton. *Annu Rev Biophys Biomol Struct* **33**: 177-198.
- MacNeil, D.J., Gewain, K.M., Ruby, C.L., Dezeny, G., Gibbons, P.H., and MacNeil, T. (1992) Analysis of *Streptomyces avermitilis* genes required for avermectin biosynthesis utilizing a novel integration vector. *Gene* **111**: 61-68.
- Margolin, W. (2005) FtsZ and the division of prokaryotic cells and organelles. *Nat Rev Mol Cell Biol* **6**: 862-871.
- Marston, A.L., Thomaides, H.B., Edwards, D.H., Sharpe, M.E., and Errington, J. (1998) Polar localization of the MinD protein of *Bacillus subtilis* and its role in selection of the mid-cell division site. *Genes Dev* **12**: 3419-3430.
- Matsuhashi, M., Wachi, M., and Ishino, F. (1990) Machinery for cell growth and division: penicillin-binding proteins and other proteins. *Res Microbiol* **141**: 89-103.
- Mazza, P., Noens, E.E., Schirmer, K., Grantcharova, N., Mommaas, A.M., Koerten, H.K., Muth, G., Flardh, K., van Wezel, G.P., and Wohleben, W. (2006) MreB of *Streptomyces coelicolor* is not essential for vegetative growth but is required for the integrity of aerial hyphae and spores. *Mol Microbiol* **60**: 838-852.
- McCormick, J.R., Su, E.P., Driks, A., and Losick, R. (1994) Growth and viability of *Streptomyces coelicolor* mutant for the cell division gene *ftsZ*. *Mol Microbiol* **14**: 243-254.
- McCormick, J.R., and Losick, R. (1996) Cell division gene *ftsQ* is required for efficient sporulation but not growth and viability in *Streptomyces coelicolor* A3(2). *J Bacteriol* **178**: 5295-5301.
- Mendez, C., Brana, A.F., Manzanal, M.B., and Hardisson, C. (1985) Role of substrate mycelium in colony development in *Streptomyces*. *Can J Microbiol* **31**: 446-450.
- Merrick, M.J. (1976) A morphological and genetic mapping study of bald colony mutants of *Streptomyces coelicolor*. *J Gen Microbiol* **96**: 299-315.
- Michie, K.A., and Lowe, J. (2006) Dynamic filaments of the bacterial cytoskeleton. *Annu Rev Biochem* **75**: 467-492.
- Molle, V., and Buttner, M.J. (2000) Different alleles of the response regulator gene *bldM* arrest *Streptomyces coelicolor* development at distinct stages. *Mol Microbiol* **36**: 1265-1278.
- Molle, V., Palframan, W.J., Findlay, K.C., and Buttner, M.J. (2000) WhiD and WhiB, homologous proteins required for different stages of sporulation in *Streptomyces coelicolor* A3(2). *J Bacteriol*.. *Mar* **182**: 1286-1295.

- Nguyen, K.T., Tenor, J., Stettler, H., Nguyen, L.T., Nguyen, L.D., and Thompson, C.J. (2003) Colonial differentiation in *Streptomyces coelicolor* depends on translation of a specific codon within the *adpA* gene. *J Bacteriol* **185**: 7291-7296.
- Nodwell, J.R., McGovern, K., and Losick, R. (1996) An oligopeptide permease responsible for the import of an extracellular signal governing aerial mycelium formation in *Streptomyces coelicolor*. *Mol. Microbiol* **22**: 881-893.
- Nodwell, J.R., and Losick, R. (1998) Purification of an extracellular signaling molecule involved in production of aerial mycelium by *Streptomyces coelicolor*. *J Bacteriol* **180**: 1334-1337.
- Nodwell, J.R., Yang, M., Kuo, D., and Losick, R. (1999) Extracellular complementation and the identification of additional genes involved in aerial mycelium formation in *Streptomyces coelicolor*. *Genetics* **151**: 569-584.
- Noens, E.E., Mersinias, V., Traag, B.A., Smith, C.P., Koerten, H.K., and van Wezel, G.P. (2005) SsgA-like proteins determine the fate of peptidoglycan during sporulation of *Streptomyces coelicolor*. *Mol Microbiol* **58**: 929-944.
- Noens, E.E., Mersinias, V., Willemse, J., Traag, B.A., Laing, E., Chater, K.F., Smith, C.P., Koerten, H.K., and van Wezel, G.P. (2007) Loss of the controlled localisation of growth stage-specific cell wall synthesis pleiotropically affects developmental gene expression in an ssgA mutant of *Streptomyces coelicolor*. *Mol Microbiol* **64**: 1244-59.
- Nothaft, H., Dresel, D., Willimek, A., Mahr, K., Niederweis, M., and Titgemeyer, F. (2003) The phosphotransferase system of *Streptomyces coelicolor* is biased for N-acetylglucosamine metabolism. *J Bacteriol* **185**: 7019-7023.
- Ohnishi, Y., and Horinouchi, S. (1999) Regulation of secondary metabolism and morphological differentiation by a microbial hormone in *Streptomyces*. *Tanpakushitsu Kakusan Koso* **44**: 1552-1561.
- Ohnishi, Y., Seo, J.W., and Horinouchi, S. (2002) Deprogrammed sporulation in *Streptomyces*. *FEMS Microbiol Lett* **216**: 1-7.
- Park, J.T. (1993) Turnover and recycling of the murine sacculus in oligopeptide permease-negative strains of *Escherichia coli*: indirect evidence for an alternative permease system and for a monolayered sacculus. *J Bacteriol* **175**: 7-11.
- Pichoff, S., and Lutkenhaus, J. (2002) Unique and overlapping roles for ZipA and FtsA in septal ring assembly in *Escherichia coli*. *Embo J* **21**: 685-693.
- Piette, A., Derouaux, A., Gerkens, P., Noens, E.E., Mazzucchelli, G., Vion, S., Koerten, H.K., Titgemeyer, F., De Pauw, E., Leprince, P., van Wezel, G.P., Galleni, M., and Rigali, S. (2005) From dormant to germinating spores of *Streptomyces coelicolor* A3(2): new perspectives from the *crg* null mutant. *J Proteome Res* **4**: 1699-1708.
- Pope, M.K., Green, B.D., and Westpheling, J. (1996) The bld mutants of *Streptomyces coelicolor* are defective in the regulation of carbon utilization, morphogenesis and cell-cell signalling. *Mol Microbiol* **19**: 747-756.
- Pope, M.K., Green, B., and Westpheling, J.R.W., IV (1998) The *bldB* gene encodes a small protein required for morphogenesis, antibiotic production, and catabolite control in *Streptomyces coelicolor*. *J. Bacteriol.* **180**: 1556-1562.
- Potuckova, L., Kelemen, G.H., Findlay, K.C., Lonetto, M.A., Buttner, M.J., and Kormanec, J. (1995) A new RNA polymerase sigma factor, sigma F, is required for the late stages of morphological differentiation in *Streptomyces* spp. *Mol Microbiol* **17**: 37-48.
- Prentki, P., and Krisch, H.M. (1984) In vitro insertional mutagenesis with a selectable DNA fragment. *Gene* **29**: 303-313.
- Prosser, J.I., and Tough, A.J. (1991) Growth mechanisms and growth kinetics of filamentous microorganisms. *Crit Rev Biotechnol* **10**: 253-274.
- Redenbach, M., Kieser, H.M., Denapaité, D., Eichner, A., Cullum, J., Kinashi, H., and Hopwood, D.A. (1996) A set of ordered cosmids and a detailed genetic and physical map for the 8 mb *Streptomyces coelicolor* a3(2) chromosome. *Mol. Microbiol* **21**: 77-96.
- Rigali, S., Schlicht, M., Hoskisson, P., Nothaft, H., Merzbacher, M., Joris, B., and Titgemeyer, F. (2004) Extending the classification of bacterial transcription factors beyond the helix-turn-helix motif as an alternative approach to discover new cis/trans relationships. *Nucleic Acids Res* **32**: 3418-3426.
- Rigali, S., Nothaft, H., Noens, E.E., Schlicht, M., Colson, S., Muller, M., Joris, B., Koerten, H.K., Hopwood, D.A., Titgemeyer, F., and van Wezel, G.P. (2006) The sugar phosphotransferase system of *Streptomyces coelicolor* is regulated by the GntR-family regulator DasR and links N-acetylglucosamine metabolism to the control of development. *Mol Microbiol* **61**: 1237-1251.
- Rothfield, L., Taghbalout, A., and Shih, Y.L. (2005) Spatial control of bacterial division-site placement. *Nat Rev Microbiol* **3**: 959-968.
- Ryding, N.J., Kelemen, G.H., Whatling, C.A., Flärdh, K., Buttner, M.J., and Chater, K.F. (1998) A developmentally regulated gene encoding a repressor-like protein is essential for sporulation in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **29**: 343-357.
- Ryding, N.J., Bibb, M.J., Molle, V., Findlay, K.C., Chater, K.F., and Buttner, M.J. (1999) New sporulation loci in *Streptomyces coelicolor* A3(2). *J. Bacteriol.* **181**: 5419-5425.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular cloning: a laboratory manual*. Cold Spring Harbor, N.Y.: Cold Spring Harbor laboratory press.
- Sanchez, M., Valencia, A., Ferrandiz, M.J., Sander, C., and Vicente, M. (1994) Correlation between the structure and biochemical activities of FtsA, an essential cell division protein of the actin family. *Embo J* **13**: 4919-4925.
- Saraste, M., Sibbald, P.R., and Wittinghofer, A. (1990) The P-loop—a common motif in ATP- and GTP-binding proteins. *Trends Biochem Sci* **15**: 430-434.
- Schlosser, A., Kampers, T., and Schrempf, H. (1997) The *Streptomyces* ATP-binding component MsiK assists in cellobiose and maltose transport. *J Bacteriol* **179**: 2092-2095.
- Schmidt, K.L., Peterson, N.D., Kustusch, R.J., Wissel, M.C., Graham, B., Phillips, G.J., and Weiss, D.S. (2004) A predicted ABC transporter, FtsEX, is needed for cell division in *Escherichia coli*. *J Bacteriol* **186**: 785-793.
- Schwedock, J., McCormick, J.R., Angert, E.R., Nodwell, J.R., and Losick, R. (1997) Assembly of the cell division protein FtsZ into ladder-like structures in the aerial hyphae of *Streptomyces coelicolor*. *Mol. Microbiol* **25**: 858.
- Sharpe, M.E., and Errington, J. (1999) Upheaval in the bacterial nucleoid. An active chromosome segregation mechanism. *Trends Genet* **15**: 70-74.
- Shih, Y.L., Le, T., and Rothfield, L. (2003) Division site selection in *Escherichia coli* involves dynamic redistribution of Min proteins within coiled structures that extend between the two cell poles. *Proc Natl Acad Sci U S A* **100**: 7865-7870.
- Smyth, G.K., and Speed, T.P. (2003) Normalization of cDNA microarray data. *Methods*: 265-273.
- Smyth, G.K. (2005) Limma: linear models for microarray data. In *Bioinformatics and Computational Biology Solutions using R and Bioconductor*. Gentleman, R., Carey, V., Dudoit, S., Irizarry, R., and Huber, W. (eds). New York: Springer, pp. 397-420.
- Slovak, P.M., Wadhams, G.H., and Armitage, J.P. (2005) Localization of MreB in *Rhodobacter sphaeroides* under conditions causing changes in cell shape and membrane structure. *J Bacteriol* **187**: 54-64.
- Soliveri, J.A., Gomez, J., Bishai, W.R., and Chater, K.F. (2000) Multiple paralogous genes related to the *Streptomyces coelicolor* developmental regulatory gene *whiB* are present in *Streptomyces* and other actinomycetes. *Microbiology* **146**: 333-343.
- Soufo, H.J., and Graumann, P.L. (2003) Actin-like proteins MreB and Mbl from *Bacillus subtilis* are required for bipolar positioning of replication origins. *Curr Biol* **13**: 1916-1920.

- Suefuji, K., Valluzzi, R., and RayChaudhuri, D. (2002) Dynamic assembly of MinD into filament bundles modulated by ATP, phospholipids, and MinE. *Proc Natl Acad Sci U S A* **99**: 16776-16781.
- Takano, E., Tao, M., Long, F., Bibb, M.J., Wang, L., Li, W., Buttner, M.J., Deng, Z.X., and Chater, K.F. (2003) A rare leucine codon in *adpA* is implicated in the morphological defect of *bldA* mutants of *Streptomyces coelicolor*. *Mol Microbiol* **50**: 475-486.
- Takano, E., Kinoshita, H., Mersinias, V., Bucca, G., Hotchkiss, G., Nihira, T., Smith, C.P., Bibb, M., Wohlleben, W., and Chater, K. (2005) A bacterial hormone (the SCB1) directly controls the expression of a pathway-specific regulatory gene in the cryptic type I polyketide biosynthetic gene cluster of *Streptomyces coelicolor*. *Mol Microbiol* **56**: 465-479.
- Tan, H., Yang, H., Tian, Y., Wu, W., Whatling, C.A., Chamberlin, L.C., Buttner, M.J., Nodwell, J., and Chater, K.F. (1998) The *Streptomyces coelicolor* sporulation-specific sigma WhiG form of RNA polymerase transcribes a gene encoding a ProX-like protein that is dispensable for sporulation. *Gene* **212**: 137-146.
- Tillotson, R.D., Wosten, H.A., Richter, M., and Willey, J.M. (1998) A surface active protein involved in aerial hyphae formation in the filamentous fungus *Schizophyllum commune* restores the capacity of a bald mutant of the filamentous bacterium *Streptomyces coelicolor* to erect aerial structures. *Mol Microbiol* **30**: 595-602.
- Traag, B.A., Kelemen, G.H., and Van Wezel, G.P. (2004) Transcription of the sporulation gene *ssgA* is activated by the IclR-type regulator SsgR in a whi-independent manner in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **53**: 985-1000.
- Tian, Y., Fowler, K., Findlay, K., Tan, H., and Chater, K.F. (2007) An Unusual Response Regulator Influences Sporulation at Early and Late Stages in *Streptomyces coelicolor*. *J Bacteriol* **189**: 2873-2885.
- Ueda, K., Oinuma, K., Ikeda, G., Hosono, K., Ohnishi, Y., Horinouchi, S., and Beppu, T. (2002) AmfS, an extracellular peptidic morphogen in *Streptomyces griseus*. *J Bacteriol* **184**: 1488-1492.
- Ukai, H., Matsuzawa, H., Ito, K., Yamada, M., and Nishimura, A. (1998) ftsE(Ts) affects translocation of K+-pump proteins into the cytoplasmic membrane of *Escherichia coli*. *J Bacteriol* **180**: 3663-3670.
- van den Ent, F., Amos, L.A., and Lowe, J. (2001) Prokaryotic origin of the actin cytoskeleton. *Nature* **413**: 39-44.
- van Wezel, G.P., van der Meulen, J., Kawamoto, S., Luiten, R.G., Koerten, H.K., and Kraal, B. (2000a) SsgA is essential for sporulation of *Streptomyces coelicolor* A3(2) and affects hyphal development by stimulating septum formation. *J Bacteriol* **182**: 5653-5662.
- van Wezel, G.P., van der Meulen, J., Taal, E., Koerten, H., and Kraal, B. (2000b) Effects of increased and deregulated expression of cell division genes on the morphology and on antibiotic production of streptomycetes. *Antonie Van Leeuwenhoek* **78**: 269-276.
- van Wezel, G.P., White J, Hoogvliet G, Bibb M.J. (2000c) Application of redD, the transcriptional activator gene of the undecylprodigiosin biosynthetic pathway, as a reporter for transcriptional activity in *Streptomyces coelicolor* A3(2) and *Streptomyces lividans*. *J Mol Microbiol Biotechnol* **2**:551-556.
- van Wezel, G.P., and Vijgenboom, E. (2004) Novel aspects of signaling in *Streptomyces* development. *Adv Appl Microbiol* **56**: 65-88.
- van Wezel, G.P., Mahr, K., Konig, M., Traag, B.A., Pimentel-Schmitt, E.F., Willimek, A., and Titgemeyer, F. (2005) GlcP constitutes the major glucose uptake system of *Streptomyces coelicolor* A3(2). *Mol Microbiol* **55**: 624-636.
- van Wezel, G.P., Krabben, P., Traag, B.A., Keijser, B.J., Kerste, R., Vijgenboom, E., Heijnen, J.J., and Kraal, B. (2006) Unlocking *Streptomyces* spp. for use as sustainable industrial production platforms by morphological engineering. *Appl Environ Microbiol* **72**: 5283-5288.
- Vicente, M., and Rico, A.I. (2006) The order of the ring: assembly of *Escherichia coli* cell division components. *Mol Microbiol* **61**: 5-8.
- Vicente, M., Rico, A.I., Martinez-Arteaga, R., and Mingorance, J. (2006) Septum enlightenment: assembly of bacterial division proteins. *J Bacteriol* **188**: 19-27.
- Vinella, D., Joseleau-Petit, D., Thevenet, D., Bouloc, P., and D'Ari, R. (1993) Penicillin-binding protein 2 inactivation in *Escherichia coli* results in cell division inhibition, which is relieved by FtsZ overexpression. *J Bacteriol* **175**: 6704-6710.
- Wachi, M., and Matsuhashi, M. (1989) Negative control of cell division by *mreB*, a gene that functions in determining the rod shape of *Escherichia coli* cells. *J Bacteriol* **171**: 3123-3127.
- Wang, L., Yu, Y., He, X., Zhou, X., Deng, Z., Chater, K.F., and Tao, M. (2007) Role of an FtsK-like protein in genetic stability in *Streptomyces coelicolor* A3(2). *J Bacteriol*.
- Wang, X., Possoz, C., and Sherratt, D.J. (2005) Dancing around the divisome: asymmetric chromosome segregation in *Escherichia coli*. *Genes Dev* **19**: 2367-2377.
- Ward, J.M., Janssen, G.R., Kieser, T., Bibb, M.J., and Buttner, M.J. (1986) Construction and characterisation of a series of multi-copy promoter-probe plasmid vectors for *Streptomyces* using the aminoglycoside phosphotransferase gene from Tn5 as indicator. *Mol Gen Genet* **203**: 468-478.
- Wardell, J.N., Stocks, S.M., Thomas, C.R., and Bushell, M.E. (2002) Decreasing the hyphal branching rate of *Saccharopolyspora erythraea* NRRL 2338 leads to increased resistance to breakage and increased antibiotic production. *Biotechnol. Bioeng.* **78**: 141-146.
- Wei, Y., Havasy, T., McPherson, D.C., and Popham, D.L. (2003) Rod shape determination by the *Bacillus subtilis* class B penicillin-binding proteins encoded by *pbpA* and *pbpH*. *J Bacteriol* **185**: 4717-4726.
- White, J., and Bibb, M. (1997) *bldA* dependence of undecylprodigiosin production in *Streptomyces coelicolor* A3(2) involves a pathway-specific regulatory cascade. *J Bacteriol* **179**: 627-633.
- Wildermuth, H. (1970) Development and organization of the aerial mycelium in *Streptomyces coelicolor*. *J Gen Microbiol* **60**: 43-50.
- Wildermuth, H., and Hopwood, D.A. (1970) Septation during sporulation in *Streptomyces coelicolor*. *J Gen Microbiol* **60**: 51-59.
- Wiley, J., Santamaría, R., Guijarro, J., Geistlich, M., and Losick, R. (1991) Extracellular complementation of a developmental mutation implicates a small sporulation protein in aerial mycelium formation by *S. coelicolor*. *Cell* **65**: 641-650.
- Withey, J.H., and Friedman, D.I. (2002) The biological roles of trans-translation. *Curr Opin Microbiol* **5**: 154-159.
- Woodcock, D.M., Crowther, P.J., Doherty, J., Jefferson, S., DeCruz, E., Noyer-Weidner, M., Smith, S.S., Michael, M.Z., and Graham, M.W. (1989) Quantitative evaluation of *Escherichia coli* host strains for tolerance to cytosine methylation in plasmid and phage recombinants. *Nucleic Acids Res* **17**: 3469-3478.
- Wu, L.J., and Errington, J. (2004) Coordination of cell division and chromosome segregation by a nucleoid occlusion protein in *Bacillus subtilis*. *Cell* **117**: 915-925.
- Yamazaki, H., Ohnishi, Y., and Horinouchi, S. (2000) An A-factor-dependent extracytoplasmic function sigma factor (sigma(AdsA)) that is essential for morphological development in *Streptomyces griseus*. *J Bacteriol* **182**: 4596-4605.
- Yamazaki, H., Ohnishi, Y., and Horinouchi, S. (2003) Transcriptional switch on of *ssgA* by A-factor, which is essential for spore septum formation in *Streptomyces griseus*. *J Bacteriol* **185**: 1273-1283.
- Young, K.D. (2003) Bacterial shape. *Mol Microbiol* **49**: 571-580.
- Yu, X.C., Weihe, E.K., and Margolin, W. (1998) Role of the C terminus of FtsK in *Escherichia coli* chromosome segregation. *J Bacteriol* **180**: 6424-6428.

CURRICULUM VITAE

Elke Noens was born on the 3rd of April 1979 in Kapellen (Belgium). In 1997, she obtained the diploma for secondary education at the Sint Lambertus Instituut (Ekeren, Belgium). In the same year, she started the study Medical Biochemistry at the University of Antwerp (Belgium). During the fourth and last year, she participated in an Erasmus exchange programme for six months, completing an internship in the department of Medical Microbiology, LUMC (Leiden, The Netherlands). In 2001, she obtained her Masters degree at the University of Antwerp (Belgium). During 2001-2002, she completed an internship in the Centre for Carbohydrate Chemistry, University of East Anglia (Norwich, UK) under the supervision of Prof. R.A. Field. In the summer of 2002, she started as a Ph.D. student at Leiden University. She worked under the supervision of Dr. G.P. van Wezel in Microbial Development (formerly Genexpress), LIC and Dr. H.K. Koerten in Electron Microscopy, MCB, LUMC. Prof. C.W.A. Pleij was her promotor. The work done as a Ph.D. student is presented in this thesis.

LIST OF PUBLICATIONS

- BJ Keijser, **EE Noens**, B Kraal, HK Koerten and GP van Wezel. (2003) The *Streptomyces coelicolor ssgB* gene is required for early stages of sporulation. *FEMS Microbiol Lett.* **225** (1): 59-67.
- A Piette, A Derouaux, P Gerkens, **EE Noens**, G Mazzucchelli, S Vion, HK Koerten, F Titgemeyer, E De Pauw, P Leprince, GP van Wezel, M Galleni, S Rigali. (2005) From dormant to germinating spores of *Streptomyces coelicolor* A3(2): new perspectives from the *crp* null mutant. *J Proteome Res.* **4** (5): 1699-708.
- EE Noens**, V Mersinias, BJ Traag, CP Smith, HK Koerten and GP van Wezel. (2005) SsgA-like proteins determine the fate of peptidoglycan during sporulation of *Streptomyces coelicolor*. *Mol. Microbiol.* **58**(4):929-44. (Including cover illustration)
- P Mazza, **EE Noens**, K Schimer, N Grantcharova, AM Mommaas, HK Koerten, G Muth, K Flardh, GP van Wezel, W Wohlleben. (2006) MreB of *Streptomyces coelicolor* is not essential for vegetative growth but is required for the integrity of aerial hyphae and spores. *Mol Microbiol.* **60** (4):838-52.
- S Rigali, L Nothaft, **EE Noens**, M Schlicht, S. Colson, M Muller, B Joris, HK Koerten, DA Hopwood, F Titgenmeyer and GP van Wezel. The sugar phosphotransferase system of *Streptomyces coelicolor* is regulated by the GntR-family regulator DasR and links N-acetylglucosamine metabolism to the control of development. *Mol. Microbiol.* **61**(5):1237-1251. (Including cover illustration)
- EE Noens**, V Mersinias, BA Traag, KF Chater, CP Smith, HK Koerten, GP van Wezel. *In vivo* localization and transcriptomic studies identify SsgA as an important developmental protein that controls early stages of sporulation-specific cell division in *Streptomyces coelicolor*. *Mol. Microbiol.* **64**(5): 1244-1259. (Including cover illustration)