

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/35803> holds various files of this Leiden University dissertation

Author: Aardenburg-van Huisstede, Astrid

Title: Morbid obesity and asthma : co-morbidity or causal relationship?

Issue Date: 2015-10-13

3

Obesity and asthma: co-morbidity or causal relationship?

Van Huisstede A, Braunstahl GJ.

Monaldi Archives Chest Disease, 2010;73(3):116-23

ABSTRACT

There is substantial evidence that obesity and asthma are related. "Obese asthma" may be a unique phenotype of asthma, characterized by decreased lung volumes, greater symptoms for a given degree of lung function impairment, destabilization or lack of asthma control, lack of eosinophilic inflammation and a different response to controller medication.

Whether this relationship between obesity and asthma is causal or represents co-morbidity due to other factors is unclear. In previous reviews concerning the relationship between obesity and asthma, five hypotheses were put forth. One of these hypotheses is that a low grade systemic inflammation caused by adipokines from the fat tissue causes or enhances bronchial inflammation. In animal models, there is an increasing amount of evidence for the role of adipokines derived from fat tissue in the relationship between obesity and asthma. The data are conflicting in humans.

Since obesity is a component of the metabolic syndrome and the metabolic syndrome is also a form of systemic inflammation, it is to be expected that there is a relationship between metabolic syndrome and asthma. The few data that are available show that there is no relationship between metabolic syndrome and asthma, but there is one between the metabolic syndrome and asthma-like symptoms.

Further research is needed to confirm the relationship between obesity and asthma in humans, where a rigorous approach in the diagnosis of asthma is essential.

INTRODUCTION

Obesity is currently a major health problem and is developing into a global epidemic. Obesity is classified by the Body Mass Index (BMI) (table 1) and is defined as abnormal or excessive fat accumulation which can cause health problems. The World Health Organisation's (WHO) latest projections indicate that, globally in 2005, approximately 1.6 billion adults (age 15+) were overweight, and at least 400 million adults were obese. WHO further predicts that, by 2015, approximately 2.3 billion adults will be overweight and more than 700 million will be obese⁽¹⁾.

Because the prevalence of both obesity and asthma have increased in the recent years, many studies have examined the possibility of a causal relationship between these two conditions⁽²⁻⁵⁾. The fact that asthma improves after weight loss with either bariatric surgery or low caloric diet supports the relationship between obesity and asthma⁽⁶⁾. Moreover, obese patients with persistent asthma have significantly worse asthma-related quality of life⁽⁷⁾, less asthma control^(8, 9), more severe disease^(4, 10-12), and more asthma-related hospital admissions than asthma patients with a normal BMI.

There are multiple hypotheses that may explain the relationship between obesity and asthma. This review will discuss the different hypotheses, and will focus on the systemic inflammation, and the possible role of the metabolic syndrome.

Table 1 BMI classification

| BMI (kg/m ²) | Classification |
|--------------------------|----------------|
| <18.5 | Underweight |
| 18.5-25 | Normal weight |
| 25-30 | Overweight |
| 30-40 | Obesity |
| >40 | Morbid Obesity |

Classification of different BMI-classes. BMI, Body Mass Index

Asthma phenotype in obesity

Recent data suggest^(13, 14) that obese patients have a different phenotype of asthma, characterized by female predominance, late onset of asthma and high symptom expression.

Classical asthma is characterised by a Th-2 cell mediated allergic airway inflammation. The characteristics of airway inflammation in asthma (exhaled nitric oxide (eNO) and inflammatory cells in induced sputum) have been investigated in obesity. eNO can be seen as a marker of eosinophilic inflammation in the airways. De Winter *et al.* showed that a higher BMI was associated with a higher eNO in healthy patients⁽¹⁵⁾. On the con-

trary, Kazaks found no significant relationship between exhaled NO and BMI in adult asthma patients⁽¹⁶⁾. However, there was no correction for use of corticosteroids.

Sputum eosinophilia is a hallmark of asthma. Several studies have found no - or even an inverse - relationship between the number of sputum eosinophils and BMI⁽¹⁷⁻¹⁹⁾. Although airway eosinophil accumulation is characteristic for asthma, little is known about the influence of obesity on eosinophil migration from bone marrow to the bronchial mucosa. Calixto⁽²⁰⁾ found, in a murine model of asthma, that diet-induced obesity enhanced eosinophil trafficking from bone marrow to the airway, but delayed their transit through the airway epithelium into the airway lumen.

If the relationship between asthma and obesity could be explained by allergic airway inflammation, then one would expect a relationship between obesity and other allergic conditions, such as allergic rhinitis or allergic conjunctivitis. Although several epidemiologic studies have reported an increased risk of atopy in the overweight and/or obese^(21, 22), others have demonstrated an increased risk of developing asthma in non-allergic compared to allergic adults⁽²³⁻²⁵⁾.

In summary, there is a great deal of conflicting data concerning the signs of bronchial inflammation in obese asthmatics (eNO, sputum eosinophilia). Data suggest that asthma in obese patients is not characterized by a classic Th2 cell driven inflammation, but probably represents a different phenotype. The inflammatory pattern might be neutrophilic, instead of eosinophilic in obese patients with asthma. Allergy is probably not an important factor in the relationship between obesity and asthma.

Therapeutic implications

The main goal of asthma treatment is to achieve adequate control of the disease, as reflected by minimal symptoms and rescue bronchodilator use. Asthma has been reported to be more difficult to control in obese patients compared to individuals with a normal weight^(9, 26). Also the improvement of exhaled nitric oxide levels with inhaled corticosteroid treatment is smaller in obese asthmatics than in lean asthmatics⁽²⁷⁾. Is this because they do not respond to inhaled corticosteroids? This could be expected when one assumes that in obese asthmatics it is a neutrophilic instead of eosinophilic airway inflammation. In a recent study by Camargo⁽²⁸⁾, they found that, compared to subjects with normal BMI, the onset to peak FEV1 may require longer treatment exposure in the very obese.

Could it be that asthma in obese patients is more difficult to control because obese asthmatics have more complaints than lean asthmatics? Although obese asthmatics have the same perceptual responses to bronchoconstriction as lean asthmatics, it is to be expected that they have more complaints of dyspnoea due to their altered lung volumes, as will be discussed later.

Weight reduction from diet and exercise has been found to improve asthma symptoms, but the most striking changes have been reported in morbidly obese asthmatics following bariatric surgery⁽²⁹⁻³⁴⁾. However, these studies are frequently small and underpowered, and are often not designed with a primary outcome or rigorous characterization of asthma. Although there is still only a small body of evidence in favour of the beneficial effect of weight reduction in obese asthmatics, weight reduction does seem to be an important aspect of the treatment of these people.

Hypotheses that may explain the relationship between obesity and asthma

Since obese asthmatics might have a non-allergic phenotype of asthma, there must be another factor explaining the relationship between obesity and asthma. Although there are multiple hypotheses, no causal relationship between obesity and asthma has, as yet, been proven. Even though asthma may promote weight gain through increased sedentary life style and occasional use of oral corticosteroids, this does not fully explain the association of asthma and obesity in the majority of patients. In previous reviews concerning the relationship between obesity and asthma, five hypotheses were put forth (figure 1).

First, obese people consume food with less nutritional value, fewer vitamins and more fat. A high amount of fat intake is associated with asthma⁽³⁵⁾. Also zinc- and magnesium deficiencies are associated with asthma and bronchial hyperreactivity (BHR)⁽³⁶⁾.

Secondly, obesity and asthma may share the same genetic risk factors. This was demonstrated by a large-scale study among 1384 twins, where a strong association between asthma and BMI was found⁽³⁷⁾.

Co-morbidities, which are a risk factor for asthma as well as obesity, might be the third hypothesis. Obesity, for instance, is a risk factor for gastro-oesophageal reflux disease (GERD) which, in turn, is a risk factor for asthma⁽³⁸⁾. The same is true for Obstructive Sleep Apnoea Syndrome (OSAS); the prevalence of OSAS is higher in severe asthma patients as well as in obese patients⁽³⁹⁾. A large-scale, questionnaire-based Northern European survey showed that after adjustments for possible confounders, obesity and nocturnal GERD were found to be independent risk factors for the onset of asthma and respiratory symptoms⁽⁴⁰⁾.

Fourthly, obesity may alter lung function parameters. Typically, obesity causes a modest reduction in total lung capacity (TLC), and a larger reduction in functional residual capacity (FRC) (figure 2)⁽⁴¹⁾. The reasons for the reduction in TLC are not known, but they are probably due to a mechanical effect of the adipose tissue. A reduction in the downward movement of the diaphragm, due to increased abdominal mass is likely to decrease TLC by limiting the space for lung expansion on inflation⁽⁴²⁾. Furthermore, obese people have a smaller tidal volume and an increased respiration rate with a decreased expiratory reserve volume (ERV)⁽⁴³⁾. The FEV1/FVC ratio is usually well preserved




Figure 1 Obesity and asthma relationship

The relationship between obesity and asthma is not yet clear. Diet, genetic factors or comorbidities which give a higher risk for asthma might also give a higher risk for obesity. Obesity gives a greater abdominal mass, which causes altered lung function parameters, which cause dyspnea. Whether or not this dyspnea is asthma is questionable. The abdominal mass also causes a dysbalance in inflammatory proteins, in favour of the pro-inflammatory proteins. This may lead to a local inflammation, which causes a low grade systemic inflammation, which, in turn, causes or enhances a local inflammation of the lung parenchyma.

or increased, which implies that the major effect of obesity is on lung volumes, with no direct effect on airway obstruction. Furthermore, weight loss significantly improves the FRC, TLC and ERV⁽⁴⁴⁾.

The decreased lung volume is associated with a decreased diameter of the peripheral airways, which leads to tidal breathing at or near closing volume⁽⁴¹⁾. Moreover, Bergeron has shown that there is a fat accumulation in the airways in obese patients with less defined cartilage ring⁽⁴⁵⁾. All this can explain why obese patients often have more complaints of dyspnea than subjects with a normal BMI. Salome showed that, although obesity reduces lung volumes, it does not alter sensitivity or maximal response to methacholine in non-asthmatic subjects⁽⁴⁶⁾. Recently, it was also shown that perceptual responses to methacholine-induced bronchoconstriction and lung hyperinflation were similar in obese and normal-weight asthmatics⁽⁴⁷⁾. Thus, obese asthmatics may have more complaints of dyspnea, but this may not be directly related to abnormal airway sensitivity, but may reflect other physiological effects of obesity.




Figure 2 Altered lung function in obesity
Obesity leads to alternations of lung volumes. ERV: expiratory reserve volume; FRC: functional residual capacity; RV: residual volume; TLC: total lung capacity

The fifth, and last, hypothesis suggests that systemic inflammation by means of fat-tissue derived adipokines may lead to asthma. The assumption is that inflammatory mediators from the fat tissue come into the systemic circulation, and find their way to the lung parenchyma, where they may cause or intensify airway inflammation.

Obesity is considered to be a state of chronic low-grade systemic inflammation, characterized by a dysbalance of pro- and anti-inflammatory proteins derived from fat tissue cells. Fat tissue was traditionally seen as an organ for storage of energy. Recently, however, it is considered to act as an endocrine organ. The fat tissue is a source of bioactive peptides, which are called adipokines⁽⁴⁸⁾. In 1993, Hotamisligil *et al* discovered that fat cells (adipocytes) in rodents excrete TNF- α . TNF- α is a pro-inflammatory cytokine⁽⁴⁹⁾. Since then, multiple inflammatory peptides have been discovered which have a relationship with obesity. Examples of adipokines are leptin, adiponectin and resistin. Also interleukin (IL)-6 and TNF- α play a role. They are produced by macrophages in the fat tissue.

Leptin (Greek leptos, thin) is excreted by fat tissue and causes the feeling of saturation and increases the metabolism. Moreover, leptin influences the T-cell immunity response and stimulates the proliferation of T-helper cells, which causes increased production of pro-inflammatory cytokines⁽⁵⁰⁾. There is an increased leptin production^(51, 52) and a relative leptin resistance in obese patients. A comparison with insulin and insulin resistance can be made: here the primary reaction of the body is to increase the production of insulin so that the same physical result is achieved. The leptin concentration is not only increased in obesity, but also in asthma^(53, 54).

Obesity is associated with necrosis of the adipocytes, causing the interstitium to be exposed to little fat drops. Macrophages will collect around this necrotizing adipocytes and become active. These macrophages will produce TNF- α and IL-6, which will cause a decrease in local production of adiponectin⁽⁵⁵⁾. Adiponectin is an insulin-regulating hormone. It also has anti-inflammatory effects: it decreases the production of pro-

inflammatory cytokines and it increases the production of IL-10 and IL-1. Plasma adiponectin levels are reversely correlated to BMI⁽⁵⁶⁾, but there is no clear relationship between adiponectin and asthma⁽⁵⁷⁻⁶⁰⁾.

Xu *et al.*⁽⁶¹⁾ have been able to show in a mice model that there is histological evidence for a significant infiltration of macrophages in the fat tissue of obese mice. Moreover, they also have shown that many inflammatory and specific macrophage genes are dramatically upregulated in the fat tissue. This suggests that macrophages in the fat tissue play an active role in obesity and macrophage-related inflammation. Weisberg *et al.*⁽⁶²⁾ showed that the macrophages are responsible for almost all expression of TNF- α and IL-6 in the fat tissue. TNF- α is a potent pro-inflammatory cytokine and is involved in many processes in the human body, in particular inflammation, cell proliferation, cell differentiation, apoptosis and fat metabolism. IL-6 regulates the production of the strong pro-inflammatory protein C-reactive protein (CRP).

In summary, in animal models the influence of the different adipokines associated with obesity seem to have an effect on the airway inflammation. In humans, however, the association is not yet confirmed. Further studies are needed to investigate whether a dysbalance in pro and anti-inflammatory cytokines in the systemic circulation may induce a change in concentration of the same cytokines in the lung tissue. This may result in an increased airway inflammation.

Metabolic syndrome

The fact that asthma improves after weight loss either with bariatric surgery or low caloric diet supports the relation between obesity and asthma. Moreover, Cottam found that obesity was manifested by a chronic inflammation (including elevation of monocytes and eosinophils), which appeared to be reversible within 6 months after bariatric surgery⁽⁶³⁾. Also, weight loss by low-energy diet was associated with a reduction in markers of systemic inflammation^(64, 65). So the hypothesis that the relationship between obesity and asthma is based on a systemic inflammation seems quite plausible. The metabolic syndrome might contribute to this low-grade systemic inflammation.

Recently, the metabolic syndrome has been postulated as a new hypothesis which might explain the relationship between obesity and asthma. Obesity, and mainly central obesity, is an important component of the metabolic syndrome. The metabolic syndrome is a term to cluster type 2 diabetes mellitus, hypertension, dyslipidemia and central obesity (table 2).

Table 2 Metabolic syndrome

| Risk factor | Objective |
|---|---|
| Abdominal obesity (waist circumference) | |
| - male | > 102 cm |
| - female | > 88 cm |
| Triglycerides | >1,7 mmol/l or treatment for hypertriglycercaemia |
| HDL-cholesterol | |
| - male | <1,03 mmol/l, or treatment for low HDL-C |
| - female | < 1,3 mmol/l or treatment for low HDL-C |
| Blood pressure | ≥130/≥85 mmHg, or treatment for hypertension |
| Blood glucose (plasma) | ≥ 6,1 mmol/l, or treatment for hyperglycaemia |

Definition of the metabolic syndrome according to NCEP-ATPIII criteria on basis of clinical characteristics. Metabolic syndrome when 3 out of 5 characteristics.

Leone found an independent relationship between lung function impairment (FEV1 or FVC < lower limit of normal) and metabolic syndrome in both sexes, predominantly due to abdominal obesity⁽⁶⁶⁾. Lee *et al*⁽⁶⁷⁾ found that the metabolic syndrome was associated with asthma symptoms. The evidence for the relationship between metabolic syndrome and asthma is scarce, mainly because it is a new idea. More research is needed to make a definitive conclusion.

The metabolic syndrome is a way to group different diseases. If there might be a relationship between the metabolic syndrome and asthma, there might also be a relationship between the separate compounds of the metabolic syndrome and asthma. In a cross-sectional population-based study, obesity and insulin resistance were associated with an increased risk of aeroallergen sensitization and asthma (allergic and non-allergic)⁽⁶⁸⁾. Thuesen⁽⁶⁹⁾ found, in a recent population-based, prospective study among 1443 adults, that insuline-resistance is a stronger risk factor for asthma symptoms than obesity. Hyperglycemia itself may not be the important factor. Shore *et al* have demonstrated in lean wild-type and leptin receptor-deficient, diabetic, obese mice that administration of metformin did not affect ozone-induced airway inflammation⁽⁷⁰⁾.

More research is needed to see whether the metabolic syndrome, or it separate components, are the "missing link" between obesity and asthma. Also here the same remarks as for the fat tissue derived adipokines can be made. If it is so that the metabolic syndrome contributes to the low grade systemic inflammation, the question remains whether this low grade systemic inflammation has its local effects on the lung parenchyma. And whether this systemic inflammation causes or enhances the bronchial inflammation.

Critical notes

Many of the articles discussed in this review are epidemiologic studies among patients with asthma, where asthma is defined as either physician-diagnosed or self-reported. Very few studies performed (extensive) lung function tests to confirm the diagnosis of asthma. Often the diagnosis is based on symptoms and medication use. The question is whether these patients truly have asthma. In a retrospective study among children, a strong correlation was found between general practitioner and specialist diagnosed asthma, irrespective of BMI⁽⁷¹⁾. In an extensive study among adult patients, the diagnosis of asthma was revisited. In 31.8% of the obese, and 28.7% of the non-obese group ($p=0.46$) the diagnosis of asthma was rejected. The percentage of misdiagnosis is the same in the obese and non-obese group, so the relationship between asthma and obesity cannot be explained by the fact that there is an overdiagnosis of asthma in the obese population⁽⁷²⁾. In a recent study Pakhale *et al.*⁽⁷³⁾ found a misdiagnosis of asthma in up to 30% of patients. They used a rigorous diagnostic testing algorithm to confirm the diagnosis of asthma. They found that misdiagnosis of asthma was associated with gender, a recent diagnosis, older age and increased FEV₁, but not with obesity. Also, obese subjects who made urgent healthcare visits for respiratory symptoms were more likely to be misdiagnosed with asthma. It is possible that these symptoms could be the result of increased effort in breathing associated with obesity, rather than airway inflammation or airway hyperresponsiveness.

Another disadvantage of most studies is that they are based on self-reported weight. A BMI based on self-reported height and weight is not accurate for either children or adults⁽⁷⁴⁾. The weight will be underreported, and the length overreported. There is however, an algorithm to correct for this.

Some previous studies of the association of adiposity and lung function have used only BMI as a marker of obesity, without taking body composition and fat distribution into account. The waist circumference - or even better, a MRI-based visceral fat quantification - is possibly a more accurate measure than BMI. The distribution of body fat may be an important determinant of lung function, and this may account for the more pronounced association between central adiposity and obstructive lung function in men as compared to women⁽⁷⁵⁾. Steele *et al.* performed a study to examine the relationship among body fatness, fat distribution, and lung function, adjusted for physical activity, energy expenditure and aerobic fitness⁽⁷⁶⁾. They found that obesity is inversely associated with lung function (FEV₁ and FVC) in adults. Central fat distribution, however, appears to have a stronger relationship to respiratory mechanics (FEV₁) in men than in women. These associations were independent of the degree of physical activity in this cohort.

CONCLUSION

From the literature discussed, it can be concluded that there is a relationship between obesity and asthma. "Obese asthma" may be a unique phenotype of asthma, characterized by decreased lung volumes, greater symptoms for a given degree of lung function impairment, destabilization or lack of asthma control, lack of eosinophilic inflammation and a different response to controller medication. Therefore, the clinical evaluation of an obese patient with asthma must require a more rigorous and objective approach.

Whether this relationship is really causal or represents parallel co-morbidities is unclear. In animal models, there is an increasing amount of evidence for the role of adipokines derived from fat tissue in the relationship between obesity and asthma. These adipokines cause a low grade systemic inflammation, which might cause or enhance bronchial inflammation. The data are conflicting in humans. However, the fact that weight loss improves asthma control and normalizes the concentration of adipokines in the serum implies that there must be some role for adipokines in the relationship between obesity and asthma.

Since obesity is a component of the metabolic syndrome and the metabolic syndrome is also a form of systemic inflammation, it is to be expected that there is a relationship between metabolic syndrome and asthma. The few data that are available show that there is no relationship between metabolic syndrome and asthma, but there is one between the metabolic syndrome and asthma-like symptoms.

Further research is needed to confirm the relationship between obesity and asthma in humans, where a rigorous approach in the diagnosis of asthma and use of good control groups is essential. Intervention-based longitudinal research might find a definitive answer to the question of what the basis of the relationship between obesity and asthma could be.

ACKNOWLEDGEMENTS

We wish to thank Mrs. Reijnhart for editing the manuscript.

REFERENCES

1. Organisation WH. Obesity and overweight, fact sheet No 311. 2006 [cited; Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/print.html>]
2. James AL, Knuiman MW, Divitini ML, Hui J, Hunter M, Palmer LJ, Maier G, Musk AB. Changes in the prevalence of asthma in adults since 1966: the Busselton health study. *Eur Respir J* 2010;35(2): 273-8.
3. Chen Y, Rennie D, Cormier Y, Dosman J. Atopy, obesity, and asthma in adults: the Humboldt study. *J Agromedicine* 2009;14(2):222-7.
4. Taylor B, Mannino D, Brown C, Crocker D, Twum-Baah N, Holguin F. Body mass index and asthma severity in the National Asthma Survey. *Thorax* 2008;63(1):14-20.
5. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med* 2007;175(7):661-6.
6. Camargo CA, Jr., Weiss ST, Zhang S, Willett WC, Speizer FE. Prospective study of body mass index, weight change, and risk of adult-onset asthma in women. *Arch Intern Med* 1999;159(21):2582-8.
7. Lavoie KL, Bacon SL, Labrecque M, Cartier A, Ditto B. Higher BMI is associated with worse asthma control and quality of life but not asthma severity. *Respir Med* 2006;100(4):648-57.
8. Saint-Pierre P, Bourdin A, Chanez P, Daures JP, Godard P. Are overweight asthmatics more difficult to control? *Allergy* 2006;61(1):79-84.
9. Clerisme-Beaty EM, Karam S, Rand C, Patino CM, Bilderback A, Riekert KA, Okelo SO, Diette GB. Does higher body mass index contribute to worse asthma control in an urban population? *J Allergy Clin Immunol* 2009;124(2):207-12.
10. Mosen DM, Schatz M, Magid DJ, Camargo CA, Jr. The relationship between obesity and asthma severity and control in adults. *J Allergy Clin Immunol* 2008;122(3):507-11 e6.
11. Akerman MJ, Calacanis CM, Madsen MK. Relationship between asthma severity and obesity. *J Asthma* 2004;41(5):521-6.
12. Varraso R, Siroux V, Maccario J, Pin I, Kauffmann F. Asthma severity is associated with body mass index and early menarche in women. *Am J Respir Crit Care Med* 2005;171(4):334-9.
13. Haldar P, Pavord ID, Shaw DE, Berry MA, Thomas M, Brightling CE, Wardlaw AJ, Green RH. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med* 2008;178(3):218-24.
14. Lugogo NL, Kraft M, Dixon AE. Does Obesity Produce a Distinct Asthma Phenotype? *J Appl Physiol* 2010;108(3):729-34.
15. De Winter-de Groot KM, Van der Ent CK, Prins I, Tersmette JM, Uiterwaal CS. Exhaled nitric oxide: the missing link between asthma and obesity? *J Allergy Clin Immunol* 2005;115(2):419-20.
16. Kazaks A, Uriu-Adams JY, Stern JS, Albertson TE. No significant relationship between exhaled nitric oxide and body mass index in people with asthma. *J Allergy Clin Immunol* 2005;116(4):929-30; author reply 930.
17. Todd DC, Armstrong S, D'Silva L, Allen CJ, Hargreave FE, Parameswaran K. Effect of obesity on airway inflammation: a cross-sectional analysis of body mass index and sputum cell counts. *Clin Exp Allergy* 2007;37(7):1049-54.
18. Sutherland TJ, Cowan JO, Young S, Goulding A, Grant AM, Williamson A, Brassett K, Herbison GP, Taylor DR. The association between obesity and asthma: interactions between systemic and airway inflammation. *Am J Respir Crit Care Med* 2008;178(5):469-75.
19. van Veen IH, Ten Brinke A, Sterk PJ, Rabe KF, Bel EH. Airway inflammation in obese and nonobese patients with difficult-to-treat asthma. *Allergy* 2008;63(5):570-4.

20. Calixto M, Lintomen L, Schenka A, Saad MJ, Zanesco A, Antunes E. Obesity enhances eosinophilic inflammation in a murine model of allergic asthma. *Br J Pharmacol* 2013;159(3):617-25.
21. Hancox RJ, Milne BJ, Poulton R, Taylor DR, Greene JM, McLachlan CR, Cowan JO, Flannery EM, Herbison GP, Sears MR. Sex differences in the relation between body mass index and asthma and atopy in a birth cohort. *Am J Respir Crit Care Med* 2005;171(5):440-5.
22. Schachter LM, Peat JK, Salome CM. Asthma and atopy in overweight children. *Thorax* 2003;58(12):1031-5.
23. Chen Y, Dales R, Jiang Y. The association between obesity and asthma is stronger in nonallergic than allergic adults. *Chest* 2006;130(3):890-5.
24. Jarvis D, Chinn S, Potts J, Burney P. Association of body mass index with respiratory symptoms and atopy: results from the European Community Respiratory Health Survey. *Clin Exp Allergy* 2002;32(6):831-7.
25. Loerbroks A, Apfelbacher CJ, Amelang M, Sturmer T. Obesity and adult asthma: potential effect modification by gender, but not by hay fever. *Ann Epidemiol* 2008;18(4):283-9.
26. Camargo CA, Jr., Sutherland ER, Bailey W, Castro M, Yancey SW, Emmett AH, Stempel DA. Effect of increased body mass index on asthma risk, impairment and response to asthma controller therapy in African Americans. *Curr Med Res Opin* 2010;26(7):1629-35.
27. Sutherland ER, Lehman EB, Teodorescu M, Wechsler ME. Body mass index and phenotype in subjects with mild-to-moderate persistent asthma. *J Allergy Clin Immunol* 2009;123(6):1328-34 e1.
28. Camargo CA, Jr., Boulet LP, Sutherland ER, Busse WW, Yancey SW, Emmett AH, Ortega HG, Ferro TJ. Body mass index and response to asthma therapy: fluticasone propionate/salmeterol versus montelukast. *J Asthma*;47(1):76-82.
29. Aaron SD, Fergusson D, Dent R, Chen Y, Vandemheen KL, Dales RE. Effect of weight reduction on respiratory function and airway reactivity in obese women. *Chest* 2004;125(6):2046-52.
30. Stenius-Aarniala B, Poussa T, Kvarnstrom J, Gronlund EL, Ylikahri M, Mustajoki P. Immediate and long term effects of weight reduction in obese people with asthma: randomised controlled study. *Bmj* 2000;320(7238):827-32.
31. Eneli IU, Skybo T, Camargo CA, Jr. Weight loss and asthma: a systematic review. *Thorax* 2008;63(8):671-6.
32. Dixon JB, Chapman L, O'Brien P. Marked improvement in asthma after Lap-Band surgery for morbid obesity. *Obes Surg* 1999;9(4):385-9.
33. Hakala K, Stenius-Aarniala B, Sovijarvi A. Effects of weight loss on peak flow variability, airways obstruction, and lung volumes in obese patients with asthma. *Chest* 2000;118(5):1315-21.
34. Nguyen NT, Hinojosa MW, Smith BR, Gray J, Varela E. Improvement of restrictive and obstructive pulmonary mechanics following laparoscopic bariatric surgery. *Surg Endosc* 2009;23(4):808-12.
35. Weiss ST. Obesity: insight into the origins of asthma. *Nat Immunol* 2005;6(6):537-9.
36. Tantisira KG, Weiss ST. Complex interactions in complex traits: obesity and asthma. *Thorax* 2001; 56 Suppl 2:ii64-73.
37. Hallstrand TS, Fischer ME, Wurfel MM, Afari N, Buchwald D, Goldberg J. Genetic pleiotropy between asthma and obesity in a community-based sample of twins. *J Allergy Clin Immunol* 2005; 116(6):1235-41.
38. Kiljander TO, Harding SM, Field SK, Stein MR, Nelson HS, Ekelund J, Illueca M, Beckman O, Sostek MB. Effects of esomeprazole 40 mg twice daily on asthma: a randomized placebo-controlled trial. *Am J Respir Crit Care Med* 2006;173(10):1091-7.
39. Julien JY, Martin JG, Ernst P, Olivenstein R, Hamid Q, Lemiere C, Pepe C, Naor N, Olha A, Kimoff RJ. Prevalence of obstructive sleep apnea-hypopnea in severe versus moderate asthma. *J Allergy Clin Immunol* 2009;124(2):371-6.

40. Gunnbjornsdottir MI, Omenaas E, Gislason T, Norrman E, Olin AC, Jogi R, Jensen EJ, Lindberg E, Bjornsson E, Franklin K, Janson C, Gulsvik A, Laerum B, Svanes C, Toren K, Tunsater A, Lillienberg L, Gislason D, Blondal T, Bjornsdottir US, Jorundsdottir KB, Talvik R, Forsberg B, Franklin K, Lundback B, Soderberg M, Ledin MC, Boman G, Norback D, Wieslander G, Spetz-Nystrom U, Cashelunge KS, Ryden E. Obesity and nocturnal gastro-oesophageal reflux are related to onset of asthma and respiratory symptoms. *Eur Respir J* 2004;24(1):116-21.
41. Beuther DA, Weiss ST, Sutherland ER. Obesity and asthma. *Am J Respir Crit Care Med* 2006;174(2):112-9.
42. Salome CM, King GG, Berend N. Physiology of obesity and effects on lung function. *J Appl Physiol* 2010;108(1):206-11.
43. Beuther DA. Obesity and asthma. *Clin Chest Med* 2009;30(3):479-88.
44. Weiner P, Waizman J, Weiner M, Rabner M, Magadle R, Zamir D. Influence of excessive weight loss after gastroplasty for morbid obesity on respiratory muscle performance. *Thorax* 1998;53(1):39-42.
45. Bergeron C, Boulet LP, Hamid Q. Obesity, allergy and immunology. *J Allergy Clin Immunol* 2005;115(5):1102-4.
46. Salome CM, Munoz PA, Berend N, Thorpe CW, Schachter LM, King GG. Effect of obesity on breathlessness and airway responsiveness to methacholine in non-asthmatic subjects. *Int J Obes (Lond)* 2008;32(3):502-9.
47. Deesomchok A, Fisher T, Webb KA, Ora J, Lam YM, Loughheed MD, O'Donnell DE. Effects of Obesity on Perceptual and Mechanical Responses to Bronchoconstriction in Asthma. *Am J Respir Crit Care Med* 2009.
48. Ronti T, Lupattelli G, Mannarino E. The endocrine function of adipose tissue: an update. *Clin Endocrinol (Oxf)* 2006;64(4):355-65.
49. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science* 1993;259(5091):87-91.
50. Lord GM, Matarese G, Howard JK, Baker RJ, Bloom SR, Lechler RI. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* 1998;394(6696):897-901.
51. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, Ohannesian JP, Marco CC, McKee LJ, Bauer TL, et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med* 1996;334(5):292-5.
52. Couillard C, Mauriege P, Imbeault P, Prud'homme D, Nadeau A, Tremblay A, Bouchard C, Despres JP. Hyperleptinemia is more closely associated with adipose cell hypertrophy than with adipose tissue hyperplasia. *Int J Obes Relat Metab Disord* 2000;24(6):782-8.
53. Guler N, Kirerleri E, Ones U, Tamay Z, Salmayenli N, Darendeliler F. Leptin: does it have any role in childhood asthma? *J Allergy Clin Immunol* 2004;114(2):254-9.
54. Mai XM, Bottcher MF, Leijon I. Leptin and asthma in overweight children at 12 years of age. *Pediatr Allergy Immunol* 2004;15(6):523-30.
55. Bruun JM, Lihn AS, Verdich C, Pedersen SB, Toubro S, Astrup A, Richelsen B. Regulation of adiponectin by adipose tissue-derived cytokines: in vivo and in vitro investigations in humans. *Am J Physiol Endocrinol Metab* 2003;285(3):E527-33.
56. Engeli S, Feldpausch M, Gorzelnik K, Hartwig F, Heintze U, Janke J, Mohlig M, Pfeiffer AF, Luft FC, Sharma AM. Association between adiponectin and mediators of inflammation in obese women. *Diabetes* 2003;52(4):942-7.
57. Kim KW, Shin YH, Lee KE, Kim ES, Sohn MH, Kim KE. Relationship between adipokines and manifestations of childhood asthma. *Pediatr Allergy Immunol* 2008;19(6):535-40.

58. Sutherland TJ, Sears MR, McLachlan CR, Poulton R, Hancox RJ. Leptin, adiponectin, and asthma: findings from a population-based cohort study. *Ann Allergy Asthma Immunol* 2009;103(2):101-7.
59. Sood A, Cui X, Qualls C, Beckett WS, Gross MD, Steffes MW, Smith LJ, Jacobs DR, Jr. Association between asthma and serum adiponectin concentration in women. *Thorax* 2008;63(10):877-82.
60. Jartti T, Saarikoski L, Jartti L, Lisinen I, Jula A, Huupponen R, Viikari J, Raitakari OT. Obesity, adipokines and asthma. *Allergy* 2009;64(5):770-7.
61. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 2003;112(12):1821-30.
62. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003;112(12):1796-808.
63. Cottam DR, Schaefer PA, Shaftan GW, Velcu L, Angus LD. Effect of surgically-induced weight loss on leukocyte indicators of chronic inflammation in morbid obesity. *Obes Surg* 2002;12(3):335-42.
64. Esposito K, Pontillo A, Di Palo C, Giugliano G, Masella M, Marfella R, Giugliano D. Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *Jama* 2003;289(14):1799-804.
65. Bastard JP, Jardel C, Bruckert E, Blondy P, Capeau J, Laville M, Vidal H, Hainque B. Elevated levels of interleukin 6 are reduced in serum and subcutaneous adipose tissue of obese women after weight loss. *J Clin Endocrinol Metab* 2000;85(9):3338-42.
66. Leone N, Courbon D, Thomas F, Bean K, Jengo B, Leynaert B, Guize L, Zureik M. Lung function impairment and metabolic syndrome: the critical role of abdominal obesity. *Am J Respir Crit Care Med* 2009;179(6):509-16.
67. Lee EJ, In KH, Ha ES, Lee KJ, Hur GY, Kang EH, Jung KH, Lee SY, Kim JH, Lee SY, Shin C, Shim JJ, Kang KH, Yoo SH. Asthma-like symptoms are increased in the metabolic syndrome. *J Asthma* 2009;46(4):339-42.
68. Husemoen LL, Glumer C, Lau C, Pisinger C, Morch LS, Linneberg A. Association of obesity and insulin resistance with asthma and aeroallergen sensitization. *Allergy* 2008;63(5):575-82.
69. Thuesen BH, Husemoen LL, Hersoug LG, Pisinger C, Linneberg A. Insulin resistance as a predictor of incident asthma-like symptoms in adults. *Clin Exp Allergy* 2009;39(5):700-7.
70. Shore SA, Williams ES, Zhu M. No effect of metformin on the innate airway hyperresponsiveness and increased responses to ozone observed in obese mice. *J Appl Physiol* 2008;105(4):1127-33.
71. Lang JE, Feng H, Lima JJ. Body mass index-percentile and diagnostic accuracy of childhood asthma. *J Asthma* 2009;46(3):291-9.
72. Aaron SD, Vandemheen KL, Boulet LP, Mclvor RA, Fitzgerald JM, Hernandez P, Lemiere C, Sharma S, Field SK, Alvarez GG, Dales RE, Doucette S, Fergusson D. Overdiagnosis of asthma in obese and nonobese adults. *CMAJ* 2008;179(11):1121-31.
73. Pakhale S, Doucette S, Vandemheen K, Boulet LP, Mclvor RA, Fitzgerald JM, Hernandez P, Lemiere C, Sharma S, Field SK, Alvarez GG, Dales RE, Aaron SD. A comparison of obese and nonobese asthmatics: Exploring an asthma-obesity interaction. *Chest* 2010;137(6):1316-1323.
74. Gorber SC, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007;8(4):307-26.
75. Harik-Khan RI, Wise RA, Fleg JL. The effect of gender on the relationship between body fat distribution and lung function. *J Clin Epidemiol* 2001;54(4):399-406.
76. Steele RM, Finucane FM, Griffin SJ, Wareham NJ, Ekelund U. Obesity is associated with altered lung function independently of physical activity and fitness. *Obesity (Silver Spring)* 2009;17(3):578-84.

