



Universiteit  
Leiden  
The Netherlands

## Drug-target residence time : a case for the adenosine A1 and A2A receptors

Guo, D.

### Citation

Guo, D. (2014, June 25). *Drug-target residence time : a case for the adenosine A1 and A2A receptors*. Retrieved from <https://hdl.handle.net/1887/26833>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/26833>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/26833> holds various files of this Leiden University dissertation.

**Author:** Guo, Dong

**Title:** Drug-target residence time : a case for the adenosine A1 and A2A receptors

**Issue Date:** 2014-06-25

# Chapter 2

## *Drug-target residence time A case for G protein-coupled receptors*

Guo D, Hillger JM, IJzerman AP, Heitman LH.  
Adapted from: *Med Res Rev.* 2014, DOI: 10.1002/med.21307



## About this chapter

A vast number of marketed drugs act on G protein-coupled receptors (GPCRs), the most successful category of drug targets to date. These drugs usually possess high target affinity and selectivity, and such combined features have been the driving force in the early phases of drug discovery. However, attrition has also been high. Many investigational new drugs eventually fail in clinical trials due to a demonstrated lack of efficacy. A retrospective assessment of successfully launched drugs revealed that their beneficial effects in patients may be attributed to their long drug-target residence times. Likewise, for some other GPCR drugs short residence time could be beneficial to reduce the potential for on-target side effects. Hence, the compounds' kinetics behavior might in fact be the guiding principle to obtain a desired and durable effect *in vivo*. We therefore propose that drug-target residence time should be taken into account as an additional parameter in the lead selection and optimization process. This should ultimately lead to an increased number of candidate drugs moving to the pre-clinical development phase and on to the market. This chapter contains examples of the kinetics behavior of GPCR ligands with improved *in vivo* efficacy and summarizes methods for assessing drug-target residence time.

## 2.1 Introduction

Drug-target residence time (RT) is an emerging concept in the drug research community. Its definition, coined by Copeland in 2006, stands for an experimental measure of the lifetime of a drug-target complex, reflected by its dissociation rate ( $k_{\text{off}}$ ).<sup>1</sup> This is an important pharmacological feature as a drug is only effective when bound to, and influencing the activity of its physiological target.<sup>1,2</sup> Recently, several notable reviews including Swinney *et al.*,<sup>2-5</sup> Copeland *et al.*,<sup>1,6,7</sup> Vauquelin *et al.*,<sup>8-11</sup> and Zhang and Monsma,<sup>12,13</sup> to name a few, have emphasized the pivotal role of RT in the early phases of drug discovery. These discussions led to increasing recognition that drug-target residence time may even be of greater importance for a drug's effect in the patient than its affinity, since the body, unlike an *in vitro* setting, is an open system where the concentration of free drug fluctuates over time.<sup>13,14</sup> The dynamic flow *in vivo*—absorption, distribution, metabolism, and excretion (ADME)—often prevents the free drug from reaching equilibrium conditions that are otherwise readily obtained in a test tube.<sup>13,14</sup> In this sense, optimizing drug candidates based on equilibrium-derived parameters alone, such as affinity ( $K_i$ ) / potency ( $IC_{50}$ ), may not be ideal.<sup>15</sup> Drug-target residence time may be a useful additional parameter as it is thought to represent a surrogate marker of drug clinical efficacy: the longer the drug occupies the receptor, the more profound the drug may exert its effect.<sup>1,12</sup> Although this reasoning most certainly is too simplistic, there is substantial evidence, as witnessed in a survey of 50 drugs on 12 different drug targets, that about 70% of long residence time therapeutics displayed higher efficacy than comparable faster dissociating drugs.<sup>5</sup>

It needs to be pointed out that the criteria for 'long' or 'short' residence time may vary for different targets and for divergent clinical indications.<sup>16</sup> For therapies requiring prolonged target occupancy, a long RT drug offers advantages as it remains bound to the target and continuously exerts its pharmacological effect even when most of the free drug has already been eliminated from circulation.<sup>17</sup> Importantly, this simultaneously implies that a slowly dissociating drug will only offer a noticeably increased effect over a faster dissociating drug, when its dissociation half-life ( $t_{1/2}$ ) outlasts its pharmacokinetic half-life, or when there are large fluctuations in the endogenous competitor (e.g., hormone or neurotransmitter) concentrations.<sup>11</sup> Further proof for the importance of RT comes from numerous examples

of drugs that are best-in-class due to slow dissociation from their target receptor. One example is the well-known muscarinic  $M_3$  receptor antagonist, tiotropium, which is a long-acting, 24 hour, anticholinergic bronchodilator used in the management of chronic obstructive pulmonary disease.<sup>29</sup> On the other hand, there are cases where the mechanism-based toxicity outweighs the therapeutic advantages by long receptor occupancy. In this situation, a fast dissociating compound displaying short-lived intervention is preferred. This may be the case for the antipsychotic  $D_2$  dopamine receptor antagonists, where long RT is not desired due to on-target toxicity associated with strong extrapyramidal motor effects. We will discuss this in a later part of this chapter.

## 2.2 Mathematical definitions of drug-target residence time

Drug-target residence time is equal to the reciprocal of the dissociation rate constant ( $RT = 1 / k_{\text{off}}$ ).<sup>1</sup> Its value is interchangeable with the dissociation half-life ( $t_{1/2} = \ln 2 / k_{\text{off}}$ ) with a factor of  $\ln 2$ , i.e.,  $RT = t_{1/2} / \ln 2$ . Both can be used to represent the duration of the drug-target binary complex. For the convenience of discussion, we chose to use the dissociation half-life to describe the drug-target residence time in the following part of this chapter, since most of the kinetic values in literature were reported with  $t_{1/2}$  values.

As mentioned above, it is necessary to determine the dissociation rate constant to calculate the RT or  $t_{1/2}$ . However, the definition of  $k_{\text{off}}$  depends on the specific mechanism of ligand-receptor interaction.<sup>1,14</sup> In general, three situations are well known. Equation 1 represents a simple one-step association and dissociation process, where the receptor (R) and ligand (L) encounter each other to form a binary complex (RL) with association rate  $k_1$  ( $k_{\text{on}}$ ) and dissociation rate  $k_2$  ( $k_{\text{off}}$ ), respectively.



In contrast, Equation 2 represents a more complicated process. Upon the binding of the ligand to the receptor, the initial complex (RL) undergoes a conformation isomerization leading to a much higher-affinity final complex

(RL\*). This is also known as the ‘induced-fit’ mechanism. In this case,  $k_{\text{off}}$  entails both forward ( $k_3$ ) and reverse steps ( $k_4$ ) in receptor isomerization as well as the on- and off-rate constants for the free ligand bound to and unbound from the free receptor, such that  $k_{\text{off}} = k_2 \cdot k_4 / (k_2 + k_3 + k_4)$ . This isomerization step enhances the interaction between the ligand and the target, hence resulting in an extended overall  $t_{1/2}$ .



A third mechanism is represented in Equation 3, where the receptor is in equilibrium between two conformational states (R and R\*) in the absence of ligand. Of these conformational states, the ligand specifically binds to one, R\*, rather than the other (R), to form the R\*L complex. This is also known as the ‘conformation-selection’ mechanism. The interconversion between conformational states R and R\* is slow relative to the binding of the ligand to the R\* state, thus it represents the rate-limiting step for the formation of the binary complex. For the dissociation rate constant, its value is equal to  $k_4$ .



Among these three mechanisms, the first situation is most commonly encountered in ligand-GPCR binding kinetics, while the second mechanism can be the case for high-affinity ligand-receptor interactions, especially for enzyme inhibitors and receptor antagonists;<sup>1,6,14</sup> the third situation is rarely observed in ligand-receptor interactions.<sup>14</sup>

## 2.3 Experimental strategies to assess drug-target residence time


Measurement of drug-target complex lifetime, reflected by  $k_{\text{off}}$  or  $t_{1/2}$ , is not a new phenomenon *per se*. To date, there are several strategies available to examine the binding kinetics of ligand-GPCR interactions. The advantages and disadvantages / limitations of these experimental strategies to access drug-target residence time are briefly outlined in Table 2.1. Of these



**Table 2.1 | The advantages and disadvantages / limitations of experimental strategies to access drug-target residence time.**

| Assay                                                            | Advantages                                                                                                                         | Disadvantages / limitations                                                                           | Ref.  |
|------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|-------|
| Direct kinetic radioligand binding assay                         | Straightforward;<br>Robust;<br>Quantitative measurement                                                                            | Ligand of interest needs to be radiolabeled;<br>Low throughput assay format                           | 19-21 |
| Indirect kinetic radioligand binding assay                       | Only one labeled ligand necessary;<br>Quantitative measurement;<br>Adjustable for high throughput screening                        | Need for suitable radiolabeled ligand;<br>Intermediate filtration and / or 'washout' steps are needed | 23-26 |
| Competition association assay                                    | Only one labeled ligand necessary;<br>Quantitative measurement                                                                     | Need for suitable radiolabeled ligand;<br>Low throughput assay format                                 | 27-34 |
| Dual-point competition association assay                         | Only one labeled ligand necessary;<br>High throughput assay format;<br>Easy readout for longer or shorter residence time compounds | Need for suitable radiolabeled ligand;<br>Qualitative measurement                                     | 18,33 |
| Dual-point IC <sub>50</sub> / K <sub>i</sub> value determination | Only one labeled ligand necessary;<br>Adjustable for high throughput screening                                                     | Need for suitable radiolabeled ligand;<br>Only selects long residence time compounds                  | 30,34 |
| Functional reversibility assay                                   | No need for labeled ligands;<br>Adjustable for high throughput screening                                                           | Qualitative measurement;<br>Not suitable for binding kinetics at low temperatures.                    | 35-46 |
| Label-free kinetic measurement                                   | No need for labeled ligands;<br>Quantitative measurement                                                                           | Need for protein purification and immobilization;<br>Low throughput assay format                      | 47-55 |

approaches, the most common and straightforward method is to radiolabel a compound of interest with high affinity and directly measure its on- and off-rate in a kinetic set-up of a radioligand binding experiment. Two examples, in the context of drug-target residence time, are radiolabeled aprepitant at the Neurokinin 1 (NK<sub>1</sub>) receptor and tiotropium at the muscarinic M<sub>3</sub> receptor (M<sub>3</sub>R). Both compounds have been well characterized by using this method.<sup>19,20</sup> Alternatively, a ligand can be fluorescently labeled to calculate its drug-target residence time. For instance, Keppler and coworkers have developed a so-called Tag-lite® platform, a combination of a Homogeneous Time-Resolved Fluorescence (HTRF®) detection method with a covalent labeling technology called SNAP-tag®.<sup>21</sup> One application note from Cisbio describes fluorescently labeled spiperone binding at the Tag-lite dopamine D<sub>2</sub> receptor.<sup>22</sup> Nevertheless, the process of radio- or fluorescent labeling is costly and labor intensive, which, as a consequence, confines kinetics determinations to a limited scope. Therefore, alternative approaches are highly desired (See Figure 2.1-2.4 for experimental schemes and representative results).




**Figure 2.1 | Indirect kinetic radioligand binding assays:** the receptor-bearing material is pre-incubated with a saturating concentration of unlabeled ligand, followed by an intermediate filtration and / or 'washout' step to initiate ligand dissociation. Subsequently, a fixed concentration of radioligand is added and incubated thereafter for a period before assay termination. This scheme is adapted from Vauquelin *et al.*<sup>35</sup> In this assay a slowly dissociating compound (squares) induces an apparent decrease in the amount of free receptor, compared to a fast dissociating compound (triangles), which allows the binding of a subsequently added radioligand; Control: no pre-incubation (spheres).

## A. Indirect kinetic radioligand binding assays

Contrary to the abovementioned, one-step, direct measurement of radiolabeled ligands' dissociation rates, indirect approaches usually contain two consecutive parts (Figure 2.1). Firstly, the receptor-bearing material is pre-incubated with a saturating concentration of unlabeled ligand (to occupy the majority of receptors), followed by an intermediate filtration and/or 'washout' step—methods vary in different labs<sup>23-26</sup>—to initiate ligand dissociation. Secondly, a fixed concentration of radioligand is added and incubated thereafter for a short period before assay termination. Thus, the unlabeled competitor's dissociation rate could be inferred from the recovery of radioligand binding. Furthermore, this method can be adopted and modified for high-throughput screening; an example is at the dopamine  $D_2$  receptor ( $D_2R$ ).<sup>26</sup>


## B. (Dual-point) competition association assay

The competition association assay has been shown to be highly accurate in determining the binding kinetics of unlabeled ligands at the beta-



**Figure 2.2 | (Dual-point) Competition association assay:** an unlabeled competitor is co-incubated with a radioligand in a kinetic association experiment, and followed for two or more time points. **a:** a radioligand association curve without co-incubation of an unlabeled competitor. **b:** a co-incubated unlabeled competitor dissociates slower than the radioligand used. **c:** a co-incubated unlabeled competitor dissociates faster than the radioligand used.  $B_{t_1}$ : specific radioligand binding at the first time point ( $t_1$ );  $B_{t_2}$ : specific radioligand binding at the second time point ( $t_2$ ). Kinetic rate index (KRI) is defined as  $B_{t_1} / B_{t_2}$ . This scheme is adapted from Guo *et al.*<sup>33</sup>

adrenergic receptor,<sup>27,28</sup> and more recently, at the muscarinic  $M_3$  receptor,<sup>29</sup> gonadotropin-releasing hormone (GnRH) receptor,<sup>30</sup> histamine  $H_1$  receptor,<sup>31</sup> and adenosine receptors,<sup>32,33</sup> to name but a few. This method is based on a framework developed by Motulsky and Mahan,<sup>34</sup> where an unlabeled competitor is co-incubated with a radioligand during a kinetic association experiment (Figure 2.2). If the competitor dissociates faster from its target than the radioligand ( $k_2 < k_4$ ), the specific binding of the radioligand will slowly and monotonically approach its equilibrium in time (Figure 2.2, **c**). However, when the competitor dissociates slower ( $k_2 > k_4$ ), the association curve of the radioligand will consist of two phases starting with a typical ‘overshoot’ followed by a decline until a new equilibrium is reached (Figure 2.2, **b**).<sup>34</sup> In contrast, the association rates of either the competitor ( $k_3$ ) or the radioligand ( $k_4$ ) are usually less influential on the pattern of the curves.<sup>34</sup> Recently, the competition association assay has also been adopted and modified by Guo *et al.* for high-throughput kinetic screening studies at the adenosine  $A_1$  receptor.<sup>33</sup> Briefly, in a so-called dual-point competition association assay, two time points are selected to measure radioligand binding: 1) at which the radiolabeled ligand just reaches equilibrium under control conditions in the absence of an unlabeled ligand (Figure 2.2, **a** and  $t_1$ ), and 2) at which the incubation time is long enough for the labeled and unlabeled compound to equilibrate with the target (Figure 2.2,  $t_2$ ). Next, the ratio of the binding at the first time point ( $B_{t_1}$ ) and that at the second time point ( $B_{t_2}$ ) is calculated, which Guo *et al.* defined as the ‘kinetic rate index’ (KRI).



**Figure 2.3 | Dual-point  $IC_{50}$  /  $K_i$  value determination:** an unlabeled competitor is co-incubated with a radioligand for either a short time (in this case, e.g., 30 min) or a long time (in this case, e.g., 10 h) and their corresponding  $IC_{50}$  or  $K_i$  values are calculated. In the representative result, a slowly dissociation compound (squares) will have a more pronounced affinity shift than a fast dissociation compound (triangles).

In this manner, the compounds that quickly dissociate from their target will have a KRI below or equal to one (Figure 2.2, *c*). Conversely, compounds that dissociate slowly from their target compared to the radioligand will have a KRI value larger than one, resulting from the typical ‘overshoot’ in the association curve (Figure 2.2, *b*). Notably, the dual-point assay and KRI value can be applied to screen for candidate compounds with either long or short RT. This suggests the general applicability of this assay for other drug targets as well.


### C. Dual-point $IC_{50}$ / $K_i$ value determination

Another method enabling kinetic screening was reported by Heise *et al.* They used a scintillation proximity assay (SPA) to measure an antagonist’s affinity after 30 min or 10 h of incubation and examined their ratio ( $K_i^{30 \text{ min}} / K_i^{10 \text{ h}}$ ; Figure 2.3).<sup>30</sup> The theory behind this is based on the observation that displacement curves will shift leftward over time until a state of equilibrium is reached; the time to reach that state is determined by the dissociation rate of the slower unbinding process.<sup>34</sup> In other words, a slowly dissociating compound requires longer incubation times to reach equilibrium than a fast dissociating one, which is often neglected. This translates into markedly decreased experimental  $IC_{50}$  or  $K_i$  values over time. In this sense, the ratio of the dual-point  $IC_{50}$  or  $K_i$  values represents a surrogate of overall RT. However,

the principle of this approach also simultaneously implies its limited application, i.e. in screening long RT candidate compounds only, since for short RT compounds the state of equilibrium is quickly established and the  $K_i$  ratio derived thereof is hardly changed.

## D. Functional reversibility assay

A ligand's binding kinetics profile could also be inferred from circumstantial evidence, which is commonly based on functional assays resembling the classical 'organ-bath' experiment. More specifically, it requires pre-incubating tissues / cells with antagonists (to allow them sufficient time to reach binding equilibrium conditions) before their challenge with an agonist (Figure 2.4).<sup>35</sup> As a result, fast dissociating antagonists can only cause a rightward shift of the agonist's concentration-dependent curve, since the receptor can quickly be liberated from the antagonist's blockade and thus readily respond to the subsequently administered agonist. These types of antagonists are categorized as 'surmountable' (Figure 2.4, pre-incubation experiment).<sup>9,36,37</sup> In contrast, slowly dissociating antagonists or, at the extreme, irreversible binders can induce not only a rightward shift but also a concomitant decline in maximal response of the agonist, due to sustained receptor blockade which, on a macroscopic scale, reduces the total receptor population available for an agonist's response.<sup>38,39</sup> Such antagonists are denoted as insurmountable (Figure 2.4, pre-incubation experiment).<sup>9,36,37,40</sup> Notably, such an insurmountable pattern can also be caused by a number of different mechanisms, such as allosteric interactions, receptor internalization or postreceptor functional response blockade.<sup>41-44</sup> None of these result from competitive antagonism. Nevertheless, these mechanisms can be distinguished in a co-incubation experiment, where a depressed maximal response can only be observed by a noncompetitive mechanism, but not for competitive antagonists, no matter how fast or slowly they dissociate from the target (Figure 2.4, co-incubation experiment). Thus, it is always necessary to perform a co-incubation experiment along with the pre-incubation experiment to clarify the mechanism of the insurmountable antagonism.<sup>37,45,46</sup> It is important to mention that the extent of the insurmountable effect, e.g. a decreased maximal response, is time dependent. During a short period of agonist exposure (following pre-incubation with a slowly dissociating antagonist), not all the



**Figure 2.4 | Functional reversibility experiment:** in the upper panel, tissues/cells are pre-incubated with an antagonist before challenge with an agonist. A slowly dissociating antagonist (squares) will cause a rightward shift of the concentration-response curve and a concomitant decline in maximal response of the agonist; a fast dissociating antagonist (triangles) will cause a rightward shift of the concentration-response curve without affecting the maximal response of the agonist. If as in the lower layer an antagonist is co-incubated with an agonist; both slowly and fast dissociating competitive antagonists will cause a rightward shift of the concentration-response curve without affecting the maximal response of the agonist. Otherwise they are non-competitive to the agonist; Control: agonist only (spheres). This scheme is adapted from Vauquelin *et al.*<sup>35</sup>

receptor sites are liberated before the response is measured, a state of so-called ‘hemi-equilibrium’.<sup>38,39,44</sup> However, if the agonist exposure lasts longer, the occupancy of the receptors will readjust with time until, ultimately, both the agonist-receptor and the antagonist-receptor interactions reach equilibrium.<sup>44</sup> Evidently, the insurmountability of such slowly dissociating compounds will change if the response is measured at different time points.

## E. Label-free kinetic measurements

One of the most well-established label-free measurements for kinetics is surface plasmon resonance measurement (SPR). In SPR, the protein is immobilized

on a coated gold chip, which is then exposed to the compound of interest under continuous flow. Ligand binding to the receptor immobilized on the chip's surface induces a real time change in the refractive index on the sensor surface. As such a change is linear to the number of molecules bound, it allows quantitative characterization of binding kinetics (for review, see<sup>16,47,48</sup>).

Another emerging label-free technology enabling kinetics measurement is by surface acoustic wave biosensor (SAW).<sup>49</sup> This methodology captures real-time mass changes on the surface, which result in a shifted phase and / or changed amplitude of the sound wave signal.<sup>50</sup> Besides these two biophysical assays, other equally valid technologies have also been reported elsewhere.<sup>51</sup>

However, these biophysical techniques have not been thoroughly tested on GPCRs. The difficulty is that these receptors are integral membrane proteins, and rapidly deteriorate when taken out of their natural environment, which is often a prerequisite for these newer biophysical approaches. Several strategies were developed to overcome this inherent difficulty. One of the successful examples is the application of a protein engineering method to generate a stabilized receptor (StaR), e.g. the adenosine A<sub>2A</sub> receptor, minimally modified for thermostability.<sup>52-54</sup> The stabilized receptors can be purified in large quantities, whilst retaining correct folding, thus generating materials suitable for a broad range of structural and biophysical studies.<sup>53</sup> Another technique is to solubilize a given GPCR from whole cell pellets and capture it on antibody surfaces for analysis, exemplified by an SPR study on the CCR5 and CXCR4 chemokine receptors.<sup>55</sup> These solubilized receptor preparations could be useful materials for binding kinetics studies.

## 2.4 Revisit kinetics data on GPCRs

To date, nearly 800 genes encoding GPCRs have been identified, representing a substantial share of the human genome.<sup>56</sup> In general, all these receptors possess structurally conserved heptahelical transmembrane motifs (7TM) connected by alternating intracellular and extracellular loops, with the amino terminus located at the extracellular side and the carboxyl terminus at the intracellular side.<sup>57</sup> Despite these common features in their molecular structure, GPCRs are characterized by a relatively low overall sequence identity (less than 20%). According to phylogenetic classification, these receptors can be divided into five main families—*Rhodopsin* (Class A), *Secretin* (Class B), *Glutamate* (Class

**Table 2.2 | Overview of GPCR ligand kinetics.**

| <b>Cmpd</b>   | <b>Target<sup>a</sup></b>               | <b>t<sub>1/2</sub><sup>b</sup></b> | <b>Clinical indications</b>              | <b>Status<sup>c</sup></b> | <b>Ref.</b> |
|---------------|-----------------------------------------|------------------------------------|------------------------------------------|---------------------------|-------------|
| Ritanserlin   | 5-HT <sub>2</sub> receptor              | 160 min (37 °C)                    | Cocaine-related disorders                | Phase II                  | 160         |
| Risperidone   | 5-HT <sub>2</sub> receptor              | Irreversible (37 °C)               | Schizophrenia; Alcohol abuse             | Marketed                  | 161         |
| Clonidine     | α <sub>2A</sub> -autoreceptor           | 27.6 s (37 °C)                     | ADHD                                     | Marketed                  | 99          |
| Bopindolol    | β <sub>1</sub> -adrenoceptor            | > 4 h (rat)                        | Hypertension                             | Marketed                  | 162         |
| Olodaterol    | β <sub>2</sub> -adrenoceptor            | 17.8 h                             | COPD                                     | Marketed                  | 163         |
| Lofentanil    | μ receptor                              | 260 min (37 °C)                    | Analgesia                                | Marketed                  | 164         |
| Candesartan   | AT <sub>1</sub> receptor                | 120 min (37 °C)                    | Hypertension                             | Marketed                  | 43          |
| UK432,097     | A <sub>2A</sub> receptor                | 173 min (5 °C)                     | COPD                                     | Terminated <sup>d</sup>   | 32          |
| Taranabant    | CB <sub>1</sub> receptor                | 122 min (37 °C)                    | Obesity                                  | Terminated <sup>e</sup>   | 25          |
| Maraviroc     | CCR5                                    | > 136 h (4 °C)                     | HIV                                      | Marketed                  | 165         |
| SCH527123     | CXCR2                                   | 22 h                               | Asthma; Psoriasis                        | Phase II                  | 166         |
| PF-4850890    | CRF <sub>1</sub> receptor               | 6.79 h (rat)                       | Depression; Anxiety                      | Preclinical               | 153         |
| Clozapine     | D <sub>2</sub> receptor                 | 14.5 s (37 °C)                     | Schizophrenia                            | Marketed                  | 135         |
| AM432         | DP <sub>2</sub> receptor                | 90 min (37 °C)                     | COPD                                     | Preclinical               | 167         |
| Uracil 3      | GnRH receptor                           | > 43 h                             | Reproductive-endocrine axis disorders    | Preclinical               | 168         |
| Desloratidine | H <sub>1</sub> receptor                 | > 6 h                              | Allergy                                  | Marketed                  | 169         |
| JNJ7777120    | H <sub>4</sub> receptor                 | 62 min                             | Inflammation                             | Preclinical               | 170         |
| Tiotropium    | M <sub>3</sub> receptor                 | 27 h                               | COPD                                     | Marketed                  | 80          |
| Aprepitant    | NK <sub>1</sub> receptor                | 154 min                            | Chemotherapy-induced emesis              | Marketed                  | 19          |
| Ibodutant     | NK <sub>2</sub> receptor                | 28 min                             | Irritable Bowel Syndrome                 | Phase II                  | 171         |
| Osanetant     | NK <sub>3</sub> receptor                | 10 min                             | Schizophrenia                            | Phase II                  | 172         |
| Almorexant    | OX <sub>2</sub> receptor                | 139 min                            | Chronic primary insomnia                 | Phase III                 | 173         |
| Clopidogrel   | P <sub>2</sub> Y <sub>12</sub> receptor | Irreversible (37 °C)               | Stroke; Heart attack; Severe chest pain  | Marketed                  | 62          |
| Apafant       | PAF receptor                            | 50 min (guinea pig)                | Arrhythmias; Breast cancer; Inflammation | Preclinical               | 174         |
| GSK1562590    | UT receptor                             | > 1 h (rat)                        | Hypertension                             | Preclinical               | 175         |
| SCH500946     | Y <sub>5</sub> receptor                 | 71 min                             | Obesity                                  | Preclinical               | 176         |

<sup>a</sup> Official IUPHAR receptor name. <sup>b</sup> Dissociation half-life at room temperature and on human receptors unless mentioned otherwise. <sup>c</sup> Clinical status obtained from [ClinicalTrials.gov](https://clinicaltrials.gov).<sup>179</sup>

<sup>d</sup> Discontinued due to lack of efficacy. <sup>e</sup> Discontinued due to high level of central nervous system side effects.<sup>177,178</sup>



C), *Adhesion* and *Frizzled/Taste2*.<sup>56,58</sup> Receptors from Class A to C have been most addressed in drug discovery efforts. The natural ligands of GPCRs are highly variable ranging from small ions and photons to large glycoproteins, which indicates the important and versatile roles these proteins play in a vast number of biological processes. These are also the reasons why GPCRs have developed into a very ‘druggable’ class of targets for more than 30% of currently available drugs.<sup>59</sup> However, in the introduction we also mentioned the tremendous attrition along R&D pipelines, and suggested that detailed mechanism-based studies of drug-target interaction should be introduced in the early phases of drug discovery to prevent ‘fail late, fail expensive’ scenarios. For this we examined the gigantic ‘pool’ of GPCR ligands and retrospectively analyzed the ‘kinotype’ (kinetics features) of some cases. Without claiming to be exhaustive we have summarized a series of prototypic GPCR ligands that have known kinetics information (Table 2.2). We listed one representative (candidate) drug per target (26 targets in total) including their clinical indications. Several conclusions can be drawn from this table. Firstly, the existing kinetics data for GPCR ligands is nowhere near the number of affinity-based evaluations in the early phases of drug discovery. The sum of targets presented here is far behind the total population of ‘druggable’ GPCRs, estimated to be 300.<sup>60,61</sup> Clearly, ligand-receptor binding kinetics has been overlooked and less investigated in the past. Secondly, the range of drug-target residence times, reflected by  $t_{1/2}$  in the table, varies from seconds to hours. Most of these are in line with the clinical applications of the related drug. For instance, patients with allergy can benefit from the long-lasting effect by desloratidine ( $t_{1/2} > 6$  h) in relieving nasal congestion, or patients with COPD from taking tiotropium ( $t_{1/2} = 27$  h) for a durable bronchodilatory effect. In both cases, their clinical indications favor sustained target occupancy. On the contrary, for some marketed drugs, serious adverse effects could go along with the desired effects due to unwanted kinetic rates. This is exemplified by clopidogrel, which is an antiplatelet agent used to inhibit the formation of blood clots. However, its irreversible inhibitory effect may prolong or exacerbate bleeding.<sup>62</sup> Another case is represented by lofentanil for analgesia. Although its longer duration of action has an advantage for certain types of analgesia,<sup>63</sup> it is still unsuitable for most practical applications.<sup>64</sup> Instead, short-acting derivatives such as sufentanil or remifentanil are preferred for medical use in human surgical procedures.<sup>65,66</sup> One can speculate that such clinical limitations could be avoided or at least foreseen if kinetics were to be incorporated in the early phase of candidate selection.

## 2.5 Structure-kinetics and structure-affinity relationships

Thus far the abovementioned analyses in retrospect are still largely based on ‘hindsight’ information; hardly any structure-kinetics relationship studies (SKR), in prospect, have been reported previously. Therefore, we should like to propose a strategy for compound optimization that combines an extensive structure-kinetics relationship study (SKR) with a classical structure-affinity relationship study (SAR). It is envisioned that this added metric, i.e., binding kinetics or the ‘kinotype’ of a given compound, offers several advantages over the traditional and simple SAR paradigm:

1) A better understanding of the molecular mechanism of action. This includes detailed characterization of not only the bound states under equilibrium conditions (SAR) but also the entire drug-target interaction that comprises metastable intermediate states and transition states (SKR).<sup>14,67</sup> Such gained knowledge offers rationales for efficient drug design.

2) A more steered drug design and development. Different targets may have distinct kinetics preferences. Aiming for long or short RT compounds is a crucial consideration in early phase discovery to triage and advance the best leads towards clinical applications.<sup>16</sup>

3) An efficient translation from *in vitro* to *in vivo* settings. This may decrease the prevalence of the ‘fail late, fail expensive’ scenario, especially considering that lack of efficacy is one of the major reasons for drug attrition.<sup>68,69</sup>

In addition, a detailed SKR study also describes the association rate of the ligand-receptor interaction, which has been less investigated and is often thought to be diffusion-limited.<sup>14</sup> As a consequence, this could lead to the assumption that optimizing for a long residence time is tantamount to optimizing the binding affinity, since  $K_D = k_{\text{off}} / k_{\text{on}}$  and  $k_{\text{on}}$  reaches an (identical) limit. However, Vilums *et al.* have recently reported a thorough SKR study on the CCR2 receptor, in which it was shown that this general assumption is not necessarily valid.<sup>18</sup> They documented that the compounds’ on-rates varied significantly from  $0.01 \text{ nM}^{-1} \cdot \text{min}^{-1}$  to  $0.0003 \text{ nM}^{-1} \cdot \text{min}^{-1}$ —more than three orders-of-magnitude from the diffusion-limited value (approximately  $10^{10} \text{ M}^{-1} \cdot \text{min}^{-1}$ ).<sup>1</sup> It is also notable that the affinity difference of two diastereomers in that paper with  $K_i$  values of 3.6 nM and 289 nM, was predominantly caused by their 18-fold difference in the on-rate, rather than their very similar RTs (135 min and 77 min). Such valuable information can

only be obtained by a detailed SKR study, which has not been reported for many GPCRs so far.

Evidently, a detailed SKR offers added value to the traditional metric of affinity. This new strategy of combining SAR and SKR can be expected to further enhance the success of drug discovery and development. To further illustrate the importance of incorporating SKR into the pipeline of early phases of drug research, in the following part of this chapter we will primarily address small molecules targeting class A and B GPCRs, which have been reported to have optimized binding kinetics, more specifically dissociation kinetics. Specific evidence will be given for three 'long RT' targets and one 'short RT' target; muscarinic  $M_3$ , tachykinin  $NK_1$  and  $CRF_1$  receptor, and dopamine  $D_2$  receptors, respectively, since these targets have been (most) extensively studied in this area.

## 2.6 Drug-target residence time for class A GPCRs

### Muscarinic receptors

Muscarinic receptors, or muscarinic acetylcholine receptors (mAChRs), are a group of class A GPCRs, which are endogenously activated by acetylcholine.<sup>70</sup> To date, five subtypes of muscarinic receptors are identified, namely  $M_1$ - $M_5$ .<sup>71,72</sup> These receptors can be further categorized into two broad groups based on their primary coupling efficiency to G proteins;  $M_2$  and  $M_4$  muscarinic receptors couple to the pertussis toxin-sensitive  $G_{i/o}$  proteins, while  $M_1$ ,  $M_3$  and  $M_5$  muscarinic receptors to  $G_{q/11}$  proteins.<sup>70,73</sup> Of all five muscarinic subtypes,  $M_3$  receptor is located in the smooth muscles of blood vessels throughout the body as well as in the lung. Hyperactivation of the lung  $M_3$  receptor can cause over constriction of smooth muscle, such as that observed during bronchoconstriction. Thus, antagonizing  $M_3$  receptor can lead to relaxation of airway smooth muscles and result in symptom relief of chronic obstructive pulmonary disease (COPD).<sup>74,75</sup> Additionally, for the treatment of chronic illness, long duration of action (preferably 24 h) is an important feature of drugs that enables both prolonged efficacy and patient compliance.<sup>76</sup> Therefore, the treatment of COPD can benefit from a long-




Figure 2.5 | Structures of small molecule M<sub>3</sub> muscarinic receptor ligands.

acting muscarinic antagonist (LAMA), as evidenced by tiotropium and other  $M_3$  antagonists summarized below.

## 1. Tiotropium

Tiotropium is a long-acting anticholinergic bronchodilator, which is currently marketed under the trade name Spiriva<sup>TM</sup> (Figure 2.5, Compound 1).<sup>77-79</sup> Given its long *in vivo* duration of action, several groups, in retrospect, performed *in vitro* time-dependent  $IC_{50}$  value analysis to obtain the accurate affinity and binding kinetics of tiotropium.<sup>29,80</sup> It was found that equilibrium is reached only after 18-20 h in a radioligand binding assay. This phenomenon is consistent with the theory by Motulsky and Mahan that a long incubation time is needed for a radioligand to reach equilibrium in the presence of a slowly dissociating competitor.<sup>34</sup> Thus, it also explains that different affinities were published by different groups; after a period of 20 h incubation, the  $K_i$  value of tiotropium was 8 pM,<sup>29,80</sup> while after only 4 h incubation, its  $K_D$  value was 0.33 nM.<sup>78</sup> The  $t_{1/2}$  of [<sup>3</sup>H]-tiotropium was 34.6 h at room temperature, which was much longer than ipratropium (Figure 2.5, Compound 2;  $t_{1/2}$  = 0.26 h, Table 2.3), a relatively short-acting agent requiring up to four doses per day.<sup>74,78</sup> Similarly, a kinetics study performed by Dowling and Charlton using a competition association assay has also demonstrated that tiotropium has a very long  $t_{1/2}$  (7.7 h) at room temperature,<sup>29</sup> which, however, is shorter than the value (34.6 h) reported by Disse *et al.*<sup>78</sup> This may be due to the difference in their buffer for membrane preparation and / or the binding buffer for kinetic measurements, for instance, the presence of sodium ions. Likewise, tiotropium has an even shorter dissociation half-life (46.2 min) under mock physiological conditions at 37 °C, where the ionic strength (including  $Na^+$ ,  $Ca^{2+}$ ,  $K^+$ ,  $Mg^{2+}$  and  $Cl^-$  ions) of the assay buffer was increased.<sup>81</sup> This example furthermore highlights the importance of careful choices in assay development for kinetic measurements. Another important pharmacological feature of tiotropium is its 'kinetic receptor subtype selectivity', namely a longer dissociation half-life at the  $M_3$  receptor, 34.7 h, than at the  $M_1$  and  $M_2$  receptors (14.6 h and 3.6 h, respectively).<sup>78</sup> In contrast, tiotropium's affinities at these receptors were similar and in the subnanomolar concentration

**Table 2.3 | Binding kinetics of M<sub>3</sub> muscarinic receptor ligands.**


| Nr  | Cmpd <sup>a</sup> | Affinity (K <sub>i</sub> or pK <sub>i</sub> ) <sup>b</sup> | Dissociation half-life (t <sub>1/2</sub> ) <sup>c</sup> | Ref   |
|-----|-------------------|------------------------------------------------------------|---------------------------------------------------------|-------|
| 1.  | Tiotropium        | 8 pM                                                       | 34.6 h                                                  | 29,78 |
| 2.  | Ipratropium       | 0.69 nM                                                    | 0.26 h                                                  | 78    |
| 3.  | Acidinium         | 18 pM                                                      | 10.7 h                                                  | 80    |
| 4.  | PF-335659         | 132 pM                                                     | > 24 h                                                  | 96    |
| 5.  | Glycopyrronium    | 10.04                                                      | 6.3 h                                                   | 80    |
| 6.  | Acetylcholine     | 4.87                                                       | 7.8 s                                                   | 98    |
| 7.  | Carbachol         | 4.09                                                       | 6.3 s                                                   | 98    |
| 8.  | Methacholine      | 4.52                                                       | 6.0 s                                                   | 98    |
| 9.  | Oxotremorine-M    | 4.61                                                       | 7.2 s                                                   | 98    |
| 10. | Oxotremorine      | 5.61                                                       | 2.6 s                                                   | 98    |
| 11. | Bethanechol       | 3.71                                                       | 3.7 s                                                   | 98    |
| 12. | Pilocarpine       | 4.78                                                       | 3.0 s                                                   | 98    |

<sup>a</sup> Compound names and synonyms used in original references. <sup>b</sup> Affinity, expressed as K<sub>i</sub> values or pK<sub>i</sub> as reported in original references; result on human receptor unless mentioned otherwise. <sup>c</sup> Dissociation half-life at room temperature.

range.<sup>78</sup> Taken together, these features make tiotropium the most widely used LAMA worldwide so far.<sup>81,82</sup>

To understand the molecular basis of the exceedingly long RT of tiotropium, Tautermann *et al.* performed a ‘kinetic mapping’ of its binding sites and the ‘exit’ channel from the M<sub>3</sub> receptor by testing the kinetics of tiotropium at mutant M<sub>3</sub> receptors.<sup>83</sup> They found that the slowly dissociating profile of tiotropium was attributed to interactions in the binding site particularly with N508<sup>6,52</sup> (N507<sup>6,52</sup> on the rat M<sub>3</sub> receptor, see Figure 2.6). Importantly, this residue appeared to ‘anchor’ tiotropium at the receptor via directly interacting with its hydroxyl-group and ester-group so as to restrict the movement of the ligand towards the exit channel. Consequently, tiotropium’s RT was 45-fold longer on the wild type M<sub>3</sub> receptor than that of its analogue lacking the hydroxyl-group, while little difference was observed at the N508<sup>6,52</sup>A mutant.

Furthermore, a recent structural biology study of the M<sub>3</sub> receptor (from rat) co-crystallized with tiotropium (PDB code: 4DAJ) provided a clear image for the molecular mechanism behind its long RT profile.<sup>84</sup> As presented in



**Figure 2.6 | The co-crystal structure of tiotropium in the rat  $M_3$  receptor.**<sup>84</sup> This figure was generated with ICM Browser v3.7 (Molsoft) from PDB code: 4DAJ. Tiotropium (black) binds within the pocket. Three tyrosine residues, Y148<sup>3,33</sup>, Y506<sup>6,51</sup> and Y529<sup>7,39</sup>, together prevent the ligand from moving out of the receptor. N507<sup>6,52</sup> interacts with the carbonyl and hydroxyl groups through H-bonds, while the ligand's typical quaternary ammonium group interacts with D147<sup>3,32</sup>.

Figure 2.6, tiotropium binds deeply in the receptor core and is covered by an aromatic 'lid' comprised of three conserved tyrosines—Y148<sup>3,33</sup>, Y506<sup>6,51</sup> and Y529<sup>7,39</sup>. This aromatic 'lid' nearly occludes the ligand from the solvent thus preventing it from being 'wetted' by water molecules and then 'washed' off the binding pocket. In addition, the abovementioned N507<sup>6,52</sup> interacts with the carbonyl and hydroxyl groups of tiotropium through H-bonds, while the ligand's typical quaternary amine interacts with D147<sup>3,32</sup>.

The elucidation of the co-crystal structure of tiotropium in the rat  $M_3$  receptor also facilitated molecular dynamic simulations to characterize the pathway by which tiotropium binds to and dissociates from the rat  $M_3$  receptor.<sup>84</sup> In contrast to a simple one-step dissociation process, the ligand appeared to firstly leave the orthosteric ligand binding pocket and then 'pause' at an extracellular 'vestibule', forming a loosely connected intermediate state, before tiotropium finally dissociated from the receptor. During this process,

the first step formed a large energy barrier that involved the aromatic 'lid' (Y148<sup>3,33</sup>, Y506<sup>6,51</sup> and Y529<sup>7,39</sup>) around the charged ligand head group opening up, while the second step posed a smaller energetic barrier. In this sense, the RT of tiotropium is greatly determined by the first step, which is rate-limiting for the ligand dissociation process.

As expected from the kinetic binding profile of tiotropium, an insurmountable antagonist behavior was observed in a functional reversibility experiment. This was detailed by Casarosa *et al.* during the preclinical evaluation of several long- and short-acting muscarinic antagonists, where tiotropium was pre-incubated for 15 min before the addition of the agonist muscarine in the inositol phosphate accumulation assay.<sup>80</sup> Fitting the dose ratios over a large range of tiotropium concentrations ( $10^{-10}$ - $10^{-5}$  M) generated a Schild plot with a slope of 1.29, higher than unity. Such a value indicated the insurmountable M<sub>3</sub>R antagonism of tiotropium, which was a result of a hemi-equilibrium due to its long M<sub>3</sub>R residence time profile.<sup>37</sup>

Taken together, the *in vitro* results mentioned above combined with further evidence unveil the molecular basis of the long RT profile of tiotropium. These results correlate with this compound's long *in vivo* duration of bronchoprotection (> 24 h) as determined in a model of acetylcholine-induced bronchoconstriction in anesthetized dogs.<sup>80</sup> Moreover, despite the possible influence of drug metabolism and pharmacokinetic factors,<sup>85</sup> the clinical profile of tiotropium in COPD highlights an inextricable link between long receptor RT and long duration of action that enables a simple, once-daily dosage regime.<sup>77,86,87</sup>

## 2. Aclidinium

Aclidinium (Figure 2.5, Compound 3) is a long-acting, inhaled muscarinic antagonist recently approved by the FDA as a maintenance treatment for COPD.<sup>88</sup> The synthesis and biological evaluation of aclidinium and its analogues were elaborately reported by Prat *et al.* and Gavalda *et al.*<sup>20,89</sup> This antagonist was specifically designed for a maintained bronchodilatory efficacy with fewer unwanted systematic anticholinergic effects, such as dry mouth (reported in 11.6% of patients treated with tiotropium bromide versus 3.5%



of patients treated with placebo), glaucoma, constipation, increased heart rate, and urinary retention.<sup>90</sup> According to Prat *et al.*, acclidinium had a rapid hydrolysis in human plasma, hence minimizing the class-related systematic side effect. However, such a fast pharmacokinetics profile of acclidinium did not influence its long duration of action (over 24 h), as evaluated in an animal model.<sup>20</sup> This phenomenon was attributed to acclidinium's slowly dissociating profile from the  $M_3$  receptor.

Compared to tiotropium, acclidinium had a similar affinity for the  $M_3$  receptor ( $K_i = 18$  pM), while its  $t_{1/2}$  (10.7 h) was approximately two-fold shorter (Table 2.3).<sup>80</sup> In another study, Gavalda *et al.* used radiolabeled acclidinium and found a  $K_D$  value of 0.32 nM; combined with  $t_{1/2}$  of 29.24 h, which was shorter than [ $^3H$ ]-tiotropium (62.19 h).<sup>20</sup>

*In vitro* functional activity and *in vivo* duration of bronchoprotection of acclidinium was also examined by Gavalda *et al.*<sup>20</sup> More specifically, they used the isolated guinea pig trachea and treated it with carbachol to generate dose-dependent contraction. A 60 min pretreatment with acclidinium or tiotropium shifted the concentration-response curves of carbachol to the right. Along with the potency shift, tiotropium also suppressed the maximal agonist effect while acclidinium did not. Such a result is indicative of the relatively shorter RT of acclidinium than that of tiotropium at the  $M_3$  receptor.

In addition to the characterization of acclidinium's functional activity, the onset of action of this compound was also studied in the carbachol-induced contraction assay. Notably, acclidinium displayed a faster onset of action ( $t_{1/2} = 6.8$  min,  $t_{max} = 35.9$  min) than tiotropium ( $t_{1/2} = 13.6$  min,  $t_{max} = 61.2$  min). For the *in vivo* duration of bronchoprotection, the effect half-life of acclidinium (defined as the time taken to reduce the maximal bronchoconstriction by 50%) was 29 h. This duration of action was considerably longer than that of ipratropium (8 h) yet shorter than tiotropium (64 h). Likewise, Casarosa *et al.* reported that acclidinium and tiotropium displayed different levels of bronchoprotection at 24 h after equi-effective dose administration (tiotropium = 35%, acclidinium = 21%).<sup>80</sup> Clearly, these *in vivo* results are in line with the *in vitro* binding kinetics.

Furthermore, compared to tiotropium's once-daily dose regime, acclidinium required a twice-daily dose regime for a 24 h bronchodilation effect in clinical trials.<sup>91-95</sup> Such a difference appears to be derived from their divergent RTs at the  $M_3$  receptor. In addition, acclidinium's faster onset of action, mostly due to its fast association rate, can provide quicker symptom relief as compared to tiotropium.

### 3. PF-3635659

The synthesis and biological evaluation of PF-3635659 (Figure 2.5, Compound 4) and its analogues was reported by Glossop *et al.*<sup>96</sup> This study nicely exemplifies a systematic combination of both a structure-affinity and structure-kinetics relationship study for early-phase drug design and development. More specifically in this study, Glossop *et al.* demonstrated that a *gem*-dimethyl substitution adjacent to the basic amine moiety of PF-3635659 exerted a profound effect on the dissociation rate. As evidence, the  $t_{1/2}$  of the initial *gem*-dimethyl-free compound (38 min) was extended to over 1440 min, indicating the importance of the *gem*-dimethyl substitution for attaining ligand receptor RT. Further chemical modifications on the heterocyclic ring and the right side of the phenyl ring induced corresponding changes in ligand affinity and kinetics, which eventually led to the discovery of the candidate compound, i.e., PF-3635659 (Compound 47 in the original paper). Following-up binding kinetics characterization of this compound in a competition association assay revealed that it had an on-rate of  $1.05 \times 10^7 \text{ M}^{-1} \cdot \text{min}^{-1}$ , slower than tiotropium and ipratropium ( $k_{\text{on}} = 2.12 \times 10^8 \text{ M}^{-1} \cdot \text{min}^{-1}$  and  $1.00 \times 10^9 \text{ M}^{-1} \cdot \text{min}^{-1}$ ) and an off-rate of  $1.39 \times 10^{-4} \text{ min}^{-1}$ , translating into a  $t_{1/2}$  of more than 24 h (Table 2.3).

Importantly, the *in vivo* onset and duration of action data for PF-3635659 were in line with its *in vitro* binding kinetics. The profiling of PF-3635659 in a conscious dog model of bronchoconstriction illustrated a slower onset of action compared to ipratropium and tiotropium, with yet a longer duration of action (> 24 h) than ipratropium.<sup>96</sup> This compound has now progressed into a phase II clinical trial. Data from healthy volunteers demonstrated that PF-3635659 provided efficacious bronchodilation over 24 h from a single inhaled dose, thus confirming the suitability of PF-3635659 as a novel once-daily inhaled muscarinic  $M_3$  receptor antagonist for the treatment of COPD.

### 4. Glycopyrronium

Glycopyrronium (Figure 2.5, compound 5) is the cation and active moiety of glycopyrronium bromide, of which the dry-powder formulation is also known as NVA237.<sup>81</sup> *In vitro* binding kinetics profiling of this compound

in a radioligand competition association assay revealed a dissociation rate of  $0.11 \pm 0.02 \text{ h}^{-1}$  from the  $M_3$  receptor, equal to a dissociation  $t_{1/2}$  of 6.3 h (Table 2.3). This value was much longer than the short-acting  $M_3$  antagonist ipratropium (0.22 h) but shorter than tiotropium (27 h).<sup>80</sup> However, glycopyrronium had a more rapid rate of onset of action (3- to 4.8-fold) than tiotropium, as determined in an *in vitro* calcium assay and rat tracheal strip assay.<sup>81</sup> This profile was similar to that obtained with ipratropium bromide in preparations of guinea-pig and human airways.<sup>97</sup>

Notably, glycopyrronium also displayed higher equilibrium binding and kinetic selectivity for  $M_3$  versus  $M_2$  receptors as compared to both tiotropium and ipratropium.<sup>80,81,97</sup> Its kinetic selectivity resulted from a 16.5-fold longer receptor RT at the  $M_3$  receptor than at the  $M_2$  receptor, which was higher than tiotropium (10.4-fold) and ipratropium (7.3-fold). Such a property may provide a better therapeutic window.

## 5. $M_3$ receptor agonists

The RTs of seven  $M_3$  agonists (Figure 2.5, Compound 6-12) were recently reported by Sykes *et al.*<sup>98</sup> These agonists are acetylcholine, carbachol, methacholine, oxotremorine-M, oxotremorine, bethanechol and pilocarpine, and are not very selective. The data, however, were only collected on the  $M_3$  receptor subtype. Among the seven agonists, oxotremorine displayed the shortest  $t_{1/2}$  at the  $M_3$  receptor (2.6 s), while acetylcholine had the longest receptor RT ( $t_{1/2} = 7.8 \text{ s}$ ). In general, these agonists' off-rates were much faster than that of the abovementioned  $M_3$  antagonists (Table 2.3). Further *in vitro* functional characterization of the  $M_3$  agonists demonstrated that they had different relative agonist efficacies. Notably, Sykes *et al.* found that there was no relationship between agonist efficacy and the equilibrium binding affinity of each agonist. Interestingly, when efficacy was compared with the dissociation rate constant, the two were highly correlated. This result suggests a relationship between the duration of agonist binding at the receptor and the intrinsic efficacy, a finding also reported by Guo *et al.* and Hoeren *et al.* at the adenosine  $A_{2A}$  receptor and at the  $\alpha_{2A}$ -adrenoceptor.<sup>32,99</sup> Importantly, such a finding again emphasizes that equilibrium models and assays alone are not sufficient to describe a dynamic signaling system; instead, a combination of kinetic models alongside the traditional affinity determination might be

preferred to better capture the nature of efficacy and to develop efficacious drugs for GPCRs.<sup>98</sup>

In summary, several case studies of long acting  $M_3$  receptor antagonists suggest their sustained pharmacological effect lies in these compounds' long receptor occupancy time. The design and development of long residence time antagonists may provide further candidate drugs with once-daily application for COPD. We also included the kinetics data of several  $M_3$  agonists, since they provide important information that agonist efficacy may be correlated to its drug-target residence time rather than the commonly tested equilibrium affinity. This further indicates the importance of combining both binding and kinetics information in the early phase of drug design and development.

## Tachykinin receptors

Tachykinin receptors comprise three different subtypes, namely the  $NK_1R$ ,  $NK_2R$  and  $NK_3R$ , which can be activated by tachykinin peptides, including Substance P (SP), neurokinin A (NKA) and neurokinin B (NKB).<sup>100-102</sup> The three members of the neurokinin receptor class are differentiated by their varying affinities for these three tachykinins, which are  $SP > NKA > NKB$  for the  $NK_1R$ ,  $NKA > NKB > SP$  for  $NK_2R$  and  $NKB > NKA > SP$  for  $NK_3R$  (for reviews, see<sup>100-102</sup>). All three subtypes are functionally coupled to  $G_{q/11}$ , which leads to the activation of phospholipase C- $IP_3$ /DAG signaling and results in elevation of intracellular  $Ca^{2+}$  levels.<sup>102</sup> The localization of the tachykinin receptors varies between subtypes in the human body: the  $NK_1R$  and  $NK_3R$  are prominent both in the central nervous system (CNS) and in peripheral tissues, while the  $NK_2R$  is widely expressed in the smooth muscle of the gastrointestinal, respiratory, and urinary tracts and to a much lesser extent in the CNS.<sup>102-104</sup> A large body of preclinical evidence has suggested that pharmacological or gene manipulation of the  $NK_1R$  has potential clinical utility in the treatment of chronic pain, migraine, neurogenic inflammation and emesis.<sup>105-107</sup> These diseases could benefit from therapeutics with a long duration of  $NK_1R$  antagonism.




Figure 2.7 | Structures of small molecule NK<sub>1</sub> receptor antagonists.

## 1. Aprepitant

Aprepitant, also known as L-754,030, MK0869 or Emend (Figure 2.7, Compound 1),<sup>19,108-110</sup> is the only tachykinin receptor antagonist on the market to date. It is a selective, slowly dissociating NK<sub>1</sub> antagonist with a  $t_{1/2}$  of 154 min at 22 °C (Table 2.4).<sup>19</sup> In comparison, it was found by Cascieri *et al.* that the dissociation rate of the radiolabeled NK<sub>1</sub>'s natural ligand SP (i.e., <sup>125</sup>I-[Tyr<sup>8</sup>]-SP) was too rapid to be measured accurately at a comparable temperature.<sup>111</sup> At 15 °C, this peptide dissociates with  $t_{1/2} = 46$  min and 0.2 min for the G protein-coupled and -uncoupled receptor states, respectively.<sup>112</sup> Evidently, aprepitant had a much longer residence time on the NK<sub>1</sub> receptor than the natural ligand which has been speculated to be the reason for its high potency and *in vivo* efficacy.

As expected from a slowly dissociating ligand, aprepitant's activity was found to be insurmountable in functional assays compared to other, fast dissociating NK<sub>1</sub>R antagonists. This functional effect was associated with its long-lasting *in vivo* efficacy, a finding shared by many authors.<sup>19,108,110</sup> One of the most elaborate studies reported on aprepitant was conducted by

**Table 2.4 | Binding kinetics of NK<sub>1</sub> receptor antagonists.**

| Nr | Cmpd <sup>a</sup> | Affinity (K <sub>i</sub> , nM) <sup>b</sup> | Dissociation half-life (t <sub>1/2</sub> ) <sup>c</sup> | Ref. |
|----|-------------------|---------------------------------------------|---------------------------------------------------------|------|
| 1  | Aprepitant        | 0.019                                       | 154 min (22 °C)                                         | 19   |
| 2  | L-742,694         | 0.037                                       | 27 min (22 °C)                                          | 111  |
| 5  | Vestipitant       | 0.027                                       | 59.7 min (37 °C)                                        | 119  |

<sup>a</sup> Compound names and synonyms used in original references. <sup>b</sup> Affinity, expressed as K<sub>i</sub> values that were reported in original references; results on human receptor unless mentioned otherwise. <sup>c</sup> Dissociation half-life at different temperatures indicated in brackets.

Lindström *et al.*<sup>108</sup> Several functional experiments (i.e., pre-incubation, co-incubation and washout experiments) were used to reaffirm that aprepitant was a competitive yet pseudo-irreversible antagonist with slow functional reversibility. Next to the measurement of its functional interactions *in vitro*, the authors also examined the pharmacokinetic/pharmacodynamics relationship of aprepitant *in vivo* by using the gerbil foot tap (GFT) assay, a model reflecting central NK<sub>1</sub>R activation.<sup>113</sup> It was found that the duration of GFT inhibition by aprepitant outlasted the time course of its plasma or brain concentrations, indicating that its long duration of action was indeed more likely due to a sustained receptor blockade rather than a slow pharmacokinetic half-life. In contrast, two other structurally unrelated NK<sub>1</sub> antagonists, CP-99994 and ZD6021, demonstrated rapid functional reversibility that was in line with their pharmacokinetic profile.<sup>108</sup> Furthermore, these authors also examined *ex vivo* receptor occupancy by autoradiography, reaffirming that time-dependent inhibition of GFT by aprepitant correlated well with NK<sub>1</sub> occupancy in the gerbil brain, a finding that was shared by Duffy *et al.* in 2002.<sup>109</sup> Taken together, all *in vitro* and *in vivo* results provide strong evidence that aprepitant is a slowly dissociating NK<sub>1</sub>R antagonist.

## 2. L-742,694

Aprepitant was originally developed from L-742,694 (Figure 2.7, Compound 2),<sup>109,111,112</sup> which was shown to be a pseudo-irreversible yet competitive antagonist by Cascieri *et al.* in 1997.<sup>111</sup> The t<sub>1/2</sub> of [<sup>3</sup>H]-L-742,694 (27 min, Table 2.4) at 22 °C is considerably shorter than the half-life of aprepitant but longer than the natural ligand SP.<sup>19,111,112</sup> Its functional interaction was

further characterized by Cascieri *et al.* by using both human and rat NK<sub>1</sub> receptor.<sup>111</sup> Interestingly, its nature of NK<sub>1</sub> receptor antagonism was species related as L-742,694 was insurmountable on the hNK<sub>1</sub> receptor but not on the rNK<sub>1</sub> receptor. A follow-up mutational analysis revealed that mutating H197<sup>5,39</sup> in TM5 into Serine reduced L-742,694 affinity and simultaneously abolished its functional insurmountability at the hNK<sub>1</sub> receptor, making it a key amino acid for L-742,694 interaction and function. When testing several structural analogues of L-742,694, the authors found that the addition of the triazoline moiety (as in L-742,694) converted a surmountable antagonist into an insurmountable one, possibly by slowing the dissociation rate.<sup>111</sup>

### 3. Compound 22

Aprepitant has formed the basis for the discovery of more potentially long acting NK<sub>1</sub>R antagonists. Young *et al.* developed a series of pyrrolidine-carboxamides and oxadiazoles based on aprepitant, aiming for feasible clinical candidates for a once-daily dosing regimen.<sup>114</sup> Their screening strategy was to find compounds with enduring functional activity *in vitro* and long duration of action *in vivo* at lower drug levels as to obtain an improved safety profile. Overall, Compound 22 (Figure 2.7, Compound 3) was found to be the best compound with efficacious functional activity in an IP-1 functional assay and long duration of action (over 24 h) in the GFT model. During the course of compound optimization, the authors were also able to establish a kinetically informed structure-activity relationship. They concluded that both receptor binding affinity and functional insurmountability are highly sensitive to both absolute and relative stereochemistry,<sup>114</sup> which was reaffirmed for another compound series by Morriello *et al.*<sup>115</sup>

### 4. Compound 20

Morriello *et al.* employed virtually the same screening technique as described by Young *et al.*<sup>114,115</sup> to develop both a series of long acting 5,5-fused pyrrolidines<sup>115-117</sup> and later a series of 5,5,5-fused tricyclic antagonists with even slower dissociation rates, with cyclopentane compound 20 (Figure 2.7,

Compound 4) being the most active.<sup>118</sup> The authors found that increasing the overall compound rigidity as well as modifying some other specific molecular components, such as by the addition of a substituent on the pyrrolidine nitrogen, increased functional insurmountability.

## 5. Vestipitant and its analogues

Vestipitant (Figure 2.7, Compound 5) has a  $t_{1/2}$  of 59.7 min (Table 2.4, 37 °C) from the human NK<sub>1</sub>R.<sup>119</sup> It is an insurmountable yet competitive antagonist with long duration of action (over 8 h) in the *in vivo* GFT model.<sup>119</sup> Compared to other structural analogues of vestipitant, it appeared that certain features such as the benzylic substitution patterns and its stereochemistry were important for vestipitant's mode of antagonism,<sup>119</sup> which was reaffirmed in another publication by Di Fabio *et al.*<sup>120</sup> Surprisingly, even a small, hydrophobic substituent, for instance a methyl substituent, played an important role in determining the magnitude of an antagonist's insurmountability and duration of action.<sup>120</sup>

## 6. MEN1149

This peptidomimetic compound (Figure 2.7, Compound 6) set the example that long acting small molecule antagonists can be developed from short acting peptide antagonists, in this case from FK888, by simple structural modifications. The main modification introduced, originally to increase metabolic stability, was the  $\beta$ -aminocycloalkyl carboxylic acid residue replacing FK888's prolyl group.<sup>121</sup> Interestingly, these modifications turned the purely surmountable FK888 into the insurmountable MEN1149 with much slower reversibility during organ bath washout assays and a longer duration of action *in vivo* of over 3 h.<sup>121</sup> The authors speculated that introduction of the  $\beta$ -amino cycloalkyl carboxylic acid caused a slower rate of dissociation which was likely to be responsible for the change in type antagonism, but only provided circumstantial evidence and no direct kinetics measurements.<sup>121</sup>



Overall, the examples summarized above provide a strong indication that NK<sub>1</sub>R are amongst the targets for which slowly dissociating antagonists could offer therapeutic advantages.

## Dopamine receptors

Dopamine receptors are a group of class A GPCRs, which are named after their endogenous ligand dopamine.<sup>122</sup> To date, five subtypes of dopamine receptors have been identified, namely D<sub>1</sub> to D<sub>5</sub>.<sup>123</sup> These receptors can be further categorized into D<sub>1</sub>-like (D<sub>1</sub> and D<sub>5</sub>) and D<sub>2</sub>-like (D<sub>2</sub>, D<sub>3</sub> and D<sub>4</sub>) subfamilies, based on their similarities in sequence, pharmacology and ability to stimulate or inhibit adenylyl cyclase activity: the D<sub>1</sub>-like subfamily couples to G<sub>s</sub> and mediates excitatory neurotransmission, while the D<sub>2</sub>-like (D<sub>2</sub>, D<sub>3</sub> and D<sub>4</sub>) subfamily couples to G<sub>i/o</sub> and mediates inhibitory neurotransmission.<sup>124</sup> Dopamine receptors, particularly D<sub>1</sub> and D<sub>2</sub> receptors, are prominent in the vertebrate CNS,<sup>123,125</sup> indicating their functional importance in neuronal signaling (for reviews, see<sup>126-131</sup>). To date, numerous drugs targeting the D<sub>2</sub> receptor are available on the market. Antagonizing the dopamine D<sub>2</sub> receptor has been reported as the molecular foundation of schizophrenia treatment.<sup>132</sup> Currently, all known marketed anti-schizophrenia drugs bind to the D<sub>2</sub> receptor and are supposed to modulate dopamine hyperfunction.<sup>26,133</sup> In general, these drugs can be categorized into two groups. The first generation or the so-called typical antipsychotics (e.g., haloperidol) can effectively treat positive schizophrenia symptoms (i.e., hallucination and delusions), yet they invoke concomitant on-target adverse effects, such as extrapyramidal symptoms (EPS) and hyperprolactinaemia, through excessive blockade of the D<sub>2</sub> receptor. The second generation, also known as 'atypical' antipsychotics, are defined according to their 'clozapine-like' properties, i.e., low incidence of the on-target/mechanism-based side effects, which are mostly attributed to their short residence times at the D<sub>2</sub> receptor.<sup>134</sup> Other theories have also been proposed to explain 'atypicality'. One of these is the multi-receptor hypothesis or polypharmacology.<sup>180</sup> Many atypical antipsychotics were found to have high affinity at other neurotransmitter receptors in addition to D<sub>2</sub> receptors, in particular at serotonin 5-HT<sub>2A</sub> receptors, while typical antipsychotics, like haloperidol, bind more specifically to D<sub>2</sub> receptors.<sup>135</sup> However, fully occupying serotonin 5-HT<sub>2A</sub> receptors with different antipsychotics at




Figure 2.8 | Structures of small molecule  $D_2$  receptor antagonists.

clinically relevant doses did not induce EPS to the same level.<sup>134</sup> Hence, the atypicality is more likely connected to the compounds' short residence times at the D<sub>2</sub> receptor than their antagonism at the 5-HT<sub>2A</sub> receptors.

Additionally, it is worth mentioning that interaction of atypical antipsychotics with additional receptors may cause undesired off-target side-effects—distinct from its on-target side-effect (e.g., EPS and hyperprolactinaemia), such as 5HT<sub>2C</sub> receptors (weight gain),  $\alpha_1$ -adrenoceptors (orthostatic hypotension, reflex tachycardia, and hypnosedation),  $\alpha_2$ -adrenoceptors (tachycardia), histamine H<sub>1</sub> receptors (sedation and weight gain), and muscarinic receptors (blurred vision, dry mouth, constipation, and cognitive impairment).<sup>135</sup> These side-effects can logically be minimized by decreasing the compound's affinity or residence time on the off-targets.

Contrary to aforementioned cases of other GPCRs, where long RT ligands are desired for optimal clinical outcomes, short residence time D<sub>2</sub> receptor antagonists are preferred, since they will supposedly induce less on-target 'mechanism-based toxicology'.<sup>25,35</sup> In the following part of this section we will focus on published binding kinetics data of dopamine D<sub>2</sub> antagonists and the related benefit from short RT in clinical applications.

## 1. Haloperidol and other slowly dissociating D<sub>2</sub> receptor antagonists

Haloperidol (Figure 2.8, Compound 1) was approved by the U.S. Food and Drug Administration (FDA) in 1967 and marketed as Haldol in U.S. and other countries. Haloperidol exhibited high-affinity antagonism for human dopamine D<sub>2</sub> receptor (1.8 nM) with a  $k_{\text{off}}$  of 0.01 s<sup>-1</sup>, equal to a  $t_{1/2}$  of 72.4 s (Table 2.5).<sup>135</sup> Notably, its  $t_{1/2}$  was substantially changed when the kinetics were tested at rat striatal membranes (i.e.,  $t_{1/2}$  = 42 min), indicating significant species differences.<sup>134</sup> One might argue that the absolute value of haloperidol's  $t_{1/2}$  is much smaller than other classical slowly dissociating GPCR antagonists, such as aforementioned NK<sub>1</sub>R antagonist aprepitant ( $t_{1/2}$  = 154 min). It should be realized that the time scale for endogenous dopamine concentration burst can be within a millisecond to second range, which is much shorter than the  $t_{1/2}$  of haloperidol.<sup>136</sup> In this sense, the receptor blockade by haloperidol is already too lengthy to allow normal physiological responses to fast and small changes in the neurotransmitter concentration.

**Table 2.5 | Binding kinetics of D<sub>2</sub> receptor antagonists.**

| Nr | Cmpd <sup>a</sup> | Affinity (K <sub>i</sub> , nM) <sup>b</sup> | Dissociation half-life (t <sub>1/2</sub> ) <sup>c</sup> | Ref |
|----|-------------------|---------------------------------------------|---------------------------------------------------------|-----|
| 1  | Haloperidol       | 1.8                                         | 72.4 s                                                  | 135 |
| 2  | Nemonapride       | 0.025 (rat)                                 | 355 min (rat)                                           | 134 |
| 3  | Spiperone         | 0.1 (rat)                                   | 200 min (rat)                                           | 134 |
| 4  | Sertindole        | 1.2 (rat)                                   | 47 min (rat)                                            | 134 |
| 5  | Chlorpromazine    | 1.3 (rat)                                   | 35 min (rat)                                            | 134 |
| 6  | Raclopride        | 1.6 (rat)                                   | 28 min (rat)                                            | 134 |
| 7  | Olanzapine        | 6.4 (rat)                                   | 18 min (rat)                                            | 134 |
| 8  | Clozapine         | 137                                         | 14.5 s                                                  | 135 |
| 9  | Quetiapine        | 67.6 (rat)                                  | 23 s (rat)                                              | 139 |
| 10 | JNJ-37822681      | 158                                         | 6.5 s                                                   | 135 |

<sup>a</sup> Compound names and synonyms used in original references. <sup>b</sup> Affinity, expressed as K<sub>i</sub> values that reported in original references; result on human receptor unless mentioned otherwise. <sup>c</sup> Dissociation half-life at room temperatures and on human receptors unless mentioned otherwise.

Next to the result from the direct kinetic binding assay at both human and rat D<sub>2</sub> receptor preparations, haloperidol was found to have long-lasting D<sub>2</sub> blockade by using an indirect kinetic measurement. Only 48% (lower than that of clozapine) of the total receptor population was recovered and able to bind to subsequently added [<sup>3</sup>H]-spiperone after pre-incubating haloperidol for one hour with the receptor.<sup>135</sup>

The binding kinetics of other slowly dissociating D<sub>2</sub> receptor antagonists were elaborately studied by Kapur and Seeman.<sup>134</sup> The authors investigated the affinity and the kinetics of nine typical and atypical antipsychotics (main results are included in Table 2.5; compound structures in Figure 2.8). Among these compounds, nemonapride and spiperone had a t<sub>1/2</sub> of 355 min and 200 min, respectively, indicative of D<sub>2</sub> receptor blockade for more than 3 h; these two compounds were followed by sertindole (47 min), haloperidol (40 min), chlorpromazine (35 min), raclopride (28 min) and olanzapine (18 min) in terms of their t<sub>1/2</sub> values. In contrast, the two atypical antipsychotics clozapine and quetiapine displayed faster dissociation half-lives, i.e., less than 1 min. Furthermore, it is important to note that the k<sub>off</sub> values varied a 1000-fold from 0.002 min<sup>-1</sup> to 3.013 min<sup>-1</sup>, yet less variation was found in k<sub>on</sub>. Apparently, the rate at which antipsychotic agents come off the receptor (k<sub>off</sub>)

causes the variation in their affinity for the  $D_2$  receptor, while differences in  $k_{on}$  do not account for that. Moreover, since atypical antipsychotics display relatively faster dissociation than typical ones, it can be inferred that the atypical antipsychotic effects are mostly attributed to their high  $k_{off}$  values for the  $D_2$  receptor.

## 2. Clozapine

Clozapine was the first atypical antipsychotic drug on the market (Figure 2.8, Compound 8). Compared to haloperidol, clozapine had a relatively low affinity for the  $D_2$  receptor ( $K_i = 137$  nM) and a 5-fold shorter  $t_{1/2}$  of approximately 14.5 s (Table 2.5).<sup>135</sup> This (partially) explained the observation by Kapus and Seeman, that clozapine reached equilibrium 100 times faster than haloperidol in the displacement assay.<sup>134</sup> Such a fast molecular interaction enabled clozapine to have a fast-on and then fast-off blockade of the  $D_2$  receptor. Therefore, the receptor was readily responsive to the endogenous dopamine burst upon physical stimulation, hence lowering the risk of EPS or hyperprolactinaemia. This mechanism was further supported by the *in vivo* observation that clozapine caused a surmountable blockade of striatal dopamine receptors in contrast to cataleptogenic neuroleptics (the insurmountable control).<sup>137</sup> Moreover, clinical positron emission tomography (PET) analysis of a subject receiving clozapine (400 mg) revealed transiently high  $D_2$  receptor occupancy (i.e., 80%) that rapidly declined to low levels (i.e., 15%) within 48 h. In comparison,  $D_2$  receptor occupancy by haloperidol (5 mg) peaked at approximately 80%, but it declined more slowly over time, showing  $D_2$  occupancy of over 10% even 3–5 days later.<sup>138</sup> This reaffirmed that fast kinetics may be favorable for a low incidence of mechanism-based, on-target, toxicology.

## 3. Quetiapine

Quetiapine (Figure 2.8, Compound 9), also known as Seroquel, Xeroquel or Ketipinor, is another marketed atypical antipsychotic approved for

schizophrenia treatment. Similar to the molecular properties of clozapine, Caboni *et al.* found that quetiapine displayed low  $D_2$  receptor affinity ( $K_i = 67.6$  nM) and a fast dissociation rate ( $1.8 \text{ min}^{-1}$ ), equal to a  $t_{1/2}$  of 23 s, at a rat striatum homogenate (Table 2.5).<sup>139</sup> In the subsequent *in vivo* profiling, Caboni *et al.* performed time-dependent analysis of plasma prolactin levels, a surrogate of peripheral  $D_2$  receptor blockade, after orally treating rats with vehicle or different dose of quetiapine or haloperidol. They found that compared to haloperidol treatment quetiapine displayed a short-lived increase of prolactin level, which was in line with its short RT profile at the  $D_2$  receptor. Moreover, serial PET scans evaluating the  $D_2$  receptor occupancy of quetiapine demonstrated that this compound dissociated very rapidly from the  $D_2$  receptor.<sup>140</sup> Taken together, all *in vitro* and *in vivo* results provided strong evidence that quetiapine is a fast dissociating  $D_2$  receptor antagonist.

#### 4. JNJ-37822681

JNJ-37822681 (Figure 2.8, Compound 10) was developed by Janssen Pharmaceuticals. One of the most elaborate studies on JNJ-37822681 has been reported by Langois *et al.*<sup>135</sup> The compound was initially selected from an indirect kinetic screening assay (for method, see<sup>23,135</sup>), where JNJ-37822681 was estimated to have a  $k_{\text{off}}$  similar to clozapine. Subsequently, a kinetic [ $^3\text{H}$ ]-JNJ-37822681 binding experiment revealed that this compound had an extremely short  $t_{1/2}$  (6.5 s, Figure 2.5) at the  $D_2$  receptor, which was approximately 2-fold and 11-fold shorter than clozapine and haloperidol, respectively. In the follow-up *in vivo* profiling, JNJ-37822681 displayed a larger specific margin (compared with haloperidol) between the inhibition and blockade of apomorphine-induced behaviors, which represents a model for psychosis. This result was further reaffirmed in phase IIb clinical trials, where JNJ-37822681 displayed an efficacious antipsychotic effect but low propensity of EPS and prolactin liability.<sup>141,142</sup>

## 5. Overall structural determinants for binding kinetics at the dopamine D<sub>2</sub> receptor

Compared to the residence times of M<sub>3</sub>R or NK<sub>1</sub>R antagonists, D<sub>2</sub> receptor antagonists are preferred to have a short RT to circumvent on-target side effects. It is therefore highly desired to analyze the structural determinants that increase ligand dissociation from the receptor for future rational drug design. For this reason, Tresadern *et al.* performed a comprehensive statistical analysis of the binding kinetics of over 1800 D<sub>2</sub> receptor antagonists and examined their physicochemical properties that potentially discriminate fast dissociating compounds from slowly dissociating ones.<sup>26</sup> They discovered that faster dissociating antagonists were in general less lipophilic and had lower molecular weight. This seems to hold true for other fast dissociating GPCR ligands as well, a finding shared by Miller *et al.*<sup>143</sup> However, it is also important to note that other descriptors, for instance polar surface area, partial positive charge or the amount of hydrogen bond donor and acceptor groups, caused different effects on the ligand's dissociation, depending on the particular cluster of chemicals that was analyzed. This suggested that local effects of a specific chemical scaffold on the binding kinetics could be dominant over other physiochemical descriptors. Thus, instead of intuitively modifying a compound's lipophilicity, molecular weight or other molecular properties, a more specific structure-kinetics relationship study is needed to direct future drug design and discovery.

In summary, examples presented here demonstrate the benefit of fast dissociating D<sub>2</sub> receptor antagonists for lowering the risk of mechanism-based toxicology. Such an advantage over long RT antagonists is supported by evidence from preclinical and clinical aspects of the typical and atypical antipsychotics. This case study on the D<sub>2</sub> receptor further emphasizes the necessity of characterizing a ligand's binding kinetics profile in the early phases of candidate drug selection.

## 2.7 Drug-target residence time for class B GPCRs

### Corticotrophin releasing factor type-1 receptor

Corticotrophin releasing factor (CRF) is a 41 amino-acid peptide that exerts its effects through activation of CRF receptors, which are members of the family B GPCRs.<sup>144</sup> The CRF receptor subfamily has two subtypes, namely CRF<sub>1</sub> and CRF<sub>2</sub>. Both receptors are primarily coupled to G<sub>s</sub> protein, resulting in the activation of adenylate cyclase and increased cAMP levels.<sup>145,146</sup> Since CRF acts on the pituitary gland to regulate the hypothalamic-pituitary-adrenal axis and the CNS to modulate behavior responses to stress, it has caught considerable attention in drug discovery for psychiatric diseases.<sup>144-147</sup> In particular, antagonism of the CRF<sub>1</sub> receptor has been proposed for rebalancing a dysfunctional stress axis.<sup>147-149</sup> Although highly potent *in vitro* and efficacious in animal models, early reported compounds such as CP-154,526 and NBI-27914 are highly lipophilic, which leads to compound accumulation in tissues or long elimination half-life, and thus high risks of toxicity.<sup>150</sup> For this reason, efforts have been specifically directed towards discovering less lipophilic, fast clearance CRF<sub>1</sub> receptor antagonists to decrease the incidence of toxicity but with maintained drug efficacy. A long residence time CRF<sub>1</sub> receptor antagonist is therefore highly desired, which could provide the benefit of a long duration of action and at the same time less accumulation on the unwanted targets. Such compounds would exemplify new mechanisms for depression treatment beyond classical monoamine modulators.<sup>148,151</sup>

#### 1. The binding kinetics of CRF<sub>1</sub>R antagonists and their *in vivo* pharmacodynamics

One of the most elaborate binding kinetics studies at the CRF<sub>1</sub> receptor was performed by Fleck *et al.*<sup>151</sup> In this article, the authors investigated a series of CRF<sub>1</sub> receptor antagonists regarding their *in vitro* binding kinetics and *in vivo* pharmacodynamics. They found that the  $k_{\text{off}}$  and  $k_{\text{on}}$  values of 12 CRF<sub>1</sub>






Figure 2.9 | Structures of small molecule CRF<sub>1</sub> receptor antagonists.

receptor compounds varied 170-fold and 13-fold, respectively, which resulted in a maximal change of around 510-fold in the kinetically derived affinity ( $K_D = k_{\text{off}} / k_{\text{on}}$ ) at 37 °C. Notably, this finding was in stark contrast with the previous reported  $K_i$  values measured in radioligand displacement assays, where little difference was observed between compounds. This discrepancy was mostly attributed to insufficient incubation time to reach equilibrium for compounds with long residence time profile. For instance, less than 40% of NBI 30775 dissociated from the CRF<sub>1</sub> receptor after 2 h, far from equilibrium, hence the  $K_i$  values were underestimated. Furthermore, several lead compounds displayed divergent dissociation half-lives; NBI 35965 with 16 min, NBI 34041 with 53 min and NBI 30775 with 130 min (Figure 2.9, Compound 1-3; Table 2.6). This ranking of their binding kinetics, in fact, was consistent with the ranking of their *in vivo* pharmacodynamic profiles, at least in adrenalectomized rats, where NBI 34041 and NBI 30775 produced sustained suppression of adrenocorticotropin for 4 h and 6 h, respectively, while NBI 35965 induced a more transient effect (less than 2 h). In addition, all three compounds (NBI 35965, NBI 34041 and NBI 30775)

**Table 2.6 | Binding kinetics of CRF<sub>1</sub> receptor antagonists.**

| <b>Nr</b> | <b>Cmpd<sup>a</sup></b> | <b>Affinity (K<sub>i</sub> nM)<sup>b</sup></b> | <b>Dissociation half-life (t<sub>1/2</sub>)<sup>c</sup></b> | <b>Ref</b> |
|-----------|-------------------------|------------------------------------------------|-------------------------------------------------------------|------------|
| 1.        | NBI 35965               | 2.3                                            | 16 min (37°C)                                               | 151        |
| 2.        | NBI 34041               | 1.7                                            | 53 min (37°C)                                               | 151        |
| 3.        | NBI 30775               | 0.36                                           | 130 min (37°C)                                              | 151        |
| 4.        | DMP-904                 | 1.5                                            | 5.5 h                                                       | 152        |
| 5.        | R121919                 | 12                                             | 12.8 h                                                      | 152        |
| 6.        | Cmpd 25 (Pfizer)        | 11                                             | 10.7 h                                                      | 152        |
| 7.        | SN003                   | 3.64 (rat)                                     | 0.35 h (rat)                                                | 153        |
| 8.        | PF-4325743              | 8.22 (rat)                                     | 0.37 h (rat)                                                | 153        |

<sup>a</sup> Compound names and synonyms used in original references. <sup>b</sup> Affinity, expressed as K<sub>i</sub> values that were reported in original references; result on human receptor unless mentioned otherwise. <sup>c</sup> Dissociation half-life at room temperature and on human receptor unless mentioned otherwise.

had similar pharmacokinetics. This confirmed that the unique sustained adrenocorticotropin suppression evoked by NBI 30775 was derived from its prolonged CRF<sub>1</sub> receptor occupancy rather than its pharmacokinetics.


Functional reversibility assays were also performed to kinetically profile CRF<sub>1</sub> receptor antagonists.<sup>152,153</sup> Briefly, cells expressing recombinant CRF<sub>1</sub> receptor were incubated with CRF in either the presence or absence of an antagonist to measure the suppression of the maximal cell response to CRF by the tested antagonist. Although CRF binds to a different site on the receptor than the non-peptide small molecule antagonists, the insurmountability of the non-competitive antagonists still correlated to their off-rates. In particular, Miller *et al.* plotted the antagonists' dissociation half-lives against their suppression of the maximal CRF<sub>1</sub> receptor response to its endogenous ligand. They found a good correlation between the two parameters, although the exact molecular mechanism of this link is yet to be discovered.<sup>152</sup> By using this assay, at least in a qualitative manner, Miller *et al.* investigated several other long RT compounds reported in literature. They found that DMP-904 and R121919 almost completely suppressed the maximal CRF response (Figure 2.9, Compound 4 and 5). Both compounds were found to have slow dissociation half-lives, i.e., 5.5 h and 12.8 h, respectively, which were calculated from k<sub>off</sub> values determined in a competition association assay.<sup>152</sup> In comparison, shorter residence time CRF<sub>1</sub> receptor antagonists (Figure 2.9,

Compound 7 and 8) such as SN003 ( $t_{1/2} = 0.35$  h, Table 2.6) or PF-4325743 ( $t_{1/2} = 0.37$  h, Table 2.6) did not significantly decrease the maximal CRF response.<sup>153</sup>

## 2. Overall structural determinants for binding kinetics at the CRF<sub>1</sub> receptor

A detailed structure-kinetics relationship (SKR) study was performed by Fleck *et al.*, in which the authors performed a retrospective binding kinetics investigation on a set of CRF<sub>1</sub> receptor ligands including the initial, low-affinity chemical starting points up to the lead compounds. Key features of the high affinity or long residence time CRF<sub>1</sub> receptor ligand are represented in Figure 2.10A (adapted from the models developed by Gilligan *et al.*, Kehne and De Lombaert, and Fleck *et al.*<sup>151,154,155</sup>) and the SKRs are summarized as follows: 1) *ortho*-position substitution ( $R_1$ ) on the low ring was necessary to enforce a twisted bioactive conformation to enhance ligand-receptor interactions. This can be illustrated by the conformation of CP-376395 in the recently elucidated CRF<sub>1</sub>R crystal structure (PDB code: 4K5Y, Figure 2.10B). Apparently, the pose of the compound's trimethylphenoxy moiety is rotated to fit in a hydrophobic pocket formed by F284<sup>5,51</sup>, L287<sup>5,54</sup>, T316<sup>6,42</sup>, L319<sup>6,45</sup> and L320<sup>6,46</sup> with the pyridine nitrogen at the core to form a hydrogen bond with N283<sup>5,50</sup>.<sup>156</sup> In contrast, the association rate of an *ortho*-substituent free analogue decreased 33-fold compared to its chloro-substituted analogue, suggesting a substantial barrier for the free rotating ligand to fit into the binding pocket. 2) Adding a methyl moiety at  $R_2$  resulted in an increased  $k_{on}$  and decreased  $k_{off}$ , indicating a raised probability of successful collision between this group and a rigid binding pocket on the receptor, combined with a stabilized ligand-receptor conformation. This finding was in agreement with other published results which suggested that the methyl-group interacted with a hydrophobic pocket in the receptor.<sup>154,155,157,158</sup> 3) Chemical modifications on the upper aliphatic groups at  $R_3$  demonstrated varying effects on the binding kinetics depending on their size, branching pattern and the stereochemical configuration, which may affect the compound's interaction with F203<sup>3,44</sup> and Y327<sup>6,53</sup>, two aromatic residues gating the small access channel from the extracellular part (Figure 2.10B).<sup>156</sup>

In another SKR study, Miller *et al.* started from a bicyclic purine core and



**Figure 2.10** | (A) Diagram of the putative functional groups implicated in structure-kinetics relationships of non-peptide antagonists. This diagram was adapted from literature,<sup>160,162,163</sup> based on the comprehensive analysis of ligand SAR. (B) The co-crystal structure of CP-376395 in the human CRF<sub>1</sub> receptor. This figure was generated with ICM Browser v3.7 (Molsoft) from PDB code: 4K5Y. CP-376395 (black) binds deeply within the pocket. Two aromatic residues, F203<sup>3.44</sup> and Y327<sup>6.53</sup>, together prevent the ligand from moving out of the receptor. The pose of the compound's trimethylphenoxy moiety is rotated to fit in a hydrophobic pocket formed by F284<sup>5.51</sup>, L287<sup>5.54</sup>, T316<sup>6.42</sup>, L319<sup>6.45</sup> and L320<sup>6.46</sup> with its pyridine nitrogen at the core to form a hydrogen bond with N283<sup>5.50, 156</sup>

performed a series of chemical modifications on the substitution patterns at the low ring and the upper branched groups. Follow-up functional investigations *in vitro* led to the identification of compound 25 (Figure 2.9, Compound 6) which displayed 79% suppression of the maximal CRF response. Binding kinetics characterization of this compound revealed that it had a slow  $t_{1/2}$  of 10.7 h, correlating well with its insurmountable antagonist behavior.<sup>152</sup>

## 2.8 Concluding remarks

We reviewed the existing kinetics data for a number of GPCRs. The examples presented make a strong case for the value of measuring drug-target residence time, or, more generally speaking, the kinetic profile of interaction for

compound optimization in the early phases of the drug discovery process. Thus, a strategy is warranted that comprises both a structure-kinetics relationship study (SKR) and a classical structure-affinity relationship study (SAR). This may offer added value to the traditional metric of affinity-based drug evaluation.

This chapter further summarized the factors influencing drug-target residence time, in particular, to find a universal molecular determinant or a certain ‘hot spot’ on GPCRs for designing kinetics-favorable ligands. Several kinetics-related statistical analyses indicated that manipulating physicochemical properties such as lipophilicity or molecular weight can lead to tuned drug-target residence times.<sup>26,143</sup> Alternatively, this could be achieved by modulating the rigidity / flexibility of a drug candidate.<sup>117</sup> However, these ‘intuitive’, ‘one size-fits-all’ approaches are very likely too simple and may increase the concern of binding to collateral targets, thereby resulting in unwanted side effects.<sup>67,159</sup> In this sense, fine-tuning drug-target residence time should probably be more specific. Despite this challenge, progress can still be made by gaining knowledge from both the ligand and target perspectives, such as by ligand modifications with subtle changes to probe relevant pharmacophores, mutation studies, co-crystallization and molecular dynamic studies to establish the corresponding ‘anchoring’ sites on the receptor.

We also summarized the kinetic methodologies in this chapter. Several screening schemes are available and new assays are emerging, which enable quick identification of compounds of interest. Quantitative characterization of a compound’s binding kinetics can be obtained in a kinetic binding assay or competition association assay. Functional reversibility experiments prevail in drug discovery as well, which offer an alternative way to interpret a ligand’s kinetics profile. In addition, *ex vivo* receptor occupancy experiments or *in vivo* duration of action experiments can translate the *in vitro* binding kinetics into the effect in an *in vivo* setting, which provides further evidence for a compound’s drug-target residence time. Notably, it is also necessary to take into account the PK profile of the candidate compound, particular for one aiming at long drug-target residence time, since RT can only prolong the duration of action only when the RT outlasts the PK.<sup>2,15</sup>

The compounds summarized in the present chapter are predominantly GPCR antagonists. This reflects a lack of kinetic information on agonists in the literature. However, more kinetics studies focused on the latter are emerging. For instance the case studies at the M<sub>3</sub> and A<sub>2A</sub> receptors

explored the molecular basis of the compounds' intrinsic efficacy from a kinetic perspective.<sup>32,98</sup> In addition to examining an agonist's mechanism of action, one can also specifically optimize an agonist's kinetic properties for therapeutic benefit. This can be exemplified by the administration of long-acting agonists at the GnRH receptor to intentionally desensitize the GnRH receptor and therefore antagonize gonadotropin secretion in hormone-dependent cancers.<sup>181</sup> Such a strategy could be a therapeutic substitute for targets where no antagonist is available. On the other hand, one needs to take into consideration that agonist-induced desensitization or internalization by prolonged or repeated administration can cause drug tolerance, for instance opiate drugs targeting  $\mu$ -opioid receptors.<sup>182</sup> In such a case, it might be tempting to search for shorter residence time agonists to prevent their target's internalization. Thus, a study that compares agonist binding kinetics and GPCR desensitization / internalization kinetics would be of great interest.

Taken together, a rapid expansion of the field of GPCR binding kinetics may be anticipated in the near future, which hopefully will lead to the identification of novel and better-in-class drugs for clinical applications.

## References:

1. Copeland RA, Pompliano DL, Meek TD. Drug-target residence time and its implications for lead optimization. *Nat Rev Drug Discov* 2006;5(9):730-739.
2. Swinney DC. Can binding kinetics translate to a clinically differentiated drug? From theory to practice. *Lett Drug Des Discov* 2006;3(8):569-574.
3. Swinney DC. Biochemical mechanisms of New Molecular Entities (NMEs) approved by United States FDA during 2001-2004: mechanisms leading to optimal efficacy and safety. *Curr Top Med Chem* 2006;6(5):461-478.
4. Swinney DC, Anthony J. How were new medicines discovered? *Nat Rev Drug Discov* 2011;10(7):507-519.
5. Swinney DC. Biochemical mechanisms of drug action: what does it take for success? *Nat Rev Drug Discov* 2004;3(9):801-808.
6. Copeland, RA. *Evaluation of enzyme inhibitors in drug discovery: a guide for medicinal chemists and pharmacologists*. New York: John Wiley & Sons; 2005. 153 p.
7. Copeland RA. Conformational adaptation in drug-target interactions and residence time. *Future Med Chem* 2011;3(12):1491-1501.
8. Vauquelin G, Charlton SJ. Long-lasting target binding and rebinding as mechanisms to prolong *in vivo* drug action. *Br J Pharmacol* 2010;161(3):488-508.
9. Vauquelin G, Fierens F, Verheijen I, Vanderheyden P. Insurmountable AT<sub>1</sub> receptor antagonism: the need for different antagonist binding states of the receptor. *Trends Pharmacol Sci*

- 2001;22(7):343-344.
10. Vauquelin G, Morsing P, Fierens FL, De Backer JP, Vanderheyden PM. A two-state receptor model for the interaction between angiotensin II type 1 receptors and non-peptide antagonists. *Biochem Pharmacol* 2001;61(3):277-284.
11. Vauquelin G, Van Liefde I. Slow antagonist dissociation and long-lasting *in vivo* receptor protection. *Trends Pharmacol Sci* 2006;27(7):356-359.
12. Zhang R, Monsma F. The importance of drug-target residence time. *Curr Opin Drug Discov Devel* 2009;12(4):488-496.
13. Zhang R, Monsma F. Binding kinetics and mechanism of action: toward the discovery and development of better and best in class drugs. *Expert Opin Drug Discov* 2010;5(11):1023-1029.
14. Tummino PJ, Copeland RA. Residence time of receptor-ligand complexes and its effect on biological function. *Biochemistry* 2008;47(20):5481-5492.
15. Dahl G, Akerud T. Pharmacokinetics and the drug-target residence time concept. *Drug Discov Today* 2013;18(15-16):697-707.
16. Nunez S, Venhorst J, Kruse CG. Target-drug interactions: first principles and their application to drug discovery. *Drug Discov Today* 2012;17(1-2):10-22.
17. Swinney DC. The role of binding kinetics in therapeutically useful drug action. *Curr Opin Drug Discov Devel* 2009;12(1):31-39.
18. Vilums M, Zweemer AJ, Yu Z, de Vries H, Hillger JM, Wapenaar H, Bollen IA, Barmare F, Gross R, Clemens J, Krenitsky P, Brussee J, Stamos D, Saunders J, Heitman LH, IJzerman AP. Structure-kinetic relationships-an overlooked parameter in hit-to-lead optimization: a case of cyclopentylamines as chemokine receptor 2 antagonists. *J Med Chem* 2013;56(19):7706-7714.
19. Hale JJ, Mills SG, MacCoss M, Finke PE, Cascieri MA, Sadowski S, Ber E, Chicchi GG, Kurtz M, Metzger J, Eiermann G, Tsou NN, Tattersall FD, Rupniak NM, Williams AR, Rycroft W, Hargreaves R, MacIntyre DE. Structural optimization affording 2-(R)-(1-(R)-3, 5-bis(trifluoromethyl) phenylethoxy)-3-(S)-(4-fluoro)phenyl-4- (3-oxo-1,2,4-triazol-5-yl)methylmorpholine, a potent, orally active, long-acting morpholine acetal human NK-1 receptor antagonist. *J Med Chem* 1998;41(23):4607-4614.
20. Gavalda A, Miralpeix M, Ramos I, Otal R, Carreno C, Vinals M, Domenech T, Carcasona C, Reyes B, Vilella D, Gras J, Cortijo J, Morcillo E, Llenas J, Ryder H, Beleta J. Characterization of acridinium bromide, a novel inhaled muscarinic antagonist, with long duration of action and a favorable pharmacological profile. *J Pharmacol Exp Ther* 2009;331(2):740-751.
21. Keppler A, Gendreizig S, Gronemeyer T, Pick H, Vogel H, Johnsson K. A general method for the covalent labeling of fusion proteins with small molecules *in vivo*. *Nat Biotechnol* 2003;21(1):86-89.
22. Pierre N. Determination of association ( $k_{on}$ ) and dissociation ( $k_{off}$ ) rates of spiperone on the dopamine D<sub>2</sub> receptor using a platform for GPCR applications. Volume 2013. Bedford: *American Laboratory*; 2011.
23. Leysen JE, Gommeren W. The dissociation rate of unlabelled dopamine antagonists and agonists from the dopamine-D<sub>2</sub> receptor, application of an original filter method. *J Recept Res* 1984;4(7):817-845.
24. Malany S, Hernandez LM, Smith WF, Crowe PD, Hoare SR. Analytical method for simultaneously measuring *ex vivo* drug receptor occupancy and dissociation rate: application to (R)-dimethindene occupancy of central histamine H<sub>1</sub> receptors. *J Recept Signal Transduct Res* 2009;29(2):84-93.
25. Packeu A, Wennerberg M, Balendran A, Vauquelin G. Estimation of the dissociation rate of

- unlabelled ligand-receptor complexes by a 'two-step' competition binding approach. *Br J Pharmacol* 2010;161(6):1311-1328.
26. Tresadern G, Bartolome JM, Macdonald GJ, Langlois X. Molecular properties affecting fast dissociation from the D<sub>2</sub> receptor. *Bioorg Med Chem* 2011;19(7):2231-2241.
  27. Affolter H, Hertel C, Jaeggi K, Portenier M, Staehelin M. (-)-S-[<sup>3</sup>H]CGP-12177 and its use to determine the rate constants of unlabeled beta-adrenergic antagonists. *Proc Natl Acad Sci U S A* 1985;82(3):925-929.
  28. Contreras ML, Wolfe BB, Molinoff PB. Kinetic analysis of the interactions of agonists and antagonists with beta adrenergic receptors. *J Pharmacol Exp Ther* 1986;239(1):136-143.
  29. Dowling MR, Charlton SJ. Quantifying the association and dissociation rates of unlabelled antagonists at the muscarinic M<sub>3</sub> receptor. *Br J Pharmacol* 2006;148(7):927-937.
  30. Heise CE, Sullivan SK, Crowe PD. Scintillation proximity assay as a high-throughput method to identify slowly dissociating nonpeptide ligand binding to the GnRH receptor. *J Biomol Screen* 2007;12(2):235-239.
  31. Slack RJ, Russell LJ, Hall DA, Luttmann MA, Ford AJ, Saunders KA, Hodgson ST, Connor HE, Browning C, Clark KL. Pharmacological characterization of GSK1004723, a novel, long-acting antagonist at histamine H<sub>1</sub> and H<sub>3</sub> receptors. *Br J Pharmacol* 2011;164(6):1627-1641.
  32. Guo D, Mulder-Krieger T, IJzerman AP, Heitman LH. Functional efficacy of adenosine A<sub>2A</sub> receptor agonists is positively correlated to their receptor residence time. *Br J Pharmacol* 2012;166(6):1846-1859.
  33. Guo D, van Dorp EJ, Mulder-Krieger T, van Veldhoven JP, Brussee J, IJzerman AP, Heitman LH. Dual-point competition association assay: a fast and high-throughput kinetic screening method for assessing ligand-receptor binding kinetics. *J Biomol Screen* 2013;18(3):309-320.
  34. Motulsky HJ, Mahan LC. The kinetics of competitive radioligand binding predicted by the law of mass action. *Mol Pharmacol* 1984;25(1):1-9.
  35. Vauquelin G, Bostoen S, Vanderheyden P, Seeman P. Clozapine, atypical antipsychotics, and the benefits of fast-off D<sub>2</sub> dopamine receptor antagonism. *Naunyn Schmiedeberg's Arch Pharmacol* 2012;385(4):337-372.
  36. Gaddum JH, Hameed KA, Hathway DE, Stephens FF. Quantitative studies of antagonists for 5-hydroxytryptamine. *Q J Exp Physiol Cogn Med Sci* 1955;40(1):49-74.
  37. Kenakin T, Jenkinson S, Watson C. Determining the potency and molecular mechanism of action of insurmountable antagonists. *J Pharmacol Exp Ther* 2006;319(2):710-723.
  38. Paton WD, Waud DR. The margin of safety of neuromuscular transmission. *J Physiol* 1967;191(1):59-90.
  39. Paton WDM, Rang HP. *A kinetic approach to the mechanism of drug action*. In: Harper NJ, Simmonds AB, editor. *Adv. Drug Res.* London & New York: Academic Press; 1998. p57-80.
  40. Rang HP. The kinetics of action of acetylcholine antagonists in smooth muscle. *Proc R Soc Lond B Biol Sci* 1966;164(996):488-510.
  41. Fierens FL, Vanderheyden PM, De Backer JP, Vauquelin G. Insurmountable angiotensin AT<sub>1</sub> receptor antagonists: the role of tight antagonist binding. *Eur J Pharmacol* 1999;372(2):199-206.
  42. Morsing P, Vauquelin G. How can the differences among AT<sub>1</sub>-receptor antagonists be explained? *Cell Biochem Biophys* 2001;35(1):89-102.
  43. Vauquelin G, Fierens FL, Verheijen I, Vanderheyden PM. Distinctions between non-peptide angiotensin II AT<sub>1</sub>-receptor antagonists. *J Renin-angio-aldo S* 2001;2(1 suppl):S24-S31.
  44. Vauquelin G, Van Liefde I, Birzbier BB, Vanderheyden PM. New insights in insurmountable antagonism. *Fundam Clin Pharmacol* 2002;16(4):263-272.



45. Vauquelin G, Van Liefde I, Vanderheyden P. Models and methods for studying insurmountable antagonism. *Trends Pharmacol Sci* 2002;23(11):514-518.
46. Vauquelin G, Szczuka A. Kinetic versus allosteric mechanisms to explain insurmountable antagonism and delayed ligand dissociation. *Neurochem Int* 2007;51(5):254-260.
47. Shiau AK, Massari ME, Ozbal CC. Back to basics: Label-free technologies for small molecule screening. *Comb Chem High T Scr* 2008;11(3):231-237.
48. Rich RL, Myszkla DG. Higher-throughput, label-free, real-time molecular interaction analysis. *Anal Biochem* 2007;361(1):1-6.
49. Gronewold TMA, Baumgartner A, Hierer J, Sierra S, Blind M, Schafer F, Blumer J, Tillmann T, Kiwitz A, Kaiser R, Zabe-Kuhn M, Quandt E, Famulok M. Kinetic binding analysis of aptamers targeting HIV-1 proteins by a combination of a microbalance array and mass spectrometry (MAMS). *J Proteome Res* 2009;8(7):3568-3577.
50. Gronewold TMA. Surface acoustic wave sensors in the bioanalytical field: Recent trends and challenges. *Analytica Chimica Acta* 2007;603(2):119-128.
51. Rich RL, Myszkla DG. Grading the commercial optical biosensor literature—Class of 2008: 'The Mighty Binders'. *J Mol Recogn* 2010;23(1):1-64.
52. Zhukov A, Andrews SP, Errey JC, Robertson N, Tehan B, Mason JS, Marshall FH, Weir M, Congreve M. Biophysical mapping of the adenosine A<sub>2A</sub> receptor. *J Med Chem* 2011;54(13):4312-4323.
53. Robertson N, Jazayeri A, Errey J, Baig A, Hurrell E, Zhukov A, Langmead CJ, Weir M, Marshall FH. The properties of thermostabilised G protein-coupled receptors (StaRs) and their use in drug discovery. *Neuropharmacol* 2011;60(1):36-44.
54. Congreve M, Andrews SP, Dore AS, Hollenstein K, Hurrell E, Langmead CJ, Mason JS, Ng IW, Tehan B, Zhukov A, Weir M, Marshall FH. Discovery of 1,2,4-triazine derivatives as adenosine A(2A) antagonists using structure based drug design. *J Med Chem* 2012;55(5):1898-1903.
55. Navratilova I, Dioszegi M, Myszkla DG. Analyzing ligand and small molecule binding activity of solubilized GPCRs using biosensor technology. *Anal Biochem* 2006;355(1):132-139.
56. Bjarnadottir TK, Gloriam DE, Hellstrand SH, Kristiansson H, Fredriksson R, Schioth HB. Comprehensive repertoire and phylogenetic analysis of the G protein-coupled receptors in human and mouse. *Genomics* 2006;88(3):263-273.
57. Pierce KL, Premont RT, Lefkowitz RJ. Seven-transmembrane receptors. *Nat Rev Mol Cell Biol* 2002;3(9):639-650.
58. De Amici M, Dallanocce C, Holzgrabe U, Trankle C, Mohr K. Allosteric ligands for G protein-coupled receptors: a novel strategy with attractive therapeutic opportunities. *Med Res Rev* 2010;30(3):463-549.
59. Overington JP, Al-Lazikani B, Hopkins AL. Opinion - How many drug targets are there? *Nat Rev Drug Discov* 2006;5(12):993-996.
60. Russ AP, Lampel S. The druggable genome: an update. *Drug Discovery Today* 2005;10(23-24):1607-1610.
61. Chen L, Jin L, Zhou N. An update of novel screening methods for GPCR in drug discovery. *Expert Opin Drug Discov* 2012;7(9):791-806.
62. Post JM, Alexander S, Wang YX, Vincelette J, Vergona R, Kent L, Bryant J, Sullivan ME, Dole WP, Morser J, Subramanyam B. Novel P<sub>2</sub>Y<sub>12</sub> adenosine diphosphate receptor antagonists for inhibition of platelet aggregation (II): pharmacodynamic and pharmacokinetic characterization. *Thromb Res* 2008;122(4):533-540.
63. Foldes FF. Pain control with intrathecally and peridurally administered opioids and other drugs.

- Anaesthesiol Reanim* 1991;16(5):287-298.
64. Laduron PM, Janssen PFM. Axoplasmic-transport and possible recycling of opiate receptors labeled with H-3-labeled lofentanil. *Life Sci* 1982;31(5):457-462.
  65. Savoia G, Loreto M, Gravino E. Sufentanil: an overview of its use for acute pain management. *Minerva Anesthesiol* 2001;67(9 Suppl 1):206-216.
  66. Patel S, Spencer C. Remifentanil. *Drugs* 1996;52(3):417-427.
  67. Pan AC, Borhani DW, Dror RO, Shaw DE. Molecular determinants of drug-receptor binding kinetics. *Drug Discov Today* 2013;18(13-14):667-673.
  68. Kola I, Landis J. Can the pharmaceutical industry reduce attrition rates? *Nature Reviews Drug Discovery* 2004;3(8):711-715.
  69. Kola I. The state of innovation in drug development. *Clin Pharmacol Ther* 2008;83(2):227-230.
  70. Caulfield MP, Birdsall NJ. International Union of Pharmacology. XVII. Classification of muscarinic acetylcholine receptors. *Pharmacol Rev* 1998;50(2):279-290.
  71. Bonner TI, Buckley NJ, Young AC, Brann MR. Identification of a family of muscarinic acetylcholine receptor genes. *Science* 1987;237(4814):527-532.
  72. Bonner TI. The molecular basis of muscarinic receptor diversity. *Trends Neurosci* 1989;12(4):148-151.
  73. Wess J. Molecular biology of muscarinic acetylcholine receptors. *Crit Rev Neurobiol* 1996;10(1):69-99.
  74. Barnes PJ. The role of anticholinergics in chronic obstructive pulmonary disease. *Am J Med* 2004;117 Suppl 12A:24S-32S.
  75. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodriguez-Roisin R, van Weel C, Zielinski J. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176(6):532-555.
  76. Tashkin DP. Is a long-acting inhaled bronchodilator the first agent to use in stable chronic obstructive pulmonary disease? *Curr Opin Pulm Med* 2005;11(2):121-128.
  77. Disse B, Speck GA, Rominger KL, Witek TJ, Jr., Hammer R. Tiotropium (Spiriva): mechanistical considerations and clinical profile in obstructive lung disease. *Life Sci* 1999;64(6-7):457-464.
  78. Disse B, Reichl R, Speck G, Traunecker W, Rominger KL, Hammer R. Ba 679 Br, a Novel Long-Acting Anticholinergic Bronchodilator. *Life Sci* 1993;52(5-6):537-544.
  79. Haddad EB, Mak JC, Barnes PJ. Characterization of [<sup>3</sup>H]Ba 679 BR, a slowly dissociating muscarinic antagonist, in human lung: radioligand binding and autoradiographic mapping. *Mol Pharmacol* 1994;45(5):899-907.
  80. Casarosa P, Bouyssou T, Germeyer S, Schnapp A, Gantner F, Pieper M. Preclinical evaluation of long-acting muscarinic antagonists: comparison of tiotropium and investigational drugs. *J Pharmacol Exp Ther* 2009;330(2):660-668.
  81. Sykes DA, Dowling MR, Leighton-Davies J, Kent TC, Renard E, Trifilieff A, Charlton SJ. The influence of receptor kinetics on the onset and duration of action, and the therapeutic index of NVA237 and tiotropium. *J Pharmacol Exp Ther* 2012;343(2):520-528.
  82. D'Urzo A, Ferguson GT, van Noord JA, Hirata K, Martin C, Horton R, Lu YM, Banerji D, Overend T. Efficacy and safety of once-daily NVA237 in patients with moderate-to-severe COPD: the GLOW1 trial. *Resp Res* 2011;12: 156
  83. Tautermann CS, Kiechle T, Seeliger D, Diehl S, Wex E, Banholzer R, Gantner F, Pieper MP, Casarosa P. Molecular basis for the long duration of action and kinetic selectivity of Tiotropium for the muscarinic M<sub>3</sub> receptor. *J Med Chem* 2013; 56 (21): 8746-8756.

84. Kruse AC, Hu J, Pan AC, Arlow DH, Rosenbaum DM, Rosemond E, Green HF, Liu T, Chae PS, Dror RO, Shaw DE, Weis WI, Wess J, Kobilka BK. Structure and dynamics of the M<sub>3</sub> muscarinic acetylcholine receptor. *Nature* 2012;482(7386):552-556.
85. Smith DA, Jones BC, Walker DK. Design of drugs involving the concepts and theories of drug metabolism and pharmacokinetics. *Med Res Rev* 1996;16(3):243-266.
86. Tashkin DP, Celli B, Senn S, Burkhart D, Kesten S, Menjoge S, Decramer M. A 4-year trial of tiotropium in chronic obstructive pulmonary disease. *N Engl J Med* 2008;359(15):1543-1554.
87. van Noord JA, Smeets JJ, Custers FL, Korducki L, Cornelissen PJ. Pharmacodynamic steady state of tiotropium in patients with chronic obstructive pulmonary disease. *Eur Respir J* 2002;19(4):639-644.
88. Traynor K. Acclidinium bromide oral inhalation powder approved for COPD. *Am J Health Syst Pharm* 2012;69(17):1446.
89. Prat M, Fernandez D, Buil MA, Crespo MI, Casals G, Ferrer M, Tort L, Castro J, Monleon JM, Gavaldà A, Miralpeix M, Ramos I, Domenech T, Vilella D, Anton F, Huerta JM, Espinosa S, Lopez M, Sentellas S, Gonzalez M, Alberti J, Segarra V, Cardenas A, Beleta J, Ryder H. Discovery of novel quaternary ammonium derivatives of (3R)-quinuclidinol esters as potent and long-acting muscarinic antagonists with potential for minimal systemic exposure after inhaled administration: identification of (3R)-3-[[hydroxy(di-2-thienyl)acetyl]oxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane bromide (aclidinium bromide). *J Med Chem* 2009;52(16):5076-5092.
90. Kesten S, Jara M, Wentworth C, Lanes S. Pooled clinical trial analysis of tiotropium safety. *Chest* 2006;130(6):1695-1703.
91. Suppli Ulrik C. Acclidinium Bromide: Clinical benefit in patients with moderate to severe COPD. *Open Respir Med J* 2012;6:150-154.
92. Singh D, Magnussen H, Kirsten A, Mindt S, Caracta C, Seoane B, Jarreta D, Garcia Gil E. A randomised, placebo- and active-controlled dose-finding study of aclidinium bromide administered twice a day in COPD patients. *Pulm Pharmacol Ther* 2012;25(3):248-253.
93. Maltais F, Milot J. The potential for aclidinium bromide, a new anticholinergic, in the management of chronic obstructive pulmonary disease. *Ther Adv Respir Dis* 2012;6(6):345-361.
94. Gupta V, Singh D. Acclidinium bromide for the treatment of chronic obstructive pulmonary disease. *Expert Rev Respir Med* 2012;6(6):581-588.
95. Jones PW, Singh D, Bateman ED, Agusti A, Lamarca R, de Miquel G, Segarra R, Caracta C, Garcia Gil E. Efficacy and safety of twice-daily aclidinium bromide in COPD patients: the ATTAIN study. *Eur Respir J* 2012;40(4):830-836.
96. Glossop PA, Watson CA, Price DA, Bunnage ME, Middleton DS, Wood A, James K, Roberts D, Strang RS, Yeadon M, Perros-Huguet C, Clarke NP, Trevethick MA, Machin I, Stuart EF, Evans SM, Harrison AC, Fairman DA, Agoram B, Burrows JL, Feeder N, Fulton CK, Dillon BR, Entwistle DA, Spence FJ. Inhalation by design: novel tertiary amine muscarinic M<sub>3</sub> receptor antagonists with slow off-rate binding kinetics for inhaled once-daily treatment of chronic obstructive pulmonary disease. *J Med Chem* 2011;54(19):6888-6904.
97. Haddad EB, Patel H, Keeling JE, Yacoub MH, Barnes PJ, Belvisi MG. Pharmacological characterization of the muscarinic receptor antagonist, glycopyrrolate, in human and guinea-pig airways. *Br J Pharmacol* 1999;127(2):413-420.
98. Sykes DA, Dowling MR, Charlton SJ. Exploring the mechanism of agonist efficacy: a relationship between efficacy and agonist dissociation rate at the muscarinic M<sub>3</sub> receptor. *Mol Pharmacol* 2009;76(3):543-551.

99. Hoeren M, Brawek B, Mantovani M, Löffler M, Steffens M, van Velthoven V, Feuerstein TJ. Partial agonism at the human  $\alpha_{2A}$ -autoreceptor: role of binding duration. *Naunyn Schmiedebergs Arch Pharmacol* 2008;378(1):17-26.
100. Pantaleo N, Chadwick W, Park SS, Wang L, Zhou Y, Martin B, Maudsley S. The mammalian tachykinin ligand-receptor system: an emerging target for central neurological disorders. *CNS Neurol Disord Drug Targets* 2010;9(5):627-635.
101. Pennefather JN, Lecci A, Candenas ML, Patak E, Pinto FM, Maggi CA. Tachykinins and tachykinin receptors: a growing family. *Life Sci* 2004;74(12):1445-1463.
102. Almeida TA, Rojo J, Nieto PM, Pinto FM, Hernandez M, Martin JD, Candenas ML. Tachykinins and tachykinin receptors: structure and activity relationships. *Curr Med Chem* 2004;11(15):2045-2081.
103. Simonsen KB, Juhl K, Steiniger-Brach B, Nielsen SM. Novel NK<sub>3</sub> receptor antagonists for the treatment of schizophrenia and other CNS indications. *Curr Opin Drug Discov Devel* 2010;13(4):379-388.
104. Stout SC, Owens MJ, Nemeroff CB. Neurokinin(1) receptor antagonists as potential antidepressants. *Annu Rev Pharmacol* 2001;41:877-906.
105. Albert JS. Neurokinin antagonists and their potential role in treating depression and other stress disorders. *Exp Opin Inv Drug* 2004;14(10):1421-1433.
106. Lecci A, Maggi CA. Peripheral tachykinin receptors as potential therapeutic targets in visceral diseases. *Exp Opin Ther Targets* 2003;7(3):343-362.
107. Quartara L, Altamura M, Evangelista S, Maggi CA. Tachykinin receptor antagonists in clinical trials. *Exp Opin Inv Drug* 2009;18(12):1843-1864.
108. Lindstrom E, von Mentzer B, Pahlman I, Ahlstedt I, Uvebrant A, Kristensson E, Martinsson R, Noven A, de Verdier J, Vauquelin G. Neurokinin 1 receptor antagonists: correlation between *in vitro* receptor interaction and *in vivo* efficacy. *J Pharmacol Exp Ther* 2007;322(3):1286-1293.
109. Duffy RA, Varty GB, Morgan CA, Lachowicz JE. Correlation of neurokinin (NK) 1 receptor occupancy in gerbil striatum with behavioral effects of NK<sub>1</sub> antagonists. *J Pharmacol Exp Ther* 2002;301(2):536-542.
110. Malherbe P, Knoflach F, Hernandez MC, Hoffmann T, Schnider P, Porter RH, Wettstein JG, Ballard TM, Spooen W, Steward L. Characterization of RO4583298 as a novel potent, dual antagonist with *in vivo* activity at tachykinin NK<sub>1</sub> and NK<sub>3</sub> receptors. *Br J Pharmacol* 2011;162(4):929-946.
111. Cascieri MA, Ber E, Fong TM, Hale JJ, Tang F, Shiao LL, Mills SG, MacCoss M, Sadowski S, Tota MR, Strader CD. Characterization of the binding and activity of a high affinity, pseudoirreversible morpholino tachykinin NK<sub>1</sub> receptor antagonist. *Eur J Pharmacol* 1997;325(2-3):253-261.
112. Cascieri MA, Ber E, Fong TM, Sadowski S, Bansal A, Swain C, Seward E, Frances B, Burns D, Strader CD. Characterization of the binding of a potent, selective, radioiodinated antagonist to the human neurokinin-1 receptor. *Mol Pharmacol* 1992;42(3):458-463.
113. Bristow LJ, Young L. Chromodacryorrhea and repetitive hind paw tapping: models of peripheral and central tachykinin NK<sub>1</sub> receptor activation in gerbils. *Eur J Pharmacol* 1994;253(3):245-252.
114. Young JR, Eid R, Turner C, DeVita RJ, Kurtz MM, Tsao KL, Chicchi GG, Wheeldon A, Carlson E, Mills SG. Pyrrolidine-carboxamides and oxadiazoles as potent hNK<sub>1</sub> antagonists. *Bioorg Med Chem Lett* 2007;17(19):5310-5315.
115. Morriello GJ, Chicchi G, Johnson T, Mills SG, Demartino J, Kurtz M, Tsao KL, Zheng S, Tong X, Carlson E, Townson K, Wheeldon A, Boyce S, Collinson N, Rupniak N, Devita RJ. Fused tricyclic pyrrolizinones that exhibit pseudo-irreversible blockade of the NK<sub>1</sub> receptor. *Bioorg Med Chem Lett* 2010;20(19):5925-5932.

116. Morriello GJ, DeVita RJ, Mills SG, Young JR, Lin P, Doss G, Chicchi GG, DeMartino J, Kurtz MM, Tsao K-LC, Carlson E, Townson K, Wheeldon A, Boyce S, Collinson N, Rupniak N, Moore S. Fused bicyclic pyrrolizinones as new scaffolds for human NK<sub>1</sub> antagonists. *Bioorg Med Chem* 2008;16(5):2156-2170.
117. Morriello GJ, Mills SG, Johnson T, Reibarkh M, Chicchi G, DeMartino J, Kurtz M, Davies P, Tsao KLC, Zheng S, Tong X, Carlson E, Townson K, Tattersall FD, Wheeldon A, Boyce S, Collinson N, Rupniak N, Moore S, DeVita RJ. Substituted fused bicyclic pyrrolizinones as potent, orally bioavailable hNK<sub>1</sub> antagonists. *Bioorg Med Chem Lett* 2010;20(6):2007-2012.
118. Morriello GJ, Chicchi G, Johnson T, Mills SG, Demartino J, Kurtz M, Tsao KL, Zheng S, Tong X, Carlson E, Townson K, Wheeldon A, Boyce S, Collinson N, Rupniak N, Devita RJ. Fused tricyclic pyrrolizinones that exhibit pseudo-irreversible blockade of the NK<sub>1</sub> receptor. *Bioorg Med Chem Lett* 2010;20(19):5925-5932.
119. Di Fabio R, Griffante C, Alvaro G, Pentassuglia G, Pizzi DA, Donati D, Rossi T, Guercio G, Mattioli M, Cimarosti Z, Marchioro C, Provera S, Zonzini L, Montanari D, Melotto S, Gerrard PA, Trist DG, Ratti E, Corsi M. Discovery process and pharmacological characterization of 2-(S)-(4-fluoro-2-methylphenyl)piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethylphenyl)ethyl]methylamide (vestipitant) as a potent, selective, and orally active NK<sub>1</sub> receptor antagonist. *J Med Chem* 2009;52(10):3238-3247.
120. Di Fabio R, Alvaro G, Griffante C, Pizzi DA, Donati D, Mattioli M, Cimarosti Z, Guercio G, Marchioro C, Provera S, Zonzini L, Montanari D, Melotto S, Gerrard PA, Trist DG, Ratti E, Corsi M. Discovery and biological characterization of (2R,4S)-1'-acetyl-N-((1R)-1-[3,5-bis(trifluoromethyl)phenyl]ethyl)-2-(4-fluoro-2-methylphenyl)-N-methyl-4,4'-bipiperidine-1-carboxamide as a new potent and selective neurokinin 1 (NK<sub>1</sub>) receptor antagonist clinical candidate. *J Med Chem* 2011;54(4):1071-1079.
121. Cirillo R, Astolfi M, Conte B, Lopez G, Parlani M, Terracciano R, Fincham CI, Manzini S. Pharmacology of the peptidomimetic, MEN 11149, a new potent, selective and orally effective tachykinin NK<sub>1</sub> receptor antagonist. *Eur J Pharmacol* 1998;341(2-3):201-209.
122. Carlsson A, Lindqvist M, Magnusson T. 3,4-Dihydroxyphenylalanine and 5-Hydroxytryptophan as Reserpine Antagonists. *Nature* 1957;180(4596):1200-1200.
123. Le Crom S, Kapsimali M, Barome PO, Vernier P. Dopamine receptors for every species: gene duplications and functional diversification in Craniates. *J Struct Funct Genomics* 2003;3(1-4):161-176.
124. Carlsson A, Civelli O, Keabian J, Langer S, Scatton B, Schwartz JC, Sedvall G, Seeman P, Sokoloff P, Spano P, Van Tol H. Dopamine receptors. Last modified on 27/05/2013. Accessed on 29/08/2013. *IUPHAR database (IUPHAR-DB)*, <http://www.iuphar-db.org/DATABASE/FamilyMenuForward?familyId=20>.
125. Hurley MJ, Jenner P. What has been learnt from study of dopamine receptors in Parkinson's disease? *Pharmacol Therapeut* 2006;111(3):715-728.
126. Schneier FR, Liebowitz MR, Abi-Dargham A, Zea-Ponce Y, Lin SH, Laruelle M. Low dopamine D<sub>2</sub> receptor binding potential in social phobia. *Am J Psychiatry* 2000;157(3):457-459.
127. Kienast T, Heinz A. Dopamine and the diseased brain. *CNS Neurol Disord Drug Targets* 2006;5(1):109-131.
128. Fuxe K, Manger P, Genedani S, Agnati L. in: Riederer P, Reichmann H, Youdim MBH, Gerlach M. (Eds) *Parkinson's Diseases and Related Disorders*. Vienna: Springer; 2006. p 71-83.
129. Mihara K, Kondo T, Suzuki A, Yasui-Furukori N, Ono S, Sano A, Koshiro K, Otani K, Kaneko S. Relationship between functional dopamine D<sub>2</sub> and D<sub>3</sub> receptors gene polymorphisms and

- neuroleptic malignant syndrome. *Am J Med Genet B* 2003;117B(1):57-60.
130. Faraone SV, Khan SA. Candidate gene studies of attention-deficit/hyperactivity disorder. *J Clin Psychiat* 2006;67:13-20.
  131. Hummel M, Unterwald EM. D<sub>1</sub> dopamine receptor: a putative neurochemical and behavioral link to cocaine action. *J Cell Physiol* 2002;191(1):17-27.
  132. Seeman P, Weinshenker D, Quirion R, Srivastava LK, Bhardwaj SK, Grandy DK, Premont RT, Sotnikova TD, Boksa P, El-Ghundi M, O'Dowd B F, George SR, Perreault ML, Mannisto PT, Robinson S, Palmiter RD, Tallero T. Dopamine supersensitivity correlates with D<sub>2</sub>High states, implying many paths to psychosis. *Proc Natl Acad Sci U S A* 2005;102(9):3513-3518.
  133. Miyamoto S, Duncan GE, Marx CE, Lieberman JA. Treatments for schizophrenia: a critical review of pharmacology and mechanisms of action of antipsychotic drugs. *Mol Psychiatry* 2005;10(1):79-104.
  134. Kapur S, Seeman P. Antipsychotic agents differ in how fast they come off the dopamine D<sub>2</sub> receptors. Implications for atypical antipsychotic action. *J Psychiatr Neurosci* 2000;25(2):161-166.
  135. Langlois X, Megens A, Lavreysen H, Atack J, Cik M, te Riele P, Peeters L, Wouters R, Vermeire J, Hendrickx H, Macdonald G, De Bruyn M. Pharmacology of JNJ-37822681, a specific and fast-dissociating D<sub>2</sub> antagonist for the treatment of schizophrenia. *J Pharmacol Exp Ther* 2012;342(1):91-105.
  136. Schultz W. Multiple dopamine functions at different time courses. *Annu Rev Neurosci* 2007;30:259-288.
  137. Bartholi G, Haefely W, Jalfre M, Keller HH, Pletsche A. Effects of Clozapine on Cerebral Catecholaminergic Neuron Systems. *Br J Pharmacol* 1972;46(4):736-740.
  138. Jones C, Kapur S, Remington G, Zipursky RB. Transient D<sub>2</sub> dopamine receptor occupancy in low EPS-incidence drugs: PET evidence. *Biol Psychiat* 2000;47(8):112s-112s.
  139. Carboni L, Negri M, Michielin F, Bertani S, Fratte SD, Oliosi B, Cavanni P. Slow dissociation of partial agonists from the D<sub>2</sub> receptor is linked to reduced prolactin release. *Int J Neuropsychopharmacol* 2012;15(5):645-656.
  140. Kapur S, Seeman P. Does fast dissociation from the dopamine D<sub>2</sub> receptor explain the action of atypical antipsychotics?: A new hypothesis. *Am J Psychiat* 2001;158(3):360-369.
  141. Schmidt ME, de Boer P, Andrews R, Neyens M, Rossenu S, William Falteos D, Mannaert E. D<sub>2</sub> receptor occupancy measurement of JNJ-37822681, a novel fast off-rate D<sub>2</sub> receptor antagonist, in healthy subjects using positron emission tomography: single dose versus steady state and dose selection. *Psychopharmacology (Berl)*. 2012;224(4):549-557.
  142. Schmidt ME, Kent JM, Daly E, Janssens L, Van Osselaer N, Husken G, Angelescu IG, Van Nueten L. A double-blind, randomized, placebo-controlled study with JNJ-37822681, a novel, highly selective, fast dissociating D<sub>2</sub> receptor antagonist in the treatment of acute exacerbation of schizophrenia. *Eur Neuropsychopharmacol* 2012;22(10):721-733.
  143. Miller DC, Lunn G, Jones P, Sabnis Y, Davies NL, Driscoll P. Investigation of the effect of molecular properties on the binding kinetics of a ligand to its biological target. *MedChemComm* 2012;3(4):449-452.
  144. Vale W, Spiess J, Rivier C, Rivier J. Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin. *Science* 1981;213(4514):1394-1397.
  145. Dautzenberg FM, Hauger RL. The CRF peptide family and their receptors: yet more partners discovered. *Trends Pharmacol Sci* 2002;23(2):71-77.
  146. Bale TL, Vale WW. CRF and CRF receptors: Role in stress responsivity and other behaviors.

- Annu Rev Pharmacol* 2004;44:525-557.
147. Schulz DW, Mansbach RS, Sprouse J, Braselton JP, Collins J, Corman M, Dunaiskis A, Faraci S, Schmidt AW, Seeger T, Seymour P, Tingley FD, Winston EN, Chen YL, Heym J. CP-154,526: A potent and selective nonpeptide antagonist of corticotropin releasing factor receptors. *Proc Natl Acad Sci U S A* 1996;93(19):10477-10482.
  148. Grigoriadis DE. The corticotropin-releasing factor receptor: a novel target for the treatment of depression and anxiety-related disorders. *Expert Opin Ther Tar* 2005;9(4):651-684.
  149. Holsboer F. The corticosteroid receptor hypothesis of depression. *Neuropsychopharmacol* 2000;23(5):477-501.
  150. Chen C. Recent advances in small molecule antagonists of the corticotropin-releasing factor type-1 receptor-focus on pharmacology and pharmacokinetics. *Curr Med Chem* 2006;13(11):1261-1282.
  151. Fleck BA, Hoare SR, Pick RR, Bradbury MJ, Grigoriadis DE. Binding kinetics redefine the antagonist pharmacology of the corticotropin-releasing factor type 1 receptor. *J Pharmacol Exp Ther* 2012;341(2):518-531.
  152. Miller DC, Klute W, Brown AD. Discovery of potent, metabolically stable purine CRF-1 antagonists with differentiated binding kinetic profiles. *Bioorg Med Chem Lett* 2011;21(20):6108-6111.
  153. Ramsey SJ, Attkins NJ, Fish R, van der Graaf PH. Quantitative pharmacological analysis of antagonist binding kinetics at CRF<sub>1</sub> receptors *in vitro* and *in vivo*. *Br J Pharmacol* 2011;164(3):992-1007.
  154. Gilligan PJ, Robertson DW, Zaczek R. Corticotropin releasing factor (CRF) receptor modulators: Progress and opportunities for new therapeutic agents. *J Med Chem* 2000;43(9):1641-1660.
  155. Kehne J, De Lombaert S. Non-peptidic CRF<sub>1</sub> receptor antagonists for the treatment of anxiety, depression and stress disorders. *Curr Drug Targets CNS Neurol Disord* 2002;1(5):467-493.
  156. Hollenstein K, Kean J, Bortolato A, Cheng RKY, Dore AS, Jazayeri A, Cooke RM, Weir M, Marshall FH. Structure of class B GPCR corticotropin-releasing factor receptor 1. *Nature* 2013;499(7459):438-443.
  157. Arvanitis AG, Gilligan PJ, Chorvat RJ, Cheeseman RS, Christos TE, Bakthavatchalam R, Beck JP, Cocuzza AJ, Hobbs FW, Wilde RG, Arnold C, Chidester D, Curry M, He L, Hollis A, Klaczkiewicz J, Krenitsky PJ, Rescinito JP, Scholfield E, Culp S, De Souza EB, Fitzgerald L, Grigoriadis D, Tam SW, Wong YN, Huang S, Shen HL. Non-peptide corticotropin-releasing hormone antagonists: syntheses and structure-activity relationships of 2-anilinoypyrimidines and -triazines. *J Med Chem* 1999;42(5):805-818.
  158. Hodge CN, Aldrich PE, Wasserman ZR, Fernandez CH, Nemeth GA, Arvanitis A, Cheeseman RS, Chorvat RJ, Ciganek E, Christos TE, Gilligan PJ, Krenitsky P, Scholfield E, Strucely P. Corticotropin-releasing hormone receptor antagonists: framework design and synthesis guided by ligand conformational studies. *J Med Chem* 1999;42(5):819-832.
  159. Hann MM. Molecular obesity, potency and other addictions in drug discovery. *MedChemComm* 2011;2(5):349-355.
  160. Decourcelles DC, Leysen JE, Roevens P, Vanbelle H. The Serotonin-5<sub>2</sub> receptor: a receptor-transducer coupling model to explain insurmountable antagonist effects. *Drug Develop Res* 1986;8(1-4):173-178.
  161. Smith C, Rahman T, Toohey N, Mazurkiewicz J, Herrick-Davis K, Teitler M. Risperidone irreversibly binds to and inactivates the h5-HT<sub>7</sub> serotonin receptor. *Mol Pharmacol* 2006;70(4):1264-1270.



162. Hosohata Y, Sasaki K, Shen Y, Hattori K, Suzuki J, Nagatomo T. Bopindolol is a slowly dissociating  $\beta_1$ -adrenoceptor antagonist when compared to other beta-blockers. *Biol Pharm Bull* 1995;18(8):1066-1071.
163. Casarosa P, Kollak I, Kiechle T, Ostermann A, Schnapp A, Kiesling R, Pieper M, Sieger P, Gantner F. Functional and biochemical rationales for the 24-hour long duration of action of olodaterol. *J Pharmacol Exp Ther* 2011;337(3):600-609.
164. Leysen JE, Gommeren W. Drug-receptor dissociation time, new tool for drug research: receptor binding affinity and drug-receptor dissociation profiles of Serotonin- $S_2$ , Dopamine- $D_2$ , Histamine- $H_1$  Antagonists, and Opiates. *Drug Develop Res* 1986;8(1-4):119-131.
165. Watson C, Jenkinson S, Kazmierski W, Kenakin T. The CCR5 receptor-based mechanism of action of 873140, a potent allosteric noncompetitive HIV entry inhibitor. *Mol Pharmacol* 2005;67(4):1268-1282.
166. Gonsiorek W, Fan X, Hesk D, Fossetta J, Qiu H, Jakway J, Billah M, Dwyer M, Chao J, Deno G, Taveras A, Lundell DJ, Hipkin RW. Pharmacological characterization of Sch527123, a potent allosteric CXCR1/CXCR2 antagonist. *J Pharmacol Exp Ther* 2007;322(2):477-485.
167. Stock N, Volkots D, Stebbins K, Broadhead A, Stearns B, Roppe J, Parr T, Baccei C, Bain G, Chapman C, Correa L, Darlington J, King C, Lee C, Lorrain DS, Prodanovich P, Santini A, Evans JF, Hutchinson JH, Prasit P. Sodium [2'-[(cyclopropanecarbonyl-ethyl-amino)-methyl]-4'-(6-ethoxy-pyridin-3-yl)-6-methoxy-biphenyl-3-yl]-acetate (AM432): a potent, selective prostaglandin  $D_2$  receptor antagonist. *Bioorg Med Chem Lett* 2011;21(3):1036-1040.
168. Sullivan SK, Hoare SRJ, Fleck BA, Zhu YF, Heise CE, Struthers RS, Crowe PD. Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure-activity relationships and insurmountable antagonism. *Biochem Pharmacol* 2006;72(7):838-849.
169. Anthes JC, Gilchrest H, Richard C, Eckel S, Hesk D, West RE, Jr., Williams SM, Greenfeder S, Billah M, Kreutner W, Egan RE. Biochemical characterization of desloratadine, a potent antagonist of the human histamine  $H_1$  receptor. *Eur J Pharmacol* 2002;449(3):229-237.
170. Smits RA, Lim HD, van der Meer T, Kuhne S, Bessembinder K, Zuiderveld OP, Wijtmans M, de Esch IJ, Leurs R. Ligand based design of novel histamine  $H_4$  receptor antagonists; fragment optimization and analysis of binding kinetics. *Bioorg Med Chem Lett* 2012;22(1):461-467.
171. Meini S, Bellucci F, Catalani C, Cucchi P, Giolitti A, Santicoli P, Giuliani S. Multifaceted approach to determine the antagonist molecular mechanism and interaction of ibodutant ([1-(2-phenyl-1R-[[1-(tetrahydropyran-4-ylmethyl)-piperidin-4-ylmethyl]-carbamoyl]-ethylcarbamoyl)-cyclopentyl]-amide) at the human tachykinin  $NK_2$  receptor. *J Pharmacol Exp Ther* 2009;329(2):486-495.
172. Malherbe P, Kratzeisen C, Marcuz A, Zenner MT, Nettekoven MH, Ratni H, Wettstein JG, Bissantz C. Identification of a critical residue in the transmembrane domain 2 of tachykinin neurokinin 3 receptor affecting the dissociation kinetics and antagonism mode of osanetant (SR 142801) and piperidine-based structures. *J Med Chem* 2009;52(22):7103-7112.
173. Mould R, Brown J, Marshall FH, Langmead CJ. Binding kinetics differentiates functional antagonism of orexin-2 receptor ligands. *Br J Pharmacol* 2014;171(2):351-363.
174. Ring PC, Seldon PM, Barnes PJ, Giembycz MA. Pharmacological characterization of a receptor for platelet-activating factor on guinea pig peritoneal macrophages using [ $^3H$ ]apafant, a selective and competitive platelet-activating factor antagonist: evidence that the noncompetitive behavior of apafant in functional studies relates to slow kinetics of dissociation. *Mol Pharmacol* 1993;43(2):302-312.



175. Behm DJ, Aiyar NV, Olzinski AR, McAtee JJ, Hilfiker MA, Dodson JW, Dowdell SE, Wang GZ, Goodman KB, Sehon CA, Harpel MR, Willette RN, Neeb MJ, Leach CA, Douglas SA. GSK1562590, a slowly dissociating urotensin-II receptor antagonist, exhibits prolonged pharmacodynamic activity *ex vivo*. *Br J Pharmacol* 2010;161(1):207-228.
176. Mullins D, Adham N, Hesk D, Wu Y, Kelly J, Huang Y, Guzzi M, Zhang X, McCombie S, Stamford A, Parker E. Identification and characterization of pseudoirreversible nonpeptide antagonists of the neuropeptide Y Y<sub>5</sub> receptor and development of a novel Y<sub>5</sub>-selective radioligand. *Eur J Pharmacol* 2008;601(1-3):1-7.
177. Aronne LJ, Tonstad S, Moreno M, Gantz I, Erondur N, Suryawanshi S, Molony C, Sieberts S, Nayee J, Meehan AG, Shapiro D, Heymsfield SB, Kaufman KD, Amatruda JM. A clinical trial assessing the safety and efficacy of taranabant, a CB<sub>1</sub>R inverse agonist, in obese and overweight patients: a high-dose study. *Int J Obes (Lond)* 2010;34(5):919-935.
178. Kipnes MS, Hollander P, Fujioka K, Gantz I, Seck T, Erondur N, Shentu Y, Lu K, Suryawanshi S, Chou M, Johnson-Levonas AO, Heymsfield SB, Shapiro D, Kaufman KD, Amatruda JM. A one-year study to assess the safety and efficacy of the CB<sub>1</sub>R inverse agonist taranabant in overweight and obese patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12(6):517-531.
179. <http://clinicaltrials.gov/>
180. Meltzer HY. Multireceptor atypical antipsychotic drugs. In: Ellenbroek BA, Cools AR, editors. *Atypical Antipsychotics*. Basel: Birkhauser Verlag; 2000. p191-213.
181. Huirne JA, Lambalk CB. Gonadotropin-releasing-hormone-receptor antagonists. *Lancet* 2001;358(9295):1793-1803.
182. Keith DE, Anton B, Murray SR, Zaki PA, Chu PC, Lissin DV, Monteillet-Agius G, Stewart PL, Evans CJ, von Zastrow M. mu-opioid receptor internalization: Opiate drugs have differential effects on a conserved endocytic mechanism *in vitro* and in the mammalian brain. *Mol Pharmacol* 1998;53(3):377-384.

