



Universiteit
Leiden
The Netherlands

The role of apolipoprotein CI in lipid metabolism and bacterial sepsis

Berbée, J.F.P.

Citation

Berbée, J. F. P. (2007, May 24). *The role of apolipoprotein CI in lipid metabolism and bacterial sepsis*. Retrieved from <https://hdl.handle.net/1887/11973>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/11973>

Note: To cite this publication please use the final published version (if applicable).

Chapter 3

Apolipoprotein CI Causes Hypertriglyceridemia Independent of the Very-low-density Lipoprotein Receptor and Apolipoprotein CIII in Mice

Caroline C. van der Hoogt^{1,2}, Jimmy F.P. Berbée^{1,2}, Sonia M.S. Espirito Santo^{1,2}, Gery Gerritsen³, Yvonne D. Krom³, André van der Zee³, Louis M. Havekes^{1,2,4}, Ko Willems van Dijk^{2,3}, Patrick C.N. Rensen^{1,2}

From the ¹Department of Biomedical Research, TNO-Quality of Life, Gaubius Laboratory, P.O. Box 2215, 2301 CE Leiden, The Netherlands; Departments of ²General Internal Medicine, Endocrinology and Metabolic Diseases, ³Human Genetics, and ⁴Cardiology, Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands.

Biochim Biophys Acta 2006, 1761 (2): 213-220.

Abstract

Objective: We have recently shown that the predominant hypertriglyceridemia in human apolipoprotein C1 (*APOC1*) transgenic mice is mainly explained by apoCI-mediated inhibition of the lipoprotein lipase (LPL)-dependent triglyceride (TG)-hydrolysis pathway. Since the very-low-density lipoprotein receptor (VLDLr) and apoCIII are potent modifiers of LPL activity, our current aim was to study whether the lipolysis-inhibiting action of apoCI would be dependent on the presence of the VLDLr and apoCIII *in vivo*.

Methods and Results: Hereto, we employed liver-specific expression of human apoCI by using a novel recombinant adenovirus (Ad*APOC1*). In wild-type mice, moderate apoCI expression leading to plasma human apoCI levels of 12-33 mg/dL dose-dependently and specifically increased plasma TG (up to 6.6-fold, $P < 0.001$), yielding the same hypertriglyceridemic phenotype as observed in human *APOC1* transgenic mice. Ad*APOC1* still increased plasma TG in *vldlr*^{-/-} mice (4.1-fold, $P < 0.001$) and in *apoc3*^{-/-} mice (6.8-fold, $P < 0.001$) that were also deficient for the low-density lipoprotein receptor (LDLr) and LDLr-related protein (LRP) or apoE, respectively.

Conclusions: Thus, irrespective of receptor-mediated remnant clearance by the liver, liver-specific expression of human apoCI causes hypertriglyceridemia in the absence of the VLDLr and apoCIII. We conclude that apoCI is a powerful and direct inhibitor of LPL activity independent of the VLDLr and apoCIII.

Introduction

Human apolipoprotein CI (apoCI), encoded by the *APOC1* gene, is primarily expressed in the liver and secreted into the circulation¹⁻⁴. To study the function of apoCI, mice have been generated that express human apoCI⁵⁻⁷. These mice have a hyperlipidemic phenotype, with the most pronounced effect on triglycerides (TG) in VLDL. Initially, it has been suggested that apoCI exerts its hyperlipidemic effect by interfering with the apoE-mediated hepatic remnant clearance via the low-density lipoprotein (LDL) receptor (LDLr)⁸ and LDLr-related protein (LRP)⁹. However, *APOC1* mice deficient for apoE still show severe hypertriglyceridemia^{10,11}, indicating that apoCI predominantly affects a lipid clearance route other than receptor recognition. Indeed, we and others showed that apoCI directly and dose-dependently inhibits lipoprotein lipase (LPL)-mediated TG hydrolysis *in vitro* with a 60% efficiency compared with the main endogenous LPL inhibitor apoCIII^{10,12}, and we have recently demonstrated that the apoCI-induced hypertriglyceridemia is mainly explained by impaired LPL-mediated lipolytic conversion of TG-rich lipoproteins¹⁰.

The VLDL receptor (VLDLr) is required for normal LPL regulation *in vivo* as disruption of the VLDLr results in hypertriglyceridemia associated with reduced LPL activity^{13,14}. Therefore, we wondered whether the effect of apoC1 on LPL is dependent on the VLDLr. Both *vldlr*^{-/-} and *APOC1* mice are protected from diet-induced obesity on a wild-type (WT) as well as an *ob/ob* background^{13,15} because of defective TG hydrolysis leading to reduced delivery of VLDL-derived FFA into adipose tissue. Since enrichment of VLDL with apoC1 inhibits its binding to the VLDLr *in vitro*¹⁶, this mechanism may be responsible for the TG-raising effect of apoC1 *in vivo*. Likewise, LPL activity is also highly dependent on the presence of the main endogenous LPL inhibitor apoCIII^{17,18}, but a potential interaction between apoC1 and apoCIII in lipolysis is unknown. We have observed by direct comparison that apoCIII is a more potent LPL inhibitor than apoC1 *in vitro*¹⁰, but the effect of apoC1 independent of apoCIII has not yet been assessed *in vivo*.

Taken together, the hypertriglyceridemic effect of apoC1 is mainly explained by the inhibition of LPL-mediated TG clearance. However, it is unknown whether this effect results from direct inhibition of LPL or indirectly via interactions with the VLDLr and/or apoCIII. Therefore, the aim of this study was to investigate whether apoC1 inhibits LPL-mediated lipolytic conversion *in vivo* through interaction with the VLDLr or apoCIII. Hereto, we expressed human apoC1 using an adenovirus in mice that lack the VLDLr or apoCIII. Our results show that apoC1 is a powerful inhibitor of LPL activity *in vivo* independent of the VLDLr and apoCIII.

Materials and methods

Animals – MX1Cre⁺*Irfp*^{lox/lox}*ldlr*^{-/-}*vldlr*^{-/-}¹⁹, *apoe*^{-/-}, *apoe*^{-/-}*apoc3*^{-/-}¹⁸, and WT mice (C57Bl/6 background), 3–4 months of age, were used in the experiments. Mice were obtained from our breeding colonies at the Institutional Animal Facility and housed under standard conditions in conventional cages and were fed regular chow *ad libitum*. LRP deficiency was induced in MX1Cre⁺*Irfp*^{lox/lox}*ldlr*^{-/-}*vldlr*^{-/-} mice by intraperitoneal injection of polyinosinic:polycytidylic ribonucleic acid (pI: pC, Sigma, St Louis, MO, U.S.A.), which results in the complete absence of LRP protein in liver membrane extracts¹⁹. Experiments were performed after 4 h of fasting at 12:00 pm with food withdrawn at 8:00 am. Female mice were used unless indicated otherwise. All experiments have been approved by the Institutional Ethical Committee on Animal Care and Experimentation.

Construction of Recombinant Adenoviral Vector Expressing Human ApoC1 –A recombinant, replication-deficient adenoviral vector expressing human *APOC1* (Ad*APOC1*) under control of a cytomegalovirus promoter was constructed by the method of He *et al.*²⁰. Briefly, the N-terminal *KpnI*-*HindIII* (2140 bp) fragment of

human *APOC1* genomic DNA was transposed into the corresponding sites of the pAdTrack-CMV vector. Subsequently, the C-terminal *HindIII-HindIII* (2403 bp) fragment of human genomic *APOC1* DNA was cloned into the resulting plasmid. The identity of the resulting construct was verified by sequence analysis, and homologous recombination with the adenoviral backbone vector pAdEasy-1 took place in BJ5183 cells. Recombinant plasmids were transfected into the adenovirus packaging cell line PER.C6²¹ and amplified. *E. coli* strain BJ5183 and the pAdTrack-CMV and pAdEasy-1 vectors were a kind gift of Dr. Vogelstein (Johns Hopkins School of Medicine, Baltimore, MD, U.S.A.). Recombinant adenovirus was purified twice via caesium chloride gradient centrifugation and dialyzed against 25 mM Tris, 137 mM NaCl, 5 mM KCl, 0.73 mM NaH₂PO₄, 0.9 mM CaCl₂, and 0.5 mM MgCl₂, pH 7.45, followed by dialysis against the same buffer supplemented with sucrose (50 g/L). For storage, aliquots of 150 µL virus were frozen at -80°C. Routinely, virus titers of the stocks varied from 1x10¹⁰ to 1x10¹¹ pfu (plaque forming units)/mL.

Administration of Adenoviral Vectors to Mice – At least 3 days before adenovirus injection into mice, basal serum lipid values were measured (t=0). At day 0, mice were injected into the tail vein with either Ad*APOC1* or a recombinant virus expressing β-galactosidase (AdLacZ) as a control, both diluted with PBS to a total volume of 200 µL. To prevent sequestration of low doses of virus by Kupffer cells and to achieve a more linear dose-response relationship of *APOC1* expression, mice were pre-injected with 0.5x10⁹ pfu AdLacZ at 3 h before injection²².

Plasma Lipid and Lipoprotein Analysis – In all experiments blood was collected from the tail vein into chilled paraoxon (Sigma, St. Louis, MO, U.S.A.)-coated capillary tubes to prevent ongoing *in vitro* lipolysis²³. The tubes were placed on ice, centrifuged at 4°C, and the obtained plasma was assayed for total cholesterol (TC), TG, and free fatty acids (FFA) using the commercially available enzymatic kits 236691, 11488872 (Roche Molecular Biochemicals, Indianapolis, IN, U.S.A.), and NEFA-C (Wako Chemicals, Neuss, Germany), respectively. For determination of the plasma lipoprotein distribution by FPLC, 50 µL of pooled plasma per group was injected onto a Superose 6 column (Äkta System; Amersham Pharmacia Biotech, Piscataway, NJ, U.S.A.), and eluted at a flow rate of 50 µL/min with PBS, 1 mM EDTA, pH 7.4. Fractions of 50 µL were collected and assayed for TC and TG as described above. Human apoC1 was quantified by ELISA exactly as described¹⁰.

Hepatic VLDL-TG Production – Mice were fasted for 4 h and anesthetized by i.p. administration of 6.2 mg/kg of Acepromazine (Pfizer Animal Health, Capelle a/d IJssel, The Netherlands), 0.3 mg/kg of Fentanyl Bipharm (Pharma Hameln, Hameln, Germany), and 6.2 mg/kg of Midazolam (Roche, Mijdrecht, The Netherlands). Subsequently, mice were injected via the tail vein with Triton WR-1339 (500 mg/kg of body weight) to block TG-hydrolysis and hepatic lipoprotein uptake²⁴. Blood samples were drawn at 1, 30, 60, 90, and 120 min after administration, and plasma TG levels were measured as described above. The VLDL-TG production rates were calculated from the linear increase in TG in time (mM TG/min).

Hepatic mRNA Expression – Total RNA was isolated from liver samples using RNA Insta-Pure reagent (Eurogentec, Seraing, Belgium), treated with DNase I, and reverse-transcribed using random primers. cDNA levels were determined by real-time PCR on an ABI Prism 7700 Sequence Detection System (Applied Biosystems, Foster City, CA) using SYBR-Green technology. PCR primers (forward: 5'-GAGGACAAGGCTCGGGAAGT-3'; reverse: 5'-TGAAACCACTCCCGCATCT-3') were designed using Primer Express 1.5 Software with the manufacturer's default settings and obtained from Isogen Bioscience (Maarssen, The Netherlands). The expression levels of human *APOC1* were determined relative to murine HPRT.

Statistical Analysis – Data were analyzed using the unpaired Student's *t* test unless indicated otherwise. *P* values less than 0.05 were considered statistically significant.

Results

Characterization of the Effect of AdAPOC1 on Plasma Lipid Levels in WT mice – First, we addressed the effects of AdAPOC1 on lipid metabolism in WT mice. Figure 1 illustrates the time-course changes for human apoC1 and lipid levels in plasma up to 14 days after injection of a low dose (0.5×10^9 pfu) of AdAPOC1. Plasma human apoC1 protein was detected already at day 3, reached peak concentrations 5 days after injection (18 ± 9 mg/dL), and was disappeared from the plasma at day 14 (Fig. 1A). Plasma TC levels were increased only from day 5 ($170 \pm 19\%$ of initial value; $P < 0.01$), and returned to basal levels at day 10. However, apoC1 expression had the most profound effect on plasma TG levels. TG were already increased at day 3, reached peak levels at day 5 ($335 \pm 117\%$ of initial value; $P < 0.05$) and stayed significantly elevated up to day 10 (Fig. 1B).




Figure 1. Effect of AdAPOC1 on time-course changes of human apoC1, TG, and TC plasma levels in WT mice. Mice were injected with AdAPOC1 (0.5×10^9 pfu). Before injection and at indicated times after injection, fasted plasma was collected from the individual mice and assayed for (A) human apoC1 protein, (B) total cholesterol (TC; closed circles) and triglycerides (TG; open circles). Initial values before injection are set at 100% and values are expressed as percentage of the value before injection $\pm$ S.D. ($n=4$). Asterisks indicate statistically significant differences with the initial value (* $P<0.05$; ** $P<0.01$).




Figure 2. Effect of AdAPOC1 on hepatic APOC1 expression and plasma TG in WT mice. Mice were injected with either AdLacZ (3.3×10^9 pfu) or AdAPOC1 (0.3 – 3.3×10^9 pfu). After 5 days the liver was collected for mRNA isolation and human APOC1 mRNA measurement (A), and plasma was isolated to determine TG (B). Values are expressed as means $\pm$ S.D. ($n=5$).

Subsequently, we evaluated the virus dose-dependent effects of AdAPOC1 in WT mice (Fig. 2). Injection of 0, 0.3, 1.0, and 3.3×10^9 pfu of AdAPOC1 dose-dependently increased human APOC1 mRNA expression in the liver at 5 days after injection (Fig. 2A). Concomitantly, plasma human apoC1 protein levels were dose-dependently increased up to 33 ± 11 mg/dL at the highest dose (Table 1). Table 1 summarizes the plasma lipid levels in the WT mice before and 5 days

Table 1. Dose-dependent effect of AdAPOC1 administration on plasma lipid levels in WT mice.

Condition	Human apoC1 (mg/dL)	TG (mM)	TC (mM)	FFA (mM)
Before adenovirus				
	—	0.22 ± 0.03	1.8 ± 0.2	0.49 ± 0.10
After adenovirus				
Control	n.d.	0.53 ± 0.10	2.2 ± 0.2	0.54 ± 0.03
0.3 × 10 ⁹	12 ± 5	0.79 ± 0.33	2.7 ± 0.2**	0.76 ± 0.04
1.0 × 10 ⁹	23 ± 8	2.36 ± 0.78***	3.4 ± 0.9*	0.75 ± 0.20
3.3 × 10 ⁹	33 ± 11	3.52 ± 0.11***	4.0 ± 0.7***	1.00 ± 0.13***

Plasma was obtained from fasted WT mice before and 5 days after administration of either AdLacZ or AdAPOC1. Plasma lipid levels were measured and values are expressed as means ± S.D. (n=5). Statistical differences were assessed between control virus and AdAPOC1 receiving mice 5 days after injection. **P*<0.05. ***P*<0.01. ****P*<0.001. n.d., not detectable; TG, triglycerides; TC, total cholesterol.

after administration of AdAPOC1. Virus administration *per se* resulted in a slight increase in TG at day 5 (0.53±0.10 vs 0.22±0.03 mM). In mice that were given AdAPOC1 TG dose-dependently increased up to 6.6-fold at the highest dose, which reached statistical significance for 1.0×10⁹ and 3.3×10⁹ pfu of virus. The elevated TG levels were mainly confined to the VLDL and IDL/LDL fractions (Fig. 2B). AdAPOC1 also dose-dependently increased TC, however to a lower extent as compared to TG (up to 1.8-fold at the highest dose), which was also due to an increase in both VLDL and IDL/LDL (not shown). In addition to TG and TC, FFA plasma levels were also increased 1.9-fold (*P*<0.001) at the highest dose. Thus, injection of AdAPOC1 into WT mice leads to a major increase in TG levels accompanied by a mild increase in TC and FFA levels. To address the mechanism(s) underlying the hypertriglyceridemia in these mice, we determined whether the hepatic VLDL-TG production after Triton WR-1339 administration was affected. As shown in Table 2, no differences were observed between the groups of mice that received AdAPOC1 and mice that received control virus. In addition, as the mice were 4 h fasted in all experiments, the observed effect of AdAPOC1 on plasma TG levels cannot be caused by an effect on intestinal chylomicron-TG production. Taken together, the hypertriglyceridemic effect of AdAPOC1 is caused by impaired TG clearance.

Effect of AdAPOC1 on Plasma Lipid Levels in VLDLr-Deficient Mice –

Besides interfering with the peripheral TG hydrolysis by LPL, apoC1 has been postulated to inhibit the hepatic uptake of TG-rich lipoproteins via the LRP and

Table 2. Effect of AdAPOC1 administration on hepatic VLDL-TG production in WT mice.

Condition	VLDL-TG production (mM TG/min)
Control	0.082 ± 0.023
0.3 x 10 ⁹	0.080 ± 0.015
1.0 x 10 ⁹	0.094 ± 0.042
3.3 x 10 ⁹	0.068 ± 0.007

Plasma was obtained from fasted WT mice before and 5 days after administration of either AdLacZ or AdAPOC1. Plasma lipid levels were measured and values are expressed as means ± S.D. (n=5). Statistical differences were assessed between control virus and AdAPOC1 receiving mice 5 days after injection. **P*<0.05. ***P*<0.01. ****P*<0.001. n.d., not detectable; TG, triglycerides; TC, total cholesterol.

LDLr. To investigate the VLDLr-dependency of the apoCI-inhibited TG hydrolysis without potential concomitant effects of apoCI on hepatic receptor recognition, the effect of AdAPOC1 was compared in *lrp1dlr*^{-/-} double knockout mice versus *lrp1dlr*^{-/-}*vldlr*^{-/-} triple knockout mice. At a low viral dose (0.5x10⁹ pfu), AdAPOC1 resulted in moderate levels of human apoCI protein in *lrp1dlr*^{-/-} and *lrp1dlr*^{-/-}*vldlr*^{-/-} mice (17±10 and 45±25 mg/dL, respectively) (Table 3). Similar as on a WT background, APOC1 expression in both *lrp1dlr*^{-/-} and *lrp1dlr*^{-/-}*vldlr*^{-/-} mice resulted in a more pronounced relative increase of TG than of TC as compared to mice that received control virus, mainly in VLDL and IDL/LDL (Fig. 3). However, the effect of AdAPOC1 on plasma lipid levels was remarkably higher in *lrp1dlr*^{-/-}*vldlr*^{-/-} mice as compared to *lrp1dlr*^{-/-} mice. Specifically, AdAPOC1 resulted in 2-fold higher plasma TG levels in *lrp1dlr*^{-/-}*vldlr*^{-/-} mice as compared to *lrp1dlr*^{-/-} mice (14.1±4.3 vs 7.2±1.5 mM, *P*<0.05). Thus, apoCI does not require the presence of the VLDLr for inhibiting the LPL-dependent TG hydrolysis *in vivo*. On the contrary, the TG-raising effect of apoCI is even enhanced in the absence of the VLDLr.

Effect of AdAPOC1 on Plasma Lipid Levels in ApoCIII-Deficient Mice –

ApoCIII^{17,16} and apoE^{25,26} are both inhibitors of LPL *in vivo*, albeit that apoCIII is the most prominent endogenous inhibitor of LPL activity²⁷. To investigate whether the effect of human apoCI on LPL-dependent TG hydrolysis *in vivo* depends on the interaction between apoCI and apoCIII, AdAPOC1 was administered to apoE-knockout mice with or without concomitant apoCIII deficiency. Both *apoE*^{-/-} and *apoE*^{-/-}*apoc3*^{-/-} mice received a low adenoviral dose (0.5x10⁹ pfu), which resulted in human apoCI plasma levels of 45±4 and 52±10 mg/dL respectively. In *apoE*^{-/-} mice, plasma TG was increased by 5.4-fold (*P*<0.001), whereas TC and FFA were not affected (Table 4). Again, the TG-raising effect was specific for

Table 3. Effect of AdAPOC1 administration on plasma lipid levels in *lrp1dlr^{-/-}vldlr^{-/-}* mice.

Genotype	Virus	Human apoC1 (mg/dL)	TG (mM)	TC (mM)	FFA (mM)
Before adenovirus					
<i>lrp1dlr^{-/-}</i>	—	—	2.3 ± 0.8	18 ± 3	0.6 ± 0.1
<i>lrp1dlr^{-/-}vldlr^{-/-}</i>	—	—	3.2 ± 0.7	22 ± 5	0.6 ± 0.1
After adenovirus					
<i>lrp1dlr^{-/-}</i>	AdLacZ	n.d.	5.0 ± 1.0	18 ± 3	1.4 ± 0.4
<i>lrp1dlr^{-/-}</i>	AdAPOC1	17 ± 10	7.2 ± 1.5*	21 ± 1	1.6 ± 0.4
<i>lrp1dlr^{-/-}vldlr^{-/-}</i>	AdLacZ	n.d.	6.2 ± 1.8	24 ± 3	1.8 ± 0.1
<i>lrp1dlr^{-/-}vldlr^{-/-}</i>	AdAPOC1	45 ± 25	14.1 ± 4.3**	31 ± 3**	2.1 ± 0.3*

Plasma was obtained from fasted *lrp1dlr^{-/-}* and *lrp1dlr^{-/-}vldlr^{-/-}* mice before and 5 days after administration of either AdLacZ or AdAPOC1, and levels of human apoC1 and lipids were measured. Values are expressed as means ± S.D. ($n=5$). Statistical differences were assessed between control virus and AdAPOC1 receiving mice 5 days after injection. * $P<0.05$. ** $P<0.01$. n.d., not detectable; TC, total cholesterol; TG, triglycerides.




Figure 3. Effect of AdAPOC1 on plasma TG distribution in *lrp1dlr^{-/-}* and *lrp1dlr^{-/-}vldlr^{-/-}* mice. *lrp1dlr^{-/-}* (A) and *lrp1dlr^{-/-}vldlr^{-/-}* (B) mice were injected with 0.5×10^9 pfu AdLacZ (open circles) or AdAPOC1 (closed circles). After 5 days, fasted plasma was collected, pooled per group ($n=5$), and subjected to FPLC to separate lipoproteins. Fractions were assayed for TG.

VLDL (Fig. 4). In the absence of apoCIII, a 6.8-fold ($P<0.001$) increase in VLDL and IDL/LDL TG was found, in addition to a 2.0-fold ($P<0.001$) increase in TC levels. Thus, expression of human apoC1 leads to hypertriglyceridemia even in the absence of the major LPL inhibitor apoCIII, which indicates that apoC1 is an independent and direct modulator of LPL activity *in vivo*.




Figure 4. Effect of AdAPOC1 on plasma TG distribution in *apoe*^{-/-} and *apoe*^{-/-}*apoc3*^{-/-} mice. *Apoe*^{-/-} (A) and *apoe*^{-/-}*apoc3*^{-/-} (B) mice were injected with 0.5×10^9 pfu AdLacZ (open circles) or AdAPOC1 (closed circles). After 5 days, fasted plasma was collected, pooled per group ($n=5$), and subjected to FPLC to separate lipoproteins. Fractions were assayed for TG.

Discussion

Increased expression of apoCI in mice has been shown to result in combined hyperlipidemia, with a pronounced increase in TG levels in addition to elevated TC levels^{5,6,7,10}. We have recently shown that the combined hyperlipidemia in human APOC1 transgenic mice is the consequence of impaired LPL-mediated hydrolysis of VLDL-TG¹⁰. However, it was still unclear whether apoCI impairs the lipolytic conversion and clearance of lipoproteins *in vivo* by inhibiting LPL activity directly or indirectly via effects on the VLDLr and/or apoCIII, which are both modulators of LPL activity. We generated a novel adenoviral vector encoding human apoCI (AdAPOC1) as this vector allows us to study the interactions of human apoCI with the VLDLr and apoCIII without the need of elaborate crossbreeding of transgenic mouse lines. Upon administration of AdAPOC1 to WT mice, apoCI dose-dependently increased plasma lipid levels, with a preferential increase of TG as compared to TC, which is specific for VLDL and its remnants (*i.e.* IDL and LDL). Together with the observation that AdAPOC1 raises TG levels at an earlier stage (day 3) than TC levels (day 5), these data underscore our previous findings that apoCI primarily interferes with the LPL-dependent processing of VLDL rather than with the hepatic uptake of its core remnants¹⁰. The apoCI-induced hypertriglyceridemia was still apparent upon deficiency of the LDLr and LRP (involved in hepatic remnant clearance), or VLDLr and apoCIII (involved in peripheral LPL-dependent TG hydrolysis), indicating that apoCI is indeed a powerful and direct modulator of LPL activity. Since the hepatic uptake of TG-

Table 4. Effect of AdAPOC1 administration on plasma lipid levels in *apoe*^{-/-} and *apoe*^{-/-}*apoc3*^{-/-} mice.

Genotype	Virus	Human apoCI (mg/dL)	TG (mM)	TC (mM)	FFA (mM)
Before adenovirus					
<i>apoe</i> ^{-/-}	—	—	1.0 ± 0.3	13 ± 3	0.8 ± 0.1
<i>apoe</i> ^{-/-} <i>apoc3</i> ^{-/-}	—	—	0.6 ± 0.2	8 ± 2	0.5 ± 0.1
After adenovirus					
<i>apoe</i> ^{-/-}	AdLacZ	n.d.	1.2 ± 0.1	15 ± 2	1.0 ± 0.1
<i>apoe</i> ^{-/-}	AdAPOC1	45 ± 4	6.3 ± 1.3**	17 ± 2	1.0 ± 0.2
<i>apoe</i> ^{-/-} <i>apoc3</i> ^{-/-}	AdLacZ	n.d.	0.6 ± 0.1	8 ± 1	0.6 ± 0.1
<i>apoe</i> ^{-/-} <i>apoc3</i> ^{-/-}	AdAPOC1	52 ± 10	4.0 ± 1.2***	17 ± 3***	0.8 ± 0.1*

Plasma was obtained from fasted *apoe*^{-/-} and *apoe*^{-/-}*apoc3*^{-/-} mice before and 5 days after administration of either AdLacZ or AdAPOC1. Plasma human apoCI and plasma lipid levels were measured. Values are expressed as means ± S.D. (*n*=4). Statistical differences were assessed between control virus and AdAPOC1 receiving mice 5 days after injection. **P*<0.05. ***P*<0.01. ****P*<0.001. n.d., not detectable; TC, total cholesterol; TG, triglycerides.

rich lipoproteins is highly dependent on processing by LPL, thereby generating remnant particles with reduced size with an altered apolipoprotein composition, apoCI will have a secondary inhibiting effect on hepatic remnant uptake. In addition, the hepatic remnant uptake may be directly affected by apoCI.

At similar plasma human apoCI levels, the plasma lipid phenotype of AdAPOC1-treated mice is virtually identical to that of APOC1 transgenic mice^{6,7,10}. Since intravenous injection of recombinant adenoviruses leads to nearly exclusive infection of the liver, this indicates that liver-derived apoCI can be fully responsible for the hyperlipidemia in APOC1 transgenic mice. We did not observe any effect of hepatic APOC1 expression on hepatic VLDL-TG production, which is in line with our recent observations in APOC1 transgenic mice¹⁰. Furthermore, we can rule out that the hypertriglyceridemic effect of APOC1 expression is caused by an effect on TG influx into plasma by modulating intestinal lipid absorption as we used 4 h fasted mice to exclude potential influx of chylomicrons from the intestine. In addition, we have recently shown that intestinal TG absorption in transgenic APOC1 mice is not affected as compared to WT mice¹⁰. Therefore, our present experiments with AdAPOC1 confirm our previous conclusion that the hypertriglyceridemia in APOC1 transgenic mice is truly caused by a defect in peripheral lipolysis¹⁰, thereby preventing subsequent hepatic remnant uptake.

ApoCI has been shown to inhibit the binding of VLDL to the VLDLr *in vitro* and *in vivo*¹⁶. Since the VLDLr is commonly assumed to be required for normal LPL functioning *in vivo*, apoCI might indeed act by disrupting the binding of VLDL to the VLDLr, thereby inhibiting the LPL-mediated TG hydrolysis. We thus evaluated the effect of hepatic human apoCI expression in VLDLr-deficient mice. These

mice were also deficient for the LDLr and hepatic LRP to exclude a potentially additional effect of apoCI on receptor-mediated hepatic remnant clearance. Hepatic apoCI expression in LRP and LDLr double-deficient mice increased plasma VLDL-TG and VLDL-cholesterol. Remarkably, the additional absence of the VLDLr resulted in an aggravated combined hyperlipidemia, associated with higher plasma human apoCI levels, while the hepatic apoCI expression levels were similar in *lrp-ldlr^{-/-}* and *lrp-ldlr^{-/-}vldlr^{-/-}* mice. This suggests that, although apoCI reduces the binding affinity of VLDL for the VLDLr¹⁶, the VLDLr can still facilitate the lipolysis of apoCI-enriched VLDL by bringing VLDL in close proximity of LPL at the endothelial surface. The VLDLr could also be involved in the clearance of apoCI, leading to accumulation of plasma apoCI in the absence of the VLDLr, with a concomitant increase in combined hyperlipidemia. Regardless of the precise mechanism, the TG levels in LRP, LDLr, and VLDLr triple-knockout mice increased 7.9 mM after AdAPOC1 administration, illustrating that the inhibition of LPL-mediated TG hydrolysis by apoCI can indeed occur independently of the VLDLr *in vivo*.

It is commonly known that apoCIII is the most prominent physiological inhibitor of LPL activity^{12,18,28,29}. Indeed, human APOC3 transgenic mice show marked hypertriglyceridemia and hypercholesterolemia, with severely elevated VLDL levels³⁰. Reciprocally, *apoc3*-deficient mice on an *apoE^{-/-}* background show decreased VLDL-TG levels^{18,31}. Therefore, we also questioned whether the capacity of apoCI to inhibit LPL proceeds via an interaction with apoCIII. Since we and others have shown that apoE is also an inhibitor of LPL *in vivo*^{25,26}, we determined the effect of apoCI expression in apoE-knockout and apoE/apoCIII double-knockout mice. Injection of AdAPOC1 into *apoE^{-/-}* mice resulted in aggravated hyperlipidemia, which confirms previous findings that apoCI can act independently of apoE^{10,11}. Importantly, apoCI expression in *apoE^{-/-}apoc3^{-/-}* mice led to a similar increase in TG as observed in *apoE^{-/-}* only mice, establishing that the LPL-inhibiting effect of apoCI occurs without interaction of apoCI with endogenous apoCIII.

In conclusion, we have demonstrated that low to moderate liver-specific expression of human apoCI in mice results in dose-dependent hypertriglyceridemia even in the absence of the VLDLr or apoCIII, presumably by direct inhibition of LPL activity, as we have recently shown¹⁰. It is evident that LPL activity strongly determines plasma TG levels, as overexpression of LPL in mice markedly reduces VLDL-TG levels^{32,33}, whereas heterozygous deficiency of LPL results in accumulation of VLDL-TG³⁴. Furthermore, LPL is an important factor for the influx of fatty acids into adipose tissue. We have shown that an increased LPL function leads to aggravated obesity on high fat diet¹⁷ whereas a decreased LPL function results in protection against diet-induced obesity^{13,15}. Therefore, our findings

may have important implications for the treatment of hypertriglyceridemia and obesity.

Acknowledgements

This work was performed in the framework of the Leiden Center for Cardiovascular Research LUMC-TNO, and supported by the Leiden University Medical Center (Gisela Thier Fellowship to P.C.N.R.), the Netherlands Organization for Scientific Research (NWO RIDE grant 014-90-001, NWO grant 908-02-097, and NWO VIDI grant 917.36.351), and the Netherlands Heart Foundation (NHS grant 2003B136).

References

1. Allan CM, Walker D, Segrest JP, Taylor JM. Identification and characterization of a new human gene (APOC4) in the apolipoprotein E, C-I, and C-II gene locus. *Genomics* 1995;28:291-300.
2. Lauer SJ, Walker D, Elshourbagy NA, Reardon CA, Levy-Wilson B, Taylor JM. Two copies of the human apolipoprotein C-I gene are linked closely to the apolipoprotein E gene. *J Biol Chem* 1988;263:7277-7286.
3. Scott J, Knott TJ, Shaw DJ, Brook JD. Localization of genes encoding apolipoproteins CI, CII, and E to the p13-----cen region of human chromosome 19. *Hum Genet* 1985;71:144-146.
4. Smit M, Kooij-Meijis E, Frants RR, Havekes L, Klasen EC. Apolipoprotein gene cluster on chromosome 19. Definite localization of the APOC2 gene and the polymorphic Hpa I site associated with type III hyperlipoproteinemia. *Hum Genet* 1988;78:90-93.
5. Jong MC, Dahlmans VE, van Gorp PJ, van Dijk KW, Breuer ML, Hofker MH, Havekes LM. In the absence of the low density lipoprotein receptor, human apolipoprotein C1 overexpression in transgenic mice inhibits the hepatic uptake of very low density lipoproteins via a receptor-associated protein-sensitive pathway. *J Clin Invest* 1996;98:2259-2267.
6. Shachter NS, Ebara T, Ramakrishnan R, Steiner G, Breslow JL, Ginsberg HN, Smith JD. Combined hyperlipidemia in transgenic mice overexpressing human apolipoprotein CI. *J Clin Invest* 1996;98:846-855.
7. Simonet WS, Bucay N, Pitas RE, Lauer SJ, Taylor JM. Multiple tissue-specific elements control the apolipoprotein E/C-I gene locus in transgenic mice. *J Biol Chem* 1991;266:8651-8654.
8. Sehaye E and Eisenberg S. Mechanisms of inhibition by apolipoprotein C of apolipoprotein E-dependent cellular metabolism of human triglyceride-rich lipoproteins through the low density lipoprotein receptor pathway. *J Biol Chem* 1991;266:18259-18267.
9. Weisgraber KH, Mahley RW, Kowal RC, Herz J, Goldstein JL, Brown MS. Apolipoprotein C-I modulates the interaction of apolipoprotein E with beta-migrating very low density lipoproteins (beta-VLDL) and inhibits binding of beta-VLDL to low density lipoprotein receptor-related protein. *J Biol Chem* 1990;265:22453-22459.
10. Berbee JF, van der Hoogt CC, Sundaraman D, Havekes LM, Rensen PC. Severe hypertriglyceridemia in human APOC1 transgenic mice is caused by apoC-I-induced inhibition of LPL. *J Lipid Res* 2005;46:297-306.
11. Conde-Knape K, Bensadoun A, Sobel JH, Cohn JS, Shachter NS. Overexpression of apoC-I in apoE-null mice: severe hypertriglyceridemia due to inhibition of hepatic lipase. *J Lipid Res* 2002;43:2136-2145.
12. Havel RJ, Fielding CJ, Olivecrona T, Shore VG, Fielding PE, Egelrud T. Cofactor activity of protein components of human very low density lipoproteins in the hydrolysis of triglycerides by lipoproteins lipase from different sources. *Biochemistry* 1973;12:1828-1833.
13. Goudriaan JR, Tacke PJ, Dahlmans VE, Gijbels MJ, van Dijk KW, Havekes LM, Jong MC. Protection from obesity in mice lacking the VLDL receptor. *Arterioscler Thromb Vasc Biol* 2001;21:1488-1493.
14. Obunike JC, Lutz EP, Li Z, Paka L, Katopodis T, Strickland DK, Kozarsky KF, Pillarisetti S, Goldberg IJ. Transcytosis of lipoprotein lipase across cultured endothelial cells requires both heparan sulfate proteoglycans and the very low density lipoprotein receptor. *J Biol Chem*

- 2001;276:8934-8941.
15. Jong MC, Voshol PJ, Muurling M, Dahlmans VE, Romijn JA, Pijl H, Havekes LM. Protection from obesity and insulin resistance in mice overexpressing human apolipoprotein C1. *Diabetes* 2001;50:2779-2785.
 16. Jong MC, van Dijk KW, Dahlmans VE, van der Boom H, Kobayashi K, Oka K, Siest G, Chan L, Hofker MH, Havekes LM. Reversal of hyperlipidaemia in apolipoprotein C1 transgenic mice by adenovirus-mediated gene delivery of the low-density-lipoprotein receptor, but not by the very-low-density-lipoprotein receptor. *Biochem J* 1999;338 (Pt 2):281-287.
 17. Duivenvoorden I, Teusink B, Rensen PC, Romijn JA, Havekes LM, Voshol PJ. Apolipoprotein C3 deficiency results in diet-induced obesity and aggravated insulin resistance in mice. *Diabetes* 2005;54:664-671.
 18. Jong MC, Rensen PC, Dahlmans VE, van der Boom H, van Berkel TJ, Havekes LM. Apolipoprotein C-III deficiency accelerates triglyceride hydrolysis by lipoprotein lipase in wild-type and apoE knockout mice. *J Lipid Res* 2001;42:1578-1585.
 19. Espirito Santo SM, Rensen PC, Goudriaan JR, Bensadoun A, Bovenschen N, Voshol PJ, Havekes LM, van Vlijmen BJ. Triglyceride-rich lipoprotein metabolism in unique VLDL receptor, LDL receptor, and LRP triple-deficient mice. *J Lipid Res* 2005;46:1097-1102.
 20. He TC, Zhou S, da Costa LT, Yu J, Kinzler KW, Vogelstein B. A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci U S A* 1998;95:2509-2514.
 21. Fallaux FJ, Bout A, van dV, I, van den Wollenberg DJ, Hehir KM, Keegan J, Auger C, Cramer SJ, van Ormondt H, van der Eb AJ, Valerio D, Hoeben RC. New helper cells and matched early region 1-deleted adenovirus vectors prevent generation of replication-competent adenoviruses. *Hum Gene Ther* 1998;9:1909-1917.
 22. Tao N, Gao GP, Parr M, Johnston J, Baradet T, Wilson JM, Barsoum J, Fawell SE. Sequestration of adenoviral vector by Kupffer cells leads to a nonlinear dose response of transduction in liver. *Mol Ther* 2001;3:28-35.
 23. Zambon A, Hashimoto SI, Brunzell JD. Analysis of techniques to obtain plasma for measurement of levels of free fatty acids. *J Lipid Res* 1993;34:1021-1028.
 24. Jong MC, Dahlmans VE, van Gorp PJ, Breuer ML, Mol MJ, van der ZA, Frants RR, Hofker MH, Havekes LM. Both lipolysis and hepatic uptake of VLDL are impaired in transgenic mice coexpressing human apolipoprotein E*3Leiden and human apolipoprotein C1. *Arterioscler Thromb Vasc Biol* 1996;16:934-940.
 25. Huang Y, Liu XQ, Rall SC, Jr., Taylor JM, von Eckardstein A, Assmann G, Mahley RW. Overexpression and accumulation of apolipoprotein E as a cause of hypertriglyceridemia. *J Biol Chem* 1998;273:26388-26393.
 26. Rensen PC and van Berkel TJ. Apolipoprotein E effectively inhibits lipoprotein lipase-mediated lipolysis of chylomicron-like triglyceride-rich lipid emulsions in vitro and in vivo. *J Biol Chem* 1996;271:14791-14799.
 27. Gerritsen G, Rensen PC, Kypreos KE, Zannis VI, Havekes LM, Willems vD. ApoC-III deficiency prevents hyperlipidemia induced by apoE overexpression. *J Lipid Res* 2005;46:1466-1473.
 28. Brown WV and Baginsky ML. Inhibition of lipoprotein lipase by an apoprotein of human very low density lipoprotein. *Biochem Biophys Res Commun* 1972;46:375-382.
 29. Ebara T, Ramakrishnan R, Steiner G, Shachter NS. Chylomicronemia due to apolipoprotein CIII overexpression in apolipoprotein E-null mice. Apolipoprotein CIII-induced hypertriglyceridemia is not mediated by effects on apolipoprotein E. *J Clin Invest* 1997;99:2672-2681.
 30. Ito Y, Azrolan N, O'Connell A, Walsh A, Breslow JL. Hypertriglyceridemia as a result of human apo CIII gene expression in transgenic mice. *Science* 1990;249:790-793.
 31. Maeda N, Li H, Lee D, Oliver P, Quarfordt SH, Osada J. Targeted disruption of the apolipoprotein C-III gene in mice results in hypotriglyceridemia and protection from postprandial hypertriglyceridemia. *J Biol Chem* 1994;269:23610-23616.
 32. Excoffon KJ, Liu G, Miao L, Wilson JE, McManus BM, Semenkovich CF, Coleman T, Benoit P, Duverger N, Branellec D, Deneffe P, Hayden MR, Lewis ME. Correction of hypertriglyceridemia and impaired fat tolerance in lipoprotein lipase-deficient mice by adenovirus-mediated expression of human lipoprotein lipase. *Arterioscler Thromb Vasc Biol* 1997;17:2532-2539.
 33. Shimada M, Shimano H, Gotoda T, Yamamoto K, Kawamura M, Inaba T, Yazaki Y, Yamada N. Overexpression of human lipoprotein lipase in transgenic mice. Resistance to diet-induced hypertriglyceridemia and hypercholesterolemia. *J Biol Chem* 1993;268:17924-17929.
 34. Coleman T, Seip RL, Gimble JM, Lee D, Maeda N, Semenkovich CF. COOH-terminal disruption of lipoprotein lipase in mice is lethal in homozygotes, but heterozygotes have elevated triglycerides and impaired enzyme activity. *J Biol Chem* 1995;270:12518-12525.