



Universiteit
Leiden
The Netherlands

The impact of obesity on the pharmacokinetics of drugs in adolescents and adults

Rongen, A. van; Rongen A. van

Citation

Rongen, A. van. (2016, December 7). *The impact of obesity on the pharmacokinetics of drugs in adolescents and adults*. Retrieved from <https://hdl.handle.net/1887/44789>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/44789>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/44789> holds various files of this Leiden University dissertation


Author: Rongen, Anne van

Title: The impact of obesity on the pharmacokinetics of drugs in adolescents and adults

Issue Date: 2016-12-07

Chapter 3

Drug disposition in obesity: toward evidence-based dosing



Catherijne A.J. Knibbe
Margreke J.E. Brill
Anne van Rongen
Jeroen Diepstraten
Piet Hein van der Graaf
Meindert Danhof

Annu Rev Pharmacol Toxicol. 2015; 55: 149-67

ABSTRACT

Obesity and morbid obesity are associated with many physiological changes affecting pharmacokinetics, such as increased blood volume, cardiac output, splanchnic blood flow, and hepatic blood flow. In obesity, drug absorption appears unaltered, although recent evidence suggests that this conclusion may be premature. Volume of distribution may vary largely, but the magnitude and direction of changes seem difficult to predict, with extrapolation on the basis of total body weight being the best approach to date. Changes in clearance may be smaller than in distribution, whereas there is growing evidence that the influence of obesity on clearance can be predicted on the basis of reported changes in the metabolic or elimination pathways involved. For obese children, we propose two methods to distinguish between developmental and obesity-related changes. Future research should focus on the characterization of physiological concepts to predict the optimal dose for each drug in the obese population.

INTRODUCTION

Obesity represents a serious and increasing health problem worldwide. In the United States in 2009-2010, the prevalence of obesity (body mass index (BMI) $> 30 \text{ kg/m}^2$) was 35.9%, and the prevalence of morbid obesity (BMI $> 40 \text{ kg/m}^2$) was 6.3% (8.2% for women and 4.4% for men)¹. Alarming, the prevalence of overweight and obesity in children is also increasing. According to the most recent National Health and Nutrition Survey (2009-2010) 31.8% of US children and adolescents (age 2-19 years) are overweight (≥ 85 th percentile of BMI for age), 16.9% are obese (≥ 95 th percentile), and 12.3% are morbidly obese (≥ 97 th percentile)². Worldwide prevalence rates for obesity in adults and overweight and obesity rates in children are also high, exceeding 24% in, for instance, Canada, Spain, the United Kingdom, Greece, Mexico, Saudi Arabia, Egypt, Australia, New Zealand, and some parts of South America³.

Obesity increases the risk of many diseases and health conditions, such as hypertension, cardiovascular disease, dyslipidaemia, type 2 diabetes, cancer, and osteoarthritis, thereby diminishing average life expectancy⁴. In addition, obese individuals are also more likely to suffer from chronic pain⁵⁻⁶ and nosocomial infections⁷⁻⁸. Because these comorbidities often require pharmacotherapeutic or surgical and anaesthetic treatment, an important question is how to optimize the dose of drugs, particularly in light of the fact that the morbidly obese patient group is increasing. In this respect, specific attention should be paid to obese children, who are likely to become obese adults. Comorbidities associated with childhood obesity are hypertension, obstructive sleep apnoea, diabetes mellitus, and coronary artery disease, necessitating pharmacotherapeutic or even surgical or bariatric treatment⁹⁻¹⁰. Furthermore, obese children are also more likely to develop asthma or severe asthma¹¹, but their response to inhaled steroids is decreased¹². Moreover, overweight and obesity have been reported as independent predictors of the relapse risk of acute lymphoblastic leukaemia¹³. It cannot be excluded that these differences result from changes in the pharmacokinetics of chemotherapeutic agents in overweight or obese children. Therefore, it is of utmost importance to gain insight into how to adjust the dose of drugs in obese and morbidly obese children and adolescents. This issue should be viewed through the perspective of the fact that even in nonobese children, 37-80% of drugs are prescribed in an off-label or unlicensed manner¹⁴⁻¹⁶.

In this review, we provide an overview of the current knowledge on changes in drug disposition in obese patients in relation to physiological changes associated with obesity. Our ultimate goal is to direct future research aiming for individualized dosing in this growing and heterogeneous patient population. We pay specific attention to changes in drug disposition in obese children.

PHYSIOLOGICAL CHANGES ASSOCIATED WITH OBESITY

Obesity is associated with many physiological and pathophysiological changes that may affect drug disposition. Obesity and morbid obesity are not only associated with an increase in fat but also in lean body weight (LBW), which is the weight devoid of all adipose tissue. The percentage of fat mass per kilogram of total body weight increases more than LBW in obese patients, with, for instance, an increase in LBW representing 20-40% of total excess of weight in morbidly obese patients¹⁷⁻¹⁸.

To supply the excess body mass with oxygen and nutrients, blood volume, cardiac output, and capillary flow increase substantially in obese and, in particular, morbidly obese individuals¹⁹⁻²². Serum albumin and total protein concentrations are reported to be comparable in lean and obese subjects, even though concentrations of alpha-1-acid glycoprotein are increased²³. In the cardiovascular system, the increased blood volume and cardiac output eventually leads to systemic hypertension, left and right ventricular hypertrophy, and an increased risk for sudden cardiac death due to conduction disorders²⁴⁻²⁵. Pulmonary function is uniformly altered in obesity, with reduced lung volumes²⁶ and a higher incidence of obstructive sleep apnea syndrome²⁷.

Nonalcoholic steatohepatitis and histological abnormalities such as fatty infiltration in the liver are very common in morbidly obese patients²⁸⁻²⁹. Because of the accumulation of fat in the liver of obese individuals, functional morphology may be altered owing to sinusoidal narrowing³⁰⁻³¹. However, because of increased blood volume and cardiac output, liver blood flow is not necessarily reduced in obese subjects³². Although liver volume is reported to be increased in obese individuals³³, the results of studies on the influence of obesity on expression and function of CYP enzymes are inconclusive, with the exception of CYP3A and CYP2E1; the expression and function of these enzymes have been reported to be decreased and increased, respectively³⁴.

There are conflicting data on alterations in renal function. Irrespective of the presence of hypertension, investigators have reported increases in glomerular filtration rate and effective renal plasma flow³⁵⁻³⁷. However, there is also evidence of unaltered renal function³⁸. In studies in Zucker rats with genetic obesity, researchers found that, after an initial increase in glomerular filtration rate, this rate normalized and subsequently decreased in the later stages of obesity, ultimately leading to end-stage renal disease³⁹⁻⁴¹. In morbidly obese patients who presented with proteinuria, one study reported focal glomerular sclerosis, diabetic nephropathy, or both⁴². In addition, estimates of the creatinine clearance from standard formulas tend to be inaccurate in obese patients⁴³⁻⁴⁵. Even though obesity-associated renal damage may be unpredictable, the available evidence indicates that it is best to use LBW in the Cockcroft-Gault formula for estimation of creatinine clearance in obese patients^{44,46}.

With respect to the functioning of the gastrointestinal tract, studies in obese subjects have found accelerated gastric emptying of solids⁴⁷⁻⁵⁰, high splanchnic blood flow¹⁹, and increased gut wall permeability⁵¹⁻⁵². Because studies on the influence of obesity on intestinal transit time and motility have shown contradictory results, the exact impact of obesity on drug or nutrient absorption remains unclear^{50,53-54}. Wisén & Johansson⁵⁴ found that obese subjects had significantly higher absorption in the proximal small intestine. Studies on the influence of obesity on enterohepatic recirculation are lacking.

MEASURES TO QUANTIFY BODY SIZE AND OVERWEIGHT

BMI is the international metric recommended by the World Health Organization to classify obesity⁵⁵. A BMI value between 18.5 and 25 kg/m² is considered healthy. BMI values greater than 30 and 40 kg/m² indicate obesity and morbid obesity, respectively⁵⁵. As BMI does not differentiate adipose tissue from muscle mass, BMI should be considered a descriptor of body shape instead of a measure of body composition⁵⁶⁻⁵⁷. For a child's weight status (2-18 years), an age- and sex-specific percentile for BMI (BMI-for-age) is used because children's body compositions vary as they age and between boys and girls⁵⁸⁻⁵⁹. For children younger than 2 years, weight-for-length charts are used. Overweight is defined as a BMI between the 85th and 95th percentile and obesity above the 95th percentage for children of the same age and sex⁶⁰.

The value of the ideal body weight (IBW) parameter is most commonly calculated using the equation by Devine⁶¹. Similar to BMI, this measure is rarely used as the basis for the individualization of drug dosage in obese patients, except for some specific drugs such as muscle relaxants⁶²⁻⁶⁴ and remifentanyl⁶⁵. This measure may lack predictive value for the dose adjustment of other drugs because it is based on height and sex only and does not consider body weight in any way⁵⁶. Adjusted body weight is an empirical, IBW-based metric with different correction factors (0.14-0.98) that was developed after the discovery that IBW was a suboptimal parameter for drug dosing in obese subjects⁶⁶, but very little evidence supports using this as a guide for dosing⁶⁷.

Body surface area (BSA) is mainly used for dosing of anticancer drugs, a practice that has a historical rather than scientific basis. BSA can be calculated using the equations by Dubois and Dubois⁶⁸ or Mosteller⁶⁹. The equations are based on the theory of Euclidean geometry and account for height and weight⁶⁶. Remarkably, recent reports have shown that there is no evidence to reduce the dose or dose capping when BSA-adjusted doses are used in obese or morbidly obese cancer patients⁷⁰⁻⁷¹. These results may be explained by the nonlinear relation of BSA with total body weight (Figure 1), reducing the absolute increase in dose in relation to the increase in body weight.

Because of the drawbacks of the previous measures, researchers have proposed using lean body weight (LBW) as a measure of body composition⁷³. Information on body weight as well as height and gender are required to calculate LBW (Figure 1). LBW represents the weight of bones, muscles, tendons, and organs without body fat (i.e., fat-free mass). The most recent LBW equation, proposed by Janmahasatian et al.⁷², provided good predictions of the fat-free mass as measured with bioelectrical impedance analysis or dual-energy X-ray absorptiometry. The exact value of LBW as a predictor for dosing remains to be established. In this respect, it is important to note that in pharmacometric studies, this parameter was not always identified as the best predictor^{67,74-75}. Peters et al.⁷⁶ proposed a new formula to calculate LBW in children. However, researchers have very limited experience with this measure as a predictor for dosing drugs in obese children⁷⁷.




Figure 1 Lean body weight⁷² (a) and body surface area⁶⁸ (b) vs. total body weight for males of various heights.

In general, actual body weight should be used with caution as a body-size descriptor in obesity because its value is influenced by factors such as age, sex, height, muscle mass, and obesity. Nevertheless, nonlinear functions of total body weight (TBW) show good performance as predictors of clearance in several pharmacokinetic studies covering wide ranges in body weight^{74-75,78}. Similarly, in a large study on the variation in clearance and volume of distribution of 12 different drugs, total body weight appeared to be a consistent and reliable size descriptor for the prediction of these parameters in the obese⁷⁹.

THE INFLUENCE OF OBESITY ON ORAL BIOAVAILABILITY AND ABSORPTION RATE

Only six studies have directly compared the oral bioavailability and absorption rate of drugs between obese and nonobese subjects on the basis of both oral and intravenous

administration⁸⁰⁻⁸⁵. For propranolol, clearance (CL) after an intravenous dose was not different between six obese (136 ± 36 kg) and six control (67 ± 5 kg) subjects. However, oral clearance (CL/F) was lower in obese patients, indicating that the bioavailability (F) of propranolol was slightly higher for obese subjects ($35 \pm 4\%$ vs. $27 \pm 2\%$, $P > 0.05$)⁸⁰. In the discussion of their article, the authors point out that the slightly higher bioavailability reported for propranolol may also be applicable for triazolam^{80,86}. Unfortunately, in the study on triazolam, there were no observations after intravenous administration⁸⁶, which makes it impossible to draw conclusions on an eventual difference in absolute bioavailability. For midazolam, no difference in bioavailability was found between normal-weight volunteers (66 ± 2 kg, $n = 20$) and obese volunteers (117 ± 8 kg, $n = 20$) ($40 \pm 3\%$ vs. $42 \pm 4\%$, $P > 0.05$, respectively), nor was a difference found in time of maximum concentration ($T_{\max}$) or maximum concentration ($C_{\max}$) itself⁸¹. Similarly, no difference in bioavailability or oral absorption rate was found for trazodone, cyclosporine, dexfenfluramine, and moxifloxacin between obese and nonobese subjects⁸²⁻⁸⁵.

In view of the limited number of studies on oral absorption, we most recently studied midazolam bioavailability in 20 morbidly obese patients (mean body weight 144 kg (112-186 kg) and mean BMI 47 kg/m^2 ($40\text{-}68 \text{ kg/m}^2$)) and 12 healthy volunteers (76 kg (63-93 kg) and mean BMI 22 kg/m^2 ($19\text{-}26 \text{ kg/m}^2$)) (<http://clinicaltrials.gov/show/NCT01519726>). For this study, a semisimultaneous oral and intravenous administration design was chosen in which morbidly obese patients received 7.5 mg of midazolam orally followed by a 5 mg intravenous bolus dose after 159 ± 67 min. Healthy volunteers received 2-mg oral and 1-mg intravenous midazolam separated by 150 min. This study design allowed for the characterization of both clearance and bioavailability in a single pharmacokinetic study. Results of this study show an increased bioavailability ($60 \pm 13\%$ vs. $28 \pm 7\%$, $P < 0.01$) and a lower oral absorption rate ($0.057 \pm 14\% \text{ min}^{-1}$ vs. $0.13 \pm 5\% \text{ min}^{-1}$, $P < 0.01$), but no influence of obesity on systemic clearance in morbidly obese patients compared to healthy volunteers⁸⁷. Dose simulations of the final population pharmacokinetic model showed that after a 7.5-mg oral midazolam, $C_{\max}$ is only slightly lower, whereas $T_{\max}$ is increased for morbidly obese patients (Figure 2a).

The significant difference in oral bioavailability reported in this study⁸⁷ may result from the larger body weights of the subjects compared to the previous study by Greenblatt et al.⁸¹, who reported no difference in bioavailability (mean body weight of 144 kg vs. 117 kg). The observed higher bioavailability could be explained by an increased splanchnic blood flow¹⁹, which may lead to reduced contact between midazolam and intracellular CYP3A enzymes in the gut wall. Also, the increase in bioavailability may be explained by increased paracellular absorption through the gut wall, or a combination of both^{51-52,88-89}. The higher midazolam bioavailability found in morbidly obese patients, however, does not seem to result in higher $C_{\max}$ values (Figure 2a); this may be explained by the higher volume of distribution⁸⁷ which was also reported by Greenblatt et al.⁸¹. The lower absorp-

tion rate (and therefore increased $T_{\max}$) in morbidly obese patients may be the result of the difference in midazolam formulation, as healthy volunteers received an oral solution and morbidly obese patients a tablet. As midazolam effectiveness is determined by the initial midazolam concentrations after an oral dose, this study suggests that the net result of the alterations in the different pharmacokinetic parameters is that no adjustments in oral midazolam dose seem necessary for obese individuals. However, a different conclusion should be drawn for intravenous administration, given the substantially increased volumes of distribution of midazolam in morbidly obese patients (Figure 2b,c)^{81,87}.




Figure 2 Population-predicted midazolam concentrations over time based on the final pharmacokinetic model in three typical morbidly obese patients (112, 145, and 186 kg) and one healthy volunteer (76 kg) after (a) a 7.5-mg oral dose (linear scale) (b) a 5-mg intravenous bolus dose (logarithmic scale) and (c) a 2.5-mg/h continuous infusion (linear scale). Figure adapted from Reference 87 (Chapter 7) with permission.

In conclusion, there is limited information on the influence of obesity on drug pharmacokinetics after oral administration, despite the fact that most drugs are given orally. From the very small number of studies on drug absorption identified in this review, it seems that drug absorption is rather unaltered. However, this may be a premature conclusion warranting further systematic evaluations on drug absorption⁹⁰. Given the reported accelerated gastric emptying of solids⁴⁷⁻⁵⁰, increased splanchnic blood flow¹⁹, and increased gut permeability⁵¹⁻⁵² in obese subjects, changes in absorption rate

and oral bioavailability cannot be excluded. The recent study on midazolam oral and intravenous pharmacokinetics in both morbidly obese patients and healthy volunteers confirms some of these anticipated changes⁸⁷. The design of this study may be used as an example to study drug absorption because both oral and intravenous administration were evaluated within each individual. Investigators analyzing results on drug absorption from a study without data after intravenous administration risk being unable to distinguish between the influence of obesity on clearance and bioavailability (or between volume of distribution and bioavailability). Finally, the consequences of altered absorption rate and oral bioavailability should each be evaluated for their clinical relevance and impact on drug dosing in the obese population.

THE INFLUENCE OF OBESITY ON DRUG DISTRIBUTION

Volume of distribution is an important parameter that is often substantially altered in obese patients^{79,90-92}. It is particularly important to characterize changes in volume of distribution when a rapid onset of the effect is needed as the peak concentration after single-dose administration is largely determined by the volume of distribution. The same applies for the time to reach steady state and an eventual loading dose as part of a continued or repeated administration scheme. A rapid onset of effect may be clinically relevant in anaesthesia, for anticoagulation, and for antimicrobial drug effects.

In general, drug distribution depends on the physicochemical properties of the drug, such as molecular weight, lipid solubility, and protein binding, as well as the properties of the biological system^{91,93}. The latter properties may differ between subjects (obese subjects vs. healthy volunteers). In obese subjects, changes in volume of distribution may be expected to result from increased blood volume, increased cardiac output and blood flow, increased LBW, increased adipose tissue and reduced tissue perfusion^{19-22,91-92}, with only a limited influence of changes in blood proteins (i.e., albumin, alpha acid glycoprotein)^{23,94}.

From the available evidence, the values of the volume of distribution appear highly variable in obese individuals and more difficult to predict than the values of clearance^{79,90}. While intuitively more influence of obesity on lipophilic drugs than on hydrophilic drugs may be expected⁹³, Jain et al.⁹⁰ concluded, on the basis of an overview of the ratios of volume of distribution of various drugs in obese vs. nonobese individuals, that changes in volume of distribution cannot be predicted on the basis of lipophilicity alone. More specifically, they showed that, for lipophilic drugs, the values for volume of distribution normalized with body weight may be increased, unchanged, or reduced⁹⁰. Also, in our experience, volume of distribution is difficult to predict. For instance, no influence of obesity on the peripheral volumes of distribution of propofol was observed, despite the high lipophilicity of the drug^{75,77-78}. For hydrophilic drugs, unchanged or

decreased ratios of volume of distribution normalized with body weight were observed, but the magnitude of the effect of obesity was smaller than for lipophilic drugs⁹⁰.

Similarly, Mahmood⁷⁹ concluded, on the basis of a study on the pharmacokinetics of 12 different drugs, that predictions of volume of distribution in the obese from the values in normal-weight subjects were less accurate than predictions of clearance. Although total body weight appeared to be a more consistent and reliable size descriptor than other size descriptors for the prediction of volume of distribution⁷⁹, as was suggested before⁵⁶, linear scaling of volume of distribution with body weight was reported to lead to overprediction of volume of distribution in the obese for many drugs. Instead, prediction of volume of distribution by an allometric model on the basis of total body weight was more accurate. However, for the 12 drugs studied, the exponents of allometric functions were found to vary widely (0.27-2.459), illustrating the variability of changes in volume of distribution as a result of total body weight⁷⁹. As the allometric models were built on data from normal-weight subjects, Mahmood concluded that inclusion of data from the obese into these allometric models could lead to better predictions⁷⁹.

The relative impact of the obesity-related changes in volume of distribution with respect to adjusting the dose in obese individuals is illustrated below in three examples.

Example 1: Cefazolin

In a clinical microdialysis study, cefazolin concentrations in subcutaneous adipose tissue and in plasma were evaluated in morbidly obese and nonobese patients⁷⁴. Previously, no influence of morbid obesity was found on protein binding or on trough concentrations of cefazolin, whereas a modest influence of obesity was found on cefazolin peak concentrations upon an intravenous bolus administration⁹⁴. The results of the microdialysis study show that cefazolin penetration into the subcutaneous tissue over 4 h after dosing in obese patients was reduced by 30% on average (Figure 3).

These results were explained by reduced distribution of cefazolin to the subcutaneous tissue, which was found to depend on body weight, while there was no evidence for an increased peripheral volume of distribution represented by the subcutaneous tissue compartment⁷⁴. Instead, the value of the central volume of distribution was found to depend on body weight, and there was no influence of weight on clearance. Because time above the minimal inhibitory concentration at the target site is relevant for cefazolin prophylaxis, these findings have important consequences for the dosing regimen, particularly for the heaviest patients⁷⁴. In this respect, it is also important to take into account that obesity is an independent risk factor for postoperative surgical site infection^{7-8,95}.

Example 2: Nadroparin

A second example concerns anti-Xa levels, which Diepstraten et al.⁹⁶ measured to evaluate the effect of nadroparin in morbidly obese patients (107-260 kg). Prophylactic




Figure 3 Concentrations of (a) subcutaneous interstitial space fluid (ISF) cefazolin and (b) unbound plasma cefazolin in morbidly obese (blue, $n=7$ for panel a and $n=8$ for panel b) and nonobese (red, $n=7$ for both panels) patients. Figure adapted from Reference 74 with permission.

ranges have been defined for anti-Xa levels 4 h after subcutaneous dosing⁹⁷⁻⁹⁸. Volume of distribution is an essential parameter to determine the optimal dose for nadroparin. Upon subcutaneous administration, anti-Xa levels correlated best with LBW rather than BMI or total body weight, so dose adjustments on the basis of LBW are proposed⁹⁶.

An explanation for the finding that LBW should be used to dose low-molecular-weight heparins such as nadroparin could be that anti-Xa is a large, hydrophilic molecule that mainly distributes over vascular tissue and blood. Investigators have previously reported that blood volume increases with body weight in a nonlinear manner²², which probably corresponds to LBW. Also, researchers have proposed to adjust the dose for enoxaparin, another low-molecular-weight heparin, in obese individuals on the basis of LBW⁹⁹. Optimal dosing of low-molecular-weight heparins in obese individuals is particularly important because these individuals are at increased risk for venous thrombosis embolisms¹⁰⁰.

Example 3: Atracurium

As a third example, we present a pharmacodynamic study on atracurium in morbidly obese patients ($\text{BMI} > 40 \text{ kg/m}^2$, body weight 112-260 kg)⁶². Patients were randomized to receive atracurium on the basis of IBW or total body weight (TBW). Dosing on the basis of IBW resulted in a predictable profile of muscle relaxation, allowing for adequate intubation conditions and recovery of muscle strength within 60 min. In the patients for whom the dose was individualized on the basis of TBW, a dose-dependent prolongation of action was shown (Figure 4); thus, van Kralingen et al⁶² concluded that atracurium should be dosed on IBW.

In this example, changes in both pharmacokinetics (volume of distribution, clearance), and pharmacodynamics may have contributed to these results. Similar results

have previously been reported for rocuronium⁶³⁻⁶⁴. Remarkably, these results have led to an IBW-based dosing advice for rocuronium in the European label, whereas in the United States, rocuronium is still advised to be dosed on total bodyweight⁹⁰.




Figure 4 Effect of atracurium expressed as time to recovery of the twitch response of the neuromuscular train-of-four (TOF) to 5% vs. dose for morbidly obese patients dosed 0.5 mg/kg based on ideal body weight (orange squares, $n=8$) and dosed 0.5 mg/kg based on total body weight (green triangles, $n=9$). Figure adapted from Reference 62 with permission.

From this overview, it seems that the current level of understanding of the comprehensive effect of obesity on volume of distribution is limited. Although volume of distribution often changes with obesity, the direction and magnitude is not always predictable^{79,90}, despite many efforts to correlate it to physicochemical properties^{17,90-92}. When no information is available, extrapolation on the basis of total body weight with an estimated allometric exponent from results in normal-weight subjects seems preferable⁷⁹.

THE INFLUENCE OF OBESITY ON DRUG METABOLISM AND EXCRETION

Typically, there is more attention for the influence of obesity on metabolic and elimination clearance than on drug distribution^{79,101-103}. This may be explained by the fact that drug clearance is considered the most important pharmacokinetic parameter because it determines the maintenance dose of drugs.

A systematic review on reported clearance values of drugs in both obese and non-obese patients showed that the influence of obesity on drug metabolism and elimina-

tion differs between specific metabolic or elimination pathways¹⁰¹, even though the magnitude of its influence seems relatively small compared to the influence of obesity on distribution⁷⁹. Overall, the clearance of drugs primarily metabolized through the Phase II metabolism enzyme uridine diphosphate glucuronosyltransferase is reported to increase with obesity. For drugs that are eliminated through Phase I metabolism, the changes may differ depending on the pertinent enzyme. For example, an increased CYP2E1 clearance, a lower CYP3A clearance, and a trend toward higher clearance of CYP1A2, CYP2C9, CYP2C19, and CYP2D6 substrates have been reported¹⁰¹. In agreement with these literature findings, oral clearances were successfully predicted for eight drugs that are primarily cleared by CYP3A, CYP1A2, CYP2E1, and CYP2C9 on the basis of physiologically based pharmacokinetic modelling, in which known alterations in physiology resulting from obesity are implemented¹⁰³. More specifically, seven out of nine cases (involving eight drugs) were within 2-fold of the actual ratio between clearance in obese and lean patients¹⁰³. Remarkably, in this study, oral clearances of the CYP3A substrates alprazolam, midazolam, triazolam, and cyclosporine in the obese were somewhat over-predicted compared to observed oral clearance values, which were expected to be lower in the obese¹⁰³. As for midazolam, similar systemic clearance and higher bioavailability in morbidly obese patients were recently reported⁸⁷; it is emphasized that oral clearance equals CL/F and that reported differences in oral clearance in the obese may result from differences in systemic clearance, bioavailability, or both. Therefore, investigators should take care to predict systemic clearance on the basis of oral data as long as limited information is available on drug absorption in the obese.

With respect to renal clearance, higher values are reported in obese individuals^{35,101}. Recent results on the renally excreted antibiotic cefazolin in morbidly obese patients undergoing bariatric surgery did not identify an influence of body weight on cefazolin clearance, however^{74,94}. Even though this finding may be an artefact resulting from the relatively short sampling time in the study, a lack of change in glomerular filtration rate in obese individuals without microalbuminuria has been reported before³⁸, emphasizing that renal clearance of drugs may not necessarily be increased.

Concerning drug clearance mediated by liver blood flow, higher values were reported for a small number of high-extraction-ratio drugs with clearance values of more than 1.5 L/min¹⁰¹, which confirm early reports on increased hepatic flow in obese patients¹⁹.

Recently, Mahmood⁷⁹ has used an allometric equation to scale the pharmacokinetics of 12 drugs that are eliminated through different routes between healthy normal-weight subjects and obese patients. The results of this study indicate that clearances of these 12 different drugs increase in a nonlinear manner with total body weight⁷⁹, confirming a previous report⁵⁶. Clearance in the obese could be predicted with accuracy from normal-weight subjects using total body weight and simple allometry if an allometric exponent was estimated within the normal-weight population⁷⁹. In addition, allometric scaling

with a fixed exponent of 0.75 or 1.0 was found to be inferior to the allometric model in which the exponent was estimated. Mahmood⁷⁹ also states that obesity may not have an impact on clearance at all, as was the case for phenazone, carbamazepine, lithium, remifentanyl, cefazolin, and theophylline; thus, we emphasize that allometric scaling using a fixed exponent of 0.75 or 1.0 on the basis of results from normal-weight patients should not be applied unless more data become available. This argument also applies to the proposal to scale clearance with LBW with an exponent of 2/3, independent of the drug's primary route of metabolism and elimination¹⁰², as this approach assumes an increase in clearance with obesity, which may not be the case for all drugs^{79,104}.

In conclusion, for clearance, the influence of obesity seems smaller and somewhat easier to predict compared to alterations in volume of distribution, even though many questions remain on the exact quantification¹⁰¹. From the results presented here, it seems that predictions can be made on the basis of the primary pathway involved^{101,103}. When no information is available, extrapolation on the basis of total body weight with an estimated allometric exponent from results in normal-weight subjects seems preferable⁷⁹.

CHARACTERIZATION OF THE INFLUENCE OF OBESITY IN CHILDREN

Despite the increasing numbers of obese and morbidly obese children, very limited pharmacokinetic and dosing information in obese children is available¹⁰⁵⁻¹⁰⁷. A specific aspect that investigators, regulators, and prescribers should consider when determining dosing guidelines for obese children and adolescents is that, in general paediatric practice, dosing regimens are expressed in mg/kg. This linear mg/kg-based dosing is subject to debate even in normal-weight children between 0 and 18 years¹⁰⁸⁻¹¹², but an overdose may be anticipated if the dosing is based on mg/kg total body weight in overweight and, particularly, obese and morbidly children. This underscores the need to develop dedicated models for obese and morbidly obese children and adolescents⁷⁸. Performing these studies in the target population of obese individuals is even more relevant given that differences in pharmacokinetics, pharmacodynamics, or even the disease itself may exist in this population¹²⁻¹³.

In view of the limited number of pharmacokinetic studies in obese children^{101,113-114}, we present two pharmacokinetic studies in which data from overweight and obese children (and adults) of a large age range, along with their controls, are analyzed. In obese children, total body weight can be considered to be composed of both weight resulting from growth and development and weight from varying levels of obesity. This raises the question of, for instance, whether an obese 9-year-old child weighing 60 kg – in whom part of this body weight is physiological weight, i.e., body weight conforming to his age, and the other part is overweight – should receive the same dose as a normal-weight

16-year-old individual of the same weight. The distinction between physiological weight and overweight should be kept in mind when weight is studied as a covariate in children of varying ages and varying degrees of obesity.

Example 1: Propofol

For propofol, researchers performed a population pharmacokinetic meta-analysis with data from morbidly obese adults, adolescents, and children and their nonobese controls (body weight 37–184 kg, age 9–79 years)⁷⁷. In this analysis, propofol clearance was found to increase with body weight according to a power function. Age was identified and implemented as a second covariate using a bilinear function with two distinct slopes, reflecting an initial increase and, at the age of 41 years, a subsequent decrease in clearance (Figure 5).




Figure 5 Individual post hoc propofol clearance estimates vs. total body weight for morbidly obese adults and their nonobese controls (red circles) and obese adolescents and children and their nonobese controls (brown circles) ($n=94$). The blue dashed lines indicate the population clearance values for 15, 41, and 65 years. Figure adapted from Reference 77 with permission.

Example 2: Busulfan

In another study, investigators determined busulfan concentrations from a large population of underweight, normal-weight, and overweight children, adolescents, and adults (0.1–35 years)¹¹⁵. This study used a previously derived, body weight--driven, pharmacokinetic model for busulfan in children of all ages¹¹⁶. The results showed that the derived model¹¹⁶ proved equally predictive in normal-weight, underweight, and overweight children¹¹⁵. In addition, Bartelink et al. developed an exploratory model in which the body weight of each patient was considered to be composed of two parts: (a) physiological body weight related to growth (mean body weight-for-age) and (b) overweight,

i.e., body weight related to under/overweight for a certain age (body weight Z-score) (Figure 6). Despite adequate performance of this exploratory model in which weight as a result of growth and obesity was disentangled (Figure 6), the model was not superior over the simple, weight-based model¹¹⁵⁻¹¹⁶.

To capture the entire developmental change in clearance across the paediatric age range, this pharmacokinetic analysis of busulfan in over- and underweight children of all ages used an advanced power function based on body weight in which the exponent was allowed to change with body weight¹¹⁶⁻¹¹⁷. This advanced power function was needed because very young infants were also included in the busulfan analysis, whereas the propofol analysis did not consider children younger than 9 years of age⁷⁷. When this function was used for busulfan, the data were adequately described, and no influence of age could be identified. In contrast, for propofol, a bilinear, age-based function with two distinct slopes was found (Figure 5)⁷⁷. The reason for this difference may be in part that, for the busulfan analysis, no patients above 35 years were included¹¹⁵. For busulfan, these results imply that within the ranges of age and weight studied, dosing in children can be based on actual body weight, irrespective of the level of over- or underweight¹¹⁵⁻¹¹⁶.

In conclusion, although very limited pharmacokinetic and dosing information is available in obese children¹⁰⁵⁻¹⁰⁷, we present two approaches on how to analyse data from children varying in age and degree of obesity (Figures 5 and 6). Future clinical studies should focus on the pharmacokinetics and pharmacodynamics of commonly used




Figure 6 Busulfan clearance vs. mean body weight-for-age for an exploratory model of overweight and underweight for children of all ages. In the model, a function for body weight due to growth (described using mean body weight-for-age, blue line) and a function for body weight due to under- and overweight (described using the body weight Z-score, orange lines) were implemented. Orange lines represent body weight Z-scores of +1 (dark orange) and +2 (light orange), and purple lines represent body weight Z-scores of -1 (dark purple) and -2 (light purple). Figure adapted from Reference 115 with permission.

drugs in obese and morbidly obese children and adolescents to expand our knowledge in this clinically important area. Such studies should perform proper evaluations of the exact influence of weight resulting from growth, obesity, and age; these evaluations may be complicated because of the interrelation between weight and age in different manners, and they should use advanced validation frameworks, such as those described for paediatric pharmacokinetic analyses ¹¹⁸.

PERSPECTIVES

To predict the optimal dose for each drug in the obese, not only well-designed clinical studies on drug disposition in obese adults and children upon oral and intravenous administration are needed. Future research should also focus on the characterization of physiological concepts that can be used across drugs. From this overview, it is clear that for none of the parameters bioavailability, volume of distribution or clearance, a general covariate model with one size descriptor and one allometric exponent can be defined without paying attention to the nature of the compound involved, including the route of elimination. In this respect, physiologically based modelling principles that take into account both drug characteristics and physiological changes in the obese body are of large importance.

For obesity-related changes in clearance, a recently reported, semiphysiological approach applied in children, in which information for one drug was used to predict changes for another drug sharing the same metabolic or elimination pathway, may deserve attention. Using this approach, the maturation function for glucuronidation of morphine in young children ¹¹⁹⁻¹²⁰ was found to adequately predict the maturation in zidovudine glucuronidation in infants ¹²¹. As the physicochemical drug parameters were not found to affect this maturation profile, researchers concluded that this maturation function for glucuronidation can also be used for other substrates of this enzyme ¹²². This approach of between-drug predictions was also applied to renally excreted drugs in 0.5-5 kg neonates on the basis of a model derived for amikacin ¹²³. This model has recently been extended to older children and adults ¹²⁴ to obtain adequate predictions for other renally excreted drugs ¹²⁵⁻¹²⁶.

To predict volumes of distribution in the obese, investigators need to take into account both physicochemical properties and physiological changes in the obese body. Most recently, a new covariate relation that integrates body weight and LBW as covariates, with a weighting factor depending on the physicochemical properties of the drug, was proposed to predict volume of distribution at steady state ¹²⁷. Even though this approach was applied to only a limited number of obese individuals weighing below 100 kg, it deserves further exploration in the obese population, particularly because this

approach to covariate modelling led to similar results as a whole-body, physiologically based pharmacokinetic model¹²⁷.

CONCLUSION

In conclusion, although studies are particularly needed on absorption and distribution of drugs in obese individuals, some insight has been gained into changes in important metabolic and elimination pathways in obesity. For obese children, investigators need to perform clinical studies for which the proposed models^{77,115} can be used to analyze the data. Future research should focus on the characterization of physiological concepts to predict the optimal dose for each drug in the obese.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding or financial holdings that might be perceived as affecting the objectivity of this review.

REFERENCES

1. Flegal KM, Carroll MD, Kit BK, Ogden CL: Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *Jama* 2012; 307: 491-7
2. Ogden CL, Carroll MD, Kit BK, Flegal KM: Prevalence of obesity and trends in body mass index among US children and adolescents, 1999-2010. *Jama* 2012; 307: 483-90
3. World, Obes., Fed.: World map of obesity, 2014
4. Haslam DW, James WP: Obesity. *Lancet* 2005; 366: 1197-209
5. Stone AA, Broderick JE: Obesity and pain are associated in the United States. *Obesity (Silver Spring)* 2012; 20: 1491-5
6. McCarthy LH, Bigal ME, Katz M, Derby C, Lipton RB: Chronic pain and obesity in elderly people: results from the Einstein aging study. *J Am Geriatr Soc* 2009; 57: 115-9
7. Choban PS, Heckler R, Burge JC, Flancbaum L: Increased incidence of nosocomial infections in obese surgical patients. *Am Surg* 1995; 61: 1001-5
8. Huttunen R, Karppele M, Syrjanen J: Obesity and nosocomial infections. *J Hosp Infect* 2013; 85: 8-16
9. Oyeturji TA, Franklin AL, Ortega G, Akolkar N, Qureshi FG, Abdullah F, Cornwell EE, Nwomeh BC, Fullum TM: Revisiting childhood obesity: persistent underutilization of surgical intervention? *Am Surg* 2012; 78: 788-93
10. Schilling PL, Davis MM, Albanese CT, Dutta S, Morton J: National trends in adolescent bariatric surgical procedures and implications for surgical centers of excellence. *J Am Coll Surg* 2008; 206: 1-12
11. Black MH, Zhou H, Takayanagi M, Jacobsen SJ, Koebnick C: Increased asthma risk and asthma-related health care complications associated with childhood obesity. *Am J Epidemiol* 2013; 178: 1120-8
12. Forno E, Lescher R, Strunk R, Weiss S, Fuhlbrigge A, Celedon JC: Decreased response to inhaled steroids in overweight and obese asthmatic children. *J Allergy Clin Immunol* 2011; 127: 741-9
13. Gelelete CB, Pereira SH, Azevedo AM, Thiago LS, Mundim M, Land MG, Costa ES: Overweight as a prognostic factor in children with acute lymphoblastic leukemia. *Obesity (Silver Spring)* 2011; 19: 1908-11
14. Conroy S, Choonara I, Impicciatore P, Mohn A, Arnell H, Rane A, Knoeppel C, Seyberth H, Pandolfini C, Raffaelli MP, Rocchi F, Bonati M, Jong G, de Hoog M, van den Anker J: Survey of unlicensed and off label drug use in paediatric wards in European countries. *European Network for Drug Investigation in Children. Bmj* 2000; 320: 79-82
15. Ernest TB, Elder DP, Martini LG, Roberts M, Ford JL: Developing paediatric medicines: identifying the needs and recognizing the challenges. *J Pharm Pharmacol* 2007; 59: 1043-55
16. t Jong GW, Vulto AG, de Hoog M, Schimmel KJ, Tibboel D, van den Anker JN: A survey of the use of off-label and unlicensed drugs in a Dutch children's hospital. *Pediatrics* 2001; 108: 1089-93
17. Cheymol G: Clinical pharmacokinetics of drugs in obesity. An update. *Clin Pharmacokinet* 1993; 25: 103-14
18. Cheymol G: Effects of obesity on pharmacokinetics implications for drug therapy. *Clin Pharmacokinet* 2000; 39: 215-31
19. Alexander JK, Dennis EW, Smith WG, Amad KH, Duncan WC, Austin RC: Blood volume, cardiac output, and distribution of systemic blood flow in extreme obesity. *Cardiovasc Res Cent Bull* 1962; 1: 39-44

20. Licata G, Scaglione R, Barbagallo M, Parrinello G, Capuana G, Lipari R, Merlino G, Ganguzza A: Effect of obesity on left ventricular function studied by radionuclide angiocardiology. *Int J Obes* 1991; 15: 295-302
21. Herrera MF, Deitel M: Cardiac function in massively obese patients and the effect of weight loss. *Can J Surg* 1991; 34: 431-4
22. Lemmens HJ, Bernstein DP, Brodsky JB: Estimating blood volume in obese and morbidly obese patients. *Obes Surg* 2006; 16: 773-6
23. Blouin RA, Kolpek JH, Mann HJ: Influence of obesity on drug disposition. *Clin Pharm* 1987; 6: 706-14
24. Crocker DW: Lipomatous infiltrates of the heart. *Arch Pathol Lab Med* 1978; 102: 69-72
25. Bharati S, Lev M: Cardiac conduction system involvement in sudden death of obese young people. *Am Heart J* 1995; 129: 273-81
26. Jones RL, Nzekwu MM: The effects of body mass index on lung volumes. *Chest* 2006; 130: 827-33
27. Rajala R, Partinen M, Sane T, Pelkonen R, Huikuri K, Seppalainen AM: Obstructive sleep apnoea syndrome in morbidly obese patients. *J Intern Med* 1991; 230: 125-9
28. Guzzaloni G, Grugni G, Minocci A, Moro D, Morabito F: Liver steatosis in juvenile obesity: correlations with lipid profile, hepatic biochemical parameters and glycemic and insulinemic responses to an oral glucose tolerance test. *Int J Obes Relat Metab Disord* 2000; 24: 772-6
29. Moretto M, Kupski C, Mottin CC, Repetto G, Garcia Toneto M, Rizzolli J, Berleze D, de Souza Brito CL, Casagrande D, Colossi F: Hepatic steatosis in patients undergoing bariatric surgery and its relationship to body mass index and co-morbidities. *Obes Surg* 2003; 13: 622-4
30. Ijaz S, Yang W, Winslet MC, Seifalian AM: Impairment of hepatic microcirculation in fatty liver. *Microcirculation* 2003; 10: 447-56
31. Farrell GC, Teoh NC, McCuskey RS: Hepatic microcirculation in fatty liver disease. *Anat Rec (Hoboken)* 2008; 291: 684-92
32. Casati A, Putzu M: Anesthesia in the obese patient: pharmacokinetic considerations. *J Clin Anesth* 2005; 17: 134-45
33. Johnson TN, Tucker GT, Tanner MS, Rostami-Hodjegan A: Changes in liver volume from birth to adulthood: a meta-analysis. *Liver Transpl* 2005; 11: 1481-93
34. Kotlyar M, Carson SW: Effects of obesity on the cytochrome P450 enzyme system. *Int J Clin Pharmacol Ther* 1999; 37: 8-19
35. Janmahasatian S, Duffull SB, Chagnac A, Kirkpatrick CM, Green B: Lean body mass normalizes the effect of obesity on renal function. *Br J Clin Pharmacol* 2008; 65: 964-5
36. Ribstein J, du Cailar G, Mimran A: Combined renal effects of overweight and hypertension. *Hypertension* 1995; 26: 610-5
37. Marik P, Varon J: The obese patient in the ICU. *Chest* 1998; 113: 492-8
38. Anastasio P, Spitali L, Frangiosa A, Molino D, Stellato D, Cirillo E, Pollastro RM, Capodicasa L, Sepe J, Federico P, Gaspare De Santo N: Glomerular filtration rate in severely overweight normotensive humans. *Am J Kidney Dis* 2000; 35: 1144-8
39. O'Donnell MP, Kasiske BL, Cleary MP, Keane WF: Effects of genetic obesity on renal structure and function in the Zucker rat. II. Micropuncture studies. *J Lab Clin Med* 1985; 106: 605-10
40. Kasiske BL, Cleary MP, O'Donnell MP, Keane WF: Effects of genetic obesity on renal structure and function in the Zucker rat. *J Lab Clin Med* 1985; 106: 598-604
41. Schmitz PG, O'Donnell MP, Kasiske BL, Katz SA, Keane WF: Renal injury in obese Zucker rats: glomerular hemodynamic alterations and effects of enalapril. *Am J Physiol* 1992; 263: F496-502

42. Kasiske BL, Crosson JT: Renal disease in patients with massive obesity. *Arch Intern Med* 1986; 146: 1105-9
43. Pai MP: Estimating the glomerular filtration rate in obese adult patients for drug dosing. *Adv Chronic Kidney Dis* 2010; 17: e53-62
44. Demirovic JA, Pai AB, Pai MP: Estimation of creatinine clearance in morbidly obese patients. *Am J Health Syst Pharm* 2009; 66: 642-8
45. Wuerzner G, Bochud M, Giusti V, Burnier M: Measurement of glomerular filtration rate in obese patients: pitfalls and potential consequences on drug therapy. *Obes Facts* 2011; 4: 238-43
46. Lim WH, Lim EM, McDonald S: Lean body mass-adjusted Cockcroft and Gault formula improves the estimation of glomerular filtration rate in subjects with normal-range serum creatinine. *Nephrology* 2006; 11: 250-6
47. Cardoso-Junior A, Coelho LG, Savassi-Rocha PR, Vignolo MC, Abrantes MM, de Almeida AM, Dias EE, Vieira Junior G, de Castro MM, Lemos YV: Gastric emptying of solids and semi-solids in morbidly obese and non-obese subjects: an assessment using the ¹³C-octanoic acid and ¹³C-acetic acid breath tests. *Obes Surg* 2007; 17: 236-41
48. Tosetti C, Corinaldesi R, Stanghellini V, Pasquali R, Corbelli C, Zoccoli G, Di Febo G, Monetti N, Barbara L: Gastric emptying of solids in morbid obesity. *Int J Obes Relat Metab Disord* 1996; 20: 200-5
49. Wright RA, Krinsky S, Fleeman C, Trujillo J, Teague E: Gastric emptying and obesity. *Gastroenterology* 1983; 84: 747-51
50. Xing J, Chen JD: Alterations of gastrointestinal motility in obesity. *Obes Res* 2004; 12: 1723-32
51. Teixeira TF, Souza NC, Chiarello PG, Franceschini SC, Bressan J, Ferreira CL, Peluzio Mdo C: Intestinal permeability parameters in obese patients are correlated with metabolic syndrome risk factors. *Clin Nutr* 2012; 31: 735-40
52. Horton F, Wright J, Smith L, Hinton PJ, Robertson MD: Increased intestinal permeability to oral chromium (51 Cr) -EDTA in human Type 2 diabetes. *Diabet Med* 2013
53. French SJ, Murray B, Rumsey RD, Sepple CP, Read NW: Preliminary studies on the gastrointestinal responses to fatty meals in obese people. *Int J Obes Relat Metab Disord* 1993; 17: 295-300
54. Wisen O, Johansson C: Gastrointestinal function in obesity: motility, secretion, and absorption following a liquid test meal. *Metabolism* 1992; 41: 390-5
55. WHO: Global database on BMI classification World Health Organization 2006
56. Green B, Duffull SB: What is the best size descriptor to use for pharmacokinetic studies in the obese? *Br J Clin Pharmacol* 2004; 58: 119-33
57. Eleveld DJ, Proost JH, Absalom AR, Struys MM: Obesity and allometric scaling of pharmacokinetics. *Clin Pharmacokinet* 2011; 50: 751-3
58. Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, Wei R, Curtin LR, Roche AF, Johnson CL: 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat* 11 2002; 246: 1-190
59. http://www.cdc.gov/growthcharts/percentile_data_files.htm. Accessed 05-03-2014 Centers for Disease Control and Prevention
60. Barlow SE: Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* 2007; 120 Suppl 4: S164-92
61. Devine J: Gentamycin Therapy. *Ann Pharmacother* 1974; 8: 650-655

62. van Kralingen S, van de Garde EM, Knibbe CA, Diepstraten J, Wiezer MJ, van Ramshorst B, van Dongen EP: Comparative evaluation of atracurium dosed on ideal body weight vs. total body weight in morbidly obese patients. *Br J Clin Pharmacol* 2011; 71: 34-40
63. Leykin Y, Pellis T, Lucca M, Lomangino G, Marzano B, Gullo A: The pharmacodynamic effects of rocuronium when dosed according to real body weight or ideal body weight in morbidly obese patients. *Anesth Analg* 2004; 99: 1086-9,
64. Meyhoff CS, Lund J, Jenstrup MT, Claudius C, Sorensen AM, Viby-Mogensen J, Rasmussen LS: Should dosing of rocuronium in obese patients be based on ideal or corrected body weight? *Anesth Analg* 2009; 109: 787-92
65. Egan TD, Huizinga B, Gupta SK, Jaarsma RL, Sperry RJ, Yee JB, Muir KT: Remifentanyl pharmacokinetics in obese vs. lean patients. *Anesthesiology* 1998; 89: 562-73
66. Pai MP: Drug dosing based on weight and body surface area: mathematical assumptions and limitations in obese adults. *Pharmacotherapy* 2012; 32: 856-68
67. van Rongen A, Brill MJ, Diepstraten J, Knibbe CA: Applied pharmacometrics in the obese population, *Applied Pharmacometrics*. USA Springer. In press
68. Du Bois D, Du Bois EF: A formula to estimate the approximate surface area if height and weight be known. *Arch Int Med* 1916; 17: 863-71
69. Mosteller RD: Simplified calculation of body-surface area. *N Engl J Med* 1987; 317
70. Griggs JJ, Mangu PB, Anderson H, Balaban EP, Dignam JJ, Hryniuk WM, Morrison VA, Pini TM, Runowicz CD, Rosner GL, Shayne M, Sparreboom A, Sucheston LE, Lyman GH: Appropriate chemotherapy dosing for obese adult patients with cancer: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol* 2012; 30: 1553-61
71. Sparreboom A, Wolff AC, Mathijssen RH, Chatelut E, Rowinsky EK, Verweij J, Baker SD: Evaluation of alternate size descriptors for dose calculation of anticancer drugs in the obese. *J Clin Oncol* 2007; 25: 4707-13
72. Janmahasatian S, Duffull SB, Ash S, Ward LC, Byrne NM, Green B: Quantification of lean bodyweight. *Clin Pharmacokinet* 2005; 44: 1051-65
73. Han PY, Duffull SB, Kirkpatrick CM, Green B: Dosing in obesity: a simple solution to a big problem. *Clin Pharmacol Ther* 2007; 82: 505-8
74. Brill MJ, Houwink AP, Schmidt S, Van Dongen EP, Hazebroek EJ, van Ramshorst B, Deneer VH, Mouton JW, Knibbe CA: Reduced subcutaneous tissue distribution of cefazolin in morbidly obese versus non-obese patients determined using clinical microdialysis. *J Antimicrob Chemother* 2014; 69: 715-23
75. van Kralingen S, Diepstraten J, Peeters MY, Deneer VH, van Ramshorst B, Wiezer RJ, van Dongen EP, Danhof M, Knibbe CA: Population pharmacokinetics and pharmacodynamics of propofol in morbidly obese patients. *Clin Pharmacokinet* 2011; 50: 739-50
76. Peters AM, Snelling HL, Glass DM, Bird NJ: Estimation of lean body mass in children. *Br J Anaesth* 2011; 106: 719-23
77. Diepstraten J, Chidambaran V, Sadhasivam S, Blusse van Oud-Alblas HJ, Inge T, van Ramshorst B, van Dongen EP, Vinks AA, Knibbe CA: An integrated population pharmacokinetic meta-analysis of propofol in morbidly obese and nonobese adults, adolescents, and children. *CPT Pharmacometrics Syst Pharmacol* 2013; 2: e73
78. Diepstraten J, Chidambaran V, Sadhasivam S, Esslinger HR, Cox SL, Inge TH, Knibbe CA, Vinks AA: Propofol clearance in morbidly obese children and adolescents: influence of age and body size. *Clin Pharmacokinet* 2012; 51: 543-51

79. Mahmood I: Prediction of clearance and volume of distribution in the obese from normal weight subjects: an allometric approach. *Clin Pharmacokinet* 2012; 51: 527-42
80. Bowman SL, Hudson SA, Simpson G, Munro JF, Clements JA: A comparison of the pharmacokinetics of propranolol in obese and normal volunteers. *Br J Clin Pharmacol* 1986; 21: 529-32
81. Greenblatt DJ, Abernethy DR, Locniskar A, Harmatz JS, Limjuco RA, Shader RI: Effect of age, gender, and obesity on midazolam kinetics. *Anesthesiology* 1984; 61: 27-35
82. Greenblatt DJ, Friedman H, Burstein ES, Scavone JM, Blyden GT, Ochs HR, Miller LG, Harmatz JS, Shader RI: Trazodone kinetics: effect of age, gender, and obesity. *Clin Pharmacol Ther* 1987; 42: 193-200
83. Flechner SM, Kolbeinsson ME, Tam J, Lum B: The impact of body weight on cyclosporine pharmacokinetics in renal transplant recipients. *Transplantation* 1989; 47: 806-10
84. Cheymol G, Weissenburger J, Poirier JM, Gellee C: The pharmacokinetics of dexfenfluramine in obese and non-obese subjects. *Br J Clin Pharmacol* 1995; 39: 684-7
85. Kees MG, Weber S, Kees F, Horbach T: Pharmacokinetics of moxifloxacin in plasma and tissue of morbidly obese patients. *J Antimicrob Chemother* 2011; 66: 2330-5
86. Abernethy DR, Greenblatt DJ, Divoll M, Smith RB, Shader RI: The influence of obesity on the pharmacokinetics of oral alprazolam and triazolam. *Clin Pharmacokinet* 1984; 9: 177-83
87. Brill M, van Rongen A, Houwink A, Burggraaf J, van Ramshorst B, Wiezer R, Van Dongen E, Knibbe C: Midazolam pharmacokinetics in morbidly obese patients following semi-simultaneous oral and intravenous administration: a comparison with healthy volunteers. *Clin Pharmacokinet* 2014; In press
88. Yang J, Jamei M, Yeo KR, Tucker GT, Rostami-Hodjegan A: Prediction of intestinal first-pass drug metabolism. *Curr Drug Metab* 2007; 8: 676-8
89. Rostami-Hodjegan A, Tucker GT: The effects of portal shunts on intestinal cytochrome P450 3A activity. *Hepatology* 2002; 35: 1549-50; author reply 1550-1
90. Jain R, Chung SM, Jain L, Khurana M, Lau SW, Lee JE, Vaidyanathan J, Zadezensky I, Choe S, Sahajwalla CG: Implications of obesity for drug therapy: limitations and challenges. *Clin Pharmacol Ther* 2011; 90: 77-89
91. Hanley MJ, Abernethy DR, Greenblatt DJ: Effect of obesity on the pharmacokinetics of drugs in humans. *Clin Pharmacokinet* 2010; 49: 71-87
92. Blouin RA, Warren GW: Pharmacokinetic considerations in obesity. *J Pharm Sci* 1999; 88: 1-7
93. Abernethy DR, Greenblatt DJ, Divoll M, Harmatz JS, Shader RI: Alterations in drug distribution and clearance due to obesity. *J Pharmacol Exp Ther* 1981; 217: 681-5
94. van Kralingen S, Taks M, Diepstraten J, van de Garde EM, van Dongen EP, Wiezer MJ, van Ramshorst B, Vlamincx B, Deneer VH, Knibbe CA: Pharmacokinetics and protein binding of cefazolin in morbidly obese patients. *Eur J Clin Pharmacol* 2011; 67: 985-92
95. Falagas ME, Kompoti M: Obesity and infection. *Lancet Infect Dis* 2006; 6: 438-46
96. Diepstraten J, Hackeng CM, van Kralingen S, Zapletal J, van Dongen EP, Wiezer RJ, van Ramshorst B, Knibbe CA: Anti-Xa Levels 4 h After Subcutaneous Administration of 5,700 IU Nadroparin Strongly Correlate with Lean Body Weight in Morbidly Obese Patients. *Obes Surg* 2012
97. Hirsh J, Raschke R: Heparin and low-molecular-weight heparin: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126: 188S-203S
98. Nutescu EA, Spinler SA, Wittkowsky A, Dager WE: Low-molecular-weight heparins in renal impairment and obesity: available evidence and clinical practice recommendations across medical and surgical settings. *Ann Pharmacother* 2009; 43: 1064-83

99. Barras MA, Duffull SB, Atherton JJ, Green B: Individualized compared with conventional dosing of enoxaparin. *Clin Pharmacol Ther* 2008; 83: 882-8
100. Stein PD, Beemath A, Olson RE: Obesity as a risk factor in venous thromboembolism. *Am J Med* 2005; 118: 978-80
101. Brill MJ, Diepstraten J, van Rongen A, van Kralingen S, van den Anker JN, Knibbe CA: Impact of obesity on drug metabolism and elimination in adults and children. *Clin Pharmacokinet* 2012; 51: 277-304
102. McLeay SC, Morrish GA, Kirkpatrick CM, Green B: The relationship between drug clearance and body size: systematic review and meta-analysis of the literature published from 2000 to 2007. *Clin Pharmacokinet* 2012; 51: 319-30
103. Ghobadi C, Johnson TN, Aarabi M, Almond LM, Allabi AC, Rowland-Yeo K, Jamei M, Rostami-Hodjegan A: Application of a systems approach to the bottom-up assessment of pharmacokinetics in obese patients: expected variations in clearance. *Clin Pharmacokinet* 2011; 50: 809-22
104. Hall RG, 2nd, Jean GW, Sigler M, Shah S: Dosing considerations for obese patients receiving cancer chemotherapeutic agents. *Ann Pharmacother* 2013; 47: 1666-74
105. Mulla H, Johnson TN: Dosing dilemmas in obese children. *Arch Dis Child Educ Pract Ed* 2010; 95: 112-7
106. Kendrick JG, Carr RR, Ensom MH: Pharmacokinetics and drug dosing in obese children. *J Pediatr Pharmacol Ther* 2010; 15: 94-109
107. Mahmood I: Dosing in Children: A Critical Review of the Pharmacokinetic Allometric Scaling and Modelling Approaches in Paediatric Drug Development and Clinical Settings. *Clin Pharmacokinet* 2014
108. Knibbe CA, Danhof M: Individualized dosing regimens in children based on population PKPD modelling: are we ready for it? *Int J Pharm* 2011; 415: 9-14
109. Knibbe CA, Krekels EH, Danhof M: Advances in paediatric pharmacokinetics. *Expert Opin Drug Metab Toxicol* 2011; 7: 1-8
110. Admiraal R, van Kesteren C, Boelens JJ, Bredius RG, Tibboel D, Knibbe CA: Towards evidence-based dosing regimens in children on the basis of population pharmacokinetic pharmacodynamic modelling. *Arch Dis Child* 2014; 99: 267-72
111. Cella M, Knibbe C, Danhof M, Della Pasqua O: What is the right dose for children? *Br J Clin Pharmacol* 2010; 70: 597-603
112. De Cock RF, Piana C, Krekels EH, Danhof M, Allegaert K, Knibbe CA: The role of population PK-PD modelling in paediatric clinical research. *Eur J Clin Pharmacol* 2011; 67 Suppl 1: 5-16
113. Koshida R, Nakashima E, Taniguchi N, Tsuji A, Benet LZ, Ichimura F: Prediction of the distribution volumes of cefazolin and tobramycin in obese children based on physiological pharmacokinetic concepts. *Pharm Res* 1989; 6: 486-91
114. Heble DE, Jr., McPherson C, Nelson MP, Hunstad DA: Vancomycin trough concentrations in overweight or obese pediatric patients. *Pharmacotherapy* 2013; 33: 1273-7
115. Bartelink IH, van Kesteren C, Boelens JJ, Egberts TC, Bierings MB, Cuvelier GD, Wynn RF, Slatter MA, Chiesa R, Danhof M, Knibbe CA: Predictive performance of a busulfan pharmacokinetic model in children and young adults. *Ther Drug Monit* 2012; 34: 574-83
116. Bartelink IH, Boelens JJ, Bredius RG, Egberts AC, Wang C, Bierings MB, Shaw PJ, Nath CE, Hempel G, Zwaveling J, Danhof M, Knibbe CA: Body weight-dependent pharmacokinetics of busulfan in paediatric haematopoietic stem cell transplantation patients: towards individualized dosing. *Clin Pharmacokinet* 2012; 51: 331-45

117. Wang C, Peeters MY, Allegaert K, Blusse van Oud-Alblas HJ, Krekels EH, Tibboel D, Danhof M, Knibbe CA: A bodyweight-dependent allometric exponent for scaling clearance across the human life-span. *Pharm Res* 2012; 29: 1570-81
118. Krekels EH, van Hasselt JG, Tibboel D, Danhof M, Knibbe CA: Systematic evaluation of the descriptive and predictive performance of paediatric morphine population models. *Pharm Res* 2011; 28: 797-811
119. Knibbe CA, Krekels EH, van den Anker JN, DeJongh J, Santen GW, van Dijk M, Simons SH, van Lingen RA, Jacqz-Aigrain EM, Danhof M, Tibboel D: Morphine glucuronidation in preterm neonates, infants and children younger than 3 years. *Clin Pharmacokinet* 2009; 48: 371-85
120. Krekels EH, DeJongh J, van Lingen RA, van der Marel CD, Choonara I, Lynn AM, Danhof M, Tibboel D, Knibbe CA: Predictive performance of a recently developed population pharmacokinetic model for morphine and its metabolites in new datasets of (preterm) neonates, infants and children. *Clin Pharmacokinet* 2011; 50: 51-63
121. Krekels EH, Neely M, Panoilia E, Tibboel D, Capparelli E, Danhof M, Mirochnick M, Knibbe CA: From pediatric covariate model to semiphysiological function for maturation: part I-extrapolation of a covariate model from morphine to Zidovudine. *CPT Pharmacometrics Syst Pharmacol* 2012; 1: e9
122. Krekels EH, Johnson TN, den Hoedt SM, Rostami-Hodjegan A, Danhof M, Tibboel D, Knibbe CA: From Pediatric Covariate Model to Semiphysiological Function for Maturation: Part II-Sensitivity to Physiological and Physicochemical Properties. *CPT Pharmacometrics Syst Pharmacol* 2012; 1: e10
123. De Cock RF, Allegaert K, Schreuder MF, Sherwin CM, de Hoog M, van den Anker JN, Danhof M, Knibbe CA: Maturation of the glomerular filtration rate in neonates, as reflected by amikacin clearance. *Clin Pharmacokinet* 2012; 51: 105-17
124. De Cock RF, Allegaert K, Brussee JM, Sherwin CM, Mulla H, de Hoog M, van den Anker JN, Danhof M, Knibbe CA: Simultaneous pharmacokinetic modeling of gentamicin, tobramycin and vancomycin clearance from neonates to adults: towards a semi-physiological function for maturation in glomerular filtration. *Pharm Res.* 2014; In press
125. De Cock RF, Allegaert K, Sherwin CM, Nielsen EI, de Hoog M, van den Anker JN, Danhof M, Knibbe CA: A neonatal amikacin covariate model can be used to predict ontogeny of other drugs eliminated through glomerular filtration in neonates. *Pharm Res* 2014; 31: 754-67
126. Zhao W, Biran V, Jacqz-Aigrain E: Amikacin maturation model as a marker of renal maturation to predict glomerular filtration rate and vancomycin clearance in neonates. *Clin Pharmacokinet* 2013; 52: 1127-34
127. Huisinga W, Solms A, Fronton L, Pilari S: Modeling interindividual variability in physiologically based pharmacokinetics and its link to mechanistic covariate modeling. *CPT Pharmacometrics Syst Pharmacol* 2012; 1: e4

