



Universiteit
Leiden
The Netherlands

CHECK'D?! : determinants of participation in a two-stage cardiometabolic screening among underserved groups

Groenenberg, I.

Citation

Groenenberg, I. (2017, May 11). *CHECK'D?! : determinants of participation in a two-stage cardiometabolic screening among underserved groups*. Retrieved from <https://hdl.handle.net/1887/48563>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/48563>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/48563> holds various files of this Leiden University dissertation

Author: Groenenberg, Iris

Title: CHECK'D?! : determinants of participation in a two-stage cardiometabolic screening among underserved groups

Issue Date: 2017-05-11

4

Determinants of participation in a cardiometabolic health check among underserved groups

Iris Groenenberg¹, Mathilde R. Crone¹, Sandra van Dijk², Jamila Ben Meftah¹, Barend J.C. Middelkoop¹, Willem J.J. Assendelft³, and Anne M. Stiggelbout⁴

¹ Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, the Netherlands.

² Department of Social Sciences, Institute for Psychology, Health, Medical, and Neuropsychology, Leiden University, Leiden, the Netherlands.

³ Department of Primary and Community Care, Radboud University Medical Center, Nijmegen, the Netherlands.

⁴ Department of Medical Decision Making, Leiden University Medical Center, Leiden, the Netherlands.

Preventive Medicine Reports (2016), 4:33-43

ABSTRACT

Cardiometabolic diseases affect underserved groups disparately. Participation in health checks is also lower, widening health inequalities in society. Two-stage screening (non-invasive health risk assessment (HRA) and GP consultations for high-risk individuals) seems cost-effective, provided that drop-out rates are low in both steps. We aimed to explore the process of decision-making regarding HRA participation among underserved groups (45–70y): native Dutch with a lower socioeconomic status (SES), Turkish, Moroccan, and Surinamese participants. We conducted a cross-sectional questionnaire study. The questionnaire comprised the following determinants: a self-formulated first reaction, a structured set of predefined determinants, and the most important barrier(s) and facilitator(s) for HRA completion. We used univariable and (stepwise) multivariate logistic regression analyses to assess which determinants were associated with HRA completion. Of the 892 participants in the questionnaire, 78% (n=696) also completed the HRA. Moroccans and patients from GP practices with a predominantly non-Western population less often completed the HRA. A lower SES score, wanting to know one's risk, not remembering receiving the invitation (thus requiring a phone call), fear of the test result and/or adjusting lifestyle, perceived control of staying healthy, wanting to participate, and perceiving no barriers were associated with completing the HRA. We conclude that our 'hard-to-reach' population may not be unwilling to participate in the HRA. A more comprehensive approach, involving key figures within a community informing people about and providing help completing the HRA, would possibly be more suitable. Efforts should be particularly targeted at the less acculturated immigrants with an external locus of control.

INTRODUCTION

Cardiometabolic diseases (cardiovascular disease, diabetes, and kidney disease) are leading causes of death in high-income countries (1). An increased risk of cardiometabolic disease is associated with a lower socioeconomic status (SES) and ethnicity (2, 3). Among ethnic minorities in the Netherlands, cardiovascular disease is particularly prevalent among Surinamese and Turkish people (4 - 6). Turkish, Moroccan, and especially Hindustani Surinamese people have a higher risk of developing diabetes (7). To early identify individuals with an increased risk of cardiometabolic disease, health checks are implemented worldwide (8 - 10). Several studies concluded that two-stage screening could be a cost-effective strategy (11, 12). Two-stage screening usually refers to a non-invasive risk stratification tool, followed by a blood test during an assessment by a healthcare professional. The Dutch cardiometabolic health check imbedded in primary care follows this two-stage approach, comprising a short health risk assessment (HRA) to be completed at home, and two prevention consultations (PCs) with the GP for high-risk individuals according to the HRA (13). This approach implies that patients can refrain from participation on two separate occasions (14). High drop-out rates may induce an even greater problem among underserved groups, as ethnicity and SES are inversely related to health check attendance (15). These groups usually have greater difficulties in making an informed decision about participation (16). Presumably, higher participation rates in stage one (as a result of more informed decision-making) lead to higher participation rates in stage two. To increase informed decision-making about HRA completion, insight into its determinants plays a pivotal role. Few studies specifically investigated reasons for participation in cardiometabolic health checks of underserved groups. Studies reporting determinants in these populations exclusively focus on physical assessments at a doctor's office, not two-stage screening with risk stratification as a first step. Therefore, we conducted prior qualitative research on determinants of hypothetical HRA completion (17). These determinants were mainly of a cognitive nature and included (flawed) risk perceptions, health negligence, (health) illiteracy, and language barriers. With the current study we aim to explore the process of actual decision-making about HRA completion. Research questions were: (1) what are participants' self-formulated first reactions regarding the invitation?; (2) what predefined determinants play a role in completing the HRA?; (3) what are participants' most important barriers and

facilitators?; and (4) which of the aforementioned determinants are associated with actual HRA completion?

METHODS

Design and study population

This cross-sectional study is part of a larger study investigating reach and participation of underserved populations in the Dutch cardiometabolic health check.

Between May 2012 and December 2013, patients from six general practices were invited to participate. The six practices were located in The Hague and surroundings, and encompassed both large group as well as solo practices, and urban as well as rural environments. Patients had to be native Dutch with a lower SES or Turkish, Moroccan, or Surinamese. Ethnicity is not registered by GPs in the Netherlands, this was estimated by the researchers based on family name, and was subsequently checked by the GP. The GP also selected the native Dutch patients with a lower SES, which was afterwards corroborated with a neighbourhood SES score (average income, proportion of individuals with a low income, with a low education, and without a paid job) (18). These attributes are captured in one parameter: the socioeconomic status (SES) score and has been shown to be associated with deprivation in a community (19). This score is assessed every four years by interviewing persons representing nearly each street in the Netherlands. The average SES score in the Netherlands is 0.17.

Categorization of the SES scores was as follows: average to higher SES (score ≥ 0); lower to average SES (score 0 till -1.9); lower SES (score -2 till -3.9); lowest SES (score ≤ -4).

Patients had to be 45–70 years old except for the Hindustani Surinamese. Their lower age limit was 35 years because of their genetically increased risk of diabetes. Exclusion criteria were: having (had) cardiometabolic disease, using drugs against cardiometabolic disease, or having had a complete cardiometabolic risk inventory less than a year ago (Appendix A). All patients who met the eligibility criteria ($n=1644$) were invited.

Three culturally targeted and personalized invitation steps were tested following an increasingly (cost-)intensive ‘funnelled’ design: (1) all patients received a postal invitation; (2) non-reached were approached by telephone; (3) finally non-reached were approached face-to-face by their GP (Appendix B). The latter step was not included as participation rates were very low. Postal materials were provided in Dutch, and in Turkish/Arabic for

Turkish/Moroccan patients, and included the questionnaire and the HRA simultaneously in one package. Patients were called by Turkish, Arabic, and Berber (oral-only language) speaking research assistants.

Ethical approval was given by the Committee Medical Ethics from the Leiden University Medical Center. The study followed an 'opt-out procedure' where patients could sign a response form when not interested in participation. The design and results of the larger study have been described in detail elsewhere (20).

The study population of this study consisted of those patients who completed the determinants questionnaire (n = 892), divided into two groups: HRA completers and non-completers. Postal responders filled out a self-administered written questionnaire and telephone responders answered the questionnaire by phone.

The questionnaire

The postal- and telephone-administered questionnaires followed the same structure and were based on our previous work (17). This qualitative study was embedded in a theoretical framework based on the I-change model (Fig. 1), which aims to explain health behaviours and has been applied in studies among native and immigrant populations (21 - 24). The most important determinants in the qualitative study were turned into (simply formulated) questions. The questionnaire was pilot-tested among the target population. We incorporated three steps in the questionnaire: (1) a self-formulated first reaction regarding the invitation for the health check, (2) a structured set of predefined determinants that the participant could indicate to be of importance to his/her HRA completion, (3) most important barrier(s) and most important facilitator(s) regarding HRA completion (Appendix C).




Figure 1. The I-change model, from <http://www.maastricht-university.eu/hein.devries/interests/change>

Step one: self-formulated first reaction

The questionnaire started with one (open answer) question prompting participants to express their thoughts about the invitation. Any reaction was possible: from positive to negative attitudes regarding the initiative, and from practical barriers to positive social influences. This and the open answer questions of step three were coded inductively and converted into a code tree. Coding was performed by the first author and randomly double coded by the second author. Both authors categorized the codes in the code tree conform the I-change model constructs (Appendix D). Codes and constructs were entered into SPSS. The first reactions were then computed into dichotomized variables representing the different reactions (0 = not mentioned, 1 = mentioned).

Step two: structured set of predefined determinants

The questionnaire continued with a structured set of predefined (multiple-choice) determinant questions. These determinants were categorized under the appropriate I-change construct (see Results, Table 3). Most questions consisted of three answer categories (mostly ‘no’, ‘a little’, ‘yes’), which were dichotomized for a better distribution.

Step three: most important barriers and facilitators

The final two (open answer) questions aimed to unravel what participants perceived to be the most important barrier(s) and facilitator(s) for HRA completion. The telephone questions were rephrased to match the willingness to complete the HRA: e.g. when the participant indicated to be willing to complete the HRA, the barrier(s) question was rephrased as ‘what could be a disadvantage for you of completing the HRA?’. Coding of these questions was performed as described under step one.

Measures

Primary outcome measure was HRA completion (no/yes). Patient characteristics were: gender, ethnicity, age, and SES score. We also looked at the predominant patient population of a GP practice: native Dutch with a lower SES, mixed, or non-Western immigrants.

Data analyses

Descriptive analyses were used to describe the patient population. Differences regarding sociodemographic characteristics between the patients in the postal versus the telephone step were assessed by means of t-tests and ANOVA.

(Univariable) Logistic regression analyses were performed to explore the associations with HRA completion. First, we assessed the influence of the first reactions variable as a categorical variable (reference group: not having provided a reaction); second, the influence of the structured set of predefined determinants; third, the most important barriers and facilitators. The facilitators and barriers were included jointly in the regression model as patients could report more than one barrier or facilitator. Significant associations with HRA-completion ($p < 0.05$) from these initial analyses were included in a final stepwise multivariate logistic regression model. By adding the various constructs in a stepwise manner, we investigated which associations remained significant when adjusting for each other. The first step included the relevant sociodemographic variables (model 1). Each consecutive step added the significant determinants from respectively self-formulated first reactions (model 2), predefined determinants (model 3), and most important barrier(s) and facilitator(s) (model 4).

Because the number of telephone participants who did not complete the HRA was very low, stratified analyses for the telephone step could not be performed. Only strategy-dependent or very notable differences between postal and telephone participants are highlighted in the text.

RESULTS

Demographics

Of the 1644 eligible patients, 1125 responded to the invitation (response rate: 68%) by either completing the HRA or answering that they did not want to participate. Of those who responded to the invitation, 892 participated in the questionnaire (participation rate: 79%). Among the postal responders ($n = 681$), this percentage was 92% ($n = 624$); among the telephone responders ($n = 444$), it was 60% ($n = 268$) (Fig. 2). Not surprisingly, the questionnaire participants differed from the non-participants regarding all demographic factors except for age (data not shown). The non-participants were more often male ($p < 0.001$), Moroccan or Surinamese ($p < 0.001$), from a GP practice with a predominantly non-Western patient population, and with a lower SES score ($p = 0.039$).

Telephone participants were more often from a GP practice with a predominantly non-Western patient population and had a lower SES score than postal participants (Table 1). Gender, ethnicity, and age were similarly distributed between postal and telephone participants.

Of the questionnaire participants, 78% completed the HRA. Among the postal questionnaire participants, the completion rate was 71%, among the telephone questionnaire participants, it was 94%. A minority of the patients (7%) who did not fill out the questionnaire did complete the HRA.




Figure 2. Flowchart of individuals reached by mail or telephone and their participation in the questionnaire and in the health risk assessment (HRA).

The decision making process

Step one: self-formulated first reaction to the invitation

Table 2 presents the association of the self-formulated first reactions with HRA completion. In this table, we describe the frequencies of the different first reactions. The first reaction variables were entered into a univariable regression analysis (reference group: not having provided a reaction) to explore which reactions were most important for HRA completion. In total, 13% of the participants did not provide a first reaction, most often when they did not complete the HRA. When a first reaction was provided, usually it was a positive or rational attitude ('good' or 'useful'). A positive attitude towards the invitation was associated with HRA completion, as well as expressing the intention to complete the HRA. Paradoxically, those who mentioned negative information factors (not remembering having received an invitation) and barriers to participating (having no time) were more likely to complete the HRA. This was mainly due to the telephone participants who did not remember the postal invitation or said to have had no time to participate, but were apparently able or willing to answer it when approached by phone (separate analyses on postal participants only, ORs not significant anymore: negative information factors $p = 0.627$, barriers $p = 0.477$, data not shown).

Table 1. Sociodemographic characteristics of all questionnaire participants, and stratified by those reached by postal and telephone invitation

	Total (n=892)		Postal (n=624)		Telephone (n=268)		p value
	n (%)	mean (SD)	n (%)	mean (SD)	n (%)	mean (SD)	
Gender							
Male	441 (49)		314 (50)		127 (47)		.207
Female	451 (51)		310 (50)		141 (53)		
Ethnicity							
Dutch	264 (30)		193 (31)		71 (27)		
Turkish	207 (23)		148 (24)		59 (22)		.179
Moroccan	169 (19)		120 (19)		49 (18)		
Surinamese	252 (28)		163 (26)		89 (33)		
Age (years)		50 (6.7)		51 (6.6)		49 (6.5)	.247
30-44	111 (12)		60 (10)		51 (19)		
45-49	344 (39)		235 (38)		109 (41)		
50-54	226 (25)		162 (26)		64 (24)		
55-59	115 (13)		90 (14)		25 (9)		
60-64	63 (7)		51 (8)		12 (5)		
65+	33 (4)		26 (4)		7 (3)		
Predominant GP practice population							
Dutch	226 (25)		165 (26)		61 (23)		
Mix	127 (14)		99 (16)		28 (10)		.024
Non-Western	539 (60)		360 (58)		179 (67)		
SES score		-2.0 (2.5)		-1.9 (2.4)		-2.2 (2.6)	.043
Average to higher (≥ 0)	283 (32)		204 (33)		79 (29)		
Lower to average (0 till -1.9)	202 (23)		144 (23)		58 (22)		
Lower (-2 till -3.9)	130 (15)		94 (15)		36 (13)		
Lowest (≤ -4)	277 (31)		182 (29)		95 (35)		

Table 2. Descriptive and multivariate logistic regression analyses regarding step one: self-formulated first reaction in response to the HRA invitation, for postal and telephone participants (n=892), coded according to I-change model constructs

I-change construct	HRA completers, %n with this reaction n=696	HRA non-completers, %n with this reaction n=196	OR (95% CI) of completing the HRA	Main exemplification
Not answered	10.3	23.5	1.00	
Information factors; positive	13.2	8.2	1.89 (1.08-3.32)*	'Initiative GP/researcher/other'
Information factors; negative	4.6	1.0	6.44 (1.49-27.90)*	'Cannot remember/have not received invitation'
Awareness factors; positive	12.9	10.2	1.39 (0.83-2.35)	'Obtain insight into risks', 'importance prevention', 'health/healthy aging'
Awareness factors; negative	3.0	3.6	1.06 (0.44-2.58)	'More useful for others'
Motivation factors; positive/rational attitude	57.8	55.1	1.53 (1.06-2.20)*	'Good', 'positive', 'important'
Motivation factors; negative/emotional attitude	2.9	4.6	0.88 (0.38-2.06)	'Unnecessary'
Motivation factors; positive social influences	1.1	0	NA	'Action linkage: help from others'
Intention state; precontemplation	1.7	3.6	0.69 (0.26-1.88)	'Not wanting to participate'
Intention state; contemplation	14.4	8.2	2.30 (1.29-4.08)**	'Wanting to participate'
Barriers	4.2	0.5	13.88 (1.85-104.21)*	'No time'
Ability factors	3.7	3.1	2.10 (0.82-5.40)	'Having no barriers'
Other	4.3	2.6	2.33 (0.87-6.26)	Not specified

CI: Confidence interval. NA: Not applicable. *p<.05. **p<.01

Step two: structured set of predefined determinants

Table 3 presents the influence of the predefined determinants on HRA completion, analysed by means of univariable logistic regression. Missing values varied from $n = 4$ (locus of control question) to $n = 34$ (social influences question), but were usually limited. The HRA completers were more likely than the non-completers of wanting to know their risk, of thinking that staying healthy can be controlled, and of having others finding it important for them to participate. Postal participants having one or more health complaints less often completed the HRA (separate analyses on postal participants only: OR 0.68, 95% CI 0.48–0.97, data not shown), while for the group as a whole (postal and telephone participants) we found no association.

Step three: most important barrier(s) and facilitator(s)

Table 4 presents the most important barrier(s) and facilitator(s) and their relation with HRA completion. A regression analysis was conducted including both the barriers and facilitators to assess which of them were significantly related to HRA-completion. Almost half of the participants did not answer the most important barrier(s) question and were less likely to complete the HRA. Those who did answer most often reported having no barriers (37%) or having a negative or emotional attitude (29%). Positive awareness factors ('obtaining insight into risks') were most often mentioned as important facilitator(s) (77%).

Participants who indicated fear (mainly for the test result) to be their most important barrier or who perceived no barriers (hence, ability factor) more often completed the HRA.

Participants who expressed obtaining insight into risks and finding it important as their most important facilitator more often completed the HRA.

Table 3. Descriptive and univariable logistic regression analyses of step two: structured set of predefined determinants of HRA completion, for postal and telephone participants (n=892)

	HRA completers, % n agreement with factor, n= 696	HRA non-completers, % n agreement with factor, n=196	OR (95% CI) of completing the HRA	Corresponding question ^a
Information factors; guidance	74.6	70.1	1.25 (0.88-1.78)	'Do you trust you will get the guidance you need if you have an increased risk?'
Awareness factors; prior experience	25.2	28.4	0.85 (0.59-1.21)	'Have you ever done a health check for cardiovascular disease, diabetes, or kidney disease before?'
Awareness factors; health complaints	52.7	60.2	0.74 (0.53-1.02)	'Do you have one or more health complaints at the moment?'
Awareness factors; perceived risk	54.6	59.6	0.82 (0.59-1.13)	'Do you think the chance of getting these diseases is average to high?'
Awareness factors; knowing risk	91.2	85.6	1.73 (1.07-2.80) [*]	'Do you want to know what your chance of these diseases is?'
Motivation factors; fear test result	23.4	28.5	0.77 (0.54-1.10)	'Are you afraid of the test result?'
Motivation factors; fear adjusting lifestyle	25.9	32.0	0.74 (0.53-1.05)	'Are you afraid that you have to adjust your lifestyle habits?'
Motivation factors; locus of control	59.2	46.2	1.69 (1.23-2.33) ^{***}	'Do you think you can control staying healthy?'
Motivation factors; social influences	67.7	59.9	1.40 (1.00-1.96) [*]	'Do others find it important for you to participate?'

CI: Confidence interval. NA: Not applicable. ^{*}Occasional slight adjustments in formulation so all questions are answered with yes/no. ^{**}p<.05. ^{***}p<.001.

Table 4. Descriptive and multivariate logistic regression analyses regarding step three: most important barrier(s) and facilitator(s) of HRA completion, for postal and telephone participants (n=892).

	HRA completers, %n with this barrier / facilitator n=696	HRA non-completers, %n with this barrier / facilitator n=196	OR (95% CI) of completing the HRA	Main exemplification
Not answered	38.2	62.2	1.00	
Information factors; negative	3.3	4.1	1.10 (0.48-2.55)	'Privacy'
Awareness factors; negative	12.5	11.2	1.62 (0.97-2.86)	'No health complaints'
Motivation factors; negative/emotional attitude	18.4	8.2	3.39 (1.94-5.38)***	'Fear for the test result', 'Afraid to have to adjust lifestyle habits'
Motivation factors; negative social influences	0.4	0	NA	'Action linkage: no help from others'
Intention state; precontemplation	0.6	0.5	0.95 (0.10-9.38)	'Not wanting to participate'
Barriers	4.0	5.1	1.06 (0.50-2.26)	'No time'
Ability factors	23.6	10.2	3.57 (2.15-5.95)***	'Having no barriers'
Other	2.2	2.0	1.56 (0.51-4.79)	
Not answered	11.6	24.5	1.00	
Information factors; positive	4.5	5.1	1.18 (0.53-2.60)	'Knowledge development GP/researcher/ other', 'Trust in guidance'
Awareness factors; positive	68.0	57.1	1.67 (1.16-2.39)**	'Obtain insight into risks', 'healthy aging'
Awareness factors; negative	4.0	4.6	1.22 (0.55-2.72)	'Already health complaints / already ill / receiving treatment'
Motivation factors; positive/rational attitude	18.1	12.8	1.63 (1.01-2.62)*	'Important'
Motivation factors; negative/emotional attitude	1.4	0.5	3.64 (0.46-28.97)	'Worries about health'
Motivation factors; positive social influences	1.9	0.5	3.82 (0.49-29.68)	'Important for offspring'
Motivation factors; negative social influences	0.1	1.0	0.20 (0.02-2.39)	'Social pressure'
Intention state; contemplation	7.9	9.2	0.80 (0.45-1.42)	'Wanting to participate'
Ability factors	2.9	4.6	0.85 (0.37-1.95)	'Having no facilitators'
Other	2.3	3.1	0.90 (0.34-2.40)	

CI: Confidence interval. NA: Not applicable. * p<.05. ** p<.01. *** p<.001

Stepwise model of determinants for HRA completion

Significant associations with HRA-completion ($p < 0.05$) from the initial analyses were included in a final stepwise multivariate logistic regression model. Step one of the stepwise model showed that the probability of HRA completion was highest among participants with a lower SES (score -2 till -3.9), and lowest among Moroccan participants and patients from non-Western GP practices (Table 5). Each consecutive model added the significant determinants from respectively self-formulated first reactions (model 2), predefined determinants (model 3), and most important barrier(s) and facilitator(s) (model 4). In model 2, the self-formulated first reactions significant in the initial analyses remained significantly associated with HRA completion when adjusted for characteristics of model 1, except for the positive/rational attitude. In model 3, the significant association between HRA completion and wanting to know one's risk and perceived control over staying healthy remained significant. Positive social influences were no longer significantly associated with HRA completion. In the final model the negative information factors and the contemplation state remained the only significant first reactions associated with HRA completion. The significant associations from model 3 were supplemented with fear regarding the test result and reporting no barriers, which both remained positively associated with HRA completion. The Nagelkerke R square indicates that 24% of the variance was explained by the final model.

Table 5. Stepwise multivariate logistic regression analyses presenting associations with HRA completion for postal and telephone participants, OR (95% CI)

Variable	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Ethnicity Dutch ^a	1.00	1.00	1.00	1.00
Turkish	0.58 (0.30-1.10)	0.51 (0.26-1.01)	0.47 (0.23-0.95)*	0.61 (0.29-1.27)
Moroccan	0.44 (0.250-0.78)**	0.40 (0.22-0.73)**	0.35 (0.19-0.66)**	0.38 (0.20-0.74)**
Surinamese	0.89 (0.46-1.73)	0.80 (0.40-1.58)	0.68 (0.34-1.38)	0.82 (0.39-1.72)
GP practice Dutch ^a	1.00	1.00	1.00	1.00
Mix	0.70 (0.37-1.33)	0.70 (0.36-1.35)	0.69 (0.35-1.34)	0.67 (0.33-1.34)
Non-Western	0.45 (0.23-0.88)*	0.44 (0.22-0.88)*	0.43 (0.21-0.87)*	0.33 (0.16-0.69)**
SES score Average to higher (> 0) ^a	1.00	1.00	1.00	1.00
Lower to average (0 till -2)	1.54 (0.92-2.57)	1.41 (0.83-2.39)	1.52 (0.88-2.60)	1.69 (0.96-2.96)
Lower (-2 till -3.9)	1.82 (1.01-3.29)*	1.87 (1.02-3.43)*	1.87 (1.01-3.46)*	2.13 (1.12-4.07)*
Lowest (≤ -4)	1.14 (0.69-1.88)	1.11 (0.66-1.88)	1.12 (0.66-1.91)	1.20 (0.69-2.09)
Self-formulated first reaction ^b :				
Not answered		1.00	1.00	1.00
Information factors; positive		2.24 (1.23-4.06)**	2.13 (1.17-3.88)*	1.76 (0.94-3.31)
Information factors; negative		6.75 (1.53-29.79)*	6.98 (1.56-31.20)*	6.57 (1.29-33.43)*
Motivation factors; positive/rational attitude		1.45 (0.98-2.15)	1.38 (0.92-2.05)	1.22 (0.80-1.88)
Intention state; contemplation		2.34 (1.27-4.30)**	2.24 (1.22-4.12)*	2.19 (1.16-4.13)*
Barriers		9.19 (1.20-70.22)*	7.95 (1.04-61.04)*	4.56 (0.57-36.70)
Predefined determinants:				
Awareness factors; knowing risk			2.36 (1.28-4.34)**	1.99 (1.02-3.88)*
Motivation factors; locus of control			1.61 (1.13-2.30)**	1.63 (1.12-2.36)*
Motivation factors; social influences			1.25 (0.86-1.82)	1.20 (0.81-1.77)
Most important barrier(s) ^b :				
Not answered				1.00
Motivation factors; negative/emotional attitude				3.25 (1.69-6.25)***
Ability factors				3.85 (2.14-6.95)***
Most important facilitator(s) ^b :				
Not answered				1.00
Awareness factors; positive				1.12 (0.72-1.74)
Motivation factors; positive/rational attitude				1.38 (0.81-2.35)
Nagelkerke R square	0.07	0.15	0.17	0.24

OR: Odds ratio. CI: Confidence interval. ^aReference category. ^bAll answer categories are included, only those significant in first step are presented. * p<.05. ** p<.01. *** p<.001

DISCUSSION

Conclusion

More than three quarters of the questionnaire participants also completed the HRA, mostly patients in the lower SES group. Those who did not were more often of Moroccan origin or from a non-Western GP practice. The self-formulated first reactions were generally positive and many participants expressed the wish to participate. Barriers formulated as first reaction were mostly not remembering receiving the postal invitation or not having (had) time, thus requiring a phone call. Although positive at first, when HRA non-completers further considered participation they more often did not want to know their risk and were less certain of their ability to control staying healthy. Most of the completers reported having no barriers at all. Some of them ventilated fear for the test result, but this did not prevent them from completing the HRA.

Strengths and weaknesses

To our knowledge, this is the first study exploring the decision making process of underserved populations regarding their completion in the first stage (HRA) of a cardiometabolic health check. The main strength of the study is the exertion to study both HRA completers and non-completers in the context of actual decision-making, reducing potential hypothetical bias. Given the lower levels of (health) literacy levels among these underserved groups, questionnaire missings were limited. This study was embedded in an encompassing theoretical framework. We chose this model as health check attendance can be seen as a health behavior and in that sense be studied with this comprehensive model. In addition to our positive experiences in explaining determinants of hypothetical HRA completion (17) we conclude that the I-change model is also valuable in explaining determinants of actual HRA completion.

Some limitations should be noted. First, given the lower (health) literacy levels of our populations, it is debatable whether the participants were able to formulate their answers in a way that really reflected their opinions. Those with the lowest literacy levels may have more often skipped questions that were difficult for them to fill out (i.e. the open answer questions). The open answer questions were also more often skipped by the HRA non-completers. Related to this is the fact that we tried to make the postal and telephone questionnaire as similar as possible. It cannot be denied, however, that during the telephone

questionnaire our populations may have more easily elaborated on their answers when compared to the self-administered questionnaire. Also, our rephrasing of the barriers/facilitators question may have resulted in slight differences in information obtained, but not rephrasing would have led to unnatural conversations as the research assistant would not react to the willingness already expressed by the participant. Second, all GP practices were located in The Hague and surroundings. It is, therefore, debatable whether the findings from these practices are generalizable to other practices in the Netherlands, let alone other countries. Nevertheless, we do not expect major differences with other practices with similar lower SES/ethnic groups. We expect the major differences to occur between ethnic groups, and between socioeconomic strata. Increasing the chance of generalizability was the fact that we included both large group as well as solo practices, and urban as well as rural environments. Third, potential residual confounding may have been present due to possible errors in the estimation of ethnicity based on last name and because the SES score was a neighbourhood score and not an individual score. Fourth, in the larger study we have not reached 30% of the patients, of whom we have no information on determinants of their HRA non-completion whatsoever. Aiming to reduce health inequalities, obtaining insight into the determinants of particularly these non-participants is important. Fifth, we used a cross-sectional study design. This means that we cannot conclude that the correlates we found caused the HRA completion. We can conclude that some differences exist between HRA completers and non-completers. Finally, it is possible that our simultaneously sending a questionnaire for research purposes led to distrust among some individuals, as we have shown before to be a potential problem for these vulnerable groups (17). With this simultaneous sending and our structured design of the questionnaire we might also have enabled participants to reflect on participation and to make an informed decision about HRA completion. However, we have no control group, nor a measure for informed decision making, allowing us to draw a conclusion about whether we have succeeded.

Comparison with other studies

We aimed to explore the process of decision making regarding HRA completion among underserved populations. Most first reactions provided were positive, possibly because patients felt obliged to react positively or to provide some excuse for not having completed the HRA. Which raises the question on the value of these reactions for explaining HRA participation. In the final model, most of the associations of these positive first reactions with

participation disappeared. The only negative first reaction (not remembering/receiving the invitation) remained significantly associated with HRA participation. This implies that this determinant may not have been used as an excuse, and follow-up by telephone may actually be crucial.

Despite the generally positive self-formulated first reactions, still about a quarter did not complete the HRA. A qualitative investigation among non-responders to the NHS health check concluded that participants viewed the health check positively, but lacked personal relevance (25). They concluded that, perhaps, people considered themselves in good health or had regular contact with primary care for their health complaints. To promote personal relevance and more informed decision making, it has been proposed to provide more personalized risk communication and using telephone/verbal methods (25, 26). In the current study, patients with health complaints less often participated in the postal HRA, but not so in the telephone HRA. This indicates that personalized risk communication by telephone may increase personal relevance. It has been found that up to 40% of non-participants in cardiovascular screening would reconsider their participation decision when given additional information (27).

The argument of not wanting to know one's risk is in line with findings of other studies (28, 29). The current study adds that participation in a health check is influenced by a perceived lack of control over staying healthy. This external locus of control may also imply that the HRA non-completers were the less acculturated minorities, as they more often feel that the doctor, God, or a higher power could help prevent cardiometabolic disease, rather than they themselves (30). Indeed, HRA completion was lowest in non-Western GP practices, situated in neighbourhoods with stronger non-Western communities, with generally lower acculturation rates. Additionally, Turkish and Moroccan immigrants had the lowest HRA completion rates and have been found to be less acculturated than Surinamese immigrants, and less often participate in Dutch society (31). Efforts should, therefore, be particularly targeted at the less acculturated immigrants, emphasizing the modifiability of cardiometabolic disease through lifestyle changes and boosting the confidence in their own abilities.

Interestingly, the most important barrier expressed by the completers was fear of the test result. However, as the results of the predefined set of determinants showed, many were not actually scared of the test result. This barrier may, thus, have been a potential barrier imagined to be applicable to others in the same situation. On the other hand, these participants may have participated despite of their anxiety so that in case of a high-risk test

result, they would at least know that they would experience the benefits of early diagnosis (32, 33). Indeed, a large majority of these HRA completers expressed their most important facilitator to be obtaining insight into risks.

Implications and future research

To explore the influence of perceived personal relevance on informed decision making, future studies should focus on personalized/verbalized cardiometabolic risk communication, emphasizing the modifiability of cardiometabolic risk factors and boosting the confidence of these underserved groups. Most of our underserved participants, often depicted as ‘hard-to-reach’, had a positive attitude towards the cardiometabolic health check. This, combined with the strongest correlates being not remembering having received an invitation and having had no time (thus requiring a phone call), gives rise to the idea that these groups may not be hard-to-reach in the sense that they are unwilling to complete the HRA. The results of this pragmatic intervention provide interesting leads for follow-up by means of a controlled study. Special efforts should then be made at those ‘hardest-to-reach’. A more comprehensive approach, including the involvement of key figures within a community informing people about and providing help with the HRA (reducing the amount of time needed) would possibly be more suitable for these groups. Efforts should be particularly aimed at the less acculturated immigrants.

ACKNOWLEDGMENTS

Conflicts of interest

The authors declare that there are no conflicts of interest.

Authors' contributions

MC, SD, WG, and AS filed the proposal for this study. The design, execution, and analysis were mainly done by IG and MC, in close collaboration with the research team. The paper was written by IG and critically revised by all authors. All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis. IG is guarantor.

Funding

This study was funded by the Dutch Heart Foundation, the Dutch Diabetes Foundation, and the Dutch Kidney Foundation in a collaboration called *LekkerLangLeven* [Living nice and long] (2008.20.005). The views presented in this manuscript are those of the authors and do not necessarily reflect those of the study funders.

Ethical approval

This study was approved by the Medical Ethical Committee of Leiden University Medical Center (P11.151).

Transparency document

The Transparency document associated with this article can be found, in online version.

Acknowledgments

The authors thank all GP practices for their cooperation in conducting the study at their locations, and the patients in particular for their participation.

APPENDIX A

Exclusion criteria

Having (had) one or more of the following diseases (in ICD codes):

- o K74 ANGINA PECTORIS
- o K75 ACUTE MYOCARDIAL INFARCTION
- o K76 OTHER CHRONIC ISCHEMIC HEART DISEASES
- o K77 CONGESTIVE HEART FAILURE
- o K78 ATRIAL FIBRILLATION/-FLUTTER
- o K79 PAROXYSMAL TACHYCARDIA
- o K82 COR PULMONARY
- o K83 VALVE DISEASE NOT RHEUMATIC/NOS
- o K84 OTHER HEART DISEASES
- o K86 HYPERTENSION WITHOUT ORGAN DAMAGE.
- o K87 HYPERTENSION WITH ORGAN DAMAGE.
- o K89 TRANSIENT CEREBRAL ISCHEMIA/TIA
- o K90 CEREBROVASCULAR ACCIDENT (CVA) [EX.TIA]
- o K91 ATHEROSCLEROSIS [EX.CORON.,CEREBR.]
- o K92 OTHER DISEASES PERIPHERAL ARTERIES
- o T90 DIABETES
- o T93 LIPID DISORDER
- o U88 GLOMERULONEPHRITIS/NEFROSIS
- o U99 OTHER DISEASES URINARY TRACT

Use of one of the following drugs (in ATC-classifications):

- o A10 ANTIDIABETICS
- o B01/C01/C02/C03/C07/C08/C09 ANTIHYPERTENSIVES
- o C10 ANTILIPAEIMICS

Complete risk profile with a maximum of one year old with a known measurement for all of the following factors:

- o Smoking status
- o Comments on characteristics of diet
- o Physical activity
- o Alcohol use
- o BMI
- o Waist circumference
- o Systolic blood pressure
- o Fasting glucose
- o LDL

APPENDIX B




Figure B1. Results of response and participation in three culturally targeted and personalized invitation steps following an increasingly (cost-)intensive ‘funnelled’ design

APPENDIX C

Question	Answer options
1. What is your first reaction about this invitation?	Open
2. Have you ever done a health check for cardiovascular disease, diabetes, or kidney disease before?	No (or for something else) - Yes, <i>more</i> than 1 year ago - Yes, <i>less</i> than 1 year ago
3. Are you afraid of the result?	No - A little - Yes
4. Are you afraid that you have to adjust your lifestyle habits?	No - A little - Yes
5. Do you trust you will get the guidance you need if you have an increased risk?	No - A little - Yes
6. Do you have health complaints at the moment?	None - One - A couple
7. How high do you think your chance is of getting these diseases?	Small - Average - High
8. Do you want to know what your chance of these diseases is?	No - Dubiety - Yes
9. Do you think you can control staying healthy?	No - Partly - Yes
10. Do others find it important for you to participate?	No - Little - Yes
11. What is the most important reason for you <i>not</i> to participate in the HRA? Of course, this may be another reason than what has been mentioned above. <i>Also fill out this question if you do participate in the HRA!</i>	Open
12. What is the most important reason for you to <i>do</i> participate in the HRA? Of course, this may be another reason than what has been mentioned above. <i>Also fill out this question if you do not participate in the HRA!</i>	Open

APPENDIX D

Code tree assigned to I-change constructs

Predisposing factors

None.

Information factors; positive

Healthcare professional

- o Initiative GP/researcher/other
- o Attention from GP/researcher/other
- o Knowledge development GP/researcher/other
- o Trust in guidance
- o Relationship with GP

Information factors

- o Had understood it
- o Taking the target population into account
- o (Remembered) Having received it

Information factors; negative

Healthcare professional

- o No treatment options
- o Privacy issues

Obligation

- o Feeling of obligation
- o No Feeling of obligation

Information factors

- o Had not understood it
- o Language barrier
- o (Health) Illiteracy
- o Not (remembering) having received it
- o Doubts about content aspects
- o Judgment about materials

Awareness factors; positive

Importance prevention

Health status

- o Obtain insight into risks
- o Obtain certainty about health
- o (Being) Health(y)
- o Healthy aging
- o Believes to be low-risk and wants to know risk
- o Believes to be high-risk and wants to know risk
- o Decrease risks
- o Risk perception with regard to family history
- o Risk perception with regard to lifestyle
- o Never too old
- o Previous experience with a health check
- o No previous experience with a health check

Awareness factors; negative

Health status

- o Too old

- o Already health complaints/already ill/receiving treatment
 - o No health complaints
 - o Convinced of own health
- Social environment
- o More useful for others

Motivation factors; positive/rational attitude

Positive

- o Nice
- o Good
- o Useful
- o Okay
- o Surprising
- o Interesting
- o Happy
- o Relieved
- o Can do no harm
- o Curious
- o Sensible
- o Important
- o Necessary
- o Satisfied
- o Enthusiastic
- o Grateful
- o Positive

Normal/neutral

Health status

- o Not afraid to have to adjust lifestyle habits

Locus of control

- o Believes to control staying healthy

Fear

- o No fear
- o No fear for the test result

Motivation factors; negative/emotional attitude

Negative

- o Not good
- o Not important
- o Unreliable
- o Not interested
- o Don't feel like it
- o Not necessary
- o Strange
- o Doubt
- o Negative

Health status

- o Afraid to have to adjust lifestyle habits
- o Worries about health
- o Ignoring/denial

Healthcare professional

- o Guinea pig

Locus of control

- o Believes not to control staying healthy
- o Religious beliefs

Fear

- o Fear for the test result
- o Fear for the consequences of the test result
- o Fear for doctors/hospitals

- o Panic/agitation
 - o Fear
- Own responsibility

Motivation factors; positive social influences

- Social environment
- o Action linkage: help from others
 - o Important for offspring
 - o Others find it important

Motivation factors; negative social influences

- Social environment
- o Social pressure
 - o Gossip

Intention state; precontemplation

- (Non-)Participation
- o Not wanting to participate
 - o Doubts about participation

Intention state; contemplation

- (Non-)Participation
- o Wanting to participate
 - o Need

Ability factors

- Time
- o Takes little time
- Not applicable/none

Barriers

- Time
- o No time
 - o Other priorities
 - o Job
 - o Forgot it
 - o Holidays/in home country/sick
- Money issues

Other

- Health status
- o Pregnant
- Psychological problems
- Other

REFERENCES

1. World Health Organization. Fact sheet number 310: The top ten causes of death 2008. 2011. http://www.who.int/mediacentre/factsheets/fs310_2008.pdf.
2. Mackenbach JP, Stirbu I, Roskam AJ, Schaap MM, Menvielle G, Leinsalu M et al. Socioeconomic inequalities in health in 22 European countries. *N Engl J Med* 2008;358(23):2468–81.
3. Mackenbach JP, Kunst AE, Cavelaars AE, Groenhouf F, Geurts JJ. Socioeconomic inequalities in morbidity and mortality in western Europe. The EU Working Group on Socioeconomic Inequalities in Health. *Lancet* 1997;349(9066):1655–59.
4. Bos V, Kunst AE, Keij-Deerenberg IM, Garssen J, Mackenbach JP. Ethnic inequalities in age- and cause-specific mortality in The Netherlands. *Int J Epidemiol* 2004;33(5):1112–19.
5. van Leest LATM, van Dis SJ, Verschuren WMM. Hart- en vaatziekten bij allochtonen in Nederland, een cijfermatige verkenning naar leefstijl- en risicofactoren, ziekte en sterfte [Cardiovascular diseases in non-Western immigrants in the Netherlands. An exploratory study into lifestyle, risk factors, morbidity, and mortality]. Bilthoven, the Netherlands: RIVM; 2002.
6. Dijkshoorn H, Uitenbroek DG, Middelkoop BJ. Prevalentie van diabetes mellitus en hart- en vaatziekten onder Turkse, Marokkaanse en autochtone Nederlanders [Prevalence of diabetes mellitus and cardiovascular disease among immigrants from Turkey and Morocco and the indigenous Dutch population]. *Ned Tijdschr Geneeskd* 2003;147(28):1362–66.
7. Kunst AE, Mackenbach JP, Lamkaddem M, Rademakers J, Devillé W. Overzicht en evaluatie van resultaten van wetenschappelijk onderzoek naar etnische verschillen in gezondheid, gezondheidsrisico's en zorggebruik in Nederland [Overview and evaluation of results from scientific research on ethnic differences in health, health risks, and health care use in the Netherlands]. Utrecht, the Netherlands: NIVEL; 2008.
8. Dalton AR, Bottle A, Okoro C, Majeed A, Millett C. Uptake of the NHS Health Checks programme in a deprived, culturally diverse setting: cross-sectional study. *J Public Health* 2011;33(3):422–29.

9. Brunner-Ziegler S, Rieder A, Stein KV, Koppensteiner R, Hoffmann K, Dorner TE. Predictors of participation in preventive health examinations in Austria. *BMC Public Health* 2013;13:1138.
10. Amoroso C, Harris MF, Ampt A, Laws RA, McKenzie S, Williams AM et al. The 45 year old health check - feasibility and impact on practices and patient behaviour. *Aust Fam Physician* 2009;38(5):358–362.
11. Khunti K, Gillies CL, Taub NA, Mostafa SA, Hiles SL, Abrams KR et al. A comparison of cost per case detected of screening strategies for Type 2 diabetes and impaired glucose regulation: modelling study. *Diabetes Res Clin Pract* 2012;97(3):505–13.
12. Pandya A, Weinstein MC, Salomon JA, Cutler D, Gaziano TA. Who needs laboratories and who needs statins?: comparative and cost-effectiveness analyses of non-laboratory-based, laboratory-based, and staged primary cardiovascular disease screening guidelines. *Circ Cardiovasc Qual Outcomes* 2014;7(1):25–32.
13. Dekker JM, Alsema M, Janssen PGH, van der Paardt PGH, Festen CCS, van Oosterhout et al. NHG-Standaard Het PreventieConsult module Cardiometabool [Guideline for Dutch GPs Prevention consultation Cardiometabolic module]. *Huisarts Wet* 2011;54(3):138–55.
14. Assendelft WJ, Nielen MM, Hettinga DM, van der Meer V, van Vliet M, Drenthen AJ et al. Bridging the gap between public health and primary care in prevention of cardiometabolic diseases; background of and experiences with the Prevention Consultation in The Netherlands. *Fam Pract* 2012;29 Suppl 1:i126–i31.
15. Dryden R, Williams B, McCowan C, Themessl-Huber M. What do we know about who does and does not attend general health check? Findings from a narrative scoping review. *BMC Public Health* 2012;12:723.
16. Smith SK, Simpson JM, Trevena LJ, McCaffery KJ. Factors associated with informed decisions and participation in bowel cancer screening among adults with lower education and literacy. *Med Decis Mak* 2014;34(6):756–72.
17. Groenberg I, Crone MR, van Dijk S, Gebhardt WA, Ben Meftah J, Middelkoop et al. ‘Check it out!’ Decision-making of vulnerable groups about participation in a two-stage cardiometabolic health check: a qualitative study. *Patient Educ Couns* 2015;98(2):234–44.
18. Sociaal en Cultureel Planbureau [Social and Cultural Planning Office]. Status scores 2014. The Hague, the Netherlands.

19. Reijneveld SA. The impact of individual and area characteristics on urban socioeconomic differences in health and smoking. *Int J Epidemiol* 1998;27(1):33–40.
20. Groenenberg I, Crone MR, van Dijk S, Ben Meftah J, Middelkoop BJ, Assendelft WJ et al. Response and participation of underserved populations after a three-step invitation strategy for a cardiometabolic health check. *BMC Public Health* 2015;15:854.
21. Knops-Dullens T, de Vries N, de Vries H. Reasons for non-attendance in cervical cancer screening programmes: an application of the Integrated Model for Behavioural Change. *Eur J Cancer Prev* 2007;16(5):436–45.
22. de Vries H, Mesters I, van de Steeg H, Honing C. The general public's information needs and perceptions regarding hereditary cancer: an application of the Integrated Change Model. *Patient Educ Couns* 2005;56(2):154–65.
23. Nierkens V, Stronks K, de Vries H. Attitudes, social influences and self-efficacy expectations across different motivational stages among immigrant smokers: replication of the O pattern. *Prev Med* 2006;43(4):306–11.
24. Nierkens V, Stronks K, van Oel CJ, de Vries H. Beliefs of Turkish and Moroccan immigrants in The Netherlands about smoking cessation: implications for prevention. *Health Educ Res* 2005;20(6):622–34.
25. Ellis N, Gidlow C, Cowap L, Randall J, Iqbal Z, Kumar J. A qualitative investigation of non-response in NHS health checks. *Arch Public Health* 2015;73(1):14.
26. Burgess C, Wright AJ, Forster AS, Dodhia H, Miller J, Fuller F et al. Influences on individuals' decisions to take up the offer of a health check: a qualitative study. *Health Expect* 2014;18(6):2437–48.
27. Sogaard R, Lindholt J, Gyrd-Hansen D. Individual decision making in relation to participation in cardiovascular screening: a study of revealed and stated preferences. *Scand J Public Health* 2013;41(1):43–50.
28. McAndrew LM, Musumeci-Szabo TJ, Mora PA, Vileikyte L, Burns E, Halm EA et al. Using the common sense model to design interventions for the prevention and management of chronic illness threats: from description to process. *Br J Health Psychol* 2008;13(Pt 2):195–204.
29. Farrimond H, Saukko PM, Qureshi N, Evans PH. Making sense of being at 'high risk' of coronary heart disease within primary prevention. *Psychol Health* 2010;25(3):289–304.
30. Edelman D, Christian A, Mosca L. Association of acculturation status with beliefs, barriers, and perceptions related to cardiovascular disease prevention among racial and ethnic minorities. *J Transcult Nurs* 2009;20(3):278–85.

31. Nap A, van Loon A, Peen J, van Schaik DJ, Beekman AT, Dekker JJ. The influence of acculturation on mental health and specialized mental healthcare for non-western migrants. *International J Soc Psychiatry* 2015;61(6):530-8.
32. Kaltsa A, Holloway A, Cox K. Factors that influence mammography screening behaviour: a qualitative study of Greek women's experiences. *Eur J Oncol Nurs* 2013;17(3):292–301.
33. Hennelly MO, Sly JR, Villagra C, Jandorf L. Narrative message targets within the decision-making process to undergo screening colonoscopy among Latinos: a qualitative study. *J Cancer Educ* 2015;30(2):268-76.

