



Universiteit
Leiden
The Netherlands

Clinical outcomes of modern lamellar keratoplasty techniques

Dijk, K. van

Citation

Dijk, K. van. (2018, January 16). *Clinical outcomes of modern lamellar keratoplasty techniques*. Retrieved from <https://hdl.handle.net/1887/59337>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/59337>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/59337> holds various files of this Leiden University dissertation

Author: Dijk, Korine van

Title: Clinical outcomes of modern lamellar keratoplasty techniques

Date: 2018-01-16

Chapter 1

General introduction

2

3

4

5

6

7

8

9

10

A

THE HUMAN CORNEA

The cornea, the dome-shaped 'window of the eye', covers the anterior chamber, iris and pupil. It has a diameter of approximately 11.5 millimeters and is normally about 500 to 600 micrometers (μm) thick.¹ Its function is to protect the inner ocular structures as well as to provide about 2/3 of the eye's refractive power. It is one of the most innervated and sensitive tissues of the body, with unmyelinated nerve endings derived from the ophthalmic branch of the trigeminal nerve. At the same time, the absence of blood and lymphatic vessels contributes to corneal clarity and optimal optical performance, while contributing to the cornea's immunologic privilege.^{1,2} The human cornea consists of five layers: epithelium, Bowman layer (BL), stroma, Descemet membrane (DM) and endothelium (Figure 1).




Figure 1. Schematic representation of the human cornea.

Epithelium

The epithelium is the outermost part of the cornea, comprising about 10 percent (around $53\mu\text{m}$ in thickness) of total corneal thickness.³ It is a stratified, non-keratinized structure characterized by extreme uniformity from limbus to limbus, composed of 5 to 6 layers of cells, tightly and orderly arranged without intercellular spaces, and increasingly flattened as it reaches the surface. Corneal epithelial cells have an average lifespan of 7 to 10 days, and routinely undergo orderly involution, apoptosis, and desquamation,¹ in which deeper cells replace the desquamating superficial cells in an orderly, apically directed fashion.⁴ An important source of new corneal epithelial cells are the epithelial stem cells, localized at the corneal limbus. As these cells migrate to the central cornea, they differentiate into basal cells, forming the deepest epithelial layer, which is connected to the underlying epithelial basement membrane by hemi-desmosomes. Just anterior to the basal cell layer are 2 to 3 layers of wing cells, which are covered by desquamating superficial cells. These desquamating superficial cells are tightly connected to their neighboring cells through tight junctional complexes, which prohibits tears, toxins and microbes from entering the deeper corneal layers.^{1,5} Despite the high turnover rate of the epithelial cells, a highly controlled, steady thickness profile, and a tight arrangement of cells is maintained to guarantee a constant corneal power, with

a good optical quality.^{3,6} Furthermore, the superficial flattened epithelial cells interact with the mucous layer of the tear film to ensure a smooth, stable optical surface that resists bacterial adherence.^{5,7}

Bowman Layer

The BL, situated directly underneath the epithelial basement membrane, is an 8 to 14 μm thin acellular layer with the front being smoother than the more irregular posterior part; it thins with age.^{8,9} The BL contains randomly-organized tightly-woven collagen fibrils (mostly types I and V collagen), which differ from the thicker collagen fibrils of the underlying stroma that run in alignment across the cornea in characteristic lamellae.^{10,11} Thus far, the function of this firm dense corneal layer remains unclear. It has been hypothesized that the BL may act as a biological barrier against pathogen infiltration, especially viruses.^{1,11,12} The BL may also have a role in modulating epithelial-stromal wound-healing, since subepithelial-stromal scarring seems not to occur in the presence of an intact BL.^{13,14} Additionally, given its strong rigidity, it may be valuable for maintaining the corneal shape or strength.^{8,11,15}

Stroma

About 90% of the total cornea is made up by stroma. It consists primarily of water (78%), collagen (15%), and non-collagenous proteins (7%) which are supported by scattered keratocytes,¹⁶ predominantly situated in the anterior stroma. Collagen fibers (mainly types I and V) are arranged in parallel bundles, i.e. fibrils, which are packed in parallel arranged layers or lamellae. Each lamella is positioned at right angles relative to fibers in adjacent lamellae.¹⁷ In the central cornea, the stroma comprises about 200 lamellae, which are more densely packed in the anterior region.¹⁸ The anterior lamellae are highly interlaced,¹⁹ inserting into the BL.²⁰ Also the mid-stromal lamellae are considerably interwoven,²¹ whereas the posterior lamellae seem to have less interlacing. As such, the posterior stroma swells easily while the more interwoven anterior stroma does not.²² The peripheral stroma is thicker than the central, and the collagen fibrils run more circumferentially as they approach the limbus.²³ Furthermore, stromal interweaving seems to extend to the deeper posterior lamellae in this region of the cornea.¹⁹

The lamellar arrangement within the stroma helps to maintain the cornea's overall shape, while the unique structure and organization of the collagen fibrils within the lamellae allow corneal transparency.²⁴ Specifically, uniformity in the collagen fibrils' diameters and the restriction in the range of distances between adjacent collagen fibrils, maintained by the interfibrillar proteoglycans, may be important determinants.²⁵⁻²⁷ Additionally, cytoplasmic molecules, the so-called 'corneal crystallins', within the keratocytes seem to be responsible for reducing backscatter of light from the keratocytes, supporting corneal transparency by matching the refractive index of the cell cytoplasm

to the extracellular matrix.²⁸ Dysfunction in any of these components can reduce corneal clarity, and consequently may result in functional loss.

Descemet membrane

Beneath the stroma lies the DM, an extracellular matrix, mainly composed of type IV collagen fibrils and laminin that serves as a basement membrane for the corneal endothelium. Apparently, endothelial cells create this membrane by secreting the different components.^{29,30} Three different zones can be distinguished: a 0.3 μm thin anterior non-banded zone, adjacent to the posterior stroma, an anterior banded zone (2-4 μm) and a posterior amorphous non-banded zone (>4 μm), which thickens with age.³⁰ Just anterior to the DM, a very thin, strong pre-DM layer, may exist.³¹

Endothelium

The endothelium is the corneal innermost monolayer of approximately 4 μm thin in adulthood. The endothelial layer consists of closely-packed hexagonal cells that are essential in keeping the cornea transparent, by regulating fluid and solute transport between the aqueous humor and stroma. The cells serve as selective barriers allowing leakage of solutes and nutrients from the aqueous humor to the cornea. On the other hand, a net flux of ions from the stroma to the aqueous humor is ensured through Na-K ATPase pumps, found in the basolateral endothelial cell membranes, and intracellular carbonic anhydrase, which create an osmotic gradient that causes passive diffusion of fluid in the same direction.^{1,32-34} In a healthy cornea, the endothelium maintains a perfect balance between fluid moving into and being pumped out of the cornea,^{1,32} maintaining the stroma in a relative state of deturgescence, while dysfunction of either the barrier or the pumps results in corneal edema.

Endothelial cells do not regenerate, but instead stretch to compensate for lost cells. As this process occurs, the remaining cells grow in size (polymegathism) and lose their hexagonal shape (pleomorphism).¹ The endothelial cell density is around 6000 cell/ mm^2 at birth and 3500 cells/ mm^2 by the age of 5 years.³² This number decreases gradually throughout life at an average rate of 0.6% per year.¹ Accelerated cell loss may occur after ocular surgery, such as cataract extraction and corneal transplantation, probably due to surgical trauma and postoperative inflammation.^{1,33,35} Corneal transparency may be maintained with an endothelial cell density as low as 500 cells/ mm^2 (range 750-250 cells/ mm^2), after which a proper corneal fluid balance may no longer be preserved and stromal swelling with subsequent loss of corneal transparency may occur.³³

Corneal optical properties

The visual performance of the cornea depends on its shape, transparency and surface regularity. As described, a highly-organized matrix of corneal collagen fibrils maintains

corneal clarity by minimizing light scatter. Anything that interferes with this matrix or affects the corneal surface may compromise the corneal optical quality.

In an average human eye, the cornea forms a positive prolate-shaped (steepest curvature at the corneal apex, progressively flatter curvature toward the periphery) meniscus lens.⁷ The anterior corneal surface has an approximate refractive power of +48 D, while the refractive power of the posterior corneal surface is -6 D. To convert radii of curvature into corneal power it is often assumed that the entire cornea has a uniform refractive index of 1.376, whereas refractive indices of 1.3375 (keratometric index) and 1.0 (air) are normally used to determine the anterior keratometric corneal power. In young individuals, the anterior corneal surface tends to have with-the-rule-astigmatism (steepest meridian is vertical), but astigmatism becomes more against-the-rule (steepest meridian is horizontal) with increasing age.³⁶⁻³⁸ The posterior cornea tends to be steeper in the vertical meridian, inducing against-the-rule astigmatism, that shows only a minimal variation with age.³⁹ The relationship between the anterior and posterior corneal surface is important, and normally the posterior surface compensates approximately 30% of the anterior corneal astigmatism.⁴⁰

Together with the lens, the cornea plays a major role in the shape and amount of the total human eye optical aberrations,⁴¹ i.e. any light rays misdirected from their intended image point. In the normal population, the dominant optical aberrations include the ordinary second-order spherocylindrical errors, which are called lower-order aberrations or refractive errors (myopia, hyperopia, and regular astigmatism). Lower-order aberrations are typically correctable with spectacles, whereas higher-order aberrations (HOAs), which normally comprise about 10% of total aberrations, cannot.⁴² Total eye aberrations are shown to be less than those of the anterior cornea alone, which may indicate that aberrations of both the posterior cornea and crystalline lens partly compensate for the anterior corneal surface aberrations, although this compensation mechanism seems to decrease with age.⁴⁰

A useful representation to describe total eye aberrations as well as the aberrations of both corneal surfaces is the use of Zernike Polynomials (Figure 2). The polynomials describe typical optical properties or imperfections, using varying radial orders and angular frequencies, demonstrated as different wavefront error maps.^{41,43-45} Each polynomial is named according to the image defects it represents (e.g. astigmatism, coma or spherical aberration), affects the optical quality in a specific way, and can be positive or negative (Figure 2). Generally, the magnitude of contribution decreases with increasing Zernike order, and polynomials near the center of the Zernike tree (e.g. coma, spherical aberration) tend to affect vision more than those near the edge of the tree (e.g. trefoil) (Figure 2).⁴³ Furthermore, the different polynomials interact to either increase or decrease visual function, depending on their relative contributions and how they are combined.⁴¹




Figure 2. Chart showing the Zernike tree with the common Zernike polynomials. An aberration-free flat plane, known as piston, represents a theoretically perfect eye.

A healthy clear cornea is able to transmit almost all incident light in the visible part of the spectrum.^{24,25} Loss of clarity, or increased haziness, indicates a degradation of the corneal tissue,^{46,47} and may be a sign of a pathological process, such as infection, corneal dystrophy or degeneration. Corneal haze (back-scattered light) most likely represents a mixture of light scattered from small particles and specular reflections from adjacent tissues with different refractive indices.⁴⁶ Back-scattered light itself cannot compromise the vision, but is associated with corneal changes that typically increase forward-scattered light, which degrades the periphery of the point-spread function. As such, disability glare (reduced visibility due to light sources in the visual field), impaired contrast sensitivity and decreased visual performance in unfavorable ambient light conditions may arise.⁴⁸⁻⁵⁰

Corneal imaging

Various diagnostic imaging devices are currently available to analyze the cornea, including corneal topography, corneal tomography, anterior segment optical coherence tomography, specular microscopy and confocal microscopy.

With corneal topography, the anterior corneal surface can be evaluated, most commonly by reflecting a series of concentric rings, known as Placido rings, on the cornea. A digital camera then captures the reflected pattern, after which the shape of the cornea is processed and calculated by the device's software. The principle of corneal topography




Figure 3. Corneal topography maps showing (A) a prolate-shaped cornea with the steepest curvature (warmer colors) at the corneal apex, (B) an oblate-shaped cornea after Laser in-situ keratomileusis with a flatter central curvature (cooler colors) compared to a steeper peripheral shape, and (C) an irregular-shaped cornea with the steepest curvature in the inferior corneal meridian as in keratoconus.

is based upon the assumption that the cornea is prolate, and therefore non-prolate corneas or irregular corneal surfaces are often misdiagnosed (Figure 3).⁵¹ Furthermore, to obtain a clear image, the epithelial surface as well as the tear film must be intact. The newest color light-emitting diode (LED) corneal topography devices, in which a color-coded checkerboard is projected on the cornea instead of Placido rings, may allow for higher predictability in central irregular corneas as compared to Placido-based topographers.^{52,53}

Whereas corneal topographers can only measure the anterior corneal surface, corneal tomography devices can evaluate both the anterior and posterior cornea, as well as determine corneal thickness (pachymetry), anterior chamber depth and lens parameters.⁵⁴ Three types of corneal tomographers are currently available: slit scanning devices (i.e. Orbscan II), Scheimpflug imaging devices (i.e. Pentacam), and devices that combine Scheimpflug cameras with Placido based topography (i.e. Galilei, Sirius and TMS-5/Tomey). A slit scanning device projects optical slits at a fixed 45-degree angle along multiple points across the cornea. A digital video camera captures the reflections of the slits, after which the reflections can be analyzed to construct representations of the anterior and posterior cornea.⁵⁴ With Scheimpflug-based corneal imaging, a rotating Scheimpflug camera captures multiple cross-sectional slits in a two-second scan across the cornea, while a stationary camera, centered at the pupil, aligns the images and monitors ocular fixation. The three-dimensional images are then analyzed to compose anterior and posterior corneal topographies based on height data.^{54,55} Some Scheimpflug-based tomographers (e.g. Pentacam) also provide corneal wavefront and densitometry analysis (Figure 4).

Anterior segment optical coherence tomography (OCT) uses optical interferometry to produce high-resolution, cross-sectional images of the cornea and other anterior segment structures. It is being employed more and more often to picture anterior segment pathologies, and for anatomical imaging for surgical purposes.⁵⁶ Two main OCT




Figure 4. Screen display of Scheimpflug-based (Pentacam) corneal densitometry assessment.

categories can be distinguished; time-domain (TD) vs. spectral-domain (SD, also called Fourier-domain). TD-OCT's utilize a moveable reference mirror, which moves for each scan to determine the ocular structure depth. SD-OCT's use a fixed reference mirror and a Fourier transformation algorithm of the interferogram to measure depth, resulting in faster acquisition and better image quality as compared to TD-OCT's.⁵⁷

Specular microscopy allows for visualization of the corneal endothelium, by projecting light onto the cornea, and capturing the image reflected from the optical interface between the endothelium and the aqueous humor. Modern specular microscopes allow for analysis of the endothelial cell size, shape and density.⁵⁸

Corneal confocal microscopy is a relatively novel and rapidly evolving clinical technique enabling real-time, in vivo, microstructural analysis of every layer of the cornea. In confocal microscopy, a single point of tissue is both optically illuminated (using white light or a focused laser beam) and simultaneously imaged by a point detector, both having the same focal point, or being "confocal". To optimize the optical resolution, the light source and the detector use pinholes that work 'in tandem'. To increase the field of view, the instrument instantaneously illuminates and synchronously images a small corneal region with numerous tiny light spots or slits, which can be reconstructed to create a functional field of view with high resolution and magnification.⁵⁹⁻⁶⁰

CORNEAL TRANSPLANTATION

Corneal transplantation, also known as corneal grafting or keratoplasty, is a surgical procedure in which a damaged or diseased cornea is replaced by corneal donor tissue (the graft). It remains the main method to restore vision in eyes with irreversible affection of corneal clarity. In addition, globe preservation, pain reduction and cosmetic appearance of the eye may be reasons for keratoplasty. While indications differ between locations and institutes, in developed western countries, corneal transplantation is most often performed for disorders in which the endothelium is impaired, such as Fuchs endothelial dystrophy, bullous keratopathy, and failed previous corneal grafts, or ectatic disorders, which are mostly due to keratoconus.⁶¹

History of corneal transplantation

The first suggestion to replace an opaque cornea for manufactured transparent material in order to restore vision originated from the French surgeon Guillaume Pellier de Quengsy in 1789.⁶² This was actually the first inspiration for what is nowadays known as keratoprosthesis. However, it lasted until 1818 when Franz Reisinger initiated experimental corneal transplantation in animals. Reisinger also introduced the term 'keratoplasty'.⁶³ At the end of the 19th century, around 1890, professor Arthur von Hippel promoted anterior lamellar corneal transplants, since he understood that corneal transparency mainly depended on the integrity of the endothelium and DM.⁶⁴

In 1905, Eduard Zirm performed the first successful human corneal transplantation, a full-thickness keratoplasty on a 45-year-old male with severe bilateral alkali burns. The operation was done under general anesthesia and 'strict' aseptic conditions.⁶⁵ Vladimir Filatov, a Russian eye surgeon, followed with his first attempts in full-thickness corneal transplantation in 1912, which eventually resulted in a successful transplantation using corneal tissue from a deceased human donor in 1931. Filatov is considered the father of eye banking and is credited for popularizing the use of cadaveric human donor corneas for transplantation.⁶⁶ Another pioneer was Ramon Castroviejo from Spain, who did his first penetrating transplantation in 1936 in an advanced keratoconus patient, achieving significant improvement in vision. He also devised numerous instruments, which were very important for further improvement of corneal surgery in general.^{67,68}

Further development and standardization of transplantation techniques followed, while research concentrated on tissue preservation and preparation. In 1944, a first eye bank was established in New York by Townley Paton and, in 1961, the Eye Bank Association of America (EBAA) was founded.⁶⁹ This organization developed the standards for obtaining, preserving, storing and using corneal donor tissue.

Penetrating keratoplasty

The full-thickness replacement of the cornea by donor tissue (penetrating keratoplasty, PK), regardless of the healthy layers, was the preferred surgical treatment for all corneal disorders. Although initial poor graft survival rates after PK resulted in renewed interest in lamellar corneal transplants (lamellar keratoplasty, LK) around the 1950's,⁷⁰⁻⁷³ the development of corticosteroids for the treatment of allograft rejection, the introduction of antibiotics and operating microscopes, and the evolution of microsurgical instruments and newer suture materials all contributed to the increased popularity of PK. Since its introduction, different suturing techniques and graft-sizing practices have been described, but the largest advance in PK has been the recent introduction of the femtosecond laser to trephine the recipient and donor tissues, theoretically providing better apposition and faster healing.⁷⁴⁻⁷⁶

Technically, PK is often successful, initially providing a clear cornea. However, the postoperative course may be compromised due to complications derived from corneal sutures, ocular surface problems or wound dehiscence associated with incomplete wound healing.⁷⁷⁻⁸¹ As a result, PK is often accompanied by slow, insufficient, and unsatisfactory visual rehabilitation, which may be even further complicated by refractive problems such as anisometropia and high (irregular) astigmatism.^{77,80,82,83} Furthermore, complications such as continuous and increased endothelial cell decline and allograft rejection, which may both be provoked by other complications (e.g. sutures, corneal vascularization, glaucoma), may eventually result in graft failure.^{80,84-88}

The long-term graft survival rate after PK ranges from around 50% to 80% at 10 years postoperatively, depending on indication and complications.⁸⁴⁻⁸⁶ For uncomplicated Fuchs endothelial dystrophy or keratoconus, the likelihood of having a functional graft in the long run is relatively good,^{84,85} whereas corneas transplanted for pseudophakic or aphakic bullous keratopathy, infectious corneal ulcers and re-grafts may have a lower graft survival rate.^{80,86,88-90}

Lamellar keratoplasty

Problems frequently encountered with PK eventually led to renewed interest, further developments, and popularization of LK (Figure 5). In contrast to PK, the fundamental hypothesis behind LK was to replace only the diseased part of the cornea, leaving the recipient's healthy corneal layers intact and to resect the least amount of tissue, with less risk of rejection and more success. Anterior LK potentially improves graft survival rates by selectively replacing the diseased anterior corneal layers and retaining the healthy endothelium.^{87,91} The development of posterior LK procedures has dramatically enhanced the predictability and speed of visual rehabilitation in endothelial disorders.^{77,92} Posterior and anterior LK procedures will be separately discussed in part 1 and 2 of this thesis.




Figure 5. Schematic overview displaying (A) a virgin cornea, (B) penetrating keratoplasty (PK), and (C-E) the different lamellar keratoplasty techniques, i.e. (C) Deep anterior lamellar keratoplasty (DALK), (D) Descemet stripping (automated) endothelial keratoplasty (DS(A)EK), and (E) Descemet membrane endothelial keratoplasty (DMEK). (Source: van Dijk K, et al. Contact lenses after keratoplasty what to expect and what to look for with contact lens management in post-keratoplasty corneas. Contact Lens Spectrum 2014, August Issue).

PART I: SELECTIVE, MINIMAL INVASIVE TREATMENT OF ENDOTHELIAL DISORDERS

Endothelial disorders

The various types of endothelial disorders share the presence of posterior corneal alterations, i.e. in the endothelium and DM, which may eventually cause corneal swelling and

opacification and impair visual acuity. Different corneal endothelial dystrophies with their own unique pathophysiologic mechanism may be distinguished. In addition to the endothelial dystrophies, various insults to the endothelium may affect the function of the endothelium. In this thesis, the most common “endothelial” indications for corneal transplantation will be described.

Fuchs endothelial dystrophy

Fuchs endothelial dystrophy is the most common corneal dystrophy, frequently resulting in visual loss. It is a slowly progressive, bilateral, corneal disease and was first described a century ago by the Austrian ophthalmologist Ernst Fuchs.⁹³ Primarily, it is a disorder of the posterior cornea with the formation of focal excrescences of DM, termed ‘guttae’, which appear as round dark areas within the cellular monolayer on specular microscopy (Figure 6). This is accompanied by endothelial cell density decrease and abnormally enlarged, polymorphic cells surrounding the guttae.

With advancing disease, loss of endothelial cells results in progressive stromal edema and subsequent loss of vision, especially in the morning, as an initial symptom. In end-stage disease, the cornea will slowly generate painful epithelial bullae, evolving into subepithelial fibrosis and finally ending up with corneal vascularization.^{94,95}

Two clinical subtypes have been identified: early-onset Fuchs endothelial dystrophy, which is rare, presents within the first decade of life and progresses through the second to third decade; and the more common late-onset Fuchs endothelial dystrophy that




Figure 6. Specular microscopy images of advancing Fuchs endothelial dystrophy (left to right). (Source: Ham L. Descemet Membrane Endothelial Keratoplasty: Donor Tissue Preparation and Clinical Outcomes. Optima Grafische Communicatie Rotterdam, 2011).

starts at the second to third decade and becomes symptomatic during the fifth to sixth decade of life.^{94,96,97} Early-onset Fuchs endothelial dystrophy is accompanied by a thickened anterior banded DM zone. In late-onset Fuchs endothelial dystrophy, multiple layers of basement membrane-like material have been found on the posterior nonbanded part of DM.⁹⁷ Both subtypes appear to have a similar progression from disease onset until corneal decompensation, and have a female predominance at a ratio of around 3:1.⁹⁷

The pathological mechanism behind Fuchs endothelial dystrophy is not yet completely understood. The genetic basis appears complex and heterogeneous, demonstrating variable expressivity and incomplete penetrance. Autosomal dominant inheritance of Fuchs endothelial dystrophy occurs; however, in most cases, Fuchs endothelial dystrophy has not yet been found to be associated with a specific genetic mutation.⁹⁷ Additionally, environmental factors such as oxidative stress may play a role in the development or progression of the disorder by accelerating cell damage and apoptosis.⁹⁸⁻¹⁰²

Bullous keratopathy

Bullous keratopathy refers to the development of irreversible corneal edema due to endothelial decompensation as a complication secondary to trauma (most often surgical trauma, such as complicated cataract extraction), glaucoma or congenital abnormalities.¹⁰³ It may manifest in the immediate post-traumatic period, or symptoms may not present for many years. With increasing corneal edema, (sub)epithelial bullae may form and rupture, causing painful micro-defects in the corneal surface. Moreover, in advanced bullous keratopathy, subepithelial fibrosis, with or without BL-disruption and superficial vascularization, may frequently be observed.¹⁰⁴⁻¹⁰⁶

Corneal graft failure

Corneal graft failure can be defined as an irreversible loss of corneal clarity caused by a directly dysfunctional corneal graft (primary graft failure), or occurring years after corneal transplantation (secondary graft failure).¹⁰⁷ Risk factors include continuous and increased postoperative endothelial cell decline and allograft rejection episodes, which in turn can be provoked by complications such as (secondary) glaucoma, ocular surface problems, suture-related problems and corneal vascularization.^{80,81,84-86,88,108} Furthermore, transplantation indication, socio-economic and geographical factors play a role in the possible development of graft failure.¹⁰⁸⁻¹¹⁰

Management of endothelial dysfunction

Although the management of endothelial dysfunction may be somewhat different between entities and individual cases, in general, endothelial disorders may initially be treated with topical sodium chloride solutions, topical corticosteroids and reduction of

intraocular pressure.^{111,112} Furthermore, soft bandage lenses may be useful to relieve the pain in case of recurrent erosions caused by epithelial bullae.¹¹³ However, since corneal endothelial cells show minimal or no regeneration,¹ most cases ultimately require corneal transplant surgery. Traditionally, the standard surgical approach was to replace the full thickness of the cornea with donor tissue, as in PK. More recently, various types of posterior LK techniques have been introduced, among which is Descemet membrane endothelial keratoplasty (DMEK), the latest innovation in that field,^{114,115} and the subject of this thesis (part 1).

Posterior lamellar keratoplasty

The concept of posterior LK, also called endothelial keratoplasty (EK), was established in the 1950s and 1960s by Charles Tillett and Jose Barraquer.^{73,116} However, their attempts of posterior lamellar exchange underneath a manually-dissected stromal flap were often unsuccessful and were at that time perceived as technically more difficult than a PK. In 1998, Gerrit Melles came up with a breakthrough idea, as he demonstrated that a posterior donor lamella could adhere to the posterior recipient stroma with only an air bubble in the anterior chamber as support. The procedure, called posterior LK, required excision of a 7.0–7.5-mm diameter recipient posterior stromal button with attached endothelium, after which a similar size posterior donor button - containing the same tissue layers (stroma and endothelium) - could be inserted through a 9-mm limbal incision.¹¹⁷ The initial technique was modified in 2000 by Melles et al, using a 5-mm incision and folding the donor tissue like a ‘taco’ to enable insertion.¹¹⁸ Meanwhile, Mark Terry popularized both techniques as deep lamellar endothelial keratoplasty (DLEK).^{119,120} Promising results however were tempered by technical difficulty to manually dissect both donor and host stromal beds.

In 2002, Melles et al. further simplified the procedure by removing only the host DM and its endothelial cells by a scraping movement with a reversed Sinskey hook, called descemetorhexis. Descemetorhexis combined with a DLEK-graft is referred to as Descemet stripping endothelial keratoplasty (DSEK).¹²¹ When the donor tissue then was dissected with a microkeratome, as developed by Mark Gorovoy, the procedure was called Descemet stripping automated endothelial keratoplasty (DSAEK).¹²²

Adoption of EK accelerated after eye banks began to prepare the donor tissue for surgeons,¹²³ and in the last decennium, DSEK/DSAEK has replaced PK as the standard surgical technique for the treatment of corneal endothelial disorders. Compared to PK, EK resulted in a faster visual rehabilitation, with more predictable refractive outcomes and less postoperative astigmatism, while suture and wound-related problems could be avoided. At the same time, EK provided a better retained structural integrity of the eye, reducing the risk of catastrophic eye loss from suprachoroidal haemorrhage, both intra- and postoperatively.^{92,115}

Variability in visual outcomes after DSEK/DSAEK, however, weakened these advantages to some extent. Full visual recovery was not always achieved, despite having otherwise healthy eyes and clear corneas.^{92,114,115,124,125} Possible explanations include stromal scarring and fibrosis secondary to the underlying pathology and increased light scatter.^{48-50,126-129} In addition, optical degradation at the graft-recipient interface, increased corneal thickness and increased HOAs have been studied as causes for suboptimal visual outcomes after DSEK/DSAEK.¹³⁰⁻¹³⁶ While the anterior corneal surface is left uncompromised, posterior corneal HOAs after DSEK/DSAEK seem to be comparable to or even larger than post-PK.^{137,138} This may be secondary to graft decentration, curvature-mismatch between the recipient and the graft, graft wrinkling or uneven graft thickness from asymmetric trephination.^{132,133,139} It has been theorized that thinner posterior lamellar grafts would result in a more regular posterior graft surface, while better conforming to the recipient posterior corneal curvature and, consequently, their use could improve the visual outcome.^{132,134,139}

Descemet membrane endothelial keratoplasty

Melles et al. realized that EK could be further refined by selectively implanting a donor DM and endothelium without any stroma (such as in a DSEK/DSAEK graft).¹⁴⁰ The first successful case report of a technique, nowadays referred to as DMEK, was presented in 2006.¹⁴¹

As with other keratoplasty techniques, DMEK-graft preparation may be performed by the surgeon at the time of surgery or in an eye bank setting 1 to 2 weeks prior to the surgery. Precut tissue may save surgical time, avoids surgical postponement caused by unsuccessful tissue preparation and requires fewer investments in equipment and preparation skills. Furthermore, eye banks can perform all routine quality checks, such as endothelial cell layer evaluation and sterility testing before shipping the tissue.

The DM can be relatively easily stripped off due to the fragile interconnections with the posterior stroma, making it feasible to prepare isolated grafts of DM and its endothelium.^{142,143} When the DM is peeled from a donor corneo-scleral rim, it rolls up with the endothelium on its outer surface, probably owing to the elastic properties of the tissue.^{142,143}

The initially-described DMEK-graft preparation technique consisted of stripping a 9.5-mm diameter DM from a corneo-scleral rim submerged in saline.^{142,143} The method proved safe and reproducible, with less than 5% tissue loss and an acceptable endothelial cell decrease of 4% to 7%.¹⁴² In 2012, the preparation process became safer and easier with the introduction of the so called 'no-touch' DMEK-graft preparation procedure, in which a rim of trabecular meshwork tissue is left in-situ, and the DMEK-graft is trephined on an underlying soft contact lens.¹⁴⁴

A standardized 'no-touch' surgical technique for DMEK was published in 2011 by Dapena et al.¹⁴⁵ The technique comprised the creation of three side-ports. The recipient's DM was then scored and stripped from the posterior stroma, performing a 9.0-mm diameter descemetorhexis, with the anterior chamber filled with air. Subsequently, the DMEK-roll was thoroughly rinsed with balanced salt solution, stained with 0.06% trypan blue solution, and drawn into a custom-made injector for injection into the recipient anterior chamber through a 3.0-mm limbal tunnel incision at the 12:00 o'clock position. The graft was oriented with the endothelial side down (donor DM facing the recipient posterior stroma) by indirect manipulation with air and BSS, then gently spread out over the iris, after which it was positioned onto the recipient posterior stroma with an air bubble injected underneath.¹⁴⁵ At the end of the surgery, the anterior chamber was left completely filled with air for a period of one hour, followed by an air-liquid exchange to pressurize the eye, leaving a 30% to 50% air-bubble in the anterior chamber. Patients were asked to remain supine for 48 to 72 hours after surgery.¹⁴⁵

DMEK aims to restore the normal corneal anatomy by selective replacement of the diseased corneal layers only and, as such, is a truly disease-specific form of keratoplasty (Figure 7). Initial DMEK outcomes were promising with a relatively low complication rate (early graft detachment being the main complication), good visual results in the majority of patients, and endothelial cell survival similar to earlier keratoplasty techniques.^{115,146-151}




Figure 7. Slit-lamp photographs of an eye at 3 months after Descemet membrane endothelial keratoplasty for Fuchs endothelial dystrophy. Note the 'normal' anatomical restoration of the transplanted cornea as well as its clarity.

PART II: NEW TREATMENT OPTION FOR ADVANCED KERATOCONUS

Keratoconus

Keratoconus is the most common ectatic corneal disorder, often described as a bilateral, often asymmetrical, non-inflammatory, progressive disorder characterized by thinning and protrusion of the cornea, causing a compromised optical performance. The disorder generally becomes apparent during the second decade of life, during puberty, and progresses until the fourth decade, when it usually stabilizes.^{152,153} The established prevalence of keratoconus among the general population is approximately 1/2000,¹⁵⁴ but much higher rates have been mentioned in some parts of the world.^{155,156}

The pathogenesis of keratoconus, although not well understood, may include genetic, environmental and mechanical factors. Possibly, a genetic predisposition requires an environmental event to provoke progressive disease.¹⁵⁷⁻¹⁵⁹ Different associations have been described; specifically atopy, eye rubbing, positive family history, and several syndromes and diseases, such as Down syndrome, Ehlers-Danlos syndrome and Leber congenital amaurosis.^{152,153,157,158,160-164} Furthermore, new evidence suggests an overexpression of inflammatory mediators, including cytokines and interleukin 6 (IL-6), in cases with keratoconus.¹⁶⁵⁻¹⁶⁷ In addition, oxidative stress may play an important role in the pathogenesis of keratoconus.^{98,167}

Depending on the severity of keratoconus, all layers of the cornea may become involved in the pathological process.^{152,153} The corneal epithelial basal cells may degenerate and grow towards the BL. The BL often shows breakages, which are then filled with stromal collagen. In the stroma, a decrease in the number of lamellae and keratocytes, degradation of fibroblasts, changes in lamellar organization, and uneven distribution of the collagen fibrils have been observed.^{152,153,168,169} The DM and endothelium are usually unaffected. However, breakage of DM may occur in severe cases and elongation of endothelial cells pointing towards the cone has been reported.^{152,153}

The clinical presentation of keratoconus varies depending on disease severity. While early keratoconus can go unnoticed, keratoconus progression will cause subjective visual loss and photophobia, caused by corneal deformation and astigmatism. In moderate to advanced keratoconus cases, stromal thinning may accompany corneal ectasia, which is greatest at the apex of the cone (Figure 8). Vogt striae, fine vertical parallel lines in the posterior stroma and DM, and a Fleischer ring, iron deposits on the epithelial basal membrane, may be visible on slit-lamp examination. Furthermore, BL breakage may cause corneal scarring. In severe cases, a so-called 'Munson sign', a V-shaped distortion of the lower eyelid in down gaze, may be visible and corneal hydrops, causing excessive corneal edema, may occur due to sudden breaks of DM.^{152,153,170}

To grade keratoconus, the Amsler-Krumeich classification system is one of the oldest and still the most commonly used.¹⁷¹ With this classification, which is often used for




Figure 8. Slit-lamp photograph of an eye with advanced keratoconus. Note the deformation of the corneal curvature, as well as the thinning at the apex of the cone

research purposes and is integrated in the Pentacam software, the severity of keratoconus is graded from stage I to IV using spectacle refraction, central keratometry and pachymetry, and the presence or absence of corneal scarring.^{55,171} However, the Amsler-Krumeich scale relies on relatively 'outdated' parameters and it fails to address currently available information and technological advances. Furthermore, it does not deal with objective and subjective visual function. Therefore, various modifications to better diagnose or characterize the severity of keratoconus have been suggested, such as the use of corneal topometric irregularity indices, corneal HOAs, corneal thickness maps, anterior and posterior corneal elevation, and corneal biomechanical properties.¹⁷²⁻¹⁷⁸ Preferably, a classification system should combine clinical findings such as visual performance with corneal topometric and tomographic parameters.¹⁷⁹

Management of keratoconus

The first step in the management of keratoconus comprises optimizing patient's vision by means of optical correction. In early stages of the disorder, spectacles or soft contact lenses are a viable option, while in moderate to advanced cases, specially designed soft lenses, hybrid and rigid gas-permeable lenses are indicated. Scleral lenses become desirable to achieve visual rehabilitation in cases in which wearing the more traditional contact lens designs is not possible anymore (e.g. poor centration, instability, or low tolerability).¹⁸⁰

For patients with unacceptable vision, also while wearing a contact lens, in which the vision cannot be adequately restored by other means, corneal transplantation remains the standard of care. Two main types of corneal transplantation are available for the treatment of keratoconus: PK, and deep anterior lamellar keratoplasty (DALK). In the latter only the anterior corneal layers are replaced, preserving the patient's own endothelial cell layer.¹⁸¹

Anterior lamellar keratoplasty

Since the renewed interest in lamellar corneal transplants around the 1950's, there has been a group of ophthalmic surgeons, including Paufigue,¹⁸² Malbran,¹⁸³ and Anwar,¹⁸⁴ who performed anterior LK instead of PK for visual improvement in corneal diseases with normal corneal endothelium, such as keratoconus. However, it took until the mid-1980s before innovative procedures were developed and widely adopted to remove diseased corneal anterior stroma up to the DM,¹⁸⁵⁻¹⁸⁷ as in DALK. Different lamellar dissection techniques to achieve removal of all, or almost all, of the corneal stroma have been described, including intrastromal air injection,¹⁸⁵ big-bubble technique,¹⁸⁶ hydrodissection,¹⁸⁸ viscoelastic dissection,¹⁸⁹ and manual dissection.¹⁹⁰

Most of the currently used DALK techniques are based on the Anwar big bubble or the Melles manual dissection technique. The Anwar technique comprises a partial-depth trephination of the cornea, followed by forcibly injecting air deep in the stroma (creating a "big-bubble") to detach DM from the stroma.¹⁸⁶ With the Melles manual dissection technique the stroma is manually dissected away from the underlying DM using a series of a different sizes curved spatulas.¹⁹⁰ In the latter technique, the depth of dissection can be determined after filling the anterior chamber with air, by using the air-to-endothelium interface as a reference plane.¹⁹¹ A drawback of both techniques may be that perforation of DM is relatively frequent (4 – 30%),^{91,190,192-195} and, depending on the size of perforation, conversion to PK may be required. Because with the Melles manual DALK technique the dissection is more 'controlled' and, in the event of a perforation, the perforation site tends to be small, the procedure can often be completed. Otherwise, since no corneal surface incisions have been made at the time of the dissection, the operation can be aborted and reattempted at a later date.^{190,191}

A major advantage of DALK compared to PK is the maintenance of the recipient endothelium which results in much lower rates of endothelial cell density decrease and practically eliminates endothelial allograft rejection, suggesting a long lifetime for a DALK graft.¹⁹⁶ Furthermore, the retained recipient corneal endothelium and DM allow for better preservation of the ocular integrity, permitting earlier suture removal due to faster wound recovery and, consequently, fewer wound healing-related problems.^{77,91} As a consequence, DALK provides a somewhat faster visual rehabilitation, although final visual outcomes may be at best equal to PK, provided that stromal dissection reaches the level of DM.^{91,197-200} Similar to PK, DALK grafts require fixation by sutures, which may give rise to several complications such as cheese wiring, neovascularization, and suture loosening, potentially leading to a sequence of adverse events with disappointing visual results and graft failure.

Treatments to avoid corneal transplantation

Clinical observation, supported by recent research studies, suggests that eyes with advanced keratoconus may be prone to a sequence of (inflammatory) ocular surface reactions,^{152,201-204} which may render DALK and PK high-risk procedures. Both surgeries tend to worsen any existing ocular surface problems, as both involve surface incisions, corneal denervation, and placement of long-lasting sutures. Furthermore, keratoconus patients are generally young, and may possibly survive their corneal transplant. Consequently, it seems reasonable that many of these patients require more than one corneal graft during their life, with re-grafts having inferior clinical outcomes and worse graft survival than the initial corneal transplant.^{205,206} Another limitation of PK/DALK for the treatment of keratoconus may be that these treatment options do not halt progression of keratoconus, possibly due to ongoing ectatic progression of the recipient corneal rim, increasing graft-host interface misalignment, or recurrent disease in the donor button.²⁰⁷⁻²⁰⁹

As an alternative, UV-crosslinking has recently been introduced as a treatment for progressing early to moderate keratoconus stages, with the aim to strengthen the stromal collagenous corneal matrix and thereby delaying or avoiding further ectasia progression.^{210,211} The initially described protocol, the so called Dresden protocol, entails 30 minutes of soaking a de-epithelized cornea with the vitamin and photosensitizer Riboflavin, followed by UV-A irradiation for 30 minutes.²¹⁰ Currently, UV-crosslinking is indicated in cases with documented disease progression, a clear central cornea, and a corneal thickness of at least 400 μm after removal of the epithelium.²¹¹ Although techniques are being developed to also treat thinner corneas (such as the use of hypotonic Riboflavin to swell the cornea), at present there have been limited studies of the efficacy and safety of UV-crosslinking in thin corneas, with relatively few included eyes with severe thinning (<350 μm).²¹²⁻²¹⁵ Furthermore, the risk of complications or treatment failure seems higher in steeper keratoconus corneas (>58D).^{216,217}

Another possibility to potentially postpone corneal transplantation in ectatic eyes may be by reshaping the cornea using intracorneal ring segments (ICRS), alone or in combination with UV-crosslinking.²¹⁸⁻²²⁰ ICRS are made of polymethylmethacrylate plastic and are available in numerous arc-lengths, thicknesses, and designs. The segments are placed into corneal stromal tunnels which may be created manually, or automatically with a femtosecond laser.²²⁰ By normalizing the corneal contour, ICRS may enable a contact lens-intolerant patient to become contact lens-tolerant again.^{218,221,222} Beyond this, (un)corrected visual acuity may show a modest improvement.^{220,223-225} Still, eyes with severe corneal thinning or steepening seem currently ineligible for ICRS secondary to the relatively higher rate of complications and poorer visual outcomes.²²⁶

Advanced keratoconus eyes, showing continuous progression, but being ineligible for either UV-crosslinking or ICRS, may still reach satisfactory vision with well fitted contact

lenses, most often scleral lenses. These eyes may therefore similarly benefit from stabilizing the cornea to preserve the vision and to enable continued contact lens wear, and consequently postpone or avoid the need for PK/DALK. However, a procedure that could halt keratoconus progression in these corneas was not available (Figure 9). Prevention of disease (progression) may be the first step in medical care, hence developing a treatment to also halt progression in this group of keratoconus patients was a main objective.




Figure 9. Keratoconus management decision tree. To potentially avoid or postpone the need for penetrating or deep anterior lamellar keratoplasty (PK/DALK), UV-crosslinking may be the preferred treatment for progressive keratoconus in cases with still acceptable vision. Furthermore, intracorneal ring segments (ICRS) seem viable to optimize vision and/or contact lens tolerance. However, very thin corneas seem ineligible for either UV-crosslinking or ICRS and are therefore allowed to continue to progress, with the risk that eventually PK or DALK becomes essential.

(ICRS indicates intracorneal ring segments; PK, penetrating keratoplasty; DALK, deep anterior lamellar keratoplasty; TPT, thinnest point thickness, measured after removal of the epithelium; μm , micrometers; *hypo-tonic Riboflavin should be used if TPT is between 350 - 400 μm .²²⁷)

OUTLINE OF THE THESIS

This thesis evaluates the feasibility and clinical outcomes of DMEK for managing endothelial disorders, and the use of BL grafts, i.e. Bowman layer transplantation, in the management of advanced keratoconus.

The first part concerns the outcomes of DMEK. We hypothesized that selective transplantation of only a DM and its endothelium would provide a fast and unprecedented high rate of optical recovery. In order to test this hypothesis, the first part of this research has been designed to accomplish the following objectives:

In Chapter 2, the feasibility and efficacy of DMEK for the management of Fuchs endothelial dystrophy, bullous keratopathy or previous corneal transplant failure is assessed by evaluating the clinical outcome before and up to 6 months after surgery and by documenting intra- and postoperative complications in a large consecutive series of DMEK eyes.

In Chapter 3, the incidence and causes of anterior corneal surface irregularities associated with visual complaints after successful DMEK surgery are reported and the efficacy of contact lens fitting in these cases is evaluated.

In Chapter 4, corneal higher-order aberrations, backscattered light, and their correlation with visual acuity outcome in a large series of DMEK eyes before and 6 months after surgery for Fuchs endothelial dystrophy are evaluated and compared with an age-matched control group of healthy subjects with no history of ocular disease.

In Chapter 5, the two-year refractive outcomes after DMEK in a larger series of pseudophakic eyes, undergoing DMEK for Fuchs endothelial dystrophy, are monitored. Furthermore, factors influencing the pre- to postoperative refractive changes and the time point of stabilization are determined.

The second part concerns the outcomes of BL transplantation, a newly designed procedure for advanced keratoconus cases, ineligible for UV-crosslinking or ICRS. We hypothesized that partial restoration of the corneal anatomy in an advanced keratoconus cornea might be obtained through implantation of an isolated BL graft, since fragmentation of the recipient's BL is one of the pathognomonic features in advanced keratoconus. BL transplantation aims to stabilize the ectasia and, at the same time, to preserve the patient's vision and enable continued contact lens wear. The procedure should be free from short- and long-term complications that both PK and DALK frequently entail. This part includes the following objectives:

In Chapter 6, the surgical approach for midstromal implantation of an isolated BL graft is described. Furthermore, the ability of this surgical approach to reduce ectasia and to preserve patient's vision is evaluated.

In Chapter 7, the clinical outcome of the first 22 mid-stromal BL transplantations is evaluated and the potential impact of preoperative corneal characteristics on the anatomic effect of the surgery is determined.

In Chapter 8, the clinical outcome up to 5 years after BL transplantation is evaluated to determine whether stabilization of ectasia in advanced keratoconus may be achieved long term.

The different outcomes are discussed in Chapters 9 and 10.

REFERENCES

1. DelMonte DW, Kim T. Anatomy and physiology of the cornea. *J Cataract Refract Surg* 2011; 37:588-598
2. Streilein JW. New thoughts on the immunology of corneal transplantation. *Eye* 2003;17:943-948
3. Reinstein DZ, Archer TJ, Gobbe M, Silverman RH, Coleman DJ. Epithelial thickness in the normal cornea: three-dimensional display with Artemis very high-frequency digital ultrasound. *J Refract Surg* 2008;24:571-581
4. Thoft RA, Friend J. The X, Y, Z hypothesis of corneal epithelial maintenance. Letter to the editor. *Invest Ophthalmol Vis Sci* 1983;24:1442-1443
5. McDermott ML. The corneal epithelium. In: Podos SM and Yanoff M (Ed.). *Textbook of Ophthalmology*. London: Mosby- Europe Ltd, 1994. Vol 8
6. Dohlman, C.H. The function of the corneal epithelium in health and disease. *Invest. Ophthalmol* 1971;10:383-407
7. Navarro R. The Optical Design of the Human Eye: a Critical Review. *J Optom.* 2009;2:3–18
8. Marshall J. The 2014 Bowman Lecture - Bowman's and Bruch's: a tale of two membranes during the laser revolution. *Eye (Lond)* 2015;29:46-64
9. Germundsson J, Karanis G, Fagerholm P, Lagali N. Age-related thinning of Bowman's layer in the human cornea in vivo. *Invest Ophthalmol Vis Sci* 2013;54:6143-6149
10. Gordon MK, Foley JW, Birk DE, Fitch JM, Linsenmayer TF. Type V collagen and Bowman's membrane. Quantitation of mRNA in corneal epithelium and stroma. *J Biol Chem* 1994;269:24959-24966
11. Wilson SE, Hong JW. Bowman's layer structure and function: critical or dispensable to corneal function? A hypothesis. *Cornea* 2000;19:417-420
12. Lie J, Droutsas K, Ham L. Isolated Bowman layer transplantation to manage persistent subepithelial haze after excimer laser surface ablation. *J Cataract Refract Surg* 2010;36:1036-1041
13. Melles GRJ, Binder PS, Moore MN, Anderson JA. Epithelial-stromal interactions in human keratotomy wound healing. *Arch Ophthalmol* 1995;113:1124-1130
14. Netto MV, Mohan RR, Sinha S, Sharma A, Dupps W, Wilson SE. Stromal haze, myofibroblasts, and surface irregularity after PRK. *Exp Eye Res* 2006; 82:788-797
15. Abahussin M, Hayes S, Knox Cartwright NE, Kamma-Lorger CS, Khan Y, Marshall J, Meek KM. 3D collagen orientation study of the human cornea using X-ray diffraction and femtosecond laser technology. *Invest Ophthalmol Vis Sci* 2009;50:5159-5164
16. Maurice DM. The cornea and sclera, In: Davidson H (Ed.). *The Eye*. New York Academic, 1984
17. Komai Y, Ushiki T. The three-dimensional organisation of collagen fibrils in the human cornea and sclera. *Invest. Ophthalmol. Vis. Sci* 1991;32:2244-2258
18. Bergmanson JP, Horne J, Doughty MJ, Garcia M, Gondo M. Assessment of the number of lamellae in the central region of the normal human corneal stroma at the resolution of the transmission electron microscope. *Eye Contact Lens* 2005;31:281-287
19. Radner W, Zehetmayer M, Aufreiter ., Mallinger R. Interlacing and crossangle distribution of collagen lamellae in the human cornea. *Cornea* 1998;17:537-543
20. Morishige N, Petroll WM, Nishida T, Kenney MC, Jester JV. Non-invasive corneal stromal collagen imaging using two-photon-generated second-harmonic signals. *J. Cataract Refract. Surg* 2006;32:1784-1791
21. Radner W, Mallinger R. Interlacing of collagen lamellae in the midstroma of the human cornea. *Cornea* 2002;21:598-601

22. Müller LJ, Pels E, Vrensen GFJM. The specific architecture of the anterior stroma accounts for maintenance of corneal curvature. *Br. J. Ophthalmol* 2001;85:437-443
23. Meek KM, Boote C. The organization of collagen in the corneal stroma. *Exp Eye Res* 2004;78:503-512
24. Meek KM, Leonard DW, Connon CJ, Dennis S, Khan S. Transparency, swelling and scarring in the corneal stroma. *Eye (Lond.)* 2003;17:927-936
25. Meek KM, Knupp C. Corneal structure and transparency. *Prog Retin Eye Res* 2015;49:1-16
26. Edelhauser HF. The balance between corneal transparency and edema: the Proctor Lecture. *Invest Ophthalmol Vis Sci* 2006;47:1754-1767
27. Hassell JR, Birk DE. The molecular basis of corneal transparency. *Exp Eye Res* 2010;91:326-335
28. Chen Y, Thompson DC, Koppaka V, Jester JV, Vasiliou V. Ocular aldehyde dehydrogenases: protection against ultraviolet damage and maintenance of transparency for vision. *Prog Retin Eye Res* 2013;33:28-39
29. Bourne WM, Johnson DH, Campbell RJ. The ultrastructure of Descemet's membrane. III. Fuchs' dystrophy. *Arch Ophthalmol* 1982;100:1952-1955
30. Johnson DH, Bourne WM, Campbell RJ. The ultrastructure of Descemet's membrane. I. Changes with age in normal corneas. *Arch Ophthalmol* 1982;100:1942-1947
31. Dua HS, Faraj LA, Said DG, Gray T, Lowe J. Human corneal anatomy redefined: a novel pre-Descemet layer (Dua's layer). *Ophthalmology* 2013;120:1778-1785.
32. Mergler S, Pleyer U. The human corneal endothelium: New insights into electrophysiology and ion channels. *Progress in Retinal and Eye Research* 2007;26:359-378
33. Dawson DG, Geroski DH, Edelhauser HF. Corneal endothelium: structure and function in health and disease. In: Brightbill FS (Ed). *Corneal Surgery: Theory, Technique and Tissue*, ed 4. St Louis, Mosby Elsevier; 2009:57-70
34. Elhalis H, Azizi B, Jurkunas UV. Fuchs endothelial corneal dystrophy. *Ocul Surf* 2010;8:173-184
35. Gordon SR. Cytological and immunocytochemical approaches to the study of corneal endothelial wound repair. *Prog Histochem Cytochem* 1994;28:1-64.
36. Sicam VA, Dubbelman M, van der Heijde RG. Spherical aberration of the anterior and posterior surfaces of the human cornea. *J Opt Soc Am A Opt Image Sci Vis* 2006;23:544-549
37. Ho JD, Liou SW, Tsai RJ, Tsai CY. Effects of aging on anterior and posterior corneal astigmatism. *Cornea* 2010;29:632-637
38. Ueno Y, Hiraoka T, Beheregaray S, Miyazaki M, Ito M, Oshika T. Age-related changes in anterior, posterior, and total corneal astigmatism. *J Refract Surg* 2014;30:192-197
39. Koch DD, Ali SF, Weikert MP, Shirayama M, Jenkins R, Wang L. Contribution of posterior corneal astigmatism to total corneal astigmatism. *J Cataract Refract Surg* 2012;38:2080-2087
40. Dubbelman M, Sicam VA, Van der Heijde GL. The shape of the anterior and posterior surface of the aging human cornea. *Vision Res* 2006;46:993-1001
41. Pepose JS, Applegate RA. Making sense out of wavefront sensing. *Am J Ophthalmol* 2005;139:335-343
42. Lawless MA, Hodge C. Wavefront's role in corneal refractive surgery. *Clin Exp Ophthalmol* 2005;33:199-209
43. Applegate RA, Sarver EJ, Khemsara V. Are all aberrations equal? *J Refract Surg* 2002; 18:S556-S562
44. Applegate RA, Ballentine C, Gross H, Sarver EJ, Sarver CA. Visual acuity as a function of Zernike mode and level of root mean square error. *Optom Vis Sci* 2003; 80:97-105
45. Maeda N. Clinical applications of wavefront aberrometry: a review. *Clin Exp Ophthalmol* 2009;37:118-129

46. McLaren JW, Bourne WM, Patel SV. Standardization of corneal haze measurement in confocal microscopy. *Invest Ophthalmol Vis Sci* 2010;51:5610-5616
47. Hillenaar T, van Cleynenbreugel H, Remeijer L. How normal is the transparent cornea? Effects of aging on corneal morphology. *Ophthalmology* 2012;119:241-248
48. McLaren JW, Patel SV. Modeling the effect of forward scatter and aberrations on visual acuity after endothelial keratoplasty. *Invest Ophthalmol Vis Sci* 2012;53:5545-5551
49. Patel SV, Baratz KH, Hodge DO, Maguire LJ, McLaren JW. The effect of corneal light scatter on vision after Descemet stripping with endothelial keratoplasty. *Arch Ophthalmol* 2009;127:153-160
50. Baratz KH, McLaren JW, Maguire LJ, Patel SV. Corneal haze determined by confocal microscopy 2 years after Descemet stripping with endothelial keratoplasty for Fuchs corneal dystrophy. *Arch Ophthalmol* 2012;130:867-874
51. Klein SA. Axial curvature and the skew ray error in corneal topography. *Optom Vis Sci* 1997;74:931-944
52. Vos FM, Groen FCA, van Stokkum IHM, et al. A new instrument to measure the shape of the cornea based on pseudorandom color coding. *IEEE Trans Instrum Meas* 1997;46:794-797
53. Arni V, Sicam P, Van Der Heijde RG. Topographer reconstruction of the non-rotation symmetric anterior corneal surface features. *Optom Vis Sci* 2006;83:910-918
54. Oliveira CM, Ribeiro C, Franco S. Corneal imaging with slit scanning and Scheimpflug imaging techniques. *Clin Exp Optom* 2011;94:33-42
55. Oculus Optikgeräte GmbH. Oculus Pentacam instruction manual: measurement and evaluation system for the anterior segment of the eye. Wetzlar, Germany: Oculus Optikgeräte GmbH;2005
56. Ramos JL, Li Y, Huang D. Clinical and research applications of anterior segment optical coherence tomography-a review. *Clin Experiment Ophthalmol* 2009;37:81-89
57. Kiernan DF, Mieler WF, Hariprasad SM. Spectral-domain optical coherence tomography:a comparison of modern high resolution retinal imaging systems. *Am J Ophthalmol* 2010;149:18-31
58. McCarey B, Edelhauser HF, Lynn MJ. Review of corneal endothelial specular microscopy for FDA clinical trials of refractive procedures, surgical devices and new intraocular drugs and solutions. *Cornea*. 2008; 27:1-16
59. Efron N, Perez-Gomez I, Motalib HA, et al. Confocal microscopy of the normal human cornea. *Cont Lens Anterior Eye* 2001;24:16-24
60. Niederer RL, McGhee CNJ. Clinical in vivo confocal microscopy of the human cornea in health and disease. *Prog Retin Eye Res* 2010;29:30-58
61. Eye Bank Association of America. 2015 Eye Banking Statistical Report. Washington, D.C., United States, 2016. Available at <http://restoresight.org/wp-content/uploads/2016/03/2015-Statistical-Report.pdf>
62. Pellier de Quengsy G. Paris: Didot; 1789. Précis ou cours d'opérations sur la chirurgie des yeux par M.G. Pellier de Quengsy, fils.
63. Reisinger F. Die keratoplastik, ein versuch zur enweiterund der augenheilkunde. *Bayerische Annalem* 1824;1:207
64. von Hippel A. Eine neue Methode der Hornhauttransplantation. *Albrecht von Graefes Arch Klin Exp Ophthalmol* 1888;34:108-130
65. Zirm EK: Eine erfolgreiche totale keratoplastik. *V. Graefes Arch Ophthalmol* 1906;64:580-593
66. Filatov VP. Transplantation of the cornea. *Arch Ophthalmol*.1935;13,321-347
67. Castroviejo R. Keratoplasty *Am J Ophthalmol*. 1941;24:1-20
68. Murube J. Ramon Castroviejo centenary: a life dedicated to corneal transplantation. *Surv Ophthalmol* 2005;50:215-225

69. Paton D. The founder of the first eye bank: R. Townley Paton, MD. *Refract Corneal Surg* 1991;7:190-194
70. Hallermann W. Verschiedenes uber keratoplastik. *Klin Monstabl Augenheilkd* 1959;135:252-259
71. Hallermann W. On the technic of lamellar keratoplasty. *Klin Monstabl Augenheilkd* 1963;142:243-250
72. Barraquer JI. Lamellar keratoplasty. (Special techniques). *Ann Ophthalmol* 1972;4:437-469
73. Tillett CW. Posterior lamellar keratoplasty. *Am J Ophthalmol* 1956;41:530-533
74. Farid M, Kim M, Steinert RF. Results of penetrating keratoplasty performed with a femtosecond laser zigzag incision initial report. *Ophthalmology* 2007;114:2208-2212
75. Price FW Jr, Price MO. Femtosecond laser shaped penetrating keratoplasty: one-year results utilizing a top-hat configuration. *Am J Ophthalmol* 2008;145:210-214
76. Bahar I, Kaiserman I, McAllum P, Rootman D. Femtosecond laser-assisted penetrating keratoplasty: stability evaluation of different wound configurations. *Cornea* 2008;27:209-211
77. Tan DT, Dart JK, Holland EJ, Kinoshita S. Corneal transplantation. *Lancet* 2012;379:1749-1761
78. Elder MJ, Stack RR. Globe rupture following penetrating keratoplasty. How often, why, and what can we do to prevent it? *Cornea* 2004;23:776-780
79. Nagra PK, Hammersmith KM, Rapuano CJ, et al. Wound dehiscence after penetrating keratoplasty. *Cornea* 2006;25:132-135
80. Williams KA, Muehlberg SM, Lewis RF, Coster DJ. How successful is corneal transplantation? A report from the Australian Corneal Graft Register. *Eye* 1995;9:219-227
81. Christo CG, van Rooij J, Geerards AJ, Remeijer L, Beekhuis WH. Suture-related complications following keratoplasty: a 5-year retrospective study. *Cornea* 2001;20:816-819
82. Claesson M1, Armitage WJ, Fagerholm P, Stenevi U. Visual outcome in corneal grafts: a preliminary analysis of the Swedish Corneal Transplant Register. *Br J Ophthalmol* 2002;86:174-180
83. Kirkness CM, Ficker LA, Steele AD, et al. Refractive surgery for graft-induced astigmatism after penetrating keratoplasty for keratoconus. *Ophthalmology* 1991;98:1786-1792
84. Felipe AF, Hammersmith KM, Nottage JM, Rapuano CJ, Nagra PK, Cohen EJ, Laibson PR. Indications, visual outcome, and ectasia in clear corneal transplants 20 years old or more. *Cornea* 2013;32:602-607
85. Niziol LM, Musch DC, Gillespie BW, Marcotte LM, Sugar A. Long-term outcomes in patients who received a corneal graft for keratoconus between 1980 and 1986. *Am J Ophthalmol* 2013;155:213-219
86. Ing JJ, Ing HH, Nelson LR, Hodge DO, Bourne WM. Ten-year postoperative results of penetrating keratoplasty. *Ophthalmology* 1998;105:1855-1865
87. Borderie VM, Guilbert E, Touzeau O, Laroche L. Graft rejection and graft failure after anterior lamellar versus penetrating keratoplasty. *Am J Ophthalmol* 2011;151:1024-1029
88. Inoue K, Amano S, Oshika T, Tsuru T. Risk factors for corneal graft failure and rejection in penetrating keratoplasty. *Acta Ophthalmol Scand* 2001;79:251-255
89. Williams KA, Lowe MT, Bartlett CM, et al. The Australian corneal graft registry, 2007 report Adelaide, Australia: Flinders University Press; 2007
90. Thompson RW jr, Price MO, Bowers PJ, Price FW jr. Long-term graft survival after penetrating keratoplasty. *Ophthalmology* 2003;110:1396-1402
91. Reinhart WJ, Musch DC, Jacobs DS, et al. Deep anterior lamellar keratoplasty as an alternative to penetrating keratoplasty: a report by American Academy of Ophthalmology. *Ophthalmology* 2011;118:209-218

92. Price MO, Price FW Jr. Endothelial keratoplasty - a review. *Clin Experiment Ophthalmol* 2010;38:128-140
93. Fuchs E. Dystrophia epithelialis corneae. *Albr von Graefe's Arch für Ophthalmol* 1910;76:478-508
94. Krachmer JH, Purcell JJ Jr, Young CW, Bucher KD. Corneal endothelial dystrophy: A study of 64 families. *Arch Ophthalmol* 1978;96:2036-2039
95. Adamis AP, Filatov V, Tripathi BJ, Tripathi RC: Fuchs' endothelial dystrophy of the cornea. *Surv Ophthalmol* 1993;38:149-168
96. Vedana G, Villarreal G Jr, Jun AS. Fuchs endothelial corneal dystrophy: current perspectives. *Clinical Ophthalmology* 2016;10 321-330
97. Weiss JS, Möller HU, Aldave AJ, et al. IC3D classification of corneal dystrophies – edition 2. *Cornea* 2015;34:117–159
98. Wojcik KA, Kaminska A, Blasiak J, Szaflik J, Szaflik JP. Oxidative stress in the pathogenesis of keratoconus and Fuchs endothelial corneal dystrophy. *Int. J. Mol. Sci* 2013;14:19294-19308
99. Bruinsma M, Tong CM, Melles GRJ. What does the future hold for the treatment of Fuchs endothelial dystrophy; will 'keratoplasty' still be a valid procedure? *Eye* 2013;27:1115-1122
100. Azizi B, Ziaei A, Fuchsluger T, Schmedt T, Chen Y, Jurkunas UV. p53-regulated increase in oxidative-stress-induced apoptosis in Fuchs endothelial corneal dystrophy: a native tissue model. *Invest Ophthalmol Vis Sci* 2011;52:9291-9297
101. Jurkunas UV, Bitar MS, Funaki T, Azizi B. Evidence of oxidative stress in the pathogenesis of fuchs endothelial corneal dystrophy. *Am J Pathol* 2010;177:2278-2289
102. Borderie VM, Baudrimont M, Valle ´e A, Ereau TL, Gray F, Laroche L. Corneal endothelial cell apoptosis in patients with Fuchs' dystrophy. *Invest Ophthalmol Vis Sci* 2000;41:2501-2505
103. Morishige N, Sonoda KH. Bullous keratopathy as a progressive disease: evidence from clinical and laboratory imaging studies. *Cornea*. 2013 Nov;32 Suppl 1:S77-83
104. Gonçalves ED, Paris Fdos S, Gomes JÁ, Kanecadan L, Campos M. Bullous keratopathy. *Ophthalmology* 2011;118:2303-2303
105. Obata H, Tsuru T. Corneal wound healing from the perspective of keratoplasty specimens with special reference to the function of the Bowman layer and Descemet membrane. *Cornea* 2007;26:S82-89
106. Morishige N, Yamada N, Zhang X, Morita Y, Yamada N, Kimura K, Takahara A, Sonoda KH. Abnormalities of stromal structure in the bullous keratopathy cornea identified by second harmonic generation imaging microscopy. *Invest Ophthalmol Vis Sci* 2012;53:4998-5003
107. Baydoun L, Melles GR. Refining the Terminology of Graft Failure in Reports on Endothelial Keratoplasty Outcomes. *JAMA Ophthalmol* 2016;134:125-126
108. Borderie VM, Boëlle PY, Touzeau O, Allouch C, Boutboul S, Laroche L. Predicted long-term outcome of corneal transplantation. *Ophthalmology* 2009;116:2354-2360
109. Tabin GC, Gurung R, Paudyal G, et al. Penetrating keratoplasty in Nepal. *Cornea* 2004;23:589-596
110. Pineda R. Corneal transplantation in the developing world: Lessons learned and meeting the challenge. *Cornea* 2015;0:1-6
111. Bergmanson JP, Sheldon TM, Goosey JD. Fuchs' endothelial dystrophy: a fresh look at an aging disease. *Ophthalmic Physiol Opt* 1999;19:210-222
112. Insler MS, Benefield DW, Ross EV. Topical hyperosmolar solutions in the reduction of corneal edema. *CLAO J* 1987;13:149-151
113. Arora R, Jain S, Monga S, Narayanan R, Raina UK, Mehta DK. Efcacy of continuous wear Pure Vision contact lenses for therapeutic use. *Cont Lens Anterior Eye* 2004;27:39-43
114. Melles GRJ. Posterior lamellar keratoplasty. DLEK to DSEK to DMEK. *Cornea* 2006;25:879-881

115. Dapena I, Ham L, Melles GRJ. Endothelial keratoplasty: DSEK/DSAEK or DMEK -the thinner the better? *Curr Opin Ophthalmol* 2009;20:299-307
116. Barraquer J. Special methods in corneal surgery. In: King JH, McTigue JW, eds. *The Cornea World Congress*. Washington, DC: Rutterworths, 1965;586-604
117. Melles GR, Eggink FA, Lander F et al. A surgical technique for posterior lamellar keratoplasty. *Cornea* 1998;17:618-626
118. Melles GR, Lander F, Nieuwendaal C. Sutureless, posterior lamellar keratoplasty: A case report of a modified technique. *Cornea* 2002;21:325-327
119. Terry MA, Ousley PJ. Deep lamellar endothelial keratoplasty in the first United States patients: early clinical results. *Cornea* 2001;20:239-243
120. Terry MA, Ousley PJ. Small-incision deep lamellar endothelial keratoplasty (DLEK): Six-month results in the first prospective clinical study. *Cornea* 2005;24:59-65
121. Melles GR, Wijdh RH, Nieuwendaal CP. A technique to excise the Descemet membrane from a recipient cornea (Descemetorhexis). *Cornea* 2004;23:286-288
122. Gorovoy MS. Descemet-stripping automated endothelial keratoplasty. *Cornea* 2006;25:886-889
123. Price MO, Baig KM, Brubaker JW, et al. Randomized, prospective comparison of precut vs surgeon-dissected grafts for Descemet stripping automated endothelial keratoplasty. *Am Ophthalmol* 2008;146:36-41
124. Anshu A, Price MO, Tan DTH, Price FW Jr. Endothelial keratoplasty: a revolution in evolution? *Surv Ophthalmol* 2012;57:236-252
125. Li JY, Terry MA, Goshe J, et al. Three-year visual acuity outcomes after Descemet's stripping automated endothelial keratoplasty. *Ophthalmology* 2012;119:1126-1129
126. Patel SV, McLaren JW. In vivo confocal microscopy of Fuchs endothelial dystrophy before and after endothelial keratoplasty. *JAMA Ophthalmol* 2013;131:611-618
127. Morishige N, Chikama T, Yamada N, et al. Effect of preoperative duration of stromal oedema in bullous keratopathy on early visual acuity after endothelial keratoplasty. *J Cataract Refract Surg* 2012;38:303-308
128. Morishige N, Yamada N, Teranishi S, et al. Detection of subepithelial fibrosis associated with corneal stromal edema by second harmonic generation imaging microscopy. *Invest Ophthalmol Vis Sci* 2009;50:3145-3150
129. Patel SV, McLaren JW, Hodge DO, Baratz KH. Scattered light and visual function in a randomized trial of deep lamellar endothelial keratoplasty and penetrating keratoplasty. *Am J Ophthalmol* 2008;145:97-105
130. Patel SV, Baratz KH, Maguire LJ, et al. Anterior corneal aberrations after Descemet's stripping endothelial keratoplasty for Fuchs' endothelial dystrophy. *Ophthalmology* 2012;119:1522-1529
131. Heinzlmann S, Bohringer D, Maier PC, Reinhard T. Correlation between visual acuity and interface reflectivity measured by pentacam following DSAEK. *Acta Ophthalmol* 2014;92:e1-4
132. Dirisamer M, Parker J, Naveiras M, et al. Identifying causes for poor visual outcome after DSEK/DSAEK following secondary DMEK in the same eye. *Acta Ophthalmol* 2013;91:131-139
133. Muftuoglu O, Prasher P, Bowman RW, et al. Corneal higherorder aberrations after Descemet's stripping automated endothelial keratoplasty. *Ophthalmology* 2010;117:878-884
134. Neff KD, Biber JM, Holland EJ. Comparison of central corneal graft thickness to visual acuity outcomes in endothelial keratoplasty. *Cornea* 2011;30:388-391
135. Seery LS, Nau CB, McLaren JW, et al. Graft thickness, graft folds, and aberrations after Descemet stripping endothelial keratoplasty for Fuchs dystrophy. *Am J Ophthalmol* 2011;152:910-916

136. Shinton AJ, Tsatsos M, Konstantopoulos A, Goverdhan S, Elsahn AF, Anderson DF, Hossain P. Impact of graft thickness on visual acuity after Descemet's stripping endothelial keratoplasty. *Br J Ophthalmol* 2012;96:246-249
137. Chamberlain W, Omid N, Lin A, et al. Comparison of corneal surface higher-order aberrations after endothelial keratoplasty, femtosecond laser-assisted keratoplasty, and conventional penetrating keratoplasty. *Cornea* 2011;31:6-13
138. Koh S, Maeda N, Nakagawa T, et al. Characteristic higherorder aberrations of the anterior and posterior corneal surfaces in 3 corneal transplantation techniques. *Am J Ophthalmol* 2012;153:284-290
139. Letko E, Price DA, Lindoso EMS, et al. Secondary graft failure and repeat endothelial keratoplasty after Descemet's stripping automated endothelial keratoplasty. *Ophthalmology* 2011;118:310-314
140. Melles GR, Lander F, Rietveld FJ. Transplantation of Descemet's membrane carrying viable endothelium through a small scleral incision. *Cornea* 2002;21:415-418
141. Melles GR, Ong TS, Ververs B, van der Wees J. Descemet membrane endothelial keratoplasty (DMEK). *Cornea* 2006;25:987-990
142. Lie JT, Birbal R, Ham L, van der Wees J, Melles GRJ. Donor tissue preparation for Descemet membrane endothelial keratoplasty. *J Cataract Refract Surg* 2008;34:1578-1583
143. Lie JT, Groeneveld EA, Ham L, van der Wees J, Melles G.R.J. More efficient use of donor corneal tissue with Descemet membrane endothelial keratoplasty (DMEK). *Br J Ophthalmol* 2010;94:1265-1266
144. Groeneveld-van Beek EA, Lie JT, van der Wees J, Bruinsma M, Melles GR. Standardized 'no-touch' donor tissue preparation for DALK and DMEK: harvesting undamaged anterior and posterior transplants from the same donor cornea. *Acta Ophthalmol* 2013;91:145-150
145. Dapena I, Moutsouris K, Droutsas K, Ham L, van Dijk K, Melles GR. Standardized "no-touch" technique for Descemet membrane endothelial keratoplasty. *Arch Ophthalmol* 2011;129:88-94
146. Melles GR, Ong TS, Ververs B, van der Wees J. Preliminary clinical results of Descemet membrane endothelial keratoplasty. *Am J Ophthalmol* 2008;145:222-227
147. Ham L, Dapena I, van Luijk C, van der Wees J, Melles GR. Descemet membrane endothelial keratoplasty (DMEK) for Fuchs endothelial dystrophy: review of the first 50 consecutive cases. *Eye (Lond)* 2009;23:1990-1998
148. Ham L, Balachandran C, Verschoor CA, van der Wees J, Melles GR. Visual rehabilitation rate after isolated descemet membrane transplantation: descemet membrane endothelial keratoplasty. *Arch Ophthalmol*. 2009;127:252-255
149. Ham L, Dapena I, van der Wees J, Melles GR. Endothelial cell density after Descemet membrane endothelial keratoplasty: 1- to 3-year follow-up. *Am J Ophthalmol* 2010;149:1016-1017
150. Dapena I, Moutsouris K, Ham L, Melles GR. Graft detachment rate. *Ophthalmology* 2010;117:847
151. Dapena I, Ham L, Droutsas K, et al. Learning curve in Descemet's membrane endothelial keratoplasty: First series of 135 consecutive cases. *Ophthalmology* 2011;118:2147-2154
152. Romero-Jiménez M, Santodomingo-Rubido J, Wolffsohn JS. Keratoconus: a review. *Cont Lens Anterior Eye* 2010;33:157-166
153. Rabinowitz YS. Keratoconus. *Surv Ophthalmol* 1998;42:297-319
154. Kennedy RH, Bourne WM, Dyer JA. A 48-year clinical and epidemiologic study of keratoconus. *Am J Ophthalmol* 1986;101:267-273
155. Jonas JB, Nangia V, Matin A, et al. Prevalence and associations of keratoconus in rural maharashtra in central India: the central India eye and medical study. *Am J Ophthalmol* 2009;148:760-765

156. Gokhale NS. Epidemiology of keratoconus. *Indian J Ophthalmol* 2013;61:382-383
157. Davidson AE, Hayes S, Hardcastle AJ, Tuft SJ. The pathogenesis of keratoconus. *Eye* 2014;28:189-195
158. Abu-Amero KK, Al-Muammar AM, Kondkar AA. Genetics of keratoconus: where do we stand? *J Ophthalmol* 2014;2014:641-708
159. Vazirani J, Basu S. Keratoconus: current perspectives. *Clin Ophthalmol* 2013;7:2019-2030
160. Cullen JF, Butler HG. Mongolism (Down's syndrome) and keratoconus. *Br J Ophthalmol* 1963;47:321-330
161. Robertson I. Keratoconus and the Ehlers-Danlos syndrome: a new aspect of keratoconus. *Med J Aust* 1975;1:571-573
162. Woodward EG, Morris MT. Joint hypermobility in keratoconus. *Ophthalmic Physiol Opt* 1990;10:360-362
163. McMonnies CW. Abnormal rubbing and keratectasia. *Eye Contact Lens* 2007;33:265-271
164. Balasubramanian SA, Pye DC, Willcox MD. Effects of eye rubbing on the levels of protease, protease activity and cytokines in tears: relevance in keratoconus. *Clin Exp Optom* 2013;96:214-218
165. Jun AS, Cope L, Speck C, Feng X, Lee S, Meng H, Hamad A, Chakravarti S. Subnormal cytokine profile in the tear fluid of keratoconus patients. *PLoS One* 2011;6:e16437
166. Balasubramanian SA, Mohan S, Pye DC, Willcox MD. Proteases, proteolysis and inflammatory molecules in the tears of people with keratoconus. *Acta Ophthalmol* 2012;90:303-309
167. McMonnies CW. Inflammation and keratoconus. *Optom Vis Sci* 2015;92:35-41
168. Sherwin T, Brookes NH. Morphological changes in keratoconus: pathology or pathogenesis. *Clin Experiment Ophthalmol* 2004;32:211-217
169. Meek KM, Tuft SJ, Huang Y, Gill PS, Hayes S, Newton RH, et al. Changes in collagen orientation and distribution in keratoconus corneas. *Invest Ophthalmol Vis Sci* 2005;46:1948-1956
170. Zadnik K, Barr JT, Gordon MO, Edrington TB. Biomicroscopic signs and disease severity in keratoconus. Collaborative Longitudinal Evaluation of Keratoconus (CLEK) Study Group. *Cornea* 1996;15:139-146
171. Krumeich JH, Daniel J, Knulle A. Live-epikeratophakia for keratoconus. *J Cataract Refract Surg* 1998;24:456-463
172. Maeda N, Klyce SD, Smolek MK, Thompson HW. Automated keratoconus screening with corneal topography analysis. *Invest Ophthalmol Vis Sci* 1994;35:2749-2757
173. Alió JL, Shabayek MH. Corneal higher order aberrations: a method to grade keratoconus. *J Refract Surg* 2006;22:539-545
174. McMahon TT, Szczotka-Flynn L, Barr JT, Anderson RJ, Slaughter ME, Lass JH, et al. A new method for grading the severity of keratoconus: the Keratoconus Severity Score (KSS). *Cornea* 2006;25:794-800
175. Mahmoud AM, Roberts CJ, Lembach RG, Twa MD, Herderick EE, McMahon TT. CLMI: the cone location and magnitude index. *Cornea* 2008;27:480-487
176. Li X, Yang H, Rabinowitz YS. Keratoconus: classification scheme based on videokeratography and clinical signs. *J Cataract Refract Surg* 2009;35:1597-1603
177. Sandali O, El Sanharawi M, Temstet C, Hamiche T, Galan A, Ghouali W, et al. Fourier-domain optical coherence tomography imaging in keratoconus: a corneal structural classification. *Ophthalmology* 2013;120:2403-2412
178. Kanellopoulos AJ, Asimellis G. Revisiting keratoconus diagnosis and progression classification based on evaluation of corneal asymmetry indices, derived from Scheimpflug imaging in keratoconic and suspect cases. *Clin Ophthalmol* 2013;7:1539-1548

179. Gomes JA, Tan D, Rapuano CJ, Belin MW, Ambrósio R Jr, Guell JL, Maleceze F, Nishida K, Sangwan VS; Group of Panelists for the Global Delphi Panel of Keratoconus and Ectatic Diseases. Global consensus on keratoconus and ectatic diseases. *Cornea* 2015;34:359-369
180. Visser ES, Wisse RP, Soeters N, Imhof SM, Van der Lelij A. Objective and subjective evaluation of the performance of medical contact lenses fitted using a contact lens selection algorithm. *Cont Lens Anterior Eye* 2016;39:298-306
181. Parker JS, van Dijk K, Melles GR. Treatment options for advanced keratoconus: A review. *Surv Ophthalmol* 2015;60:459-480
182. Paufigue L. Les greffes corneennes lamellaires therapeutiques. *Bull Mem Soc Fr Ophtalmol* 1949;62:210-213
183. Malbran ES. Corneal dystrophies: a clinical, pathological, and surgical approach. 28 Edward Jackson Memorial Lecture. *Am J Ophthalmol* 1972;74:771-809
184. Anwar M. Technique in lamellar keratoplasty. *Trans Ophthalmol Soc U K* 1974;94:163-171
185. Archila EA. Deep lamellar keratoplasty dissection of host tissue with intrastromal airinjection. *Cornea* 1984-1985;3:217-218
186. Anwar M, Teichmann KD. Big-bubble technique to bare Descemet's membrane in anterior lamellar keratoplasty. *J Cataract Refract Surg* 2002;28:398-403
187. Marchini G, Mastropasqua L, Pedrotti E, et al. Deep lamellar keratoplasty by intracorneal dissection. *Ophthalmology* 2006;113:1289-1300
188. Sugita J, Kondo J. Deep lamellar keratoplasty with complete removal of pathological stroma for vision improvement. *Br J Ophthalmol* 1997;81:184-188
189. Manche, E. Holland GN, Maloney RK. Deep lamellar keratoplasty using viscoelastic dissection. *Arch Ophthalmol* 1999;117:1561-1565
190. Melles GR, Lander F, Rietveld FJ, Remeijer L, Beekhuis WH, Binder PS. A new surgical technique for deep stromal, anterior lamellar keratoplasty. *Br J Ophthalmol* 1999;83:327-333
191. Melles GR, Rietveld FJ, Beekhuis WH, Binder PS. A technique to visualize corneal incision and lamellar dissection depth during surgery. *Cornea* 1999;18:80-86
192. Caporossi A, Simi C, Licignano R, et al. Air-guided manual deep lamellar keratoplasty. *Eur J Ophthalmol* 2004;14:55-58.
193. Watson SL, Ramsay A, Dart JKG, Bunce C, Craig E. Comparison of deep lamellar keratoplasty and penetrating keratoplasty in patients with keratoconus. *Ophthalmology* 2004;111:1676-1682
194. Feizi S, Javadi MA, Jamali H, Mirbabae F. Deep anterior lamellar keratoplasty in patients with keratoconus: big-bubble technique. *Cornea* 2010;29:177-182
195. Smadja D, Colin J, Krueger RR, Mello GR, Gallois A, Mortemousque B, Touboul D. Outcomes of deep anterior lamellar keratoplasty for keratoconus: learning curve and advantages of the big bubble technique. *Cornea* 2012;31:859-863
196. Borderie VM, Sandali O, Bullet J, Gaujoux T, Touzeau O, Laroche L. Long-term results of deep anterior lamellar versus penetrating keratoplasty. *Ophthalmology* 2012;119:249-255
197. Fontana L, Parente G, Tassinari G. Clinical outcomes after deep anterior lamellar keratoplasty using the big-bubble technique in patients with keratoconus. *American journal of ophthalmology* 2007;143:117-124
198. Fontana L, Parente G, Sincich A, et al. Influence of graft-host interface on the quality of vision after deep anterior lamellar keratoplasty in patients with keratoconus. *Cornea* 2011;30:497e502
199. Funnell CL, Ball J, Noble BA. Comparative cohort study of the outcomes of deep lamellar keratoplasty and penetrating keratoplasty for keratoconus. *Eye (Lond)* 2006;20:527-532

200. Jones MN, Armitage WJ, Ayliffe W, et al. Penetrating and deep anterior lamellar keratoplasty for keratoconus: a comparison of graft outcomes in the United kingdom. *Invest Ophthalmol Vis Sci* 2009;50:5625-5629
201. Yildiz EH, Erdurmus M, Hammersmith KM, et al. Comparative study of graft rejection in keratoconus patients with and without self-reported atopy. *Cornea* 2009;28:846-850
202. Sugar J, Macsai MS. What causes keratoconus? *Cornea* 2012;31:716-719
203. Cheung IMY, McGhee CNJ, Sherwin T. A new perspective on the pathobiology of keratoconus: interplay of stromal wound healing and reactive species-associate processes. *Clin Exp Optom* 2013;96:188-196
204. Lema I, Sobrino T, Durán JA, et al. Subclinical keratoconus and inflammatory molecules from tears. *Br J Ophthalmol* 2009;6:820-824
205. Claesson M, Armitage WJ. Armitage. Clinical outcome of repeat penetrating keratoplasty. *Cornea* 2013;32:1026-1030
206. Kelly T, Coster D, Williams K. Repeat penetrating corneal transplantation in patients with keratoconus. *Ophthalmology* 2011;118:1538-1542
207. Bourges JL, Savoldelli M, Dighiero P, et al. Recurrence of keratoconus characteristics: a clinical and histologic followup analysis of donor grafts. *Ophthalmology* 2003;110:1920-1925
208. Feizi S, Javadi MA, Rezaei Kanavi M. Recurrent keratoconus in a corneal graft after deep anterior lamellar keratoplasty. *J Ophthalmic Vis Res* 2012;7:328-331
209. Patel N, Mearza A, Rostron CK, et al. Corneal ectasia following deep lamellar keratoplasty. *Br J Ophthalmol* 2003;87:799-800
210. Wollensak G, Spoerl E, Seiler T. Riboflavin/ultraviolet-ainduced collagen crosslinking for the treatment of keratoconus. *Am J Ophthalmol* 2003;135:620-627
211. Chan E, Snibson GR. Current status of corneal collagen crosslinking for keratoconus: a review. *Clin Exp Optom* 2013;96:155-164
212. Raiskup F, Spoerl E. Corneal cross-linking with hypo-osmolar riboflavin solution in thin keratoconic corneas. *Am J Ophthalmol* 2011;152:28-32
213. Hafezi F. Limitation of collagen cross-linking with hypoosmolar riboflavin solution: failure in an extremely thin cornea. *Cornea* 2011;30:917-919
214. Hafezi F, Mrochen M, Iseli HP, Seiler T. Collagen crosslinking with ultraviolet-A and hypoosmolar riboflavin solution in thin corneas. *J Cataract Refract Surg* 2009;35:621-624
215. Meek KM, Hayes S. Corneal cross-linking: a review. *Ophthalmic Physiol Opt* 2013;33:78-93
216. Ivarsen A, Hjortdal J. Collagen cross-linking for advanced progressive keratoconus. *Cornea* 2013;32:903-906
217. Koller T, Mrochen M, Seiler T. Complication and failure rates after corneal crosslinking. *J Cataract Refract Surg* 2009;35:1358-1362
218. Pinero DP, Alio JL. Intracorneal ring segments in ectatic corneal disease: A review. *Clin Experiment Ophthalmol* 2010;38:154-167
219. Khan MI, Injarie A, Muhtaseb M. Intrastromal corneal ring segments for advanced keratoconus and cases with high keratometric asymmetry. *J Cataract Refract Surg* 2012;38:129-136
220. Avni-Zauberman N, Rootman DS. Cross-linking and intracorneal ring segments: review of the literature. *Eye Contact Lens* 2014;40:365-370
221. Burris TE. Intrastromal corneal ring technology: Results and indications. *Curr Opin Ophthalmol* 1998;9:9-14
222. Rabinowitz YS. Intacs for keratoconus. *Curr Opin Ophthalmol* 2007;18:279-283

223. Rodriguez LA, Guille ´n PB, Benavides MA, et al. Penetrating keratoplasty versus intrastromal corneal ring segments to correct bilateral corneal ectasia: preliminary study. *J Cataract Refract Surg* 2007;33:488-496
224. Kymionis GD, Grentzelos MA, Diakonis VF, et al. Nine-year follow-up of intacs implantation for keratoconus. *Open Ophthalmol J* 2009;3:77-81
225. Colin J, Cochener B, Savary G, et al. INTACS inserts for treating keratoconus: one-year results. *Ophthalmology* 2001;108:1409-1414
226. Health Quality Ontario. Intrastromal corneal ring implants for corneal thinning disorders: an evidence-based analysis. *Ont Health Technol Assess Ser* 2009;9:1-90
227. Zorginstituut Nederland. Standpunt Collageen crosslinking (CXL) bij keratoconus en keratectasie 2014. Consulted in October 2016 via: <https://www.zorginstituutnederland.nl/binaries/content/documents/zinl-www/documenten/publicaties/rapporten-en-standpunten/2014/1406-standpunt-collageen-crosslinking-cxl-bij-keratoconus-en-keratectasie/Standpunt+Collageen+crosslinking+-CXL+bij+keratoconus+en+keratectasie.pdf>

