



Universiteit
Leiden
The Netherlands

Infrared spectroscopy of astrophysically relevant hydrocarbons

Doney, K.D.

Citation

Doney, K. D. (2018, June 20). *Infrared spectroscopy of astrophysically relevant hydrocarbons*. Retrieved from <https://hdl.handle.net/1887/62922>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/62922>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/62922> holds various files of this Leiden University dissertation.

Author: Doney, K.D.





Title: Infrared spectroscopy of astrophysically relevant hydrocarbons

Issue Date: 2018-06-20

The full cubic and semidiagonal quartic force field of acetylene (C_2H_2), diacetylene (C_4H_2), triacetylene (C_6H_2), and tetraacetylene (C_8H_2) are determined using CCSD(T) (coupled cluster theory with single and double excitations and augmented by a perturbative treatment of triple excitations) in combination with the atomic natural orbital (ANO) basis sets. Application of second-order vibrational perturbation theory (VPT₂) results in vibrational frequencies that agree well with the known fundamental and combination band experimental frequencies of acetylene, diacetylene, and triacetylene (average discrepancies are less than 10 cm^{-1}). Furthermore, the predicted ground state rotational constants (B_0) and vibration-rotation interaction constants (α_i) are shown to be consistent with known experimental values. New vibrational frequencies and rotational parameters from the presented theoretical predictions are given for triacetylene and tetraacetylene, which can be used to aid laboratory and astronomical spectroscopic searches for characteristic transitions of these molecules.

2.1 INTRODUCTION

Due to their reactive nature, unsaturated linear hydrocarbons, such as acetylene and polyynes (general structure HC_{2n}H ; $\tilde{X}^1\Sigma_g^+$), are prevalent in combustion chemistry (Frenklach et al., 1985; Cherchneff et al., 1992; Homann, 1998; Richter & Howard, 2000), plasma processes (Fujii & Kareev, 2001; Thejaswini et al., 2011), chemical synthesis (Shirakawa, 2001; Zhao et al., 2003; Milani et al., 2006; Ma et al., 2008), chemistry of planetary atmospheres (Kunde et al., 1981; de Graauw et al., 1997; Wilson & Atreya, 2003; Kunde et al., 2004; Vuitton et al., 2006; Burgdorf et al., 2006; Meadows et al., 2008; Waite et al., 2007; Gu et al., 2009), and interstellar gas-phase chemistry (Cernicharo et al., 2001a,b; Bernard-Salas et al., 2006; Wakelam et al., 2010; Fonfría et al., 2011; Malek et al., 2012). They are of particular interest for astronomers, because they are believed to act as the ultraviolet (UV) shield in hydrocarbon-rich atmospheres (Bandy et al., 1992, 1993; Frost et al., 1995; Bizzocchi et al., 2011), and in the formation and destruction of polycyclic aromatic hydrocarbons (PAHs) (Frenklach & Feigelson, 1989; Ekern et al., 1998; Krestinin, 2000; Waite et al., 2007; Kress et al., 2010), a major reservoir of carbon in the universe. In astronomical environments, the formation of long chain polyynes from acetylene is believed to occur through polymerization reactions (Woods et al., 2003; Gu et al., 2009; Sakai & Yamamoto, 2013),



Although long carbon chain molecules (e.g., HC_n and HC_nN for $n \leq 9$) (Cernicharo & Guélin, 1996; Bell et al., 1997; Bell et al., 1999; Vuitton et al., 2007) and small polyynes (HC_{2n}H for $n \leq 3$) have been detected in carbon-rich astronomical sources (Cernicharo et al., 2001a,b; Vuitton et al., 2007), tetraacetylene has yet to be observed. One limiting factor is that as centrosymmetric molecules, polyynes lack a permanent dipole moment, and cannot be detected by radioastronomy using pure rotational transitions, unlike, e.g., HC_nN . Therefore, ro-vibrational spectra in the infrared (IR) region are the most important spectroscopic tools to detect polyynes both in the laboratory and in space. In particular, detection of acetylene, diacetylene, and triacetylene in planetary atmospheres and protoplanetary nebulae has been realized primarily through observation of the strongest perpendicular band (ν_5 , ν_8 , and ν_{11} , respectively, at $\sim 13 - 17 \mu\text{m}$) and the second strongest parallel band ($\nu_4 + \nu_5$, $\nu_6 + \nu_8$, and $\nu_8 + \nu_{11}$, respectively, at $\sim 8 \mu\text{m}$) (Kunde et al., 1981; de Graauw et al., 1997; Cernicharo et al., 2001a,b). However, accurate line positions for tetraacetylene are lacking, from either laboratory or theoretical studies.

Extensive theoretical and experimental studies have been carried out for acetylene and diacetylene in the past few decades, including high-resolution spectroscopic studies of all the fundamental bands and a significant number of the combination bands (Plyler et al., 1963; Palmer et al., 1972; Pliva, 1972a,b; Guelachvili et al., 1984; Owen et al., 1987; McNaughton & Bruget, 1992; Arié & Johns, 1992; Vanderauwera et al., 1993; Gambogi et al., 1995; Bizzocchi et al., 2011; Zhao et al., 2014a), and high level *ab initio* calculations that take into account anharmonic effects (Temsamani & Herman, 1995; Martin et al., 1998; Thorwirth et al., 2008; Simmonett et al., 2009). The combination of these studies shows that current quantum chemical theory, particularly coupled cluster theory with single and double excitations and augmented by a perturbative treatment of triple excitations (CCSD(T)) (Raghavachari et al., 1989), is able to accurately reproduce equilibrium geometries, experimental vibrational frequencies, vibration-rotation interaction constants (α_i), and ground state rotational constants (B_0).

Triacetylene and tetraacetylene are not as thoroughly studied, notably in terms of rotational information. While all of the fundamental vibrational modes of triacetylene have been measured,

there is only rotational information for the IR active fundamental modes (Bjarnov et al., 1974), and the strongest IR combination band ($\nu_8 + \nu_{11}$) (Matsumura et al., 1988b; McNaughton & Bruget, 1991; Haas et al., 1994b,a; Doney et al., 2015). However, theoretical studies of triacetylene do give rotational information for the remaining modes from CCSD(T) calculations of the vibration-rotation interaction constants (Chang et al., 2016) and the equilibrium geometry (Sattelmeyer & Stanton, 2000). In addition, the harmonic frequencies of triacetylene were calculated using partial fourth-order many-body perturbation theory [SDQ-MBPT(4)] (Sattelmeyer & Stanton, 2000). Conversely though, to the authors' knowledge, there is almost no rotational information for tetraacetylene. There has been only one low-resolution spectroscopic study of tetraacetylene, which measured three of the fundamentals (ν_6 , ν_8 , ν_{14} at 3329.4, 2023.3, and 621.5 cm^{-1} , respectively), one combination band ($\nu_{10} + \nu_{14}$ at 1229.7 cm^{-1}), and gives an estimate for the electronic ground state rotational constant, B_0 (Shindo et al., 2001). Unfortunately, the theoretical knowledge of tetraacetylene is equally limited, with only two studies of the equilibrium geometry (at the Hartree-Fock (Toto et al., 1996) and B3LYP (Kaiser et al., 2010) level of theory), and a calculation of the harmonic vibrational frequencies at the SVWN level of theory (Shindo et al., 2001). While the two modes that are most useful for astronomical identification (ν_{14} and $\nu_{10} + \nu_{14}$) were measured, the uncertainty associated with the line positions is too large to allow for an unambiguous assignment. Moreover, some high-resolution IR searches have been attempted (Zhao et al., 2013, 2014a; Doney et al., 2015; Chang et al., 2016), but so far no transitions have been assigned to tetraacetylene.

In this paper, we report the *ab initio* calculations for acetylene, diacetylene, triacetylene, and tetraacetylene. Due to the centrosymmetric nature of these molecules, observations in the laboratory and in space are most easily accomplished through their infrared spectra. As such, the properties computed and presented here are those related to that technique: fundamental vibrational frequencies, ground state rotational constants, and intramolecular interactions. The computational approach is calibrated using the well studied acetylene and diacetylene, and then extended to make predictions for triacetylene and tetraacetylene.

2.2 COMPUTATIONAL METHODS

All calculations were carried out at the CCSD(T) level of theory, which with a sufficiently large basis set has been shown to accurately reproduce experimental values of semi-rigid molecules (Raghavachari et al., 1989; Lee & Scuseria, 1995; Bartlett, 1995; Gauss & Stanton, 1997; Gauss, 1998; Martin et al., 1998; Thorwirth et al., 2008; Simmonett et al., 2009; McCaslin & Stanton, 2013). Equilibrium geometries were determined using the large core-valence correlation-consistent quadruple- ζ basis set (cc-pCVQZ), which features [8s7p5d3fig] (non-hydrogen atoms) and [4s3p2dif] (hydrogen) of (15s9p5d3fig) and (6s3p2dif) primitive basis sets, respectively (Woon & Dunning Jr., 1995; Feller, 1996; Schuchardt et al., 2007). All electron (AE)-CCSD(T)/cc-pCVQZ has been shown to give very accurate equilibrium geometries for unsaturated hydrocarbons (Auer & Gauss, 2001; Bak et al., 2001; Zhang et al., 2007; Simmonett et al., 2009). Optimizations were done using analytic energy derivatives (Stanton & Gauss, 2000), and were considered converged when the root-mean-square (rms) gradient fell below 10^{-10} au.

However, it is well known that correlation-consistent basis sets, such as cc-pCVQZ, tend to underestimate the vibrational frequencies of symmetric bending modes (π_g) of conjugated molecules, e.g., polynes, due to their susceptibility to an intramolecular variant of basis set superposition error (BSSE) (Simandiras et al., 1988; Simmonett et al., 2009). It has been shown that one way to avoid this problem is to use basis sets with a large number of Gaussian primitives (particularly f-type), such as the atomic natural orbital (ANO) basis set (with the primitive basis set (13s8p6d4f2g) for non-hydrogen atoms and (8s6p4d2f) for hydrogen) (Almlöf & Taylor, 1987; Bauschlicher Jr. & Taylor, 1993; Martin et al., 1998). The basis set has two common truncations: [4s3p2dif] for non-hydrogen atoms and [4s2pid] for hydrogen (hereafter known as ANO1), and [5s4p3d2fig] (non-hydrogen atoms) and [4s3p2dif] (hydrogen) (hereafter known as ANO2) (Almlöf & Taylor, 1987; Feller, 1996;




Figure 2.1: AE-CCSD(T)/cc-pCVQZ equilibrium geometries (Å) for HC_{2n}H . Experimentally determined equilibrium bond lengths for acetylene (Tamassia et al., 2016), diacetylene (Thorwirth et al., 2008), and triacetylene (Tay et al., 1995) are given in italics below.

Schuchardt et al., 2007). In addition, only the valence electrons of carbon are considered in the correlation treatment, *i.e.*, standard frozen-core (fc) calculations. (fc)-CCSD(T)/ANO1 has been shown to accurately reproduce experimental frequencies and intensities for small molecules (Martin et al., 1997, 1998; Vázquez & Stanton, 2007). Using the (fc)-CCSD(T)/ANO1 optimized geometry, second-order vibrational perturbation (VPT2) theory calculations were determined from full cubic and the semidiagonal part of the quartic force fields obtained by numerical differentiation of analytic CCSD(T) second derivatives (Gauss & Stanton, 1997; Matthews et al., 2007). All calculations were performed with the development version of the CFOUR program (CFOUR, 2017).

2.3 RESULTS AND DISCUSSION

2.3.1 Equilibrium structure

The AE-CCSD(T)/cc-pCVQZ equilibrium geometries are shown in Figure 2.1, with comparison to experimentally derived values (in italics) when known (Tay et al., 1995; Tamassia et al., 2016; Thorwirth et al., 2008). The theoretical equilibrium bond lengths for acetylene, diacetylene, and triacetylene all agree within 0.5% of the structures determined from experimentally measured rotational constants. As the length of the carbon chain increases, the C-H bond lengths stay essentially the same, ~ 1.062 Å, consistent with a sp-H type C-H bond. However, the $\text{C}\equiv\text{C}$ bond lengths increase (particularly the internal $\text{C}\equiv\text{C}$ bonds), while the C-C bond lengths decrease, becoming closer to that typical of CC double bonds. This suggests that the π electrons become more delocalized over the internuclear axis, and the polyyne's configuration moves from a strict triple-single bond alternation to more of a consecutive double bond character of the CC bonds, making the overall structure more rigid as C_2 units are added, an effect that also qualitatively acts to increase the biradical character of the molecule as the size grows.

The equilibrium rotational constants, B_e , obtained from the AE-CCSD(T)/cc-pCVQZ equilibrium geometries are summarized in Table 2.1, and agree well with experimental ground state rotational constants (B_0). As such, the equilibrium rotational constants suggest that the calculations predict the correct ground state geometry, because for linear molecules with more than three atoms the summation of vibration-rotation interaction constants (α_i) is expected to be close to zero, and from

$$B_0 = B_e - \frac{1}{2} \sum_i \alpha_i \quad (2.1)$$

it becomes that $B_0 \sim B_e$.

In addition, as seen for other carbon chains (*e.g.*, HC_n , HC_{2n+1}N , and H_2C_n) the centrifugal

Table 2.1: CCSD(T)/ANO1 rotational constants (in cm^{-1}) of HC_{2n}H ($n = 1 - 4$)

	HC_2H	HC_4H	HC_6H	HC_8H
Calc.				
B_e	1.181 053	0.146 248	0.044 064	0.018 823
B_0	1.175319	0.146 167	0.044 092	0.018 844
$D_e \times 10^8$	160	1.5	0.086	0.012
Expt.				
B_0	1.176 646 32(18) (Kabbadj et al., 1991)	0.146 4123(17) (Zhao et al., 2014a)	0.044 1735(12) (Doney et al., 2015)	0.020(3) (Shindo et al., 2001)
$D_0 \times 10^8$	159.8(9) (Kabbadj et al., 1991)	1.568 25(20) (Bizzocchi et al., 2011)	0.107(7) (Doney et al., 2015)	

distortion constant (D_e) decreases with increasing molecular size, with a theoretical $D_e = 1.6 \times 10^{-6} \text{ cm}^{-1}$ for acetylene, $D_e = 1.5 \times 10^{-8} \text{ cm}^{-1}$ for diacetylene, $D_e = 8.6 \times 10^{-10} \text{ cm}^{-1}$ for triacetylene, and $D_e = 1.2 \times 10^{-10} \text{ cm}^{-1}$ for tetraacetylene (Thaddeus et al., 1998). These values are consistent with those found experimentally for the respective vibrational ground states (Table 2.1). As noted by Thaddeus et al. (1998), this behavior of increasing stiffness with chain length is a distinguishing characteristic associated with *bona fide* chains.

2.3.2 Spectroscopic properties of acetylene and diacetylene

The quality of the present calculations is checked by comparison to the experimentally well studied acetylene and diacetylene. The harmonic and VPT2 fundamental frequencies of the fundamental and combination bands are given in Tables 2.2 and 2.3, for acetylene and diacetylene respectively,

Table 2.2: Harmonic and anharmonic (VPT2) frequencies (in cm^{-1}) of HC_2H fundamental and selected combination bands

Mode	CCSD(T)/ANO1 ^a		Experimental
	ω	ν	ν
$\nu_1(\sigma_g^+)$	3514.2(0)	3375.2(0)	3372.851 (Palmer et al., 1972)
$\nu_2(\sigma_g^+)$	2001.5(0)	1964.8(0)	1974.317 (Palmer et al., 1972)
$\nu_3(\sigma_u^+)$	3414.6(84.7)	3285.9(74.8)	3288.58075 (Vanderauwera et al., 1993)
$\nu_4(\pi_g)$	600.5(0)	600.6(0)	612.871 (Pliva, 1972a)
$\nu_5(\pi_u)$	752.3(90.5)	734.7(91.7)	730.332 (Pliva, 1972a)
$\nu_4 + \nu_5(\sigma_u^+)$	1352.8	1329.2(10.8)	1328.074 (Pliva, 1972a)
$\nu_2 + \nu_5(\pi_u)$	2753.8	2698.3(0.1)	2701.907 (Pliva, 1972b)
$\nu_3 + \nu_4(\pi_u)$	4015.1	3878.5(0.5)	3882.4060 (Palmer et al., 1972)
$\nu_1 + \nu_5(\pi_u)$	4266.5	4098.9(0.5)	4091.17326 (D'Cunha et al., 1991)
$\nu_1 + \nu_3(\sigma_u^+)$	6928.7	6551.9(2.0)	6556.46 (Plyler et al., 1963)
Anharmonic ZPE = 5760.1			

a. Intensities in km/mol are given in parenthesis.

and experimental values are included for comparison. The (fc)-CCSD(T)/ANO1 VPT2 fundamental frequencies show good agreement with experimental values, with most observed-calculated deviations (o-c) being less than 5 cm^{-1} and all being less than 15 cm^{-1} .

Table 2.3: Harmonic and anharmonic (VPT2) frequencies (in cm^{-1}) of HC_4H fundamental and selected combination bands

Mode	CCSD(T)/ANO1 ^a		Experimental
	ω	ν	ν
$\nu_1(\sigma_g^+)$	3465.8(0)	3332.5(0)	3332.15476 (McNaughton & Bruget, 1992)
$\nu_2(\sigma_g^+)$	2240.2(0)	2193.1(0)	2188.9285 (Guelachvili et al., 1984)
$\nu_3(\sigma_g^+)$	891.1(0)	859.2(0)	871.9582 (Guelachvili et al., 1984)
$\nu_4(\sigma_u^+)$	3465.9(152.7)	3333.1(135.5)	3333.6634 (Zhao et al., 2014a)
$\nu_5(\sigma_u^+)$	2054.1(0.2)	2016.9(0.5)	2022.2415 (Guelachvili et al., 1984)
$\nu_6(\pi_g)$	636.3(0)	624.2(0)	625.643507 (Bizzocchi et al., 2011)
$\nu_7(\pi_g)$	479.8(0)	476.9(0)	482.7078 (Guelachvili et al., 1984)
$\nu_8(\pi_u)$	636.3(78.7)	624.1(78.8)	628.040776 (Bizzocchi et al., 2011)
$\nu_9(\pi_u)$	220.7(7.3)	219.6(7.3)	219.97713 (Arié & Johns, 1992)
$2\nu_9(\sigma_g^+)$	441.4	438.5(0)	438.47757 (Arié & Johns, 1992)
$\nu_7 + \nu_9(\sigma_u^+)$	700.5	696.3(0.8)	701.8939 (Bizzocchi et al., 2011)
$\nu_6 + \nu_9(\sigma_u^+)$	857.0	843.9(0.01)	845.655513 (Bizzocchi et al., 2011)
$\nu_8 + \nu_9(\sigma_g^+)$	857.0	843.9(0)	848.365918 (Bizzocchi et al., 2011)
$\nu_7 + \nu_8(\pi_u)$	1116.1	1103.1(0.6)	1111 (Owen et al., 1987)
$\nu_6 + \nu_8(\sigma_u^+)$	1272.6	1244.7(21.8)	1241.060828 (McNaughton & Bruget, 1992)
$2\nu_6 + \nu_8(\pi_u)$	1909.0	1864.6(0.0)	1863.2512 (Guelachvili et al., 1984)
$\nu_2 + \nu_9(\pi_u)$	2460.9	2410.0(0.04)	2406.4251 (Guelachvili et al., 1984)
$\nu_5 + \nu_7(\pi_u)$	2533.9	2489.0(0.01)	2500.6458 (Guelachvili et al., 1984)
$\nu_5 + \nu_6(\pi_u)$	2690.4	2637.0(0.04)	2643.32323 (McNaughton & Bruget, 1992)
$\nu_2 + \nu_8(\pi_u)$	2876.6	2810.9(0.4)	2805 (Owen et al., 1987)
$\nu_1 + \nu_9(\pi_u)$	3686.5	3551.6(0.1)	3551.56158159 (McNaughton & Bruget, 1992)
$\nu_1 + \nu_8(\pi_u)$	4102.1	3946.9(0.7)	3939 (Owen et al., 1987)
$\nu_4 + \nu_6(\pi_u)$	4102.3	3947.8(0.7)	
$\nu_2 + \nu_5(\sigma_u^+)$	4294.3	4194.0(0.1)	
$\nu_4 + \nu_3(\sigma_u^+)$	4357.0	4192.3(0.1)	
$\nu_1 + \nu_4(\sigma_u^+)$	6931.7	6557.2(3.4)	6565.472 (Gambogi et al., 1995)
Anharmonic ZPE = 7966.9			

a. Intensities in km/mol are given in parenthesis.

Based on previous studies of acetylene (Martin et al., 1998) and diacetylene (Thorwirth et al., 2008), the use of the ANO2 basis set was evaluated compared to the ANO1 basis set. For some of the vibrational modes, such as the ν_4 mode of acetylene [612.88 cm^{-1} (observed)] (Pliva, 1972a), Martin et al. (1998) showed CCSD(T)/ANO2 can give a slightly better agreement (o-c value of $\sim 2 \text{ cm}^{-1}$) compared to the ANO1 basis set (o-c value of $\sim 12 \text{ cm}^{-1}$). However, the study by Thorwirth et al. (2008) showed that, for diacetylene, the average o-c value with CCSD(T)/ANO2 is comparable to that for the ANO1 basis set ($\sim 6 \text{ cm}^{-1}$ and $\sim 4 \text{ cm}^{-1}$, respectively). Moreover, the time cost of (fc)-CCSD(T)/ANO2 calculations compared to (fc)-CCSD(T)/ANO1 far outweighs the minor frequency differences, and does not justify the higher computational cost of the ANO2 basis set in predicting the fundamental frequencies of longer polyynes.

The (fc)-CCSD(T)/ANO1 anharmonicity constants (x_{ij} , **supplementary material**) also accurately account for the known combination bands of acetylene and diacetylene (Tables 2.2 and 2.3, respectively). All the combination bands are within 5 cm^{-1} of their observed values. For both acetylene and diacetylene, the ANO1 basis set is able to most accurately reproduce the C-H asymmetric stretch mode (ν_3 and ν_4 , respectively). Significant is the agreement between the experimental and

Table 2.4: CCSD(T)/ANO1 vibration-rotation interaction constants^a (α_i , in cm^{-1}) of HC_{2n}H ($n = 1 - 4$)

Mode	$\text{HC}_2\text{H} \times 10^3$	$\text{HC}_4\text{H} \times 10^4$	$\text{HC}_6\text{H} \times 10^5$	$\text{HC}_8\text{H} \times 10^5$
α_1	6.85(6.904 ^b) (Temsamani & Herman, 1995)	2.16(2.153) (Zhao et al., 2014a)	2.97	0.730
α_2	6.01(6.181) (Temsamani & Herman, 1995)	6.61	15.2	4.91
α_3	5.80(5.882 ^b) (Temsamani & Herman, 1995)	3.12(3.110 ^b) (Simmonett et al., 2009)	7.44	2.55
α_4	-1.46(-1.354) (Temsamani & Herman, 1995)	2.14(2.183) (Zhao et al., 2014a)	3.82	3.76
α_5	-2.13(-2.232) (Temsamani & Herman, 1995)	3.94(3.948) (Guelachvili et al., 1984)	2.99(3.58) (Doney et al., 2015)	0.930
α_6	0.73	-0.700(-0.678) (Bizzocchi et al., 2011)	9.91(9.15) (McNaughton & Bruget, 1991)	0.730
α_7	4.06	-2.70(-2.711) (McNaughton & Bruget, 1992)	9.91	4.06
α_8	2.33	-0.647(-0.636) (Bizzocchi et al., 2011)	-1.17(-1.071) (McNaughton & Bruget, 1991)	2.33
α_9	2.05	-4.13(-4.183) (McNaughton & Bruget, 1992)	-5.83	2.05
α_{10}			-7.42(-7.88) (McNaughton & Bruget, 1991)	-0.295
α_{11}			-1.06(-1.06) (Matsumura et al., 1988b)	-1.95
α_{12}			-5.07	-1.69
α_{13}			-8.47(-8.7207) (Haas et al., 1994a)	-2.26
α_{14}				-0.295
α_{15}				-0.163
α_{16}				-2.29
α_{17}				-2.80

a. Experimental values are in parentheses.

b. Deperturbed.

our predicted frequencies of $\nu_6 + \nu_8$ [1241.060 828(38) cm^{-1} (observed) (McNaughton & Bruget, 1992) and 1244.7 cm^{-1} (theoretical)], and $2\nu_6 + \nu_8$ [1863.251 2(5) cm^{-1} (observed) (Guelachvili et al., 1984) and 1864.6 cm^{-1} (theoretical)] of diacetylene; both of which had only previously been calculated with CCSD(T)/cc-pCVQZ, and had o-c values greater than 20 cm^{-1} (Simmonett et al., 2009). This suggests that the combination band VPT2 frequencies of polyynes determined using (fc)-CCSD(T)/ANO1 are accurate to aid identification of molecules, such as in astronomical surveys.

The vibration-rotation interaction constants (Table 2.4), are also determined in the course of the VPT2 calculation, and are in good agreement with both previous theoretical studies (Martin et al., 1998; Simmonett et al., 2009) and experimentally determined values (Guelachvili et al., 1984; McNaughton & Bruget, 1992; Temsamani & Herman, 1995; Simmonett et al., 2009; Bizzocchi et al., 2011; Zhao et al., 2014a). Based on the vibration-rotation interaction constants, the ground state rotational constants (B_0) were determined using the AE-CCSD(T)/cc-pCVQZ determined B_e values (Table 2.1). For acetylene, $B_0 = 1.175\,319\,\text{cm}^{-1}$, which is a 0.1% difference compared to the experi-

mentally determined value of $B_0 = 1.176\,646\,32(18)\,\text{cm}^{-1}$ (Kabbadj et al., 1991). Diacetylene shows a similar 0.2% difference between the theoretical value of $B_0 = 0.146\,167\,\text{cm}^{-1}$, and the experimentally determined value of $B_0 = 0.146\,4123(17)\,\text{cm}^{-1}$ (Zhao et al., 2014a). The consistent accuracy of these values suggests that the method presented is clearly good enough to extrapolated to and aid high-resolution infrared spectroscopic searches for the larger polyynes.

2.3.3 *Spectroscopic properties of triacetylene*

The (fc)-CCSD(T)/ANO1 harmonic and VPT2 fundamental frequencies along with the experimental frequencies are given in Table 2.5. Comparison between theoretical VPT2 frequencies and experimental fundamentals measured with high-resolution techniques show average o-c values that are

Table 2.5: Harmonic and anharmonic (VPT2) frequencies (in cm^{-1}) of HC_6H fundamental and selected combination bands

Mode	CCSD(T)/ANO1 ^a		Experimental
	ω	ν	ν
$\nu_1(\sigma_g^+)$	3463.1(0)	3330.4(0)	3313 (Bjarnov et al., 1974)
$\nu_2(\sigma_g^+)$	2284.0(0)	2213.2(0)	2201 (Bjarnov et al., 1974)
$\nu_3(\sigma_g^+)$	2061.0(0)	2023.2(0)	2019 (Bjarnov et al., 1974)
$\nu_4(\sigma_g^+)$	616.1(0)	612.7(0)	625 (Bjarnov et al., 1974)
$\nu_5(\sigma_u^+)$	3463.1(126.4)	3329.5(175.0)	3329.0533 (Doney et al., 2015)
$\nu_6(\sigma_u^+)$	2172.2(0.0)	2130.4(0.1)	2128.91637 (McNaughton & Bruget, 1991)
$\nu_7(\sigma_u^+)$	1169.6(1.7)	1160.9(0.2)	1115.0 (Haas et al., 1994a)
$\nu_8(\pi_g)$	633.0(0)	620.9(0)	622.38 (Matsumura et al., 1988b)
$\nu_9(\pi_g)$	489.5(0)	486.2(0)	491 (Bjarnov et al., 1974)
$\nu_{10}(\pi_g)$	252.0(0)	251.1(0)	258 (Bjarnov et al., 1974)
$\nu_{11}(\pi_u)$	632.0(80.5)	619.9(83.2)	621.34011 (Haas et al., 1994b)
$\nu_{12}(\pi_u)$	444.7(1.0)	441.8(1.0)	443.5 (Haas et al., 1994a)
$\nu_{13}(\pi_u)$	106.4(4.1)	105.9(3.5)	105.038616 (Haas et al., 1994a)
$\nu_9 + \nu_{13}(\sigma_u^+)$	595.9	591.7(0.8)	
$\nu_{10} + \nu_{12}(\sigma_u^+)$	696.7	691.8(1.8)	
$\nu_8 + \nu_{12}(\sigma_u^+)$	1077.7	1063.5(0.3)	
$\nu_9 + \nu_{11}(\sigma_u^+)$	1121.5	1107.1(0.7)	
$\nu_8 + \nu_{11}(\sigma_u^+)$	1265.0	1237.4(31.4)	1232.904295 (McNaughton & Bruget, 1991)
$\nu_3 + \nu_7(\sigma_u^+)$	3230.5	3182.8(0.1)	
$\nu_2 + \nu_7(\sigma_u^+)$	3453.6	3362.2(2.5)	
$3\nu_7(\sigma_u^+)$	3508.7	3498.7(0.01)	
$\nu_1 + \nu_{13}(\pi_u)$	3569.5	3436.1(0.2)	
$\nu_5 + \nu_{10}(\pi_u)$	3715.1	3583.7(0.1)	
$\nu_4 + \nu_5(\sigma_u^+)$	4079.3	3945.8(0.1)	
$\nu_1 + \nu_{11}(\pi_u)$	4095.2	3940.1(0.8)	
$\nu_5 + \nu_8(\pi_u)$	4096.2	3943.9(0.8)	
$\nu_3 + \nu_6(\sigma_u^+)$	4233.1	4141.1(0.1)	
$\nu_2 + \nu_6(\sigma_u^+)$	4456.2	4334.7(0.1)	
$\nu_2 + \nu_5(\sigma_u^+)$	5747.2	5548.4(0.2)	
$\nu_1 + \nu_5(\sigma_u^+)$	6926.3	6555.1(4.6)	
Anharmonic ZPE = 10095.8			

a. Intensities in km/mol are given in parenthesis.

smaller than those seen for acetylene or diacetylene ($\text{o-c} \sim 2 \text{ cm}^{-1}$). For the known combination band, the (fc)-CCSD(T)/ANO1 anharmonicity constants (x_{ij} , **supplementary material**) are able to reproduce the experimental value to within 5 cm^{-1} , suggesting other combination band frequencies are of equal accuracy.

For the modes observed in low-resolution studies (e.g., ν_1 and ν_{12}), the agreement is still good with o-c values less than 20 cm^{-1} . The notable exception is the internal $\text{C}\equiv\text{C}$ asymmetric stretch mode (ν_7), which differs by 45 cm^{-1} . Since no rotationally resolved data can be found for this band, it is possible that the band observed at 1115.0 cm^{-1} (Haas et al., 1994a) was mis-assigned as the ν_7 fundamental. A more likely assignment for this band is the $\nu_9+\nu_{11}$ combination band, which has a predicted VPT2 frequency of 1107.1 cm^{-1} , a calculated intensity of 0.7 km/mol , and the same symmetry. Furthermore, the combination band is expected to be $3.5\times$ more intense than the ν_7 fundamental at 0.2 km/mol , suggesting $\nu_9+\nu_{11}$ is more likely of the two to be observed. However, rotationally resolved measurements of this band are clearly needed to confirm this speculation.

We note that, a resonance between the ν_5 fundamental and the $\nu_2+\nu_7$ and $3\nu_7$ combination bands must be addressed to achieve the very small ($< 1 \text{ cm}^{-1}$) o-c difference obtained for the C-H asymmetric stretch mode, ν_5 . The vibrational frequencies as a result of resonant interactions are calculated by a deperturbation-diagonalization technique followed by transformation of the deperturbed transition moments, as discussed in Vázquez & Stanton (2007) and Matthews et al. (2007). This combination of Fermi and Darling-Dennison interactions shifts the ν_5 predicted frequency from 3333.1 to 3329.5 cm^{-1} , which is able to reproduce the experimentally observed frequency [$3329.0533(2) \text{ cm}^{-1}$ (Doney et al., 2015)] with the same accuracy seen for diacetylene ($\text{o-c} \sim 0.5 \text{ cm}^{-1}$). The combination bands involved are similarly shifted: $\nu_2+\nu_7$ from 3329.5 to 3362.2 cm^{-1} , and $3\nu_7$ from 3526.7 to 3498.7 cm^{-1} . Since the shift is most pronounced for the two combination bands, future experimental work to observe either of these bands is required to confirm this prediction.

The vibration-rotation interaction constants for triacetylene are given in Table 2.4, and are consistent with the previous CCSD(T)/cc-pCVQZ theoretical study (Chang et al., 2016), and experimentally determined values (Matsumura et al., 1988b; McNaughton & Bruget, 1991; Haas et al., 1994a; Doney et al., 2015). Consequently, the calculated ground state rotational constant $B_0 = 0.044\,092 \text{ cm}^{-1}$ is within 0.2% of the experimentally observed $B_0 = 0.044\,1735(12) \text{ cm}^{-1}$ (Doney et al., 2015).

2.3.4 Spectroscopic properties of tetraacetylene

The (fc)-CCSD(T)/ANO1 harmonic and VPT2 frequencies of the fundamental and combination bands for tetraacetylene are given in Table 2.6, and the (fc)-CCSD(T)/ANO1 anharmonicity constants (x_{ij}) are given in the **supplementary material**. For the four experimentally observed bands, agreement of the observed and calculated frequencies is good at $< 7 \text{ cm}^{-1}$, which is comparable to the uncertainty of the low resolution measurements. Furthermore, the ANO1 VPT2 frequencies are able to reproduce the experimental frequencies far better than the previous harmonic frequency calculations, which had o-c values $\sim 20 - 100 \text{ cm}^{-1}$ (Shindo et al., 2001). Of the predicted fundamental and combination bands, there are a number of bands that are found/predicted to have sufficient intensity and/or relatively unique frequency range that could offer viable target transitions to use to search for tetraacetylene in future laboratory or astronomical spectra. For example, in the IR the $\nu_1+\nu_6$ at 6550.8 cm^{-1} or $\nu_{12}+\nu_{15}$ at 871.9 cm^{-1} combination bands have both comparable predicted intensity to measured bands of di- and triacetylene, and have transitions in relatively clean regions of the spectrum. In terms of astronomical searches, the ν_{17} mode at 60.7 cm^{-1} , offers a unique target transition, since it's low frequency makes it accessible by far-IR observations, similar to the ν_2 bending mode of C_3 (Cernicharo et al., 2000).

Based on the results discussed for the other small polyynes, the theoretical vibration-rotation interaction constants given in Table 2.4 are sufficient to assist in identification of ro-vibrational bands of tetraacetylene. The α_i results in a theoretical ground state rotational constant of $B_0 =$

0.018 844 cm^{-1} that agrees within errors with the experimentally determined value, $B_0 = 0.020(3)$ cm^{-1} (Shindo et al., 2001). Overall, for polyynes the difference between the experimental and calculated rotational constants (ΔB_0) decreases from 0.001 to 0.00008 cm^{-1} as the chain length is increased, which is consistent with the trend seen for other carbon chain molecules (*e.g.*, HC_nN , HC_n , C_nO) (Etim & Arunan, 2017). Therefore, if the trend continues as expected then the ΔB_0 for tetraacetylene is equal to or smaller than that seen for triacetylene, and the determined ground state rotational constant is a good approximation of the true value.

2.4 CONCLUSIONS

Accurate equilibrium geometries have been determined at the AE-CCSD(T)/cc-pCVQZ level of theory, and the full cubic and semidiagonal quartic force field have been determined at the (fc)-

Table 2.6: Harmonic and anharmonic (VPT2) frequencies (in cm^{-1}) of HC_8H fundamental and selected combination bands

Mode	CCSD(T)/ANO1 ^a		Experimental
	ω	ν	ν
$\nu_1(\sigma_g^+)$	3462.0(0)	3330.5(0)	
$\nu_2(\sigma_g^+)$	2263.2(0)	2208.0(0)	
$\nu_3(\sigma_g^+)$	2134.6(0)	2094.2(0)	
$\nu_4(\sigma_g^+)$	1296.4(0)	1285.8(0)	
$\nu_5(\sigma_g^+)$	470.0(0)	455.3(0)	
$\nu_6(\sigma_u^+)$	3461.6(223.1)	3328.8(214.2)	3329.4 (Shindo et al., 2001)
$\nu_7(\sigma_u^+)$	2254.7(1.0)	2227.6(0.5)	
$\nu_8(\sigma_u^+)$	2064.3(0.3)	2026.6(0.6)	2023.3 (Shindo et al., 2001)
$\nu_9(\sigma_u^+)$	911.6(3.2)	922.4(2.0)	
$\nu_{10}(\pi_g)$	632.4(0)	620.0(0)	
$\nu_{11}(\pi_g)$	489.3(0)	486.0(0)	
$\nu_{12}(\pi_g)$	422.3(0)	419.5(0)	
$\nu_{13}(\pi_g)$	158.8(0)	157.9(0)	
$\nu_{14}(\pi_u)$	632.6(79.6)	619.7(79.9)	621.5 (Shindo et al., 2001)
$\nu_{15}(\pi_u)$	474.2(0.1)	470.9(0.2)	
$\nu_{16}(\pi_u)$	267.7(3.2)	266.7(3.1)	
$\nu_{17}(\pi_u)$	61.0(2.3)	60.7(2.2)	
$\nu_{11} + \nu_{17}(\sigma_u^+)$	550.3	546.5(0.6)	
$\nu_{13} + \nu_{15}(\sigma_u^+)$	633.0	628.5(1.7)	
$\nu_{12} + \nu_{16}(\sigma_u^+)$	690.1	684.8(3.1)	
$\nu_{12} + \nu_{15}(\sigma_u^+)$	896.5	871.9(3.5)	
$\nu_{11} + \nu_{15}(\sigma_u^+)$	963.5	970.5(1.3)	
$\nu_{10} + \nu_{15}(\sigma_u^+)$	1106.6	1092.2(0.6)	
$\nu_{10} + \nu_{14}(\sigma_u^+)$	1265.1	1236.7(37.5)	1229.7 (Shindo et al., 2001)
$\nu_2 + \nu_9(\sigma_u^+)$	3174.8	3130.1(0.5)	
$\nu_4 + \nu_8(\sigma_u^+)$	3360.7	3311.4(0.5)	
$\nu_6 + \nu_{10}(\pi_u)$	4094.0	3939.1(0.8)	
$\nu_1 + \nu_{14}(\pi_u)$	4094.6	3940.6(0.8)	
$\nu_1 + \nu_6(\sigma_u^+)$	6923.6	6550.8(5.7)	
Anharmonic ZPE = 12218.2			

a. Intensities in km/mol are given in parenthesis.

CCSD(T)/ANO1 level of theory for acetylene and the three smallest polyynes. No scaling or adjustments had to be included to match theoretical values with those determined by experiment. The resulting VPT2 fundamental vibrational frequencies and vibration-rotation interaction constants agree with known experimental values, showing about a 5 cm^{-1} deviation in frequencies for bands with high-resolution infrared information. For bands with only low-resolution data, the theoretical frequencies are able to confirm mode assignments or suggest a reassignment, as in the case of the observed band at 1115.0 cm^{-1} of triacetylene to the $\nu_9+\nu_{11}$ combination band, which has previously been attributed to the ν_7 fundamental. The provisional *ab initio* method used here is also able to accurately reproduce the observed frequencies of combination bands.

The calculated fundamental frequencies for triacetylene and tetraacetylene give insight as to why tetraacetylene has not yet been observed in space. Observation of centrosymmetric molecules in astronomical environments is mainly through infrared detection of the high intensity bending modes; e.g., ν_8 [$628.040\ 776(36)\text{ cm}^{-1}$ (Bizzocchi et al., 2011)] and $\nu_6+\nu_8$ [$1241.060\ 828(38)\text{ cm}^{-1}$ (McNaughton & Bruget, 1992)] of diacetylene, or ν_{11} [$621.340\ 11(42)\text{ cm}^{-1}$ (Haas et al., 1994b)] and $\nu_8+\nu_{11}$ [$1232.904\ 295(74)\text{ cm}^{-1}$ (McNaughton & Bruget, 1991)] of triacetylene. However, the analogous modes for tetraacetylene are the ν_{14} [$621.5(5)\text{ cm}^{-1}$ (Shindo et al., 2001)] and $\nu_{10}+\nu_{14}$ [$1229.7(5)\text{ cm}^{-1}$ (Shindo et al., 2001)], and are predicted to be significantly weaker in intensity due to lower column densities (Woods et al., 2003; Sakai & Yamamoto, 2013). Consequently, at these frequencies and resolutions of the previous infrared observations where polyynes were detected (Cernicharo et al., 2001a,b; Bernard-Salas et al., 2006; Fonfría et al., 2011; Malek et al., 2012), the transitions of tetraacetylene are blended with those of triacetylene. Other bands of tetraacetylene would be more suitable for identification, such as $\nu_1+\nu_6$, $\nu_{12}+\nu_{15}$, or ν_{17} that are expected to be equally strong as bands already used to identify di- and triacetylene.

Overall, the resulting computed geometries lead to equilibrium rotational constants (B_e), which when corrected for vibrational zero-point effects give ground state equilibrium constants (B_0) that agree with experimental values ($< 0.2\%$). Based on the small o-c values for acetylene, diacetylene, and triacetylene, we are confident that the fundamental frequencies and spectroscopic constants determined here offer an accurate guide for spectroscopic searches focused on detection of ro-vibrational bands of triacetylene and tetraacetylene. Such work is underway in our laboratory.

2.5 ACKNOWLEDGEMENTS

We thank Dr. D.J. Nesbitt for helpful discussions. We acknowledge the financial support of the Netherlands Organization for Scientific Research (NWO) through a VICI grant, and the Netherlands Research School for Astronomy (NOVA). D.Z. also acknowledges financial support from the National Key R&D Program of China (2017YFA0303502) and the Fundamental Research Funds for the Central Universities of China.

2.6 BIBLIOGRAPHY

- Almlöf, J. & Taylor, P. R. 1987, *The Journal of Chemical Physics*, 86, 4070
 Arié, E. & Johns, J. W. C. 1992, *Journal of Molecular Spectroscopy*, 155, 195
 Auer, A. A. & Gauss, J. 2001, *Physical Chemistry Chemical Physics*, 3, 3001
 Bak, K. L., Gauss, J., Jørgensen, P., et al. 2001, *The Journal of Chemical Physics*, 114, 6548
 Bandy, R. E., Lakshminarayan, C., Frost, R. K., & Zwier, T. S. 1992, *Science*, 258, 1630
 Bandy, R. E., Lakshminarayan, C., Frost, R. K., & Zwier, T. S. 1993, *The Journal of Chemical Physics*, 98, 5362
 Bartlett, R. J. 1995, in *Modern Electronic Structure Theory*, Part II, ed. D. R. Yarkony (Singapore: World Scientific)
 Bauschlicher Jr., C. W. & Taylor, P. R. 1993, *Theoretica Chimica Acta*, 86, 13
 Bell, M. B., Feldman, P. A., Travers, M. J., et al. 1997, *The Astrophysical Journal Letters*, 483, L61
 Bell, M. B., Feldman, P. A., Watson, J. K. G., et al. 1999, *The Astrophysical Journal*, 518, 740
 Bernard-Salas, J., Peeters, E., Sloan, G. C., et al. 2006, *The Astrophysical Journal Letters*, 652, L29
 Bizzocchi, L., Tamassia, F., Degli Esposti, C., et al. 2011, *Molecular Physics*, 109, 2181

- Bjarnov, E., Christensen, D., Nielsen, O., et al. 1974, *Spectrochimica Acta Part A: Molecular Spectroscopy*, 30, 1255
- Burgdorf, M., Orton, G., van Cleve, J., Meadows, V., & Houck, J. 2006, *Icarus*, 184, 634
- Cernicharo, J., Goicoechea, J. R., & Caux, E. 2000, *The Astrophysical Journal Letters*, 534, L199
- Cernicharo, J. & Guélin, M. 1996, *Astronomy & Astrophysics*, 309, L27
- Cernicharo, J., Heras, A. M., Pardo, J. R., et al. 2001a, *The Astrophysical Journal Letters*, 546, L127
- Cernicharo, J., Heras, A. M., Tielens, A. G. G. M., et al. 2001b, *The Astrophysical Journal Letters*, 546, L123
- CFOUR. 2017, Coupled-Cluster techniques for Computational Chemistry, a quantum-chemical program package by J.F. Stanton, J. Gauss, M.E. Harding, P.G. Szalay with contributions from A.A. Auer, R.J. Bartlett, U. Benedikt, C. Berger, D.E. Bernholdt, Y.J. Bomble, L. Cheng, O. Christiansen, F. Engel, R. Faber, M. Heckert, O. Heun, C. Huber, T.-C. Jagau, D. Jonsson, J. Jusélius, K. Klein, W.J. Lauderdale, F. Lipparini, D.A. Matthews, T. Metzroth, L.A. Mück, D.P. O'Neill, D.R. Price, E. Prochnow, C. Puzzarini, K. Ruud, F. Schiffmann, W. Schwalbach, C. Simmons, S. Stopkowitz, A. Tajti, J. Vázquez, F. Wang, J.D. Watts and the integral packages MOLECULE (J. Almlöf and P.R. Taylor), PROPS (P.R. Taylor), ABACUS (T. Helgaker, H.J. Aa. Jensen, P. Jørgensen, and J. Olsen), and ECP routines by A. V. Mitin and C. van Wüllen. For the current version, see <http://www.cfour.de>.
- Chang, C.-H., Agarwal, J., Allen, W. D., & Nesbitt, D. J. 2016, *The Journal of Chemical Physics*, 144, 074301
- Cherchneff, I., Barker, J. R., & Tielens, A. G. G. M. 1992, *The Astrophysical Journal*, 401, 269
- D'Cunha, R., Sarma, Y., Guelachvili, G., et al. 1991, *Journal of Molecular Spectroscopy*, 148, 213
- de Graauw, T., Feuchtgruber, H., Bezard, B., et al. 1997, *Astronomy & Astrophysics*, 321, L13
- Doney, K. D., Zhao, D., & Linnartz, H. 2015, *Journal of Molecular Spectroscopy*, 316, 54
- Ekern, S. P., Marshall, A. G., Szczepanski, J., & Vala, M. 1998, *The Journal of Physical Chemistry A*, 102, 3498
- Etim, E. E. & Arunan, E. 2017, *Astrophysics and Space Science*, 362, 4
- Feller, D. 1996, *Journal of Computational Chemistry*, 17, 1571
- Fonfría, J. P., Cernicharo, J., Richter, M. J., & Lacy, J. H. 2011, *The Astrophysical Journal*, 728, 43
- Frenklach, M., Clary, D. W., Gardiner, W. C., & Stein, S. E. 1985, *Symposium (International) on Combustion*, 20, 887
- Frenklach, M. & Feigelson, E. D. 1989, *The Astrophysical Journal*, 341, 372
- Frost, R. K., Zavarin, G. S., & Zwier, T. S. 1995, *The Journal of Physical Chemistry*, 99, 9408
- Fujii, T. & Kareev, M. 2001, *Journal of Applied Physics*, 89, 2543
- Gambogi, J. E., Pearson, R. Z., Yang, X., Lehmann, K. K., & Scoles, G. 1995, *Chemical Physics*, 190, 191
- Gauss, J. 1998, in *Encyclopedia of Computational Chemistry*, ed. P. v. R. Schleyer (New York: Wiley)
- Gauss, J. & Stanton, J. F. 1997, *Chemical Physics Letters*, 276, 70
- Gu, X., Kim, Y. S., Kaiser, R. I., et al. 2009, *Proceedings of the National Academy of Sciences*, 106, 16078
- Guelachvili, G., Craig, A. M., & Ramsay, D. A. 1984, *Journal of Molecular Spectroscopy*, 105, 156
- Haas, S., Winnewisser, G., Yamada, K. M. T., Matsumura, K., & Kawaguchi, K. 1994b, *Journal of Molecular Spectroscopy*, 167, 176
- Haas, S., Yamada, K. M. T., & Winnewisser, G. 1994a, *Journal of Molecular Spectroscopy*, 164, 445
- Homann, K.-H. 1998, *Angewandte Chemie International Edition*, 37, 2434
- Kabbadj, Y., Herman, M., Di Lonardo, G., Fusina, L., & Johns, J. W. C. 1991, *Journal of Molecular Spectroscopy*, 150, 535
- Kaiser, R. I., Sun, B. J., Lin, H. M., et al. 2010, *The Astrophysical Journal*, 719, 1884
- Kress, M. E., Tielens, A. G. G. M., & Frenklach, M. 2010, *Advances in Space Research*, 46, 44
- Krestinin, A. V. 2000, *Combustion and Flame*, 121, 513
- Kunde, V. G., Aikin, A. C., Hanel, R. A., et al. 1981, *Nature*, 292, 686
- Kunde, V. G., Flasar, F. M., Jennings, D. E., et al. 2004, *Science*, 305, 1582
- Lee, T. J. & Scuseria, G. E. 1995, in *Quantum Mechanical Electronic Structure Calculations with Chemical Accuracy*, ed. S. Langhoff (Dordrecht: Kluwer Academic Publishers)
- Ma, T.-B., Hu, Y.-Z., & Wang, H. 2008, *Journal of Applied Physics*, 104, 064904
- Malek, S. E., Cami, J., & Bernard-Salas, J. 2012, *The Astrophysical Journal*, 744, 16
- Martin, J. M. L., Lee, T. J., & Taylor, P. R. 1998, *The Journal of Chemical Physics*, 108, 676
- Martin, J. M. L., Taylor, P. R., & Lee, T. J. 1997, *Chemical Physics Letters*, 275, 414
- Matsumura, K., Kanamori, H., Kawaguchi, K., Hirota, E., & Tanaka, T. 1988b, *Journal of Molecular Spectroscopy*, 131, 278
- Matthews, D. A., Vázquez, J., & Stanton, J. F. 2007, *Molecular Physics*, 105, 2659
- McCaslin, L. & Stanton, J. F. 2013, *Molecular Physics*, 111, 1492
- McNaughton, D. & Bruget, D. N. 1991, *Journal of Molecular Spectroscopy*, 150, 620
- McNaughton, D. & Bruget, D. N. 1992, *Journal of Molecular Structure*, 273, 11
- Meadows, V. S., Orton, G., Line, M., et al. 2008, *Icarus*, 197, 585
- Milani, A., Tommasini, M., Del Zoppo, M., Castiglioni, C., & Zerbi, G. 2006, *Physical Review B*, 74, 153418

- Owen, N. L., Smith, C. H., & Williams, G. A. 1987, *Journal of Molecular Structure*, 161, 33
- Palmer, K. F., Mickelson, M. E., & Rao, K. N. 1972, *Journal of Molecular Spectroscopy*, 44, 131
- Pliva, J. 1972a, *Journal of Molecular Spectroscopy*, 44, 165
- Pliva, J. 1972b, *Journal of Molecular Spectroscopy*, 44, 145
- Plyler, E. K., Tidwell, E. D., & Wiggins, T. A. 1963, *Journal of the Optical Society of America*, 53, 589
- Raghavachari, K., Trucks, G. W., Pople, J. A., & Head-Gordon, M. 1989, *Chemical Physics Letters*, 157, 479
- Richter, H. & Howard, J. 2000, *Progress in Energy and Combustion Science*, 26, 565
- Sakai, N. & Yamamoto, S. 2013, *Chemical Reviews*, 113, 8981
- Sattelmeyer, K. W. & Stanton, J. F. 2000, *Journal of the American Chemical Society*, 122, 8220
- Schuchardt, K. L., Didier, B. T., Elsethagen, T., et al. 2007, *Journal of Chemical Information and Modeling*, 47, 1045
- Shindo, F., Bénilan, Y., Chaquin, P., et al. 2001, *Journal of Molecular Spectroscopy*, 210, 191
- Shirakawa, H. 2001, *Review of Modern Physics*, 73, 713
- Simandiras, E. D., Rice, J. E., Lee, T. J., Amos, R. D., & Handy, N. C. 1988, *The Journal of Chemical Physics*, 88, 3187
- Simmonett, A. C., Schaefer III, H. F., & Allen, W. D. 2009, *The Journal of Chemical Physics*, 130, 044301
- Stanton, J. F. & Gauss, J. 2000, *International Reviews in Physical Chemistry*, 19, 61
- Tamassia, F., Cane, E., Fusina, L., & Di Lonardo, G. 2016, *Physical Chemistry Chemical Physics*, 18, 1937
- Tay, R., Metha, G. F., Shanks, F., & McNaughton, D. 1995, *Structural Chemistry*, 6, 47
- Temsamani, M. A. & Herman, M. 1995, *The Journal of Chemical Physics*, 102, 6371
- Thaddeus, P., McCarthy, M. C., Travers, M. J., Gottlieb, C. A., & Chen, W. 1998, *Faraday Discussions*, 109, 121
- Thejaswini, H. C., Majumdar, A., Tun, T. M., & Hippler, R. 2011, *Advances in Space Research*, 48, 857
- Thorwirth, S., Harding, M. E., Muders, D., & Gauss, J. 2008, *Journal of Molecular Spectroscopy*, 251, 220
- Toto, J. L., Toto, T. T., de Melo, C. P., Kirtman, B., & Robins, K. 1996, *The Journal of Chemical Physics*, 104, 8586
- Vanderauwera, J., Hurtmans, D., Carleer, M., & Herman, M. 1993, *Journal of Molecular Spectroscopy*, 157, 337
- Vázquez, J. & Stanton, J. F. 2007, *Molecular Physics*, 105, 101
- Vuitton, V., Doussin, J.-F., Bénilan, Y., Raulin, F., & Gazeau, M.-C. 2006, *Icarus*, 185, 287
- Vuitton, V., Yelle, R. V., & McEwan, M. J. 2007, *Icarus*, 191, 722
- Waite, J. H., Young, D. T., Cravens, T. E., et al. 2007, *Science*, 316, 870
- Wakelam, V., Smith, I. W. M., Herbst, E., et al. 2010, *Space Science Reviews*, 156, 13
- Wilson, E. H. & Atreya, S. K. 2003, *Planetary and Space Science*, 51, 1017
- Woods, P. M., Millar, T. J., Herbst, E., & Zijlstra, A. A. 2003, *Astronomy & Astrophysics*, 402, 189
- Woon, D. E. & Dunning Jr., T. H. 1995, *The Journal of Chemical Physics*, 103, 4572
- Zhang, X., Maccarone, A. T., Nimlos, M. R., et al. 2007, *The Journal of Chemical Physics*, 126, 044312
- Zhao, D., Doney, K. D., & Linnartz, H. 2014a, *Journal of Molecular Spectroscopy*, 296, 1
- Zhao, D., Guss, J., Walsh, A. J., & Linnartz, H. 2013, *Chemical Physics Letters*, 565, 132
- Zhao, X., Ando, Y., Liu, Y., Jinno, M., & Suzuki, T. 2003, *Physical Reviews Letters*, 90, 187401

