

Photo-activated drug delivery systems Kong, L.

Citation

Kong, L. (2018, June 7). *Photo-activated drug delivery systems*. Retrieved from https://hdl.handle.net/1887/63080

Version:	Not Applicable (or Unknown)
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	<u>https://hdl.handle.net/1887/63080</u>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The following handle holds various files of this Leiden University dissertation: http://hdl.handle.net/1887/63080

Author: Kong, L. Title: Photo-activated drug delivery systems Issue Date: 2018-06-07

Preface:

The work in this thesis focuses on the development of new stimuli-responsive drug delivery systems. All four described systems rely on light as the exclusive trigger of activation and the spatiotemporal precision afforded by this approach is demonstrated, both *in vitro* and *in vivo*, throughout this thesis. Light activation, for three of the four described systems (**chapters 2-4**), leads to dePEGylation of a nanoparticle surface. In this way, the benefits of PEGylation (*e.g.* limited non-specific cellular interactions) are maintained, while the obstacles of PEGylation (*e.g.* limited uptake by targeted cells) can be overcome on demand. In **chapter 5**, light is used to switch the surface charge of a nanoparticle *in situ* and *in vivo*. The effectiveness of this approach is demonstrated within zebrafish embryos. Despite countless reports of stimuli-responsive drug delivery systems in the literature, none have yet made it to the clinic. The work in this thesis is therefore aimed at providing potential solutions to some of the roadblocks slowing the progression of stimulus-responsive drug delivery systems.