



Universiteit
Leiden
The Netherlands

Regulation of BMP and TGF β signaling pathway in cancer progression

Ren, J.

Citation

Ren, J. (2020, June 24). *Regulation of BMP and TGF β signaling pathway in cancer progression*. Retrieved from <https://hdl.handle.net/1887/123057>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/123057>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/123057> holds various files of this Leiden University dissertation.

Author: Ren, J.

Title: Regulation of BMP and TGF β signaling pathway in cancer progression

Issue Date: 2020-06-24

Chapter 2

Invasive Behavior of Human Breast Cancer Cells in Embryonic Zebrafish

Jiang Ren*, Sijia Liu*, Chao Cui, and Peter ten Dijke

Journal of visualized experiments 2017, (122):e55459.

*These authors contributed equally

Abstract

In many cases, cancer patients do not die of a primary tumor, but rather because of metastasis. Although numerous rodent models are available for studying cancer metastasis *in vivo*, other efficient, reliable, low-cost models are needed to quickly access the potential effects of (epi)genetic changes or pharmacological compounds. As such, we illustrate and explain the feasibility of xenograft models using human breast cancer cells injected into zebrafish embryos to support this goal. Under the microscope, fluorescent proteins or chemically labeled human breast cancer cells are transplanted into transgenic zebrafish embryos, Tg (*flil:EGFP*), at the perivitelline space or duct of Cuvier (Doc) 48 h after fertilization. Shortly afterwards, the temporal-spatial process of cancer cell invasion, dissemination, and metastasis in the living fish body is visualized under a fluorescent microscope. The models using different injection sites *i.e.*, perivitelline space or Doc are complementary to one another, reflecting the early stage (intravasation step) and late stage (extravasation step) of the multistep metastatic cascade of events. Moreover, peritumoral and intratumoral angiogenesis can be observed with the injection into the perivitelline space. The entire experimental period is no more than 8 days. These two models combine cell labeling, micro-transplantation, and fluorescence imaging techniques, enabling the rapid evaluation of cancer metastasis in response to genetic and pharmacological manipulations.

Keywords: Embryonic zebrafish, Human breast cancer, Metastasis, Intravasation, Extravasation, Perivitelline space, Duct of Cuvier

Video Link: The video component of this article can be found at <https://www.jove.com/video/55459/>

Introduction

Overt cancer metastasis in the clinic comprises a series of complex and multi-step events known as the ‘metastatic cascade’. The cascade has been extensively reviewed and can be dissected into successive steps: local invasion, intravasation, dissemination, arrest, extravasation, and colonization [1, 2]. A better understanding of the pathogenesis of cancer metastasis and the development of potential treatment strategies *in vivo* require robust host models of cancer cell spread. Rodent models are well established and widely used to evaluate metastasis [3], but these approaches have low efficiency, ethical limitations, and are costly as a forefront model to determine whether a particular manipulation could affect the metastatic phenotype. Other efficient, reliable, low-cost models are needed to quickly access the potential effect of (epi)genetic changes or pharmacological compounds. Due to the high genetic homology to humans and transparency of the embryos, the zebrafish (*Danio rerio*) has emerged as an important vertebrate model and is being applied increasingly in studying developmental processes, microbe-host interactions, human disease, drug screening, *etc.* [4]. The cancer metastasis models established in zebrafish may provide ideal solutions to the shortcomings of rodent models [5, 6].

Although spontaneous neoplasia is scarcely discovered in wild zebrafish [7], there are several longstanding techniques to induce desired cancer in zebrafish. Carcinogen-induced gene mutations or signaling pathways-activation can model carcinogenesis histologically and molecularly resembling human disease in zebrafish [7-9]. By taking advantage of diverse forward and reverse genetic manipulations of oncogenes or tumor suppressors, (transgenic) zebrafish also have enabled potential studies of cancer formation and maintenance [6, 10]. The induced cancer models in zebrafish cover a broad spectrum of cancer types in digestive, reproductive, blood, nervous systems, and epithelium [6].

The utilization of zebrafish in cancer research has expanded recently due to the establishment of human tumor cell xenograft models in this organism. This was first reported with human metastatic melanoma cells that were successfully engrafted in zebrafish embryos at the blastula stage in 2005 [11]. Several independent laboratories have validated the feasibility of this pioneering work by introducing a diverse range of mammalian cancer cells lines into zebrafish at various sites and developmental stages [5]. For example, injections near the blastodisc and blastocyst of the blastula stage; injections into the yolk sac, perivitelline space,

duct of Cuvier (Doc), and posterior cardinal vein of 6-h- to 5-day old embryos; and injections into the peritoneal cavity of 30-day-old immunosuppressed larvae have been performed [5,12]. Additionally, allogeneic tumor transplantations were also reported in zebrafish [12,13]. One of the great advantages of using xenografts is that the engrafted cancer cells can be easily fluorescently labeled and distinguished from normal cells. Hence, investigations into the dynamic behaviors of microtumor formation [14], cell invasion and metastasis [15-17], tumor-induced angiogenesis [15,18], and the interactions between cancer cells and host factors [17] can be clearly visualized in the live fish body, especially when transgenic zebrafish lines are applied [5].

Inspired by the high potential of zebrafish xenograft models to evaluate metastasis, we demonstrated the transvascular extravasation properties of different breast cancer cell lines in the tailfin area of Tg (*fli:EGFP*) zebrafish embryos through Doc injections [16]. The role of transforming growth factor- β (TGF β) [16] and bone morphogenetic protein (BMP) [19] signaling pathways in pro-/anti-breast cancer cell invasion and metastasis were also investigated in this model. Moreover, we also recapitulated the intravasation ability of various breast cancer cell lines into circulation using xenograft zebrafish models with perivitelline space injections.

This article presents detailed protocols for zebrafish xenograft models based upon the injection of human breast cancer cells into the perivitelline space or Doc. Using high-resolution fluorescence imaging, we show the representative process of intravasation into blood vessels and the invasive behavior of different human breast cancer cells, which move from the blood vessels into the avascular tailfin area.

Protocol

All research using transgenic fluorescent zebrafish Tg (*fli:EGFP*) strain, which has enhanced green fluorescent protein (EGFP) labeled vasculature²⁰, including housing and experiments, was carried out according to the international guidelines and approved by the local Institutional Committee for Animal Welfare (Dier Ethische Commissie (DEC) of the Leiden University Medical Center.

NOTE: As summarized in Figure 1, the protocol is roughly dissected into four steps, embryo collection (Figure 1A), microinjection (Figure 1B), screening (Figure 1C), and analysis (Figure 1D).

1. Prepare the injection needles

1. Prepare injection needles with borosilicate glass microcapillary. Put the microcapillary in a micropipette puller device with the following settings: air pressure 500; heat 650; pull 100; velocity 200; time 40. Keep the injection needles in a needle holder plate until used for injection.




Figure 1. Main steps for investigating the invasive behavior of breast cancer cells in embryonic zebrafish. **A**, After crossing parental zebrafish overnight, Tg (*fli:EGFP*) zebrafish embryos were collected the following morning and maintained at 28 °C. **B**, The embryos were dechorionated with fine tweezers under a stereo microscope 48 h post fertilization (hpf). The labeled breast cancer cells were collected and re-suspended in a small amount of PBS. After well-preparation, suspended cells were loaded into one needle. Approximately 400 cells were injected into the duct of Cuvier (Doc) of the perivitelline space under a stereo microscope. The injected embryos were maintained at 34 °C. **C**, 2 hours post injection (hpi), the embryos were subjected to careful screening under a fluorescence stereo microscope. The embryos were maintained at 34 °C for 3 or 6 d. During the interval, embryos could be subjected to designed treatment. **D**, Cancer cell dissemination by perivitelline space injection or invasion

by Doc injection was detected, counted, and imaged by confocal microscopy 3 or 6 days post injection (dpi).

2. Prepare of the fluorescent genetically labeled breast cancer cells for injection

1. Culture human breast cancer MDA-MB-231 cells at 37 °C in DMEM-high glucose media containing L-glutamine, 10% fetal bovine serum and 1:100 Penicillin-Streptomycin (Pen-Strep).
2. Culture the breast epithelial cell line MCF10A (M1), MCF10A-Ras (M2) at 37 °C in DMEM/F12 media containing L-glutamine, with 5% horse serum, 20 ng/mL epidermal growth factor, 10 mg/mL insulin, 100 ng/mL cholera enterotoxin, 0.5 mg/mL hydrocortisone, and 1:100 Pen-Strep.
3. Produce mCherry lentivirus by co-transfecting PLV-mCherry, pCMV-VSVG [21], pMDLg-RRE (gag/pol) [22], and pRSV-REV [22] plasmids into HEK293T cells. Harvest cell supernatants 48 h after transfection and store at -80 °C.
4. Infect MDA-MB-231, M1 and M2 cells at 30% confluence for 24 h with lentiviral supernatants diluted 1:1 with normal culture medium in the presence of 5 ng/mL polybrene.
5. Select single cell clones by diluting cells in a 96-well plate, which allows the outgrowth of isolated cell clones, until obtaining the stable mCherry-expressing cell lines.
6. Culture one T75 flask of cells for injection. Harvest the cells at 80% confluence with a 0.5% trypsin-EDTA treatment. Wash the cells with 1× PBS 2-3 times.
7. Re-suspend the cells in about 200 µL PBS. Store at 4 °C for less than 5 h before injection.

3. Prepare zebrafish embryos for injection

1. Set up zebrafish breeding pairs and collect embryos as shown in a previous Jove article by Rosen *et al.* [23].
2. Select the embryos that are at 0-4 hpf by removing the unfertilized and abnormal embryos. Keep the embryos in a petri-dish filled with egg water (60 µg/mL sea salts; about 60 embryos/dish) and incubate at 28 °C.
3. Dechorionate the embryos with fine tweezers at 48 hpf.
4. Anesthetize the embryos by transferring them to 40 µg/mL Tricaine (3-aminobenzoic acid) containing egg water approximately 2 min prior to injection, but no longer than 2 h prior to injection.

NOTE: Tricaine stock solution (4 mg/mL, 100×) is prepared as 400 mg tricaine powder in 97.9 mL double-distilled water and 2.1 mL 1 M Tris-base (pH 9), adjust pH to 7.4. Store in the -20 °C freezer.

4. Inject human breast cancer cells into the perivitelline space

1. Load 15 μ L of the cell suspension into an injection needle. Mount the needle onto the micromanipulator and break off the needle tip with fine tweezers to obtain a tip opening diameter of 5-10 μ m.
2. Use a pneumatic picopump and a manipulator to perform microinjection. Adjust the picopump to inject 400 cells each time. Prior to injection, count the cell numbers manually by injecting the cells on the top of a petri-dish containing 1% agarose.
3. Line up anesthetized embryos (2-3 days post fertilization (dpf)) on a flat 1% agarose injecting plate, around 10 embryos each time.
4. Orient the injection plate by hand during injections to place the embryos in the preferred position for inserting the needle (*i.e.*, diagonally).
5. Point the needle tip to the injection site and gently insert the needle tip into the perivitelline space between the yolk sac and the periderm of the zebrafish embryo (Figure 2A).




Figure 2. Perivitelline space injection site and common errors. **A**, Approximately 400 mCherry-labeled cells (MDA-MB-231) were injected into the perivitelline space. The brightfield (upper most), green vasculature (middle upper), and red cell mass (middle lower) of injected zebrafish embryos were captured by confocal microscope. The merged image (lower most) of three channels shows the stereo location of the cell mass in the embryo. **B**, The cells did not target the perivitelline space appropriately. The yolk sac was ruptured. **C**, Injected cells below threshold (much less than 400). **D**, Injected cells above threshold (much more than 400). The cell mass was too close to the duct of Cuvier, which has a broad blood stream. Scale bar = 50 μm .

6. Inject approximately 400 mCherry-labeled tumor cells. Make sure that the yolk sac is not ruptured to avoid implantation into the yolk sac.

5. Inject human breast cancer cells into the Doc

1. Prepare injection needle and zebrafish embryos as described in protocol steps 1, 2, and 3.
2. Use a 45° needle angle so that the Doc can be approached from the dorsal side of the embryo.
3. Insert the needle into the starting point of the Doc (Figure 3A) just dorsal to where the duct starts broadening over the yolk sac and inject approximately 400 cells. The injection is correct if the volume within the duct expands directly after the pulse and the yolk sac.

NOTE: Several consecutive injections can be performed without extracting the needle.

4. Transfer the injected zebrafish embryos to egg water.

NOTE: As considerable variation exists among individual zebrafish embryos, as well as the death of embryos after injection, relatively large number of zebrafish embryos (around 100) should be injected with cancer cells.

5. Maintain the zebrafish embryos at 34 °C to accommodate the optimal temperature requirements for fish and mammalian cells.

6. Screen the injected embryos

1. Screen each fish under a fluorescence stereo microscope at 2 h post-injection (hpi) for perivitelline space injection (Figure 2) or at 2-24 hpi for Doc injection (Figure 2), to ensure all the embryos are injected with similar number of tumor cells. Remove the embryos with injection errors, such as rupture (Figure 2B) or injection (Figure 3B) of yolk sac, and pick

out embryos with injected cells below (Figure 2C and Figure 3B) or above (Figure 2D and Figure 3B) threshold. Keep only the embryos with approximately 400 cells in culture.

2. Rule out the possibility that cells are introduced directly into the circulation during the injection process by removing the embryos with cells already in the circulation from further analysis. Also remove any embryo with a cell mass close to the Doc (Figure 2D).




Figure 3. Overview of duct of Cuvier (Doc) injection. A, Schematic of Doc injection at 2 days post-fertilization (dpf) with breast cancer cells in zebrafish embryos. Arrow indicates Doc. B, Examples of positive injection with around 400 breast cancer cells, negative injections including the yolk mis-injection and incorrect number of cells injection at 4 hpi. Arrows and circles indicate injected cells.

7. Image and analyze the metastatic process

1. Collect several anesthetized embryos with a wide-tip Pasteur pipette, and transfer them onto the glass bottom of a polystyrene dish.
2. Remove excess water and keep a limited amount of egg water. Manipulate the embryo into position with a hair loop tool, and place a cover on top of the glass.

3. Use an inverted confocal microscope in combination with water-immersion or long-distance dry objectives. The embryo should be positioned so that the region of interest is as close to the objective as possible.
4. Perform imaging immediately after anesthesia to reduce death risk of embryo due to liquid evaporation.
 1. Capture signals from EGFP-labeled vasculature and mCherry labeled tumor cells at the same position of the embryos to co-register injected cells with blood vessels by merging the two imaging channels.
 2. For each zebrafish embryo, collect two different sets of images from the head region and tail region.
5. Quantify the number of disseminated cells.
 1. For perivitelline space injection, count the number of cells in each fish that disseminated from the cell mass toward the embryonic fish body within the head and tail regions^{4,15}. The regions are beyond the boundaries of the heart cavity frontally, on top of the swim bladder dorsally, and beyond the urogenital opening caudally.
 2. For Doc injection, count the number of individual cells that invade the collagen fibers of the tail fin from circulation (MDA-MB-231) or the number of clusters formed by cells collectively (M2) in the caudal hematopoietic tissue (CHT) of each zebrafish [19].
6. Study invasion and metastasis in more detail, use confocal microscopy (highly recommended).
 1. Use low magnification (4× objective) to image the whole body and obtain an overview of the tumor cell dissemination pattern.

NOTE: Higher magnification (20× and 40× objectives) is suitable for studying intra- and peri-tumoral angiogenesis and precise localization of disseminated cells in the embryo body.
 2. Use a 488-nm laser to scan the zebrafish embryo vasculature, and a 543 nm laser to scan implanted tumor cells labeled with red fluorescence. Obtain a high-quality image, by scanning each embryo in eight to ten steps. Scan and average each step six times.
7. Carefully place the embryo back into the egg water if it is required for further experiments.

8. Perform statistical analysis using one-way analysis of variance (ANOVA) followed by post Hoc analysis

Representative results

In the embryonic xenograft zebrafish model with a perivitelline space injection, the hematogenous dissemination of labeled cancer cells in the fish body is considered as active migration. This process can be detected and quantified under a fluorescent microscope, as described in the methods above. To illustrate this xenograft model, we followed the dissemination process of different breast cancer cell lines with known (or without) invasion/metastasis potential according to *in vitro* and *in vivo* mouse studies, including the benign normal breast epithelial M1 cells, HRAS-transformed premalignant M2 cells, and highly metastatic MDA-MB-231 cells, 1 day post injection (dpi) onward. A high-resolution confocal microscopy image showed that MDA-MB-231 cells (red) exhibit an aggressive phenotype, with irregular borders in the perivitelline space. Pseudopodia-like protrusions and invasive fronts were also frequently present (Figure 4A, left). A few cells disseminated into blood circulation as early as 1 dpi (Figure 4A, right). At 2 dpi, clear dissemination was observed in the distal parts of the fish (Figure 4A, right). The number of disseminated cells increased further at 3 dpi (Figure 4A, D). In contrast, when M2 cells were challenged in zebrafish, they exhibited modest spread in the fish body after 2 dpi (Figure 4B). They also showed increased dissemination after time passed (Figure 4F). As shown in Figure 4C and 4G, M1 cells infrequently disseminated into zebrafish circulation, and even active local migration within the perivitelline space was infrequent during the period of observation. The M1 cell mass was virtually detained at the original injection site. If defining positive dissemination or metastasis as >5 cells in the fish body [4], MDA-MB-231 and M2 cell metastasis was observed in 92% and 57% of fish, respectively, at 3 dpi (Figure 4G). In contrast, no positive dissemination was observed with M1 cells. Therefore, this zebrafish model of human cancer cell progression accurately reflects the relative level of metastatic potential of the different cells in mice. Neovascularization (green) that sprouted from the subintestinal plexus of the embryonic zebrafish and penetrated the MDA-MB-231 or M2 cell mass was also present after the perivitelline space injection of tumor cells followed by 3 days of incubation (Figure 4A, B, left). Consistent with the disability in dissemination, only slight neovascularization was detected upon M1 cell implantation (Figure 4C).

In the embryonic xenograft zebrafish model with mCherry-labeled MDA-MB-231 cells and the Doc injection, the labeled cancer cells in the tailfin of the zebrafish are considered representative of active extravasation. The mCherry-labeled MDA-MB-231 cells were injected at




Figure 4. Comparison of dissemination ability among various breast cell lines. Approximately 400 mCherry-labeled MDA-MB-231, MCF10Aras (M2), or MCF10A (M1) cells were injected into the perivitelline space of zebrafish embryos 48 hpf. The injected embryos were followed for 3 days. **A-C**, High-resolution micrographs showing the representative migration and dissemination process of MDA-MB-231 (**A**), M2 (**B**), and M1 (**C**) cells in individual embryonic bodies 1, 2, and 3 days post-injection (dpi). Left, cell migration in the perivitelline space (red) and the peritumoral and intratumoral vasculature (green). Yellow signals indicate the overlap of microvessels and cells. Middle, the whole image of embryo. Right, visualization of disseminated cells in the posterior of embryo. Yellow arrowheads indicate single disseminated cells. Scale bar = 50 μ m. **D-F**, Quantification of the number of disseminated cells in each embryonic body at 1, 2, 3 dpi. Results are expressed as the Mean \pm SEM. Results from one-way analysis of variance (ANOVA) followed by the post hoc analysis are shown. $P < 0.05$ was accepted as statistically significant (* $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$). **G**, Comparison of the incidence of intravasation for MDA-MB-231, M2, and M1 cells in embryonic bodies at 1, 2, 3 dpi.




Figure 5. Different behavior of MDA-MB-231 and M2 cell metastasis in zebrafish with duct of Cuvier injection. **A**, Representative confocal images of the zebrafish followed at 3, 4, 5 dpi to show the single cell migration behavior of the MDA-MB-231 cells in zebrafish. Arrows indicate invasive MDA-MB-231 cells that migrated out of the vessels to the tail fins. Scale bar = 200 μ m in the left column, 50 μ m in the right column. **B**, Representative confocal images of the zebrafish followed at 1, 2, 3 dpi to show the cell cluster migration behavior of M2 cells in zebrafish. Arrows indicate invasive M2 cells that migrated out of the vessels to the caudal hematopoietic tissue (CHT) and formed a cluster between the vessels.

2 dpf. At 3 dpi, the cells started to migrate out of the vessels to the tailfin, which is enriched with collagen. Single MDA-MB-231 cells migrated one by one, independently from the vessels, to the distant tailfin (Figure 5A). At 6 dpi, the invasion could be quantified by counting the number of cells that migrated into the tailfin tissue. In the mCherry-labeled M2 cell Doc injection model, the injection was also performed at 2 dpf. However, a clustered phenotype was observed during the active extravasation process. At 1 dpi, M2 cells started to migrate out from the vessels into the CHT of the zebrafish. At 2 dpi, the migrated M2 cells started to form a cluster between the vessels in the CHT (Figure 5B). Quantification of the M2 invasive cell cluster number in the CHT region could be conducted at 6 dpi.

Discussion

Here, we described two methods to investigate the invasive behavior of breast cancer cells in Tg (*fli1:EGFP*) zebrafish embryos, with perivitelline space and Doc injections. By injecting cancer cells labeled with chemical dye or fluorescent protein into transgenic zebrafish embryos, the dynamic and spatial characteristics of invasion and metastasis can be clearly tracked in real-time at the single-cell or cluster level under a fluorescence microscope. In most cases, the rapid progression of metastasis in zebrafish ensures that the assay can be performed within 1 week after transplantation. Moreover, powerful statistics can be obtained with large cohorts of fish.

Early and late events of the metastatic cascade could be simulated and recapitulated by injecting cancer cells into the perivitelline space or Doc, respectively. The perivitelline space is the confined space between the periderm of the fish and the yolk sac, which allows one to monitor dissemination of single tumor cells from primary sites in the living body. After implantation, the cancer cells undergo local migration and invasion within the perivitelline space (considered the primary site) and then they intravasate into blood vessels and disseminate along with the circulation. At the head and tailfin (considered distant target sites), cancer cells accumulate in narrow capillary beds and extravasate. Therefore, the number of cells that are found at the distant sites in the fish body is a measurement of metastatic capability. In addition, more extravasated cells can be observed at later time points, which is also true of the Doc injection assay.

The Doc is an enlarged common cardinal vein with an extensive blood stream [24]. Directly targeting the Doc as an injection site introduces cancer cells into the circulatory system. In practice, breast cancer cells diffuse throughout the embryonic body via the blood stream

instantly after Doc injection. The cells then arrest at the caudal vein and dorsal aorta. Extravasation, invasion, and micrometastasis formation can be observed successively within 6 days. As reported previously [16], metastatic MDA-MB-231 cells and premalignant mammary M2 cells exhibit different invasive phenotypes. MDA-MB-231 cells undergo single-cell invasion of the collagen matrix-rich tailfin. Thus, the invasion potential of MDA-MB-231 cells can be measured by counting the number of cells that have extravasated and invaded the tailfin tissue. In contrast, M2 cells form clusters of different sizes and undergo collective invasion of the CHT. Quantifying the invasion potential of M2 cells by counting the number of clusters in this protocol is difficult and is preferably performed by making a 3D image using confocal microscopy and determining the volume of clustered tumor cells.

The technical challenge in cancer cell microinjection is successfully targeting the perivitelline space or Doc. The microinjection of large numbers of embryos is a tedious procedure requiring a highly skilled and patient operator. Factors that contribute to variations in the results in individual fish include the developmental stage of the embryo when injecting, differences in the number of cells injected, and the leakage of cells into the yolk sac. Though rare, the manipulation could unintentionally penetrate the vasculature and introduce cells into the circulatory system directly, especially in the perivitelline space injection. To further reduce variation and to ensure the reliability of the analyses, microscopic examination is necessary to exclude unqualified fish at time points throughout the process. In addition, blinded analysis by a professional without knowledge of the setting is strongly suggested to achieve unbiased quantification.

In summary, the two models we introduced here shed light on visualizing the processes of cell invasion and metastasis *in vivo* without invasive procedures. Although we only studied breast cancer cells in two models regarding metastatic potential, they could be extrapolated to other types of cancer. Moreover, the models could have broader applications in determining the mechanisms and new molecular targets controlling cancer cell metastasis using (epi)genetic manipulation. Due to the higher penetrability of zebrafish embryos by small-molecule compounds as compared to the feeding or injection of rodents [25], the two presented models also have advantages in terms of the high-throughput screening of potential new anti-invasion/metastasis drugs.

Disclosures

The authors have nothing to disclose.

Acknowledgements

Studies on TGF β family members are supported by the Cancer Genomics Centre Netherlands. Sijia Liu and Jiang Ren are supported by the China Scholarship Council for 4 years of study at the University of Leiden. We thank Dr. Fred Miller (Barbara Ann Karmanos Cancer Institute, Detroit, MI, USA) for the MCF10A cell lines.

References

1. Wan L, Pantel K, Kang Y. Tumor metastasis: moving new biological insights into the clinic. *Nat Med* 2013, 19(11):1450-64.
2. Obenauf AC, Massagué J. Surviving at a distance: Organ-specific metastasis. *Trends Cancer* 2015, 1(1):76-91.
3. Saxena M, Christofori G. Rebuilding cancer metastasis in the mouse. *Mol Oncol* 2013, 7 (2):283-96.
4. Teng Y, Xie X, Walker, S. *et al.* Evaluating human cancer cell metastasis in zebrafish. *BMC cancer* 2013, 13(1):453.
5. Konantz, M, Balci, TB, Hartwig, UF, *et al.* Zebrafish xenografts as a tool for in vivo studies on human cancer. *Ann N Y Acad Sci* 2012, 1266 (1):124-37.
6. Zhao, S, Huang, J, Ye, J. A fresh look at zebrafish from the perspective of cancer research. *J Exp Clin Cancer Res* 2015, 34(1):80.
7. Stanton, MF. Diethylnitrosamine-induced hepatic degeneration and neoplasia in the aquarium fish, *Brachydanio rerio*. *J Natl Cancer Inst* 1965, 34(1):117-30.
8. Lam SH, Wu YL, Vega VB, *et al.* Conservation of gene expression signatures between zebrafish and human liver tumors and tumor progression. *Nat Biotechnol* 2006, 24(1):73-5.
9. Spitsbergen JM, Kent ML. The state of the art of the zebrafish model for toxicology and toxicologic pathology research-advantages and current limitations. *Toxicol Pathol* 2003, 31(1 suppl):62-87.
10. Stoletov K, Klemke R. Catch of the day: zebrafish as a human cancer model. *Oncogene* 2008, 27(33):4509-20.
11. Lee LM, Sefior EA, Bonde, G, *et al.* The fate of human malignant melanoma cells transplanted into zebrafish embryos: assessment of migration and cell division in the absence of tumor formation. *Dev Dynam* 2005, 233(4):1560-70.
12. Mizgirev I, Revskoy S. Generation of clonal zebrafish lines and transplantable hepatic tumors. *Nat Protoc* 2010, 5(3):383-94.

13. Mizgireuv IV, Revskoy SY. Transplantable tumor lines generated in clonal zebrafish. *Cancer Res* 2006, 66(6):3120-5.
14. Stoletov K, Montel V, Lester RD, *et al.* High-resolution imaging of the dynamic tumor cell-vascular interface in transparent zebrafish. *Proc Natl Acad Sci U.S.A* 2007, 104(44):17406-11.
15. Rouhi P, Jensen LD, Cao Z, *et al.* Hypoxia-induced metastasis model in embryonic zebrafish. *Nat Protoc* 2010, 5 (12):1911-8.
16. Drabsch Y, He S, Zhang L, *et al.* Transforming growth factor- β signalling controls human breast cancer metastasis in a zebrafish xenograft model. *Breast Cancer Res* 2013, 15(6):R106.
17. He, S, Lamers, GE, Beenakker, JW, *et al.* Neutrophil-mediated experimental metastasis is enhanced by VEGFR inhibition in a zebrafish xenograft model. *J Pathol* 2012, 227(4):431-45.
18. Nicoli S, Presta M. The zebrafish/tumor xenograft angiogenesis assay. *Nat Protoc* 2007, 2(11):2918-23.
19. Boeck M, Cui C, Mulder AA, *et al.* Smad6 determines BMP-regulated invasive behaviour of breast cancer cells in a zebrafish xenograft model. *Sci Rep* 2016, 6:24968.
20. Lawson ND, Weinstein BM. In vivo imaging of embryonic vascular development using transgenic zebrafish. *Dev Biol* 2002, 248(2):307-18.
21. Stewart SA, Dykxhoorn DM, Palliser D, *et al.* Lentivirus-delivered stable gene silencing by RNAi in primary cells. *RNA* 2003, 9(4):493-501.
22. Dull T, Zufferey R, Kelly M, *et al.* A third-generation lentivirus vector with a conditional packaging system. *J Virol* 1998, 72 (11):8463-71.
23. Rosen, JN, Sweeney MF, Mably JD. Microinjection of zebrafish embryos to analyze gene function. *J Vis Exp* 2009, (25):e1115.
24. Lawson ND, Weinstein BM. Arteries and veins: making a difference with zebrafish. *Nat Rev Genet* 2002, 3(9):674-82.
25. Zon LI, Peterson RT. *In vivo* drug discovery in the zebrafish. *Nat Rev Drug Discov* 2005, 4(1):35-44.

Materials

Name	Company	Catalog number	Comments
Agarose	MP Biomedicals	AGAF0500	
Borosilicate glass capillary	Harvard Apparatus	300038	
Cholera enterotoxin	Calbiochem	227035	
Confocal microscope	Leica	SP5 STED	
DMEM-high glucose media containing L-glutamine	ThermoFisher Scientific	11965092	
DMEM/F-12 media containing L-glutamine	ThermoFisher Scientific	21041025	
Dumont #5 forceps	Fine Science Tools Inc	11252-20	
Epidermal growth factor	Merck Millipore	01-107	
Fetal bovine serum	ThermoFisher Scientific	16140071	
Fluorescent stereo microscope	Leica	M165 FC	
HEK293T cell line	American Type Culture Collection	CRL-1573	
Hydrocortisone	SigmaAldrich	227035	
Horse serum	ThermoFisher Scientific	26050088	
Insulin	SigmaAldrich	I-6634	
MCF10A (M1) cell line			Kindly provided by Dr. Fred Miller (Barbara Ann Karmanos Cancer Institute, Detroit, MI, USA)
MCF10Aras (M2) cell line			
MDA-MB-231 cell line	American Type Culture Collection	CRM-HTB-26	
Manual micromanipulator	World Precision Instruments	M3301R	
Micropipette puller	Sutter Instruments	P-97	
Wide-tip Pasteur pipette (0,5-20 ul)	Eppendorf	F276456I	

pCMV-VSVG plasmid			Kindly provided by Prof. Dr.
pMDLg-RRE (gag/pol)			Rob Hoeben (Leiden
plasmid			University Medical Center,
pRSV-REV plasmid			Leiden, The Netherlands)
Penicillin-Streptomycin (10,000 U/mL)	ThermoFisher Scientific	15140122	
PLV-mCherry plasmid	Addgene	36084	
Pneumatic picoPump	World Precision Instruments	SYS-PV820	
Polybrene	SigmaAldrich	107689	
Prism 4 software	GraphPad Software		
Stereo microscope	Leica	MZ16FA	
Tg (<i>fli:EGFP</i>) zebrafish strain			Kindly provided by Dr. Ewa Snaar-Jagalska (Institute of Biology, Leiden University, Leiden, The Netherlands)
Tris-base	SigmaAldrich	1181427300 1	
Tricaine (3-aminobenzoic acid)	SigmaAldrich	A-5040	
Trypsin-EDTA (0.5%)	ThermoFisher Scientific	15400054	
Petri dishes, polystyrene (60 × 15 mm)	SigmaAldrich	P5481- 500EA	
Polystyrene dish with glass bottom	WillCo	GWST-5040	