



Universiteit  
Leiden  
The Netherlands

**Putting a spin on it: amyloid aggregation from oligomers to fibrils**  
Zurlo, E.

**Citation**

Zurlo, E. (2020, July 9). *Putting a spin on it: amyloid aggregation from oligomers to fibrils*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/123273>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/123273>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/123273> holds various files of this Leiden University dissertation.

**Author:** Zurlo, E.

**Title:** Putting a spin on it: amyloid aggregation from oligomers to fibrils

**Issue Date:** 2020-07-09

# Bibliography

1. Brocchieri, L. & Karlin, S. Protein length in eukaryotic and prokaryotic proteomes. *Nucleic Acids Res.* **33**, 3390–3400 (2005).
2. Petsko, G. A. & Ringe, D. *Protein structure and function*. (New Science Press, 2004).
3. Richardson, J. S. The Anatomy and Taxonomy of Protein Structure. in (eds. Anfinsen, C. B., Edsall, J. T. & Richards, F. M. B. T.-A. in P. C.) **34**, 167–339 (Academic Press, 1981).
4. Dunker, A. K. *et al.* Intrinsically disordered protein. *J. Mol. Graph. Model.* **19**, 26–59 (2001).
5. Dyson, H. J. & Wright, P. E. Intrinsically unstructured proteins and their functions. *Nat. Rev. Mol. cell Biol.* **6**, 197–208 (2005).
6. Kumar, S., Mohanty, S. K. & Udgaonkar, J. B. Mechanism of formation of amyloid protofibrils of barstar from soluble oligomers: evidence for multiple steps and lateral association coupled to conformational conversion. *J. Mol. Biol.* **367**, 1186–1204 (2007).
7. Tompa, P. Intrinsically disordered proteins: a 10-year recap. *Trends Biochem. Sci.* **37**, 509–516 (2012).
8. Wright, P. E. & Dyson, H. J. Intrinsically disordered proteins in cellular signalling and regulation. *Nat. Rev. Mol. cell Biol.* **16**, 18–29 (2015).
9. Fändrich, M. & Dobson, C. M. The behaviour of polyamino acids reveals an inverse side chain effect in amyloid structure formation. *EMBO J.* **21**, 5682–5690 (2002).
10. Greenwald, J. & Riek, R. Biology of amyloid: structure, function, and regulation. *Structure* **18**, 1244–1260 (2010).
11. Rose, A. S. *et al.* NGL viewer: web-based molecular graphics for large complexes. *Bioinformatics* **34**, 3755–3758 (2018).
12. Kirschner, D. A., Abraham, C. & Selkoe, D. J. X-ray diffraction from intraneuronal paired helical filaments and extraneuronal amyloid fibers in Alzheimer disease indicates cross-beta conformation. *Proc. Natl. Acad. Sci.* **83**, 503–507 (1986).
13. Liberta, F. *et al.* Cryo-EM fibril structures from systemic AA amyloidosis reveal the species complementarity of pathological amyloids. *Nat. Commun.* **10**, 1–10 (2019).
14. Xue, W.-F. *et al.* Fibril fragmentation enhances amyloid cytotoxicity. *J. Biol. Chem.* **284**, 34272–34282 (2009).
15. Kodali, R. & Wetzel, R. Polymorphism in the intermediates and products of amyloid assembly. *Curr. Opin. Struct. Biol.* **17**, 48–57 (2007).
16. Caughey, B. & Lansbury Jr, P. T. Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu. Rev. Neurosci.* **26**, 267–298 (2003).
17. Kirkitadze, M. D., Bitan, G. & Teplow, D. B. Paradigm shifts in Alzheimer's disease and other neurodegenerative disorders: the emerging role of oligomeric assemblies. *J. Neurosci. Res.* **69**, 567–577 (2002).
18. Haass, C. & Selkoe, D. J. Soluble protein oligomers in neurodegeneration: Lessons from the

- Alzheimer's amyloid  $\beta$ -peptide. *Nat. Rev. Mol. Cell Biol.* **8**, 101–112 (2007).
19. van Diggelen, F., Tepper, A. W. J. W., Apetri, M. M. & Otzen, D. E.  $\alpha$ -Synuclein Oligomers: A Study in Diversity. *Isr. J. Chem.* **57**, 699–723 (2017).
  20. Kaye, R. *et al.* Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science*. **300**, 486–489 (2003).
  21. Portelius, E., Mattsson, N., Andreasson, U., Blennow, K. & Zetterberg, H. Novel A $\beta$  isoforms in Alzheimer's Disease-Their Role in Diagnosis and Treatment. *Curr. Pharm. Des.* **17**, 2594–2602 (2011).
  22. Jakob-Roetne, R. & Jacobsen, H. Alzheimer's disease: from pathology to therapeutic approaches. *Angew. Chemie Int. Ed.* **48**, 3030–3059 (2009).
  23. Chopra, K., Misra, S. & Kuhad, A. Neurobiological aspects of Alzheimer's disease. *Expert Opin. Ther. Targets* **15**, 535–555 (2011).
  24. Lanctôt, K. L. *et al.* Neuropsychiatric signs and symptoms of Alzheimer's disease: New treatment paradigms. *Alzheimer's Dement. Transl. Res. Clin. Interv.* **3**, 440–449 (2017).
  25. Braak, H. *et al.* Pattern of brain destruction in Parkinson's and Alzheimer's diseases. *J. Neural Transm.* **103**, 455–490 (1996).
  26. Selkoe, D. J. Alzheimer's disease: genes, proteins, and therapy. *Physiol. Rev.* **81**, 741–766 (2001).
  27. Gralle, M. & Ferreira, S. T. Structure and functions of the human amyloid precursor protein: the whole is more than the sum of its parts. *Prog. Neurobiol.* **82**, 11–32 (2007).
  28. Serra-Batiste, M. *et al.* A $\beta$ 42 assembles into specific  $\beta$ -barrel pore-forming oligomers in membrane-mimicking environments. *Proc. Natl. Acad. Sci.* **113**, 10866 LP – 10871 (2016).
  29. Cline, E. N., Bicca, M. A., Viola, K. L. & Klein, W. L. The amyloid- $\beta$  oligomer hypothesis: Beginning of the third decade. *J. Alzheimer's Dis.* **64**, S567–S610 (2018).
  30. Hayden, E. Y. & Teplow, D. B. Amyloid  $\beta$ -protein oligomers and Alzheimer's disease. *Alzheimers. Res. Ther.* **5**, 60 (2013).
  31. Lee, S. J. C., Nam, E., Lee, H. J., Savellieff, M. G. & Lim, M. H. Towards an understanding of amyloid- $\beta$  oligomers: characterization, toxicity mechanisms, and inhibitors. *Chem. Soc. Rev.* **46**, 310–323 (2017).
  32. Parkinson, J. *An Essay on the Shaking Palsy* (London: Sherwood, Neely and Jones). (1817).
  33. Cookson, M. R.  $\alpha$ -Synuclein and neuronal cell death. *Mol. Neurodegener.* **4**, 9 (2009).
  34. De Rijk, M. C. *et al.* Prevalence of Parkinson's disease in the elderly: the Rotterdam Study. *Neurology* **45**, 2143–2146 (1995).
  35. Ozansoy, M. & Başak, A. N. The central theme of Parkinson's disease:  $\alpha$ -synuclein. *Mol. Neurobiol.* **47**, 460–465 (2013).
  36. Venda, L. L., Cragg, S. J., Buchman, V. L. & Wade-Martins, R.  $\alpha$ -Synuclein and dopamine at the crossroads of Parkinson's disease. *Trends Neurosci.* **33**, 559–568 (2010).
  37. Sun, J. *et al.* Functional cooperation of  $\alpha$ -synuclein and VAMP2 in synaptic vesicle recycling. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 11113–11115 (2019).
  38. Chen, S. W. *et al.* Structural characterization of toxic oligomers that are kinetically trapped

- during  $\alpha$ -synuclein fibril formation. *Proc. Natl. Acad. Sci.* **112**, E1994 LP-E2003 (2015).
39. Cremades, N. *et al.* Direct observation of the interconversion of normal and toxic forms of  $\alpha$ -synuclein. *Cell* **149**, 1048–1059 (2012).
  40. Merrifield, R. B. Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. *J. Am. Chem. Soc.* **85**, 2149–2154 (1963).
  41. Atherton, N. M. *Principles of electron spin resonance*. (Ellis Horwood Limited, 1993).
  42. Roessler, M. M. & Salvadori, E. Principles and applications of EPR spectroscopy in the chemical sciences. *Chem. Soc. Rev.* **47**, 2534–2553 (2018).
  43. Brustolon, M. R. *Principles and Applications of Electron Paramagnetic Resonance Spectroscopy*. (John Wiley & Sons, 2008).
  44. Briere, R., Lemaire, H. & Rassat, A. NITROXYDES 15-SYNTHESE ET ETUDE DE RADICAUX LIBRES STABLES PIPERIDINIQUES ET PYRROLIDINIQUE. *Bull. Soc. Chim. Fr.* 3273 (1965).
  45. Hubbell, W. L. & Altenbach, C. Investigation of structure and dynamics in membrane proteins using site-directed spin labeling. *Curr. Opin. Struct. Biol.* **4**, 566–573 (1994).
  46. Hubbell, W. L., Mchaourab, H. S., Altenbach, C. & Lietzow, M. A. Watching proteins move using site-directed spin labeling. *Structure* **4**, 779–783 (1996).
  47. Hubbell, W. L., Cafiso, D. S. & Altenbach, C. Identifying conformational changes with site-directed spin labeling. *Nat. Struct. Biol.* **7**, 735–739 (2000).
  48. Stoll, S. & Schweiger, A. EasySpin, a comprehensive software package for spectral simulation and analysis in EPR. *J. Magn. Reson.* **178**, 42–55 (2006).
  49. Milov, A. D., Ponomarev, A. B. & Tsvetkov, Y. D. Electron-electron double resonance in electron spin echo: Model biradical systems and the sensitized photolysis of decalin. *Chem. Phys. Lett.* **110**, 67–72 (1984).
  50. Schiemann, O. & Prisner, T. F. Long-range distance determinations in biomacromolecules by EPR spectroscopy. *Q. Rev. Biophys.* **40**, 1 (2007).
  51. Bowman, M. K., Maryasov, A. G., Kim, N. & DeRose, V. J. Visualization of distance distribution from pulsed double electron-electron resonance data. *Appl. Magn. Reson.* **26**, 23–39 (2004).
  52. Chiang, Y. W., Borbat, P. P. & Freed, J. H. The determination of pair distance distributions by pulsed ESR using Tikhonov regularization. *J. Magn. Reson.* **172**, 279–295 (2005).
  53. Jeschke, G., Panek, G., Godt, A., Bender, A. & Paulsen, H. Data analysis procedures for pulse ELDOR measurements of broad distance distributions. *Appl. Magn. Reson.* **26**, 223 (2004).
  54. Jeschke, G. *et al.* DeerAnalysis2006—a comprehensive software package for analyzing pulsed ELDOR data. *Appl. Magn. Reson.* **30**, 473–498 (2006).
  55. Jeschke, G. & Polyhach, Y. Distance measurements on spin-labelled biomacromolecules by pulsed electron paramagnetic resonance. *Phys. Chem. Chem. Phys.* **9**, 1895–910 (2007).
  56. Jeschke, G. *et al.* DeerAnalysis2006 - A comprehensive software package for analyzing pulsed ELDOR data. *Appl. Magn. Reson.* **30**, 473–498 (2006).
  57. Mchaourab, H. S., Lietzow, M. A., Hideg, K. & Hubbell, W. L. Motion of spin-labeled side chains in T4 lysozyme. Correlation with protein structure and dynamics. *Biochemistry* **35**, 7692–7704 (1996).

58. Jeschke, G. DEER Distance Measurements on Proteins. *Annu. Rev. Phys. Chem.* **63**, 419–446 (2012).
59. Fajer, P. G. Site directed spin labelling and pulsed dipolar electron paramagnetic resonance (double electron–electron resonance) of force activation in muscle. *J. Phys. Condens. Matter* **17**, S1459–S1469 (2005).
60. Shabestari, M. H., Wolfs, C. J. A. M., Spruijt, R. B., Van Amerongen, H. & Huber, M. Exploring the structure of the 100 amino-acid residue long N-terminus of the plant antenna protein CP29. *Biophys. J.* **106**, 1349–1358 (2014).
61. Drescher, M., Huber, M. & Subramaniam, V. Hunting the Chameleon: Structural Conformations of the Intrinsically Disordered Protein Alpha-Synuclein. *ChemBioChem* **13**, 761–768 (2012).
62. Petrlova, J. *et al.* A differential association of Apolipoprotein E isoforms with the amyloid- $\beta$  oligomer in solution. *Proteins* **79**, 402–16 (2011).
63. Grimaldi, M. *et al.* Membrane charge dependent states of the  $\beta$ -amyloid fragment A $\beta$  (16–35) with differently charged micelle aggregates. *Biochim. Biophys. Acta (BBA)-Biomembranes* **1798**, 660–671 (2010).
64. Shabestari, M. *et al.* The Aggregation Potential of the 1-15- and 1-16-Fragments of the Amyloid  $\beta$  Peptide and Their Influence on the Aggregation of A $\beta$ 40. *Appl. Magn. Reson.* **44**, 1167–1179 (2013).
65. Shabestari, M. H., Meeuwenoord, N. J., Filippov, D. V. & Huber, M. Interaction of the amyloid  $\beta$  peptide with sodium dodecyl sulfate as a membrane-mimicking detergent. *J. Biol. Phys.* **42**, 299–315 (2016).
66. Hashemi Shabestari, M. *et al.* Three Long-Range Distance Constraints and an Approach Towards a Model for the  $\alpha$ -Synuclein-Fibril Fold. *Appl. Magn. Reson.* **46**, 369–388 (2015).
67. Laganowsky, A. *et al.* Atomic view of a toxic amyloid small oligomer. *Science* **335**, 1228–31 (2012).
68. Georgieva, E. R., Borbat, P. P., Ginter, C., Freed, J. H. & Boudker, O. Conformational ensemble of the sodium-coupled aspartate transporter. *Nat. Struct. Mol. Biol.* **20**, 215–221 (2013).
69. Dunkel, S., Pulagam, L. P., Steinhoff, H.-J. & Klare, J. P. *In vivo* EPR on spin labeled colicin A reveals an oligomeric assembly of the pore-forming domain in *E. coli* membranes. *Phys. Chem. Chem. Phys.* **17**, 4875–4878 (2015).
70. Warshaviak, D. T., Khramtsov, V. V., Cascio, D., Altenbach, C. & Hubbell, W. L. Structure and dynamics of an imidazoline nitroxide side chain with strongly hindered internal motion in proteins. *J. Magn. Reson.* **232**, 53–61 (2013).
71. Stoller, S., Sicoli, G., Baranova, T. Y., Bennati, M. & Diederichsen, U. TOPP: A novel nitroxide-labeled amino acid for EPR distance measurements. *Angew. Chemie - Int. Ed.* **50**, 9743–9746 (2011).
72. Jeschke, G. Conformational dynamics and distribution of nitroxide spin labels. *Prog. Nucl. Magn. Reson. Spectrosc.* **72**, 42–60 (2013).
73. Fawzi, N. L. *et al.* A rigid disulfide-linked nitroxide side chain simplifies the quantitative analysis of PRE data. *J. Biomol. NMR* **51**, 105–114 (2011).
74. Schilder, J. *et al.* Protein docking using an ensemble of spin labels optimized by intra-

- molecular paramagnetic relaxation enhancement. *Phys. Chem. Chem. Phys.* **18**, (2015).
75. Di Valentin, M., Albertini, M., Zurlo, E., Gobbo, M. & Carbonera, D. Porphyrin triplet state as a potential Spin label for nanometer distance measurements by peldor spectroscopy. *J. Am. Chem. Soc.* **136**, 6582–6585 (2014).
  76. Milov, A. D. *et al.* The secondary structure of a membrane-modifying peptide in a supramolecular assembly studied by PELDOR and CW-ESR spectroscopies. *J. Am. Chem. Soc.* **123**, 3784–3789 (2001).
  77. McNulty, J. C. *et al.* Electron spin resonance of TOAC labeled peptides: Folding transitions and high frequency spectroscopy. *Biopolym. - Pept. Sci. Sect.* **55**, 479–485 (2000).
  78. Monaco, V. *et al.* Orientation and immersion depth of a helical lipopeptaibol in membranes using TOAC as an ESR probe. *Biopolymers* **50**, 239–253 (1999).
  79. Toniolo, C., Crisma, M. & Formaggio, F. TOAC, a nitroxide spin-labeled, achiral C $\alpha$ -tetrasubstituted  $\alpha$ -amino acid, is an excellent tool in material science and biochemistry. *Pept. Sci.* **47**, 153–158 (1998).
  80. Chen, K., Liu, Z., Zhou, C., Bracken, W. C. & Kallenbach, N. R. Spin relaxation enhancement confirms dominance of extended conformations in short alanine peptides. *Angew. Chemie - Int. Ed.* **46**, 9036–9039 (2007).
  81. Hansen, P. *et al.* Distinguishing Helix Conformations in Alanine Rich Peptides Using the Unnatural Amino Acid TOAC and Electron Spin Resonance. *J. Am. Chem. Soc.* **118**, 271–272 (1996).
  82. Monaco, V. *et al.* Determining the occurrence of a 310-helix and an  $\alpha$ -helix in two different segments of a lipopeptaibol antibiotic using TOAC, a nitroxide spin-labeled C( $\alpha$ )-tetrasubstituted  $\alpha$ -amino acid. *Bioorganic Med. Chem.* **7**, 119–131 (1999).
  83. Ghimire, H. *et al.* Distance measurements on a dual-labeled TOAC AChR M2 $\delta$  peptide in mechanically aligned DMPC bilayers via dipolar broadening CW-EPR spectroscopy. *J. Phys. Chem. B* **116**, 3866–3873 (2012).
  84. Marchetto, R., Schreier, S. & Nakaie, C. R. A novel spin-labeled amino acid derivative for use in peptide synthesis: (9-fluorenylmethyloxycarbonyl)-2,2,6,6-tetramethylpiperidine-N-oxyl-4-amino-4-carboxylic acid. *J. Am. Chem. Soc.* **115**, 11042–11043 (1993).
  85. Cilli, E. M., Marchetto, R., Schreier, S. & Nakaie, C. R. Correlation between the mobility of spin-labeled peptide chains and resin solvation: An approach to optimize the synthesis of aggregating sequences. *J. Org. Chem.* **64**, 9118–9123 (1999).
  86. Nakaie, C. R. *et al.* Synthesis and pharmacological properties of TOAC-labeled angiotensin and bradykinin analogs. *Peptides* **23**, 65–70 (2002).
  87. Di Valentin, M., Albertini, M., Zurlo, E., Gobbo, M. & Carbonera, D. Porphyrin triplet state as a potential Spin label for nanometer distance measurements by peldor spectroscopy. *J. Am. Chem. Soc.* **136**, 6582–6585 (2014).
  88. Di Valentin, M. *et al.* Light-Induced Porphyrin-Based Spectroscopic Ruler for Nanometer Distance Measurements. *Chem. - A Eur. J.* **22**, (2016).
  89. Milov, A. D. *et al.* Synthesis and conformational properties of a TOAC doubly spin-labeled analog of the medium-length, membrane active peptaibiotic ampullosporin as revealed by CD, fluorescence, and EPR spectroscopies. *Biopolym. - Pept. Sci. Sect.* **102**, 40–48 (2014).

90. Shabestari, M. H. *et al.* Conformation and EPR characterization of rigid, 310-helical peptides with TOAC spin labels: Models for short distances. *Biopolym. - Pept. Sci. Sect.* **102**, 244–251 (2014).
91. Hanson, P. *et al.* Electron spin resonance and structural analysis of water soluble, alanine-rich peptides incorporating TO AC. *Mol. Phys.* **95**, 957–966 (1998).
92. Polese, A. *et al.* First interchain peptide interaction detected by ESR in fully synthetic, template-assisted, two-helix bundles. *J. Am. Chem. Soc.* **121**, 11071–11078 (1999).
93. Hanson, P., Millhauser, G., Formaggio, F., Crisma, M. & Toniolo, C. ESR characterization of hexameric, helical peptides using double TOAC spin labeling. *J. Am. Chem. Soc.* **118**, 7618–7625 (1996).
94. Greenfield, N. & Fasman, G. D. Computed Circular Dichroism Spectra for the Evaluation of Protein Conformation. *Biochemistry* **8**, 4108–4116 (1969).
95. Fasman, G. D. *Circular dichroism and the conformational analysis of biomolecules.* (Springer Science & Business Media, 2013).
96. Chiti, F. & Dobson, C. M. Protein misfolding, functional amyloid, and human disease. *Annu. Rev. Biochem.* **75**, 333–366 (2006).
97. Michaels, T. C. T. *et al.* Chemical Kinetics for Bridging Molecular Mechanisms and Macroscopic Measurements of Amyloid Fibril Formation. *Annu. Rev. Phys. Chem.* **69**, annurev-physchem-050317-021322 (2018).
98. Chiti, F. & Dobson, C. M. Amyloid formation by globular proteins under native conditions. *Nat Chem Biol* **5**, 15–22 (2009).
99. Bordignon E., Steinhoff HJ. Membrane Protein Structure and Dynamics Studied by Site-Directed Spin-Labeling ESR. In: ESR Spectroscopy in Membrane Biophysics. Biological Magnetic Resonance, vol **27**. Springer, Boston, MA(2007)
100. Mchaourab, H. S., Lietzow, M. A., Hideg, K. & Hubbell, W. L. Motion of spin-labeled side chains in T4 lysozyme. Correlation with protein structure and dynamics. *Biochemistry* **35**, 7692–7704 (1996).
101. Hubbell, W. L., Mchaourab, H. S., Altenbach, C. & Lietzow, M. A. Watching proteins move using site-directed spin labeling. *Structure* **4**, 779–783 (1996).
102. Bartucci, R. *et al.* Backbone dynamics of alamethicin bound to lipid membranes: Spin-echo electron paramagnetic resonance of TOAC-spin labels. *Biophys. J.* **94**, 2698–2705 (2008).
103. Hanson, P. *et al.* Electron spin resonance and structural analysis of water soluble, alanine-rich peptides incorporating TO AC. *Mol. Phys.* **95**, 957–966 (1998).
104. Schreier, S. *et al.* Conformational basis for the biological activity of TOAC-labeled angiotensin II and bradykinin: Electron paramagnetic resonance, circular dichroism, and fluorescence studies. *Biopolymers* **74**, 389–402 (2004).
105. Schreier, S., Bozelli, J. C., Marín, N., Vieira, R. F. F. & Nakaie, C. R. The spin label amino acid TOAC and its uses in studies of peptides: Chemical, physicochemical, spectroscopic, and conformational aspects. *Biophys. Rev.* **4**, 45–66 (2012).
106. Latajka, R. *et al.* Pentapeptides containing two dehydrophenylalanine residues--synthesis, structural studies and evaluation of their activity towards cathepsin C. *J. Pept. Sci.* **14**, 1084–1095 (2008).



107. Naiki, H., Higuchi, K., Hosokawa, M. & Takeda, T. Fluorometric determination of amyloid fibrils in vitro using the fluorescent dye, thioflavine T. *Anal. Biochem.* **177**, 244–249 (1989).
108. Hashemi Shabestari, M. *et al.* Conformation and EPR characterization of rigid, 310-helical peptides with TOAC spin labels: Models for short distances. *Pept. Sci.* **102**, 244–251 (2014).
109. Murphy, L. R., Matubayasi, N., Payne, V. A. & Levy, R. M. Protein hydration and unfolding – insights from experimental partial specific volumes and unfolded protein models. *Fold. Des.* **3**, 105–118 (2005).
110. Fischer, H., Polikarpov, I. & Craievich, A. F. Average protein density is a molecular-weight-dependent function. *Protein Sci.* **13**, 2825–2828 (2004).
111. Micsonai, A. *et al.* BeStSel: a web server for accurate protein secondary structure prediction and fold recognition from the circular dichroism spectra. *Nucleic Acids Res.* **46**, W315–W322 (2018).
112. Micsonai, A. *et al.* Accurate secondary structure prediction and fold recognition for circular dichroism spectroscopy. *Proc. Natl. Acad. Sci.* **112**, E3095–E3103 (2015).
113. Biancalana, M. & Koide, S. Molecular mechanism of Thioflavin-T binding to amyloid fibrils. *Biochim. Biophys. Acta (BBA)-Proteins Proteomics* **1804**, 1405–1412 (2010).
114. Milov, A. D., Ponomarev, A. B. & Tsvetkov, Y. D. Electron-electron double resonance in electron spin echo: model biradical systems and the sensitized photolysis of decalin. *Chem. Phys. Lett.* **110**, 67–72 (1984).
115. Bode, B. E. *et al.* Counting the monomers in nanometer-sized oligomers by pulsed electron-electron double resonance. *J. Am. Chem. Soc.* **129**, 6736–6745 (2007).
116. Gu, L. *et al.* Antiparallel triple-strand architecture for prefibrillar A $\beta$ 42 oligomers. *J. Biol. Chem.* **289**, 27300–13 (2014).
117. Cohen, S. I. A. *et al.* Proliferation of amyloid-42 aggregates occurs through a secondary nucleation mechanism. *Proc. Natl. Acad. Sci.* **110**, 9758–9763 (2013).
118. Bleiholder, C., Dupuis, N. F., Wyttenbach, T. & Bowers, M. T. Ion mobility–mass spectrometry reveals a conformational conversion from random assembly to  $\beta$ -sheet in amyloid fibril formation. *Nat. Chem.* **3**, 172 (2011).
119. Seo, J. *et al.* An infrared spectroscopy approach to follow  $\beta$ -sheet formation in peptide amyloid assemblies. *Nat. Chem.* **9**, 39 (2017).
120. Alexander, N., Al-Mestarihi, A., Bortolus, M., Mchaourab, H. & Meiler, J. De novo high-resolution protein structure determination from sparse spin-labeling EPR data. *Structure* **16**, 181–195 (2008).
121. Jao, C. C., Hegde, B. G., Chen, J., Haworth, I. S. & Langen, R. Structure of membrane-bound  $\alpha$ -synuclein from site-directed spin labeling and computational refinement. *Proc. Natl. Acad. Sci.* **105**, 19666–19671 (2008).
122. Meirovitch, E., Liang, Z. & Freed, J. H. Protein dynamics in the solid-state from <sup>2</sup>H NMR lineshape analysis. III. MOMD in the presence of Magic Angle Spinning. *Solid State Nucl. Magn. Reson.* **89**, 35–44 (2018).
123. Meirovitch, E., Liang, Z. & Freed, J. H. Protein Dynamics in the Solid State from <sup>2</sup>H NMR Line Shape Analysis. II. MOMD Applied to C-D and C-CD<sub>3</sub> Probes. *J. Phys. Chem. B* **119**, 14022–14032 (2015).

124. Budil, D. E., Lee, S., Saxena, S. & Freed, J. H. Nonlinear-least-squares analysis of slow-motion EPR spectra in one and two dimensions using a modified Levenberg–Marquardt algorithm. *J. Magn. Reson. Ser. A* **120**, 155–189 (1996).
125. García de la Torre, J., Huertas, M. . & Carrasco, B. HYDRONMR: Prediction of NMR Relaxation of Globular Proteins from Atomic-Level Structures and Hydrodynamic Calculations. *J. Magn. Reson.* **147**, 138–146 (2000).
126. Perrin, F. Mouvement Brownien d'un ellipsoïde (II). Rotation libre et dépolarisation des fluorescences. Translation et diffusion de molécules ellipsoïdales. *J. Phys. le Radium* **7**, 1–11 (1936).
127. Koenig, S. H. Brownian motion of an ellipsoid. A correction to Perrin's results. *Biopolymers* **14**, 2421–2423 (1975).
128. Marsh, D. & Horváth L.I. *Advanced EPR: applications in biology and biochemistry*. (Elsevier, 2012).
129. López, C. J., Fleissner, M. R., Guo, Z., Kusnetzow, A. K. & Hubbell, W. L. Osmolyte perturbation reveals conformational equilibria in spin-labeled proteins. *Protein Sci.* **18**, 1637–1652 (2009).
130. Columbus, L., Kálai, T., Jekő, J., Hideg, K. & Hubbell, W. L. Molecular motion of spin labeled side chains in  $\alpha$ -helices: analysis by variation of side chain structure. *Biochemistry* **40**, 3828–3846 (2001).
131. Zhang, Z. *et al.* Multifrequency electron spin resonance study of the dynamics of spin labeled T4 lysozyme. *J. Phys. Chem. B* **114**, 5503–5521 (2010).
132. Langen, R., Oh, K. J., Cascio, D. & Hubbell, W. L. Crystal structures of spin labeled T4 lysozyme mutants: implications for the interpretation of EPR spectra in terms of structure. *Biochemistry* **39**, 8396–8405 (2000).
133. Fleissner, M. R., Cascio, D. & Hubbell, W. L. Structural origin of weakly ordered nitroxide motion in spin-labeled proteins. *Protein Sci.* **18**, 893–908 (2009).
134. Clayton, D. F. & George, J. M. The synucleins: a family of proteins involved in synaptic function, plasticity, neurodegeneration and disease. *Trends Neurosci.* **21**, 249–254 (1998).
135. Burré, J. *et al.*  $\alpha$ -Synuclein promotes SNARE-complex assembly in vivo and in vitro. *Science*. **329**, 1663–1667 (2010).
136. Nemani, V. M. *et al.* Increased Expression of  $\alpha$ -Synuclein Reduces Neurotransmitter Release by Inhibiting Synaptic Vesicle Reclustering after Endocytosis. *Neuron* **65**, 66–79 (2010).
137. Villar-Piqué, A., Lopes da Fonseca, T. & Outeiro, T. F. Structure, function and toxicity of alpha-synuclein: the Bermuda triangle in synucleinopathies. *J. Neurochem.* **139**, 240–255 (2016).
138. Arosio, P., Knowles, T. P. J. & Linse, S. On the lag phase in amyloid fibril formation. *Phys. Chem. Chem. Phys.* **17**, 7606–7618 (2015).
139. Ke, P. C. *et al.* Implications of peptide assemblies in amyloid diseases. *Chem. Soc. Rev.* **46**, 6492–6531 (2017).
140. van Diggelen, F., Tepper, A. W. J. W., Apetri, M. M. & Otzen, D. E.  $\alpha$ -Synuclein Oligomers: A Study in Diversity. *Isr. J. Chem.* **57**, 699–723 (2017).
141. Iyer, A. & Claessens, M. M. A. E. Disruptive membrane interactions of alpha-synuclein aggregates. *Biochim. Biophys. Acta - Proteins Proteomics* **1867**, 468–482 (2019).

142. Ilie, I. M. & Caflisch, A. Simulation studies of amyloidogenic polypeptides and their aggregates. *Chem. Rev.* **119**, 6956–6993 (2019).
143. Afitska, K., Fucikova, A., Shvadchak, V. V & Yushchenko, D. A.  $\alpha$ -Synuclein aggregation at low concentrations. *Biochim. Biophys. Acta (BBA)-Proteins Proteomics* **1867**, 701–709 (2019).
144. Cascella, R. *et al.* Probing the Origin of the Toxicity of Oligomeric Aggregates of  $\alpha$ -Synuclein with Antibodies. *ACS Chem. Biol.* **14**, 1352–1362 (2019).
145. Pérez-Pi, I. *et al.*  $\alpha$ -Synuclein-Confocal Nanoscanning (ASYN-CONA), a Bead-Based Assay for Detecting Early-Stage  $\alpha$ -Synuclein Aggregation. *Anal. Chem.* **91**, 5582–5590 (2019).
146. Xu, M., Loa-Kum-Cheung, W., Zhang, H., Quinn, R. J. & Mellick, G. D. Identification of a New  $\alpha$ -Synuclein Aggregation Inhibitor via Mass Spectrometry Based Screening. *ACS Chem. Neurosci.* **10**, acschemneuro.9b00092 (2019).
147. van Diggelen, F. *et al.* Two conformationally distinct  $\alpha$ -synuclein oligomers share common epitopes and the ability to impair long-term potentiation. *PLoS One* **14**, e0213663 (2019).
148. van Raaij, M. E., Segers-Nolten, I. M. J. & Subramaniam, V. Quantitative morphological analysis reveals ultrastructural diversity of amyloid fibrils from  $\alpha$ -synuclein mutants. *Biophys. J.* **91**, L96–L98 (2006).
149. Veldhuis, G., Segers-Nolten, I., Ferlemann, E. & Subramaniam, V. Single-Molecule FRET Reveals Structural Heterogeneity of SDS-Bound  $\alpha$ -Synuclein. *ChemBioChem* **10**, 436–439 (2009).
150. Jao, C. C., Der-Sarkissian, A., Chen, J. & Langen, R. Structure of membrane-bound  $\alpha$ -synuclein studied by site-directed spin labeling. *Proc. Natl. Acad. Sci.* **101**, 8331–8336 (2004).
151. Steigmiller, S., Börsch, M., Gräber, P. & Huber, M. Distances between the b-subunits in the tether domain of F<sub>0</sub>F<sub>1</sub>-ATP synthase from *E. coli*. *Biochim. Biophys. Acta - Bioenerg.* **1708**, 143–153 (2005).
152. Iljina, M. *et al.* Quantifying co-oligomer formation by  $\alpha$ -synuclein. *ACS Nano* **12**, 10855–10866 (2018).
153. Iljina, M. *et al.* Kinetic model of the aggregation of alpha-synuclein provides insights into prion-like spreading. *Proc. Natl. Acad. Sci.* **113**, E1206–E1215 (2016).
154. Meisl, G. *et al.* Molecular mechanisms of protein aggregation from global fitting of kinetic models. *Nat. Protoc.* **11**, 252 (2016).
155. Gaspar, R. *et al.* Secondary nucleation of monomers on fibril surface dominates  $\alpha$ -synuclein aggregation and provides autocatalytic amyloid amplification. *Q. Rev. Biophys.* **50**, (2017).
156. Cohen, S. I. A. *et al.* Proliferation of amyloid- $\beta$ 42 aggregates occurs through a secondary nucleation mechanism. *Proc. Natl. Acad. Sci.* **110**, 9758–9763 (2013).
157. Drescher, M. *et al.* Spin-label EPR on  $\alpha$ -synuclein reveals differences in the membrane binding affinity of the two antiparallel helices. *ChemBioChem* **9**, 2411–2416 (2008).
158. Reyes, J. F. *et al.* Binding of  $\alpha$ -synuclein oligomers to Cx32 facilitates protein uptake and transfer in neurons and oligodendrocytes. *Acta Neuropathol.* **138**, 23–47 (2019).
159. Quist, A. *et al.* PNAS-2005-Quist-10427-32. **102**, 1–6 (2005).
160. Glabe, C. G. Structural classification of toxic amyloid oligomers. *J. Biol. Chem.* **283**, 29639–29643 (2008).

161. Buell, A. K. *et al.* Solution conditions determine the relative importance of nucleation and growth processes in  $\alpha$ -synuclein aggregation. *Proc. Natl. Acad. Sci.* **111**, 7671–7676 (2014).
162. Pannier, M., Veit, S., Godt, A., Jeschke, G. & Spiess, H. W. Dead-time free measurement of dipole–dipole interactions between electron spins. *J. Magn. Reson.* **142**, 331–340 (2000).
163. Schiemann, O. & Prisner, T. F. Long-range distance determinations in biomacromolecules by EPR spectroscopy. **1**, 1–53 (2020).
164. Drescher, M. EPR in protein science. in *EPR Spectroscopy* 91–119 (Springer, 2011).
165. Tsvetkov, Y. D., Milov, A. D. & Maryasov, A. G. Pulsed electron–electron double resonance (PELDOR) as EPR spectroscopy in nanometre range. *Russ. Chem. Rev.* **77**, 487 (2008).
166. Schmidt, T., Wälti, M. A., Baber, J. L., Hustedt, E. J. & Clore, G. M. Long Distance Measurements up to 160 Å in the GroEL Tetradecamer Using Q-Band DEER EPR Spectroscopy. *Angew. Chemie Int. Ed.* **55**, 15905–15909 (2016).
167. Qi, M., Groß, A., Jeschke, G., Godt, A. & Drescher, M. Gd (III)-PyMTA label is suitable for in-cell EPR. *J. Am. Chem. Soc.* **136**, 15366–15378 (2014).
168. Bellapadrona, G. *et al.* Probing Protein Conformation in Cells by EPR Distance Measurements using Gd 3+ Spin Labeling. *J. Am. Chem. Soc.* **136**, 13458–13465 (2014).
169. Igarashi, R. *et al.* Distance Determination in Proteins inside *Xenopus laevis* Oocytes by Double Electron–Electron Resonance Experiments. *J. Am. Chem. Soc.* **132**, 8228–8229 (2010).
170. Krstić, I. *et al.* Long-range distance measurements on nucleic acids in cells by pulsed EPR spectroscopy. *Angew. Chemie Int. Ed.* **50**, 5070–5074 (2011).
171. Collauto, A. *et al.* Gd ( III ) complexes as paramagnetic tags : Evaluation of the spin delocalization over the nuclei of the ligand. **263**, 156–163 (2016).
172. Garbuio, L., Zimmermann, K., Häussinger, D. & Yulikov, M. Gd ( III ) complexes for electron – electron dipolar spectroscopy : Effects of deuteration , pH and zero field splitting. *J. Magn. Reson.* **259**, 163–173 (2015).
173. Yang, Y. *et al.* A Reactive, Rigid GdIII Labeling Tag for In-Cell EPR Distance Measurements in Proteins. *Angew. Chemie Int. Ed.* **56**, 2914–2918 (2017).
174. Keizers, P. H. J., Desreux, J. F., Overhand, M. & Ubbink, M. Increased paramagnetic effect of a lanthanide protein probe by two-point attachment. *J. Am. Chem. Soc.* **129**, 9292–9293 (2007).
175. Fleissner, M. R. *et al.* Structure and dynamics of a conformationally constrained nitroxide side chain and applications in EPR spectroscopy. *Proc. Natl. Acad. Sci.* **108**, 16241–16246 (2011).
176. Liu, W.-M. *et al.* A Two-Armed Lanthanoid-Chelating Paramagnetic NMR Probe Linked to Proteins via Thioether Linkages. *Chem. - A Eur. J.* **20**, 6256–6258 (2014).
177. Georgieva, E. R. *et al.* Effect of freezing conditions on distances and their distributions derived from Double Electron Electron Resonance (DEER): A study of doubly-spin-labeled T4 lysozyme. *J. Magn. Reson.* **216**, 69–77 (2012).
178. Pannier, M., Veit, S., Godt, A., Jeschke, G. & Spiess, H. . Dead-Time Free Measurement of Dipole–Dipole Interactions between Electron Spins. *J. Magn. Reson.* **142**, 331–340 (2000).
179. Clayton, J. A. *et al.* Gd3+-Gd3+ distances exceeding 3 nm determined by very high frequency continuous wave electron paramagnetic resonance. *Phys. Chem. Chem. Phys.* **19**, 5127–5136 (2017).

180. Matalon, E. *et al.* Gadolinium(III) Spin Labels for High-Sensitivity Distance Measurements in Transmembrane Helices. *Angew. Chemie Int. Ed.* **52**, 11831–11834 (2013).
181. Feintuch, A., Otting, G. & Goldfarb, D. Gd<sup>3+</sup> + Spin Labeling for Measuring Distances in Biomacromolecules: Why and How? *Methods Enzymol.* **563**, 415–457 (2015).
182. Liu, L., Quillin, M. L. & Matthews, B. W. Use of experimental crystallographic phases to examine the hydration of polar and nonpolar cavities in T4 lysozyme. *Proc. Natl. Acad. Sci.* **105**, 14406 – 14411 (2008).
183. Yagi, H. *et al.* Gadolinium tagging for high-precision measurements of 6 nm distances in protein assemblies by EPR. *J. Am. Chem. Soc.* **133**, 10418–10421 (2011).
184. Neuhaus, E. M., Almers, W. & Soldati, T. Morphology and dynamics of the endocytic pathway in *Dictyostelium discoideum*. *Mol. Biol. Cell* **13**, 1390–1407 (2002).
185. Lu, H. P. Revealing time bunching effect in single-molecule enzyme conformational dynamics. *Phys. Chem. Chem. Phys.* **13**, 6734–6749 (2011).
186. Yang, Y., Yang, F., Li, X.-Y., Su, X.-C. & Goldfarb, D. In-Cell EPR Distance Measurements on Ubiquitin Labeled with a Rigid PyMTA-Gd(III) Tag. *J. Phys. Chem. B* **123**, 1050–1059 (2019).
187. Annesley, S. J. & Fisher, P. R. *Dictyostelium discoideum*—a model for many reasons. *Mol. Cell. Biochem.* **329**, 73–91 (2009).
188. Richman, M. *et al.* In vitro and mechanistic studies of an antiamyloidogenic self-assembled cyclic d, l- $\alpha$ -peptide architecture. *J. Am. Chem. Soc.* **135**, 3474–3484 (2013).
189. Chemerovski-glikman, M., Richman, M. & Rahimipour, S. Structure-based study of antiamyloidogenic cyclic D, L -  $\alpha$ -peptides. *Tetrahedron* **70**, 7639–7644 (2014).
190. Barnes, D. E. & Yaffe, K. The projected effect of risk factor reduction on Alzheimer’s disease prevalence. *Lancet Neurol.* **10**, 819–828 (2011).
191. Roßner, S., Sastre, M., Bourne, K. & Lichtenthaler, S. F. Transcriptional and translational regulation of BACE1 expression-Implications for Alzheimer’s disease. *Prog. Neurobiol.* **79**, 95–111 (2006).
192. Bucciantini, M. *et al.* Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature* **416**, 507–511 (2002).
193. Kelly, J. W. The alternative conformations of amyloidogenic proteins and their multi-step assembly pathways. *Curr. Opin. Struct. Biol.* **8**, 101–106 (1998).
194. Dobson, C. M., Swoboda, B. E. P., Joniau, M. & Weissman, C. The structural basis of protein folding and its links with human disease. *Philos. Trans. R. Soc. B Biol. Sci.* **356**, 133–145 (2001).
195. Chiti, F. & Dobson, C. M. Protein Misfolding, Functional Amyloid, and Human Disease. *Annu. Rev. Biochem.* **75**, 333–366 (2006).
196. Chiti, F. & Dobson, C. M. Protein Misfolding, Amyloid Formation, and Human Disease: A Summary of Progress Over the Last Decade. *Annu. Rev. Biochem.* **86**, 27–68 (2017).
197. Zurlo, E., Gorroño Bikandi, I., Meeuwenoord, N. J., Filippov, D. V. & Huber, M. Tracking amyloid oligomerization with monomer resolution using a 13-amino acid peptide with a backbone-fixed spin label. *Phys. Chem. Chem. Phys.* **21**, 25187–25195 (2019).
198. Ghalebani, L., Wahlström, A., Danielsson, J., Wärmländer, S. K. T. S. & Gräslund, A. pH-dependence of the specific binding of Cu(II) and Zn(II) ions to the amyloid- $\beta$  peptide.

- Biochem. Biophys. Res. Commun.* **421**, 554–560 (2012).
199. Vassar, P. S. & Culling, C. F. Fluorescent stains, with special reference to amyloid and connective tissues. *Arch. Pathol.* **68**, 487–498 (1959).
  200. Di Valentin, M., Albertini, M., Zurlo, E., Gobbo, M. & Carbonera, D. Porphyrin triplet state as a potential Spin label for nanometer distance measurements by peldor spectroscopy. *J. Am. Chem. Soc.* **136**, (2014).
  201. Sepkhanova, I. *et al.* Monitoring Alzheimer amyloid peptide aggregation by EPR. *Appl. Magn. Reson.* **36**, 209 (2009).
  202. Chen, M., Margittai, M., Chen, J. & Langen, R. Investigation of  $\alpha$ -synuclein fibril structure by site-directed spin labeling. *J. Biol. Chem.* **282**, 24970–24979 (2007).
  203. Margittai, M. & Langen, R. Template-assisted filament growth by parallel stacking of tau. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 10278–10283 (2004).
  204. Lebel, R. G. & Goring, D. A. I. Density, Viscosity, Refractive Index, and Hygroscopicity of Mixtures of Water and Dimethyl Sulfoxide. *J. Chem. Eng. Data* **7**, 100–101 (1962).