



Universiteit
Leiden
The Netherlands

Clinical outcomes in bariatric surgery

Poelemeijer, Y.Q.M.

Citation

Poelemeijer, Y. Q. M. (2020, November 25). *Clinical outcomes in bariatric surgery*. Retrieved from <https://hdl.handle.net/1887/138403>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/138403>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/138403> holds various files of this Leiden University dissertation.

Author: Poelmeijer, Y.Q.M.

Title: Clinical outcomes in bariatric surgery

Issue date: 2020-11-25



Measuring Quality of Life in Bariatric Surgery: A Multicentre Study

Youri Q.M. Poelemeijer^{1,2}, Elise T.W. van der Knaap¹, Perla J. Marang-van de Mheen³, Ahmet Demirkiran⁴, Marinus J. Wiezer⁵, Eric J. Hazebroek⁶, Jan Willem M. Greve⁷, Ronald S.L. Liem^{8,9}

¹Dutch Institute for Clinical Auditing, Scientific Bureau, Leiden, Netherlands

²Leiden University Medical Center, Department of Surgery, Leiden, Netherlands

³Leiden University Medical Center, Department of Biomedical Data Science, Leiden, Netherlands

⁴Rode Kruis Hospital, Department of Surgery, Beverwijk, Netherlands

⁵St. Antonius Hospital, Department of Surgery, Nieuwegein, Netherlands

⁶Rijnstate, Department of Surgery, Arnhem, Netherlands

⁷Zuyderland Medical Center, Department of Surgery, Heerlen, Netherlands

⁸Groene Hart Hospital, Department of Surgery, Gouda, Netherlands

⁹Dutch Obesity Clinic, The Hague, Netherlands

Surg Endosc. 2020 Jan 28. [Epub ahead of print]

ABSTRACT

BACKGROUND

Current studies mainly focus on total weight loss and comorbidity reduction. Only a few studies compare Quality of Life (QoL) after sleeve gastrectomy (SG) and Roux-en-Y gastric bypass (RYGB). This study was conducted to examine the extent of improvement in QoL on different domains after primary bariatric surgery and compare these results to Dutch reference values.

METHODS

The study included prospectively collected data from patients who underwent primary bariatric surgery in five Dutch hospitals. The RAND-36 questionnaire was used to measure the patient's QoL; preoperatively and twelve months postoperatively. Postoperative scores were compared to Dutch reference values, standardized for age, using t-test. A difference of more than 5% was considered a minimal important difference. A multivariate linear regression analysis was used to compare SG and RYGB on the extent of improvement, adjusted for case-mix factors.

RESULTS

In total, 4864 patients completed both the pre- and postoperative questionnaire. Compared with Dutch reference values, patients postoperatively reported clinically relevant better *physical functioning* (RYGB + 6.8%), *physical role limitations* (SG + 5.6%; RYGB + 6.2%) and *health change* (SG + 77.1%; RYGB + 80.0%), but worse *general health perception* (SG - 22.8%; RYGB - 17.0%). Improvement in QoL was similar between SG and RYGB, except for *physical functioning* (β 2.758; p-value 0.008) and *general health perception* (β 2.607; p-value < 0.001) for which RYGB patients improved more.

CONCLUSIONS

SG and RYGB patients achieved a better postoperative score in *physical functioning*, *physical role limitations*, and *health change* compared to Dutch reference values, and a worse score in *general health perception*.

INTRODUCTION

International literature provides evidence that bariatric surgery can contribute to substantial weight loss and a positive effect on obesity-related comorbidities.^{1, 2} On the other hand, bariatric surgery may also lead to severe postoperative complications, as well as endocrine and metabolic complications.^{1, 3-5} Psychological consequences of bariatric surgery were also described as complex and not always entirely understood.⁶⁻⁹ Therefore, bariatric surgery requires a detailed evaluation of its impact on a patient's level and is best assessed with a Quality of Life (QoL) assessment.¹⁰

Standard clinical outcomes, such as weight loss and resolution of comorbidities, are mostly objectively measured in registries, resulting in quantitative data, which are convenient for analyses. QoL assessments, however, are primarily patient-reported measurements and may be more challenging to interpret. On the other hand, it is essential to include QoL assessments in the evaluation of health interventions of bariatric surgery as the patient perspective can provide valuable information on the effectiveness of bariatric surgery that cannot be obtained from clinical outcome measures alone.¹⁰⁻¹² QoL could be measured by using questionnaires reflecting the patient's perspective on the effects of the provided healthcare or treatment given to the patient in their daily lives.^{10, 11} Literature showed that the Short Form 36-item Health Survey® (SF-36) is the most commonly used QoL measurement in bariatric surgery.¹⁰ A nearly identical questionnaire is the RAND 36-item Health Survey (RAND-36) that evaluates the same domains as the SF-36. The difference between these two questionnaires mainly consists of the commercial fees required for using the questionnaire.^{13, 14} The RAND-36 is also the standard QoL measuring tool offered to patients in all Dutch bariatric hospitals.

Recent studies mainly focussed on clinical outcomes such as total weight loss and obesity-related comorbidity reduction.^{15, 16} The few initial studies which compared

QoL after sleeve gastrectomy (SG) and Roux-en-Y gastric bypass (RYGB) did not use the RAND-36 and included a low sample size (range 50–1703) with only two studies of more than 1000 patients.^{17, 18} Another pitfall in previously conducted studies is the low volume of postoperative respondents as well as single-centre studies, making the results on improvement after bariatric surgery less reliable and also not generalizable to other settings in daily practice.¹⁹ Furthermore, a recent study comparing RYGB and SG in three different European countries showed differences in preoperative characteristics, which may have been the reason for a different surgical approach but could also affect the outcomes including postoperative QoL after bariatric surgery.²⁰ We therefore com-

pared changes between these procedures, not only looking at statistical significance but also considering clinically relevant differences.

The aim of this study is to compare improvement in QoL after primary bariatric surgery for the two mainly performed primary bariatric procedures in the Netherlands: SG and RYGB. In addition, the study compares postoperative values with reference values for the general Dutch population. A multicentre study design is chosen for a better representation across multiple sites.

MATERIALS AND METHODS

QoL-data were prospectively collected from all patients undergoing a primary RYGB or SG in the five participating hospitals in the Netherlands between 1 January 2015 and 1 January 2017. QoL-data were linked to the national bariatric DATO-registry covering all centres providing bariatric surgery.²¹

The scientific committee, which coordinates the national DATO-registry, represents all participating bariatric centres and all members are mandated by the practising hospital where they practice. This committee approved the research proposal for the present study and manuscript for publication. A more in depth description about the scientific committee is given in an earlier scientific article.²¹

PATIENTS

In the Netherlands, patients with a body mass index (BMI) ≥ 40.0 kg/m² or with a BMI ≥ 35.0 kg/m² and one or more obesity associated comorbidities were eligible for bariatric surgery during the study period.^{21,22} These obesity associated comorbidities were type-2 diabetes mellitus (T2DM), hypertension (HT), dyslipidaemia, gastroesophageal reflux disease (GERD), obstructive sleep apnoea syndrome (OSAS) and musculoskeletal pain. Further treatment strategies, including the choice for SG or RYGB, were determined by a multidisciplinary team and by shared decision making with the patient.

Baseline characteristics in patients undergoing SG or RYGB were compared using the mean \pm standard deviation (SD) for normally distributed variables and the median with interquartile range for non-normally distributed variables. The Mann-Whitney *U* test was performed for continuous variables and χ^2 for categorical variables. The threshold for significance has been set at 0.05.

QUALITY OF LIFE (QOL)

During the development process of the nationwide DATO registry, several questionnaires were considered including the Bariatric Analysis and Reporting Outcome System (BAROS)^{23,24}, SF-36 and RAND-36 questionnaires. Given the controversy surrounding BAROS a generic QoL questionnaire was preferred. The RAND-36 and SF-36 are identical, except for different scoring algorithms for the *pain* and *general health perception* scales, resulting in the choice for the RAND-36 questionnaire.^{13,14,25,26}

RAND-36

The Dutch version of the RAND-36 is a validated and standardized translation of the original RAND-36 questionnaire.^{13,14} The questionnaire contains 36 questions within nine scales. These scales are *physical functioning*, *social functioning*, *physical role limitations*, *emotional role limitations*, *mental health*, *vitality*, *pain*, *general health perception* and *health change perception*. Previous studies have shown this to be a valid tool for the measurement of QoL among obese patients undergoing bariatric surgery.^{25,27,28}

Each patient undergoing bariatric surgery in one of the five participating bariatric centres was included in the study. The preoperative questionnaire was completed during the initial screening for bariatric surgery. The postoperative questionnaire was administered 12 months (range 9–15) after primary surgery. The questionnaires were part of the standard given care in the five participating centres.

ANALYSING THE QUESTIONNAIRE

All completed questionnaires were analysed by a predefined algorithm, provided by the RAND-36 research group and included in the original article.^{13,14} A brief summary is given below.

All scores were recoded following the provided algorithm: a high score indicates a more favourable health state (or outcome) of the patient.^{14,26} Each item was scored on a 0 to 100 range. An average of all scores in each of the nine individual scales has been calculated. Missing values were replaced with the personal mean of the specific scale if at least half of the answers on the questions of the scale were provided.^{13,14,26}

COMPARING TO THE DUTCH REFERENCE POPULATION

First, postoperative RAND-36 scores were divided into six age groups (18–24, 25–34, 35–44, 45–54, 55–64 and 65+). Second, the extent of improvement after surgery was calculated by subtracting preoperative RAND-36 scores from the postoperative RAND-36 scores, providing the delta separately for SG and RYGB. Third, the postoperative

RAND-36 scores were compared with the Dutch reference values¹³, in order to see if patients experience the same QoL postoperatively as the Dutch reference group.

Finally, for a valid comparison, the age distribution of the Dutch reference population¹³ was applied to the age-specific QoL-values of the SG or RYGB patients to prevent overall values being different because of a difference in age distribution. The age-standardized QoL scores for each of the nine scales were compared for SG and RYGB patients with the Dutch population using the *t*-test.

POSTOPERATIVE INFLUENCES ON THE QOL OUTCOMES

The Clavien–Dindo classification (CD) is used to determine whether a patient had experienced a severe postoperative complication.^{29, 30} All patients with a CD grade ≥ 3 , within 30 days after primary surgery, were registered as severe. Due to the low number of severe complications, both operative techniques have been combined and the *t*-test compared the severe complicated group with the uncomplicated group.

A distinction has also been made to see whether the achievement of Total Weight Loss (TWL) influences the QoL outcomes.^{21, 31} All patients were subdivided into patients who reached 20% TWL within 12 months postoperatively and patients who did not. There were no patients in this cohort missing preoperative or postoperative weight scores. Both groups are compared using the *t*-test.

COMPARISON BETWEEN SG AND RYGB

In order to compare between SG and RYB, we compared the extent of improvement between SG and RYGB patients adjusted for patient variables that differed at baseline using multivariate linear regression analysis reporting the β estimate and *p*-value.

Analyses were performed using R version 3.5.1 and the R-packages “Companion to Applied Regression”-package (car 3.0-2), “A Grammar of Data Manipulation”-package (dplyr 0.7.8) and “Table 1”-package (tableone 0.9.3)’ were used.

RESULTS

A total of 5574 unique patients underwent a primary SG or RYGB. Patients who were operated and who did not complete both questionnaires ($n = 710$) were excluded. A total of 4864 (87.3%) patients were eligible for analyses, having completed both a preoperative and postoperative questionnaire. Baseline characteristics were shown in **Table 1** and correspond to the national bariatric benchmark in the Netherlands.²¹

Table 1: Baseline characteristics showing preoperative measurements and prevalence of obesity-related comorbidities.

	Sleeve gastrectomy		Roux-en-Y gastric bypass		p-value
	N	%	N	%	
Number of patients	965	19.8%	3,899	80.2%	
Gender (female)	745	77.2%	3,152	80.8%	0.013*
Weight (median, kg, IQR)	133.9	(119.4 – 155.0)	123.9	(113.0 – 136.4)	<0.001*
BMI (median, kg/m ² , IQR)	45.8	(42.0 – 52.7)	43.0	(40.4 – 46.7)	<0.001*
Waist circumference (median, cm, IQR)	133	(123 – 145)	127	(120 – 136)	<0.001*
Age (mean, years, SD)	39.3	± 12.5	45.6	± 10.3	<0.001*
Type 2 diabetes mellitus	170	17.6%	1,160	29.9%	<0.001*
Hypertension	297	30.8%	1,593	40.9%	<0.001*
Dyslipidaemia	128	13.3%	902	23.1%	<0.001*
GERD	49	5.1%	233	6.0%	0.317
OSAS	200	20.7%	812	20.8%	0.965
Musculoskeletal pain	430	44.6%	2,294	58.8%	<0.001*

Abbreviations: N, number; SD, standard deviation; IQR, interquartile range; GERD, gastroesophageal reflux disease; OSAS, obstructive sleep apnoea syndrome. * statistically significant difference measured.

Patients with a SG were significantly heavier ($p < 0.001$), reflected in both higher BMI and higher waist circumference ($p < 0.001$). Statistically significant differences in obesity-related diseases were seen for T2DM, hypertension, dyslipidaemia and musculoskeletal pain. All these obesity-related diseases were seen significantly more often in the RYGB group ($p < 0.001$) compared to SG.

SLEEVE GASTRECTOMY

For patients undergoing SG, the best postoperative QoL scores were seen in relatively young bariatric patients (**Table 2a**). However, the older bariatric patients showed a larger positive delta score and therefore showed a bigger improvement compared to the younger and middle-aged patients (**Table 2b**). For example, the youngest patients scored postoperatively better on *physical functioning* and *physical role limitations*. On the other hand, the delta scores for these domains are slightly lower for the younger patients compared to the oldest group (**Table 2a, b**).

ROUX-EN-Y GASTRIC BYPASS

For RYGB patients, especially the three youngest categories showed better postoperative scores in almost all RAND-36 domains except for the domain *vitality and general health perception* (**Table 3a**). However, the largest improvement (delta) was seen in the oldest group (65+). This applies to all domains except *physical role limitations*,

Table 2a & Table 2b: Postoperative unadjusted scale scores & pre- and postoperative change in scale scores (delta) per age group after **sleeve gastrectomy**, measured with RAND-36. Abbreviations: SD, standard deviation.

Age (n)	18-24 (n=135)		25-34 (n=257)		35-44 (n=211)		45-54 (n=226)		55-64 (n=124)		65+ (n=12)		Overall (n=965)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Physical functioning	91.26	± 13.90	90.42	± 18.33	88.39	± 19.81	83.03	± 20.76	75.27	± 25.79	58.33	± 37.77	85.92	± 20.92
Social functioning	84.68	± 21.38	86.04	± 21.47	87.30	± 19.74	84.03	± 21.48	82.00	± 23.28	77.00	± 34.12	85.03	± 21.46
Physical role limitations	88.79	± 26.07	87.57	± 27.46	85.97	± 27.93	81.65	± 32.00	79.30	± 33.30	70.83	± 45.87	84.63	± 29.69
Emotional role limitations	85.37	± 31.27	86.35	± 30.38	90.28	± 26.63	85.49	± 31.24	83.04	± 31.02	83.33	± 40.82	86.42	± 30.08
Mental health	76.37	± 17.76	78.21	± 17.53	81.29	± 15.41	79.08	± 18.03	80.99	± 15.50	72.67	± 31.94	79.21	± 17.15
Vitality	61.44	± 20.74	62.91	± 18.73	67.90	± 17.58	67.25	± 19.69	68.60	± 18.44	61.67	± 32.20	65.68	± 19.20
Pain	82.68	± 24.76	84.39	± 21.85	80.48	± 26.79	75.79	± 26.33	73.16	± 27.44	61.00	± 39.18	79.44	± 25.72
General health perception	55.68	± 15.15	56.80	± 15.96	56.44	± 15.90	55.42	± 16.03	51.53	± 18.00	51.33	± 24.45	55.48	± 16.26
Health change	93.68	± 16.27	93.44	± 14.83	92.90	± 17.74	92.49	± 19.10	93.82	± 13.11	95.83	± 10.21	93.18	± 16.56
Age (n)	18-24 (n=135)		25-34 (n=257)		35-44 (n=211)		45-54 (n=226)		55-64 (n=124)		65+ (n=12)		Overall (n=965)	
Physical functioning	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD
	28.51	± 19.94	30.50	± 23.68	30.81	± 26.11	33.61	± 24.77	33.98	± 23.26	19.17	± 42.12	31.46	± 24.24
Social functioning	8.56	± 26.69	17.37	± 29.25	15.64	± 26.54	14.57	± 26.20	14.12	± 25.78	29.33	± 55.09	14.85	± 27.48
Physical role limitations	19.54	± 41.84	28.07	± 43.76	29.03	± 42.97	32.66	± 45.62	27.42	± 46.63	29.17	± 62.08	28.28	± 44.38
Emotional role limitations	0.78	± 36.75	7.85	± 40.84	6.45	± 36.14	10.60	± 38.88	4.67	± 42.51	17.00	± 18.62	6.99	± 38.94
Mental health	3.17	± 17.48	10.50	± 20.20	7.77	± 17.68	7.68	± 17.74	6.49	± 16.02	2.00	± 28.48	7.65	± 18.32
Vitality	8.16	± 19.96	10.61	± 21.36	14.74	± 21.50	18.79	± 20.83	19.09	± 22.44	0.00	± 28.28	14.31	± 21.64
Pain	13.03	± 25.40	18.66	± 26.66	18.05	± 29.37	17.42	± 26.64	20.12	± 26.03	26.33	± 42.07	17.77	± 27.17
General health perception	18.94	± 16.61	21.54	± 17.02	19.38	± 17.73	19.54	± 17.42	18.41	± 17.09	15.33	± 14.40	19.76	± 17.20
Health change	54.02	± 27.72	55.03	± 29.44	51.94	± 32.30	53.61	± 32.98	56.72	± 30.87	70.83	± 36.80	54.22	± 31.02

Table 3a & Table 3b: Postoperative unadjusted scale scores & pre- and postoperative change in scale scores (delta) per age group after Roux-en-Y gastric bypass, measured with RAND-36. Abbreviations: SD, standard deviation.

Age (n)	18-24 (n=84)		25-34 (n=537)		35-44 (n=1,040)		45-54 (n=1,427)		55-64 (n=741)		65+ (n=70)		Overall (n=3,899)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Physical functioning	94.42	± 10.54	92.90	± 13.67	91.94	± 15.58	88.19	± 18.80	82.17	± 22.11	84.44	± 20.23	88.75	± 18.35
Social functioning	88.42	± 16.77	83.89	± 22.03	87.05	± 21.24	85.07	± 22.61	84.12	± 23.07	89.06	± 20.19	85.40	± 22.14
Physical role limitations	90.42	± 24.83	86.17	± 29.91	87.23	± 28.79	84.45	± 31.37	82.22	± 31.89	87.50	± 29.44	85.18	± 30.49
Emotional role limitations	88.82	± 27.28	83.14	± 33.50	90.07	± 26.37	86.61	± 31.17	87.60	± 30.28	94.43	± 20.28	87.44	± 29.96
Mental health	81.93	± 12.64	78.02	± 17.83	81.41	± 15.96	80.37	± 17.64	80.36	± 17.16	85.26	± 15.90	80.45	± 17.06
Vitality	62.50	± 20.14	63.20	± 19.89	67.19	± 19.74	67.49	± 20.30	68.81	± 19.77	74.54	± 18.31	67.11	± 20.04
Pain	87.42	± 20.75	82.60	± 23.44	82.76	± 23.22	79.84	± 24.59	76.44	± 25.89	78.56	± 28.79	80.48	± 24.45
General health perception	57.47	± 11.38	57.71	± 15.01	60.42	± 14.81	60.14	± 14.84	59.44	± 14.79	62.81	± 15.59	59.75	± 14.82
Health change	95.42	± 13.41	94.17	± 16.92	95.37	± 15.17	94.61	± 16.13	94.47	± 16.59	95.83	± 15.16	94.77	± 16.01
Age (n)	18-24 (n=84)		25-34 (n=537)		35-44 (n=1,040)		45-54 (n=1,427)		55-64 (n=741)		65+ (n=70)		Overall (n=3,899)	
Physical functioning	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD	Delta	SD
	24.17	± 20.34	30.66	± 22.87	31.75	± 23.31	34.54	± 23.14	36.76	± 24.36	41.85	± 24.73	33.61	± 23.50
Social functioning	8.45	± 24.03	13.03	± 26.80	14.20	± 28.57	13.99	± 27.51	15.88	± 28.40	16.13	± 29.31	14.20	± 27.84
Physical role limitations	16.25	± 39.82	23.57	± 43.65	27.49	± 44.97	29.19	± 45.80	33.11	± 47.96	35.19	± 47.71	28.57	± 45.73
Emotional role limitations	4.47	± 43.26	-0.97	± 39.34	6.24	± 36.86	3.60	± 40.44	12.83	± 45.57	9.93	± 34.10	5.56	± 40.55
Mental health	7.07	± 13.77	7.54	± 18.36	8.44	± 17.88	6.84	± 18.46	7.99	± 18.36	7.26	± 20.46	7.59	± 18.23
Vitality	3.75	± 22.71	12.47	± 22.74	16.68	± 23.02	16.98	± 22.39	18.38	± 22.89	18.89	± 22.12	16.32	± 22.82
Pain	11.58	± 28.36	17.95	± 31.18	18.92	± 27.69	20.99	± 28.07	20.98	± 27.36	21.57	± 33.49	19.85	± 28.42
General health perception	15.47	± 18.15	21.40	± 18.11	22.53	± 17.34	23.53	± 17.23	22.89	± 16.30	23.93	± 18.13	22.70	± 17.28
Health change	42.50	± 30.64	51.49	± 30.40	55.08	± 29.28	57.62	± 28.81	58.16	± 29.34	57.87	± 24.68	55.91	± 29.34

emotional role limitations and *health change*. For these domains, the second-oldest group (55–64) showed the largest improvement (**Table 3b**).

OVERALL QUALITY OF LIFE

The comparison of the Dutch reference values with both the postoperative SG and RYGB values were standardized to the age distribution of the Dutch reference group.¹³ As suggested in recent scientific literature, a statistical significant QoL-score difference of > 5% is considered a minimal important difference (MID).^{32, 33}

Results showed a MID in the domains *physical functioning* (for RYGB), *physical role limitations* and *health change* (for both SG and RYGB) compared to Dutch reference values. Especially, for the domain *health change*, a large difference was observed (**Table 4a**). However, patients postoperatively still report lower scores on the domain *general health perception*. In addition, the SG scores were slightly lower than the RYGB in all RAND- 36 domains (**Table 4a**).

COMPLICATIONS AND TOTAL WEIGHT LOSS

Hypothetically, a postoperatively complicated course has a negative influence on the postoperative QoL outcomes. To make reliable calculations, both operative techniques were combined and the postoperative complicated group was compared with the uncomplicated group. **Table 4b** shows that the positive effects in the domains *physical functioning* and *physical role limitations* have disappeared (shown in **Table 4a**). A MID was seen in the domain of *social functioning*, *physical role limitations*, *emotional role limitations*, *vitality* and *pain*. Patients postoperatively still report lower scores on the domain *general health perception*, while they still report a significant *health change* (**Table 4b**).

SG VS. RYGB

Comparing the extent of improvement between SG and RYGB patients on each domain, a significant difference was seen in the domains *physical functioning* and *general health perception* when adjusted for differences in baseline characteristics; T2DM, hypertension, dyslipidaemia and musculoskeletal pain. These significant differences were mostly seen in the RYGB group (**Table 5**).

DISCUSSION

Current studies on bariatric surgery particularly focus on weight loss and improvement of obesity-related diseases but do not sufficiently take the patient's perspective into

Table 4a & Table 4b: Table 4a represents the postoperative adjusted baseline scale scores for sleeve gastrectomy and Roux-en-Y gastric bypass. Table 4b represents the postoperative scale scores for patients with a postoperative severe complication (Clavien-Dindo grade ≥ 3) and scale scores for patients with a successful postoperative total weight loss (>20 %TWL) after 1 year. All outcomes are measured with RAND-36. Percentage difference (%) and p-value are comparisons to the Dutch benchmark.

	Sleeve gastrectomy			Roux-en-Y gastric bypass			Dutch reference group			
	Mean	SD	%	p-value	Mean	SD	%	p-value	Mean	SD
	n = 965			n = 3,899						
Physical functioning	86.02	± 19.66	+3.5	<0.001	88.76	± 17.69	+6.8*	<0.001*	83.09	± 20.39
Social functioning	85.02	± 21.32	-2.7	0.002	85.40	± 22.05	-2.2	<0.001	87.35	± 19.74
Physical role limitations	84.73	± 29.21	+5.6*	<0.001*	85.19	± 30.36	+6.2*	<0.001*	80.24	± 34.84
Emotional role limitations	86.41	± 29.90	+1.9	0.128	87.43	± 29.72	+3.1	<0.001	84.79	± 31.46
Mental health	79.12	± 17.01	+2.8	<0.001	80.44	± 16.96	+4.5	<0.001	77.00	± 18.66
Vitality	65.53	± 18.98	-3.0	0.002	67.09	± 19.92	-0.7	0.241	67.59	± 19.82
Pain	79.55	± 25.15	-2.0	0.072	80.49	± 24.27	-0.8	0.217	81.14	± 24.37
General health perception	55.50	± 16.11	-22.8*	<0.001*	59.74	± 14.76	-17.0*	<0.001*	71.95	± 21.60
Health change	93.21	± 16.29	+77.1*	<0.001*	94.76	± 15.97	+80.0*	<0.001*	52.63	± 18.50
	Postoperative complication			Unsuccessful %TWL			Dutch reference group			
	n = 119			n = 696						
Physical functioning	80.58	± 27.29	-3.0	<0.001	82.42	± 21.50	-0.8	<0.001	83.09	± 20.39
Social functioning	77.90	± 30.93	-10.8*	0.003*	81.11	± 23.50	-7.4*	0.001*	87.35	± 19.74
Physical role limitations	75.64	± 39.88	-5.7*	0.005*	79.43	± 33.80	-1.0	0.002	80.24	± 34.84
Emotional role limitations	83.71	± 33.91	-1.27	0.292	79.16	± 36.18	-6.6*	<0.001*	84.79	± 31.46
Mental health	78.90	± 18.81	+2.5	0.500	76.38	± 17.57	+0.8	<0.001	77.00	± 18.66
Vitality	61.92	± 22.48	-8.4*	0.027*	63.47	± 19.98	-6.1*	0.004*	67.59	± 19.82
Pain	70.51	± 30.66	-13.1*	<0.001*	76.72	± 25.51	-5.4*	0.015*	81.14	± 24.37
General health perception	55.74	± 16.02	-22.5*	0.060*	53.46	± 16.37	-25.7*	<0.001*	71.95	± 21.60
Health change	88.46	± 23.73	+68.1*	0.001*	89.72	± 21.88	+70.5*	<0.001*	52.63	± 18.50

Abbreviations: SD, standard deviation; %, percentage. * statistically significant and clinically relevant difference measured. ** p-value measured with the postoperative uncomplicated and the successful %TWL group.

Table 5: Postoperative delta scale scores of sleeve gastrectomy and Roux-en-Y gastric bypass, measured with RAND-36. Beta's were estimated using linear regression and adjusted for T2DM, hypertension, dyslipidaemia and musculoskeletal pain. Compared with a multivariate logistic regression analysis.

	Sleeve gastrectomy		Roux-en-Y gastric bypass		Beta estimate	p-value
	n = 965		n = 3,899			
	Delta mean	SD	Delta mean	SD		
Physical functioning	31.46	± 24.24	33.61	± 23.50	2.758	0.008*
Social functioning	14.85	± 27.48	14.20	± 27.84	-0.821	0.509
Physical role limitations	28.28	± 44.38	28.57	± 45.73	-1.352	0.505
Emotional role limitations	6.99	± 38.94	5.56	± 40.55	-2.650	0.141
Mental health	7.65	± 18.32	7.59	± 18.23	0.483	0.544
Vitality	14.31	± 21.64	16.32	± 22.82	0.919	0.361
Pain	17.77	± 27.17	19.85	± 28.42	1.148	0.360
General health perception	19.76	± 17.20	22.70	± 17.28	2.607	<0.001*
Health change	54.22	± 31.02	55.91	± 29.34	-0.072	0.957

Abbreviations: SD, standard deviation; T2DM, type 2 diabetes mellitus. * statistically significant and clinically relevant difference measured.

account.^{21, 34-38} It is important to focus more on postoperative outcomes from a patient's perspective, because of the enormous increase in bariatric procedures worldwide.^{22, 39-41}

There were a few initial studies comparing QoL after SG and RYGB, but these studies had mostly a low sample size and did not use the RAND 36-item Health Survey (RAND-36).^{17, 18} In addition, these studies were almost all single-centre studies and most of them reported low postoperative response rates.¹⁹

However, there were two larger population-based studies comparing postoperative QoL after bariatric surgery. The first study from Waljee et al. had a larger sample size comparing to the previous noted studies, but does not distinguish between SG and RYGB and has a poor follow-up rate.^{42, 43} The second study from Sarwer et al. focused on the QoL and sexual functioning of patients with obesity and looked specific on the changes in these domains, but does not made a distinction between RYGB and SG either.⁴⁴ As a result, the question remained which postoperative differences in QoL could be measured between the two most commonly used surgery techniques and how these changes relate to the Dutch population. Therefore, the first multicentre study has been conducted comparing QoL between SG and RYGB with a large sample size and a postoperative response rate of more than 85%.

Results showed that bariatric patients had meaningful higher postoperative scores on *physical functioning*, *physical role limitations* and *health change* for both SG and RYGB compared to Dutch reference values, but meaningful lower scores on *general health perception*. It may be concluded that patients feel better postoperatively, but not yet fully healthy. These results could be a prelude to focus more on these domains, so that bariatric patients do not end up in social isolation and feeling healthier, similar to the national average.

Table 4b clearly showed that a postoperatively severe complicated course or failure to achieve the desired weight loss influences the QoL outcomes. Where first a meaningful positive postoperative score in *physical functioning* and *physical role limitations* was seen (**Table 4a**), a significant negative score was now seen in the severe complicated group and the unsuccessful %TWL group. In addition, a negative trend is also observed in almost all other domains. This argues for better psychological postoperative support for patients where the outcomes do not meet Textbook Outcome.⁴⁵

The two bariatric surgical techniques showed a similar QoL improvement in all domains except for *physical functioning* and *general health perception* for which RYGB patients showed a higher postoperative improvement. This difference could be explained by the underlying indication for treatment. Particularly for patients with a high BMI (> 50 kg/m²) a SG is preferably, so a second stage procedure may follow.⁴⁶ The use of the SG for morbid obese patient stems from the use of this procedure as a modification to the duodenal switch. Later on, it was used as a first part of a two-stage gastric bypass procedure on morbid obese patients. In the beginning of this century multiple studies have been published with a laparoscopic sleeve gastrectomy as an isolated bariatric procedure, with promising results.^{47, 48} During the time when this study was conducted, the RYGB was often used as a second-stage procedure in Dutch bariatric hospitals. In recent years and increase in the use of the "one anastomosis gastric bypass" (OAGB) and the "single anastomosis duodeno-ileal bypass with sleeve gastrectomy" (SADI-s) was seen. In addition, minor modifications have been applied to the existing RYGB. This makes the RYGB more successful in patients with a higher BMI. This has led to a decrease in the number of SG procedures nowadays.

Patient with a SG could experience a worse health perception compared to patients with a single RYGB operation. Despite the fact that BMI is added in the case-mix model, also the weight loss in both groups can be experienced differently.

Another significant difference was seen in the preoperative registered obesity-related diseases in the RYGB-group. Several studies suggest that the RYGB has a greater

beneficial effect on obesity-related diseases after surgery compared to SG.⁴⁹⁻⁵² This also could have an effect on the surgeon's choice for the type of bariatric surgery and therefore the differences in experienced QoL.

But the proper interpretation of these results remains a point of discussion. As has been shown for other type of surgeries and diseases, the RAND-36 is a generic questionnaire and may not be specific enough to fully analyse the QoL in bariatric patients.^{53, 54} For example, when looking at *physical functioning*, the score was calculated on the basis of ten questions. These questions relate to typical activities during the day and may be one of the key items for patients with obesity, but can only be answered with a limited number of options; limited a lot, limited a little, or not limited at all which is likely to capture only the very severe physical limitations e.g. induced by severe obesity. More subtle differences may not be measured adequately, while this is essential for obese and bariatric patients. Using a bariatric-specific quality of life questionnaire may detect more clinically relevant differences. However, as already mentioned in the introduction, there were currently no suitable bariatric questionnaires that could be applied in current scientific research.

Another point of debate is calculating and reporting the MID. Not only the baseline scores may vary by population and context, but also the differences experienced and noticed by the patient may vary. Therefore the proper interpretation of these results remains a point of discussion.⁵⁵ This means, for example, that an increase of 10 points should be interpreted differently if the baseline values differ. And also, an individual rise from 10 to 20 on a 100-point scale can be interpreted differently than a rise from 60 to 70.

As mentioned earlier in the discussion, other studies have not shown differences between SG and RYGB in QoL, while different outcomes following these two operations can be hypothesised.^{17, 18} For example, the indication for the type of bariatric surgery is mainly based on the choice and expertise of the specific surgeon. There were some studies that suggest that the RYGB has a more beneficial effect on metabolic obesity-related diseases.⁴⁶ We tried to correct this by adding these baseline differences (**Table 1**) in the case-mix model.

There are some limitations, despite the accuracy of this study. A possible response bias could be generated by excluding patients from the study without postoperative measurements. However, this study offers a high response rate, whereby it can be assumed that the influence will be very small. In addition, this study didn't focus on statistically significant differences alone but also described whether these differences

were clinically relevant (minimal important difference; MID) which at the same time will also safeguard against finding chance differences. This study does not have specific data to make a comparison between patients who participate in a multidisciplinary postoperative coaching program and patients who did not. These coaching programs could consist of participating in support groups, appointments with a dietician and psychological follow-up by trained professionals.

There is still no consensus, whether to correct for multiple testing or not. Current statements say, an adjustment is particularly required in confirmatory analyses with multiple analyses stating one final conclusion.⁵⁶ This study, on the other hand, has an exploratory meaning with the aim of not missing potentially important findings by a standard adjustment of multiple testing.⁵⁷ Therefore, the results in this study were not corrected for multiple testing.

The strength of this study was the large sample size, high response rate and the prospective study design. Most other studies collecting patient-reported outcome data struggle to get a sufficiently high response rate as a part of daily routine clinical care. Therefore, the number of requested items has been kept to a minimum, but at the cost of having limited other data to e.g. adjust for patient characteristics. Furthermore, this was a multicentre study including hospitals located in different geographic areas, therefore a representative group from the population was obtained. A limitation of the current study was the availability of only one-year follow-up data. When all Dutch hospitals have implemented the PROMs registration, the follow-up will be extended to an annual follow-up up to five years after the primary surgery in the national DATO-registry. This will further substantiate the current outcomes of this study.

CONCLUSIONS

This study showed that bariatric patients achieve better postoperative *physical functioning*, *physical role limitations*, and *health change* for both SG and RYGB compared to Dutch reference values, but worse general health perception. In addition, a larger improvement in *general health perception* was seen in patients who underwent RYGB compared to SG. Further studies are needed to develop a specific QoL-questionnaire, which focuses on the different aspects of the bariatric patient and the different inclusion criteria for a specific procedure.

REFERENCES

1. Colquitt JL, Pickett K, Loveman E, Frampton GK. Surgery for weight loss in adults. *Cochrane Database Syst Rev.* 2014(8):CD003641.
2. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA.* 2013;309(1):71-82.
3. Puzziferri N, Roshek TB, 3rd, Mayo HG, Gallagher R, Belle SH, Livingston EH. Long-term follow-up after bariatric surgery: a systematic review. *JAMA.* 2014;312(9):934-42.
4. Ramanan B, Gupta PK, Gupta H, Fang X, Forse RA. Development and validation of a bariatric surgery mortality risk calculator. *J Am Coll Surg.* 2012;214(6):892-900.
5. Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrenbach K, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA.* 2004;292(14):1724-37.
6. Jarvholm K, Karlsson J, Olbers T, Peltonen M, Marcus C, Dahlgren J, et al. Characteristics of adolescents with poor mental health after bariatric surgery. *Surg Obes Relat Dis.* 2016;12(4):882-90.
7. Herget S, Rudolph A, Hilbert A, Bluher S. Psychosocial status and mental health in adolescents before and after bariatric surgery: a systematic literature review. *Obes Facts.* 2014;7(4):233-45.
8. Ortega J, Fernandez-Canet R, Alvarez-Valdeita S, Cassinello N, Baguena-Puigcerver MJ. Predictors of psychological symptoms in morbidly obese patients after gastric bypass surgery. *Surg Obes Relat Dis.* 2012;8(6):770-6.
9. Bocchieri LE, Meana M, Fisher BL. A review of psychosocial outcomes of surgery for morbid obesity. *J Psychosom Res.* 2002;52(3):155-65.
10. Coulman KD, Abdelrahman T, Owen-Smith A, Andrews RC, Welbourn R, Blazeby JM. Patient-reported outcomes in bariatric surgery: a systematic review of standards of reporting. *Obes Rev.* 2013;14(9):707-20.
11. Weldring T, Smith SM. Patient-Reported Outcomes (PROs) and Patient-Reported Outcome Measures (PROMs). *Health Serv Insights.* 2013;6:61-8.
12. Mazer LM, Azagury DE, Morton JM. Quality of Life After Bariatric Surgery. *Curr Obes Rep.* 2017;6(2):204-10.
13. Van der Zee KI, Sanderman R. Handleiding RAND-36 (2e druk): University Medical Center Groningen; 2012 [Available from: https://www.umcg.nl/SiteCollectionDocuments/research/institutes/SHARE/assessment%20tools/handleiding_rand36_2e_druk.pdf].
14. Hays RD, Morales LS. The RAND-36 measure of health-related quality of life. *Ann Med.* 2001;33(5):350-7.
15. Parmar C, Abdelhalim MA, Mahawar KK, Boyle M, Carr WRJ, Jennings N, et al. Management of super-super obese patients: comparison between one anastomosis (mini) gastric bypass and Roux-en-Y gastric bypass. *Surg Endosc.* 2017;31(9):3504-9.
16. Madhok B, Mahawar KK, Boyle M, Carr WR, Jennings N, Schroeder N, et al. Management of Super-super Obese Patients: Comparison Between Mini (One Anastomosis) Gastric Bypass and Sleeve Gastrectomy. *Obes Surg.* 2016;26(7):1646-9.
17. Lindeskilde N, Gladstone BP, Lubeck M, Nielsen J, Clausen L, Vach W, et al. The impact of bariatric surgery on quality of life: a systematic review and meta-analysis. *Obes Rev.* 2015;16(8):639-51.
18. Andersen JR, Aasprang A, Karlsen TI, Natvig GK, Vage V, Kolotkin RL. Health-related quality of life after bariatric surgery: a systematic review of prospective long-term studies. *Surg Obes Relat Dis.* 2015;11(2):466-73.
19. Versteegden DPA, Van Himbeek MJJ, Nienhuijs SW. Improvement in quality of life after

- bariatric surgery: sleeve versus bypass. *Surg Obes Relat Dis.* 2018;14(2):170-4.
20. Poelemeijer YQM, Liem RSL, Vage V, Mala T, Sundbom M, Ottosson J, et al. Gastric Bypass Versus Sleeve Gastrectomy: Patient Selection and Short-term Outcome of 47,101 Primary Operations from the Swedish, Norwegian, and Dutch National Quality Registries. *Ann Surg.* 2019.
 21. Poelemeijer YQM, Liem RSL, Nienhuijs SW. A Dutch Nationwide Bariatric Quality Registry: DATO. *Obes Surg.* 2018;28(6):1602-10.
 22. Fried M, Yumuk V, Oppert JM, Scopinaro N, Torres A, Weiner R, et al. Interdisciplinary European guidelines on metabolic and bariatric surgery. *Obes Surg.* 2014;24(1):42-55.
 23. Oria HE, Moorehead MK. Bariatric analysis and reporting outcome system (BAROS). *Obes Surg.* 1998;8(5):487-99.
 24. Oria HE, Moorehead MK. Updated Bariatric Analysis and Reporting Outcome System (BAROS). *Surg Obes Relat Dis.* 2009;5(1):60-6.
 25. Cunningham WE, Nakazono TT, Tsai KL, Hays RD. Do differences in methods for constructing SF-36 physical and mental health summary measures change their associations with chronic medical conditions and utilization? *Qual Life Res.* 2003;12(8):1029-35.
 26. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. *Health Econ.* 1993;2(3):217-27.
 27. Al Amer R, Al Khalifa K, Alajlan SA, Al Ansari A. Analyzing the Psychometric Properties of the Short Form-36 Quality of Life Questionnaire in Patients with Obesity. *Obes Surg.* 2018;28(8):2521-7.
 28. van Nunen AM, Wouters EJ, Vingerhoets AJ, Hox JJ, Geenen R. The health-related quality of life of obese persons seeking or not seeking surgical or non-surgical treatment: a meta-analysis. *Obes Surg.* 2007;17(10):1357-66.
 29. Poelemeijer YQM, Liem RSL, Vage V, Mala T, Sundbom M, Ottosson J, et al. Peri-operative Outcomes of Primary Bariatric Surgery in North-Western Europe: a Pooled Multinational Registry Analysis. *Obes Surg.* 2018;28(12):3916-22.
 30. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240(2):205-13.
 31. van de Laar A, de Caluwe L, Dillemans B. Relative outcome measures for bariatric surgery. Evidence against excess weight loss and excess body mass index loss from a series of laparoscopic Roux-en-Y gastric bypass patients. *Obes Surg.* 2011;21(6):763-7.
 32. Ringash J, O'Sullivan B, Bezjak A, Redelmeier DA. Interpreting clinically significant changes in patient-reported outcomes. *Cancer.* 2007;110(1):196-202.
 33. Warkentin LM, Majumdar SR, Johnson JA, Agborsangaya CB, Rueda-Clausen CF, Sharma AM, et al. Weight loss required by the severely obese to achieve clinically important differences in health-related quality of life: two-year prospective cohort study. *BMC Med.* 2014;12:175.
 34. Kindel TL, Oleynikov D. The Improvement of Gastroesophageal Reflux Disease and Barrett's after Bariatric Surgery. *Obes Surg.* 2016;26(4):718-20.
 35. Bachler T, le Roux CW, Bueter M. How do patients' clinical phenotype and the physiological mechanisms of the operations impact the choice of bariatric procedure? *Clin Exp Gastroenterol.* 2016;9:181-9.
 36. Brethauer SA, Kim J, el Chaar M, Pappasavas P, Eisenberg D, Rogers A, et al. Standardized outcomes reporting in metabolic and bariatric surgery. *Surg Obes Relat Dis.* 2015;11(3):489-506.
 37. Mahawar KK. Outcome Measures in Bariatric Surgery. *Obes Surg.* 2015;25(11):2161.
 38. Dogan K, Kraaij L, Aarts EO, Koehestanie P, Hammink E, van Laarhoven CJ, et al. Fast-

- track bariatric surgery improves perioperative care and logistics compared to conventional care. *Obes Surg.* 2015;25(1):28-35.
39. Welbourn R, Pournaras DJ, Dixon J, Higa K, Kinsman R, Ottosson J, et al. Bariatric Surgery Worldwide: Baseline Demographic Description and One-Year Outcomes from the Second IFSO Global Registry Report 2013-2015. *Obes Surg.* 2018;28(2):313-22.
 40. Angrisani L, Santonicola A, Iovino P, Vitiello A, Zundel N, Buchwald H, et al. Bariatric Surgery and Endoluminal Procedures: IFSO Worldwide Survey 2014. *Obes Surg.* 2017;27(9):2279-89.
 41. Angrisani L, Santonicola A, Iovino P, Formisano G, Buchwald H, Scopinaro N. Bariatric Surgery Worldwide 2013. *Obes Surg.* 2015;25(10):1822-32.
 42. Waljee JF, Ghaferi A, Cassidy R, Varban O, Finks J, Chung KC, et al. Are Patient-reported Outcomes Correlated With Clinical Outcomes After Surgery?: A Population-based Study. *Ann Surg.* 2016;264(4):682-9.
 43. Waljee JF, Ghaferi A, Finks JF, Cassidy R, Varban O, Carlin A, et al. Variation in Patient-reported Outcomes Across Hospitals Following Surgery. *Med Care.* 2015;53(11):960-6.
 44. Sarwer DB, Lavery M, Spitzer JC. A review of the relationships between extreme obesity, quality of life, and sexual function. *Obes Surg.* 2012;22(4):668-76.
 45. Poelemeijer YQM, Marang-van de Mheen PJ, Wouters M, Nienhuijs SW, Liem RSL. Textbook Outcome: an Ordered Composite Measure for Quality of Bariatric Surgery. *Obes Surg.* 2019;29(4):1287-94.
 46. Fried M, Hainer V, Basdevant A, Buchwald H, Deitel M, Finer N, et al. Inter-disciplinary European guidelines on surgery of severe obesity. *Int J Obes (Lond).* 2007;31(4):569-77.
 47. Eisenberg D, Bellatorre A, Bellatorre N. Sleeve gastrectomy as a stand-alone bariatric operation for severe, morbid, and super obesity. *JLS.* 2013;17(1):63-7.
 48. Felberbauer FX, Langer F, Shakeri-Manesch S, Schmaldienst E, Kees M, Kriwanek S, et al. Laparoscopic sleeve gastrectomy as an isolated bariatric procedure: intermediate-term results from a large series in three Austrian centers. *Obes Surg.* 2008;18(7):814-8.
 49. Ricci C, Gaeta M, Rausa E, Asti E, Bandera F, Bonavina L. Long-term effects of bariatric surgery on type II diabetes, hypertension and hyperlipidemia: a meta-analysis and meta-regression study with 5-year follow-up. *Obes Surg.* 2015;25(3):397-405.
 50. Mingrone G, Panunzi S, De Gaetano A, Guidone C, Iaiconelli A, Nanni G, et al. Bariatric-metabolic surgery versus conventional medical treatment in obese patients with type 2 diabetes: 5 year follow-up of an open-label, single-centre, randomised controlled trial. *Lancet.* 2015;386(9997):964-73.
 51. Baskota A, Li S, Dhakal N, Liu G, Tian H. Bariatric Surgery for Type 2 Diabetes Mellitus in Patients with BMI <30 kg/m²: A Systematic Review and Meta-Analysis. *PLoS One.* 2015;10(7):e0132335.
 52. Lee WJ, Almulaifi AM, Tsou JJ, Ser KH, Lee YC, Chen SC. Duodenal-jejunal bypass with sleeve gastrectomy versus the sleeve gastrectomy procedure alone: the role of duodenal exclusion. *Surg Obes Relat Dis.* 2015;11(4):765-70.
 53. Ware JE, Jr., Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) Project. *J Clin Epidemiol.* 1998;51(11):903-12.
 54. Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE Open Med.* 2016;4:2050312116671725.
 55. King MT. A point of minimal important difference (MID): a critique of terminology and methods. *Expert Rev Pharmacoecon Outcomes Res.* 2011;11(2):171-84.
 56. Bender R, Lange S. Adjusting for multiple testing--when and how? *J Clin Epidemiol.* 2001;54(4):343-9.

57. Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology*. 1990;1(1):43-6.