

Worldwide treatment opportunities of rheumatoid arthritis Bergstra, S.A.

Citation

Bergstra, S. A. (2018, September 4). *Worldwide treatment opportunities of rheumatoid arthritis*. Retrieved from https://hdl.handle.net/1887/64997

Version:	Not Applicable (or Unknown)
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/64997

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/64997</u> holds various files of this Leiden University dissertation.

Author: Bergstra, S.A. Title: Worldwide treatment opportunities of rheumatoid arthritis Issue Date: 2018-09-04



Inequity in access to bDMARD care and how it influences disease outcomes across countries worldwide: results from the METEOR-registry.

S.A. Bergstra, J.C. Branco, D Vega-Morales, K Salomon-Escoto, N Govind, CF Allaart, RBM Landewé

Ann Rheum Dis, 2018; accepted for publication

ABSTRACT

Objective: To establish in a global setting the relationships between countries' socioeconomic status (SES), measured biologic (b)DMARD-usage and disease outcomes. To assess if prescription- and reimbursement rules and generic access to medication relates to a countries' bDMARD-usage.

Methods: Data on disease activity and drug use from countries that had contributed at least 100 patients were extracted from the METEOR database. Mean disease outcomes of all available patients at the final visit were calculated on a per-country basis. A questionnaire was sent to at least two rheumatologists per country inquiring about DMARD-prices, access to treatment and valid regulations for prescription and reimbursement.

Results: Data from 20.379 patients living in 12 different countries showed that countries' SES was positively associated with measured disease activity (meanDAS28), but not always with physical functioning (HAQ-score). A lower country's SES, stricter rules for prescription and reimbursement of bDMARDs, as well as worse affordability of bDMARDs were associated with lower bDMARD-usage. bDMARD-usage was negatively associated with disease activity (albeit not with physical functioning), but the association was moderate at best.

Conclusions: Disease activity in RA-patients as well as bDMARD-usage varies across countries worldwide. The (negative) relationship between countries' bDMARD-usage and level of disease activity is complex and under the influence of many factors, including – but not limited to- countries' SES, affordability of bDMARDs and valid prescription- and reimbursement rules for bDMARDs.

INTRODUCTION

Earlier diagnosis and treatment, the implementation of treat-to-target and new treatment options, including biologic disease modifying anti-rheumatic drugs (bDMARDs), have improved treatment and prognosis of rheumatoid arthritis (RA) patients tremendously. [1-3] Since many of these treatments are costly, patients across the world may not benefit similarly. Indeed, a lower level of welfare has been associated with higher disease activity in RA-patients in the past.[4]

One of the potentially critical factors is poorer access to bDMARDs.[2, 5] Current recommendations advise starting bDMARDs after a first csDMARD strategy has failed.[5] But such a strategy may not be feasible in greater parts of the world. In many countries there are various restrictions in the prescription and reimbursement of bDMARD.[6-9] Within Europe, differences in socioeconomic welfare are associated with differences in prescription and reimbursement of bDMARDs.[6, 10] Stricter prescription rules and reimbursement criteria of bDMARDs may result in more infrequent use of bDMARDs and in worse health outcomes.[6, 9] To date, only one study, limited to European countries, has been performed that has taken into account all currently available bDMARDs.[6] We have investigated here daily-practice data regarding bDMARD-use in different countries worldwide and have assessed if a lower country's socioeconomic status (SES) is associated with worse clinical outcomes and lower bDMARD-usage. We have also assessed if countries' bDMARD-usage was associated with stricter prescription- and reimbursement rules and worse access to medication.

METHODS

Data selection

Disease activity and medication use in RA-patients in various countries on various treatments were extracted from the METEOR registry, an international database capturing data of daily clinical practice of patients with a clinical diagnosis of RA.[11] Data were gathered retrospectively and anonymously; hence no informed consent was needed. We selected visits after 1-1-2000, from countries that had included at least 100 patients with follow-up data available.

Missing data on disease activity and function (HAQ-score) were imputed using multivariate normal imputation (30 imputations).[12] For each country average DAS28 and HAQ and the proportion of patients in DAS28-remission (DAS28<2.6) were calculated by taking the average of all patients at the last available visit. Furthermore, the proportion of patients that ever used a biological was calculated per country.

Questionnaire

Per participating country, preferably in the region of data collection, at least 2 rheumatologists answered a questionnaire, based on questionnaires used by Putrik et al.[13] In case of disagreement between rheumatologists they were contacted by email, and if necessary additional rheumatologists were contacted to also complete the questionnaire. The questionnaire included questions about availability and affordability of DMARDs, acceptability, reimbursement and prescription rules (supplementary file 1). Drug prices provided in local currency were converted into euros or international dollars at the rate of 10-1-2017. When all questions were processed, a preliminary report was sent to all collaborators, to check correctness of the data.

Outcome measures

Based on the questionnaire results 2 composite scores were calculated: a composite score for clinical eligibility criteria for the start of bDMARDs, based on 3 questions from the questionnaire and with an optimum score of 5 indicating 'least requirements', and a composite score for access to mediation, based on questions on availability, affordability and acceptability, with an optimum score of 9 indicating 'highest level of access' (table 1).[6, 13]

In addition, we calculated the average annual national price of the most frequently used csDMARDs and bDMARDs. These included the csDMARDs methotrexate, sulfasalazine, hydroxychloroquine and leflunomide and prednisone and the bDMARDs etanercept, adalimumab, infliximab, rituximab, certolizumab, tocilizumab, abatacept and golimumab. For each DMARD a most common treatment scheme was used to calculate the costs for one year usage (the annual national price, averaged over the first 2 treatment years).[13] Furthermore, an affordability index for bDMARDs was constructed by dividing the average annual national price for all bDMARDs by the gross domestic product. All medication prices reflect official manufacturer's prices per country, not taking into account local or temporary discounts.

The gross domestic product (GDP) per capita in international dollars and the minimum wage per year, the household-net-adjusted-disposable-income and the health-expenditure-per-capita in US dollars were derived from web-based sources.[14-16] Data regarding the minimum wage and the average price for csDMARDs and bDMARDs were used to calculate the days to work at the minimum wage to cover 30 days of treatment with a csDMARD or bDMARD.[13]

Statistical analyses

At a country level, associations between several indicators of SES, clinical outcomes, medication use, access to medication and prescription and reimbursement rules were assessed using univariable linear regression analyses.

Since analyses were performed at a country level and the number of included countries was limited, multivariable regression analyses were not performed. Regression results for the GDP per capita, the household-net-adjusted-disposable-income and the health-expenditure-per-capita were assessed per 10.000 Intl\$ or US\$. All analyses were performed using Stata SE 14 (StataCorp LP).

Table 1. Composite scores for the clinical eligibility criteria for the start of bDMARDs and for the access to medication.

Composite score clinical crit	eria start of bDMAR	RDs		
	0	1	2	
Is there any requirement for disease duration?	Any requirement	No requirement	NA	
Number of DMARDs to be failed	>2	2	<2	
Level of DAS28	>3.2	≤3.2	No requirement	
Composite score access to n	nedication			
Composite score access to n	nedication 0	1	2	3
Composite score access to n Number of reimbursed bDMARDs	nedication 0	1 1-5	2 6-7	3 8
Composite score access to n Number of reimbursed bDMARDs Average annual price of all reimbursed bDMARDs	nedication 0 0 Highest quartile	1 1-5 Second quartile	2 6-7 Third quartile	3 8 Lowest quartile

NA = not applicable

RESULTS

Country and database characteristics

Twelve countries with 20.379 patients were analysed: United States (state of Massachusetts), Mexico, South-Africa, Japan, Brazil, United Kingdom, Spain, Ireland, Portugal, France, India (state of Maharashtra) and the Netherlands. Data from Qatar and Italy were ultimately excluded from the analyses, since only one rheumatologist in Qatar was available to complete the questionnaire and data from Italy were mainly derived from a biologics register. The number of questionnaire responders per country is listed in supplementary file 2.

Table 2 presents average country and database characteristics. The number of patients

per country ranged from 123 (Spain) to 7.749 (India) and the number of patients ever using a bDMARD ranged from 0.9% (South-Africa) to 75% (Ireland). There were important differences in DAS28- and HAQ-scores across countries. Overall, and expectedly, DAS28 was positively associated with HAQ-score, except in India, where the average DAS28 was highest but the average HAQ-score was among the lowest of all countries. As expected, there were important differences in SES between countries, reflected – for example – by differences in GDP per capita (ranging from Intl\$ 5,733 in India to Intl\$ 61,378 in Ireland) and by large differences in the country's number of days required to work at the minimum wage to cover 30 days of treatment with a bDMARD (ranging from 562 days in India to only 19 days in France).

Average annual medication prices also substantially differed between countries (figure 1). For bDMARDs, drug prices (Intl\$) in the US (highest) were 5.9 times higher than in France (lowest) and for csDMARDs, drug prices in the US (highest) were 14.7 times higher than in the Netherlands (lowest).



Figure 1. Average annual price for csDMARDs (fig 1-A) and bDMARDs (fig 1-B) per country in international dollars (light blue) and in euros (dark blue), prices first quarter 2017.

Countries' SES and clinical outcomes

We first assessed if a lower SES was associated with worse clinical outcomes, by testing associations between GDP per capita and DAS28. Indeed, patients in countries with a higher GDP per capita had a lower average DAS28 and a higher proportion of them were in DAS28-remission [DAS28 lower by β (95% Cl) -0.32 (-0.41; -0.021) and an additional 4.2 (0.14; 8.26) percent of patients in DAS28-remission for every 10.000 Intl\$ additional GDP]. The effect was less prominent in the US and Ireland, both countries with the highest GDP per capita (figure 2A, 2C).

Then, we factored drug-prices into the 'model' by testing the association between the number of days needed to work at the minimum wage in order to afford 30 days of treatment with a bDMARD. Now the association was largely driven by two low-GDP countries (Mexico and India) (figure 2B, 2D) that yet have among the highest drug prices relative to the income. In most other countries, DAS28 and remission percentages were only slightly higher with each extra working day needed to afford bDMARDs: DAS28 higher by β (95% CI) 0.026 (0.012; 0.041) and -0.052 (-0.084; -0.020) less patients in DAS28-remission per additional minimal wage day required to afford 30 days bDMARDs. Finally, we tested health-expenditures-per-capita as well as household's-net-adjusted-disposable income as proxies for SES and assessed the associations with DAS28. In general, the effects were similar: mean DAS28 was -1.3 (-2.6; -0.015) points lower for every additional \$10.000 health-expenditure-per-capita, which culminated into 25 (-2.3; 52.0) percent more patients in DAS28-remission. Such effects were not found for household's net-adjusted disposable income (data not shown).

Overall, RA-patients from low-GDP-countries –on a per-capita basis- appear to have a higher DAS28 than patients from high-GDP-countries, regardless of countries' drug prices. It may be that in some countries drug-prices may mitigate the effects of SES on RA outcomes, (drug prices were for instance importantly lower in Brazil and South-Africa). For HAQ-score, however, the associations with all indicators of SES were less clear: e.g. -0.031 (-0.13; 0.064) lower HAQ per 10.000 Intl\$ increase in GDP per capita and 0.000034 (-0.00091; 0.00098) higher HAQ per additional minimal wage day required to afford 30 days bDMARDs.

SES and bDMARD-usage

It is attractive to assume that the inverse association between SES and DAS28 is mediated by the countries' bDMARD use (or: RA care in high-income countries is better since these can afford bDMARDs). We have sought evidence to underscore this assumption. First, we assessed whether SES was associated with bDMARD-usage per country. Indeed, a statistically significant association was found between GDP per capita and the proportion bDMARD-usage [11.2 (4.82; 17.5), fig3-A], indicating that per additional 10.000 Intl\$ GDP per capita an additional 11% of patients used a bDMARD.

	India	South-Africa	Brazil	Mexico	Portugal	Spain
Country characteristics						
Population (x1.000.000)	1311.1	10.4	207.8	127.0	10.4	46.4
GDP per capita (Intl\$)	5733	12393	14533	16490	26549	32219
Minimum wage per year (US\$)	778	2197	3660	1438	8384	10365
Household net adjusted disposable income (US\$)	NA	8712	11487	12806	19882	22007
Health expenditure per capita (US\$)	75	570	947	677	2097	2658
Days work at minimum wage to cover 30 days treatment bDMARD	562	160	223	431	53	49
Days work at minimum wage to cover 30 days treatment csDMARD	2.2	4.9	3.3	7.4	0.7	0.7
Mean price bDMARDs year / GDP per capita	8.45	1.93	3.07	3.07	0.84	0.70
Composite score access to medication	3	c	ß	1	9	5
Composite score clinical criteria	4	1	1	4	5	4
Database characteristics						
Number of patients	7749	670	189	1191	3874	123
Mean time since diagnosis at last recorded visit (days)	1304	577	4900	2898	5599	1327
% patients bDMARD use	0.95	0.90	19.6	0.6	44.5	16.3
Mean last DAS28	5.1	4.2	4.2	4.0	3.5	3.3
% patients in DAS28-remission	2.3	19.9	17.0	20.9	32.4	42.7
Mean last HAQ	0.67	1.27	1.26	0.71	1.05	0.55

Table 2. Baseline characteristics per country.

	France	Japan	лК	Netherlands	USA	Ireland
Country characteristics						
Population (x1.000.000)	66.5	127.0	65.1	16.9	321.4	4.64
GDP per capita (Intl\$)	37775	37872	38509	46354	52704	61378
Minimum wage per year (US\$)	19886	12269	21793	20673	15080	20967
Household net adjusted disposable income (US\$)	29759	27323	26687	27759	41071	22969
Health expenditure per capita (US\$)	4959	3703	3935	5694	9403	4239
Days work at minimum wage to cover 30 days treatment bDMARD	19	39	32	26	171	48
Days work at minimum wage to cover 30 days treatment csDMARD	0.3	1.8	0.6	0.2	5.4	0.4
Mean price bDMARDs year / GDP per capita	0.37	0.44	0.69	0.42	1.55	0.62
Composite score access to medication	6	4	D	9	4	4
Composite score clinical criteria	4	5	2	4	5	4
Database characteristics						
Number of patients	161	309	1291	3330	803	689
Mean time between diagnosis and last visit (days)	5375	2503	3256	3181	3513	3921
% patients bDMARD use	60.2	50.5	14.7	28.2	48.6	75.0
Mean last DAS28	2.5	3.2	3.9	3.2	3.6	3.8
% patients in DAS28-remission	61.5	38.2	26.0	39.1	30.5	28.8
Mean last HAQ	0.61	0.59	1.29	0.85	0.67	0.85
GDP = Gross Domestic Product, Intl\$ = international dollar, I MARD = conventional synthetic disease modifying anti-rheu >1 available visit. % patients bDMARD use' indicates the nu 'mean last HAO' are the DAS28 and HAO at the last available	JS\$ = United S matic drug, N mber of patie e visit in the d	states dollar, bD A = not available nts using a biolc atabase.	MARD = biolog e. 'Number of p ogical DMARD o	ic disease modif atients' indicate luring at least 1	Ying anti-rheu es the number visit. 'Mean la	matic drug, csD- of patients with st DAS28' and

When taking drug-prices into account, the picture is more obscure. Although in Mexico and India bDMARD-usage was lowest, in the countries with highest GDP per capita bDMARD-usage was highly variable (ranging from close to 10% in the UK to 75% in Ireland), [fig3-B, β (95% CI) -0.080 (-0.16; 0.0021)]. This suggests that not only GDP and drug-prices but also other mechanisms (such as limitative regulations for reimbursement) determine bDMARD-usage.

bDMARD-usage and clinical outcomes

It is questionable, however, if a higher percentage of bDMARD-usage translates automatically into better disease outcomes. We assessed whether bDMARD-usage across



Figure 2. Associations between 'GDP per capita (Intl\$)' and 'days to work at the minimum wage to cover 30 days of treatment with a bDMARD' with clinical outcomes per country.



Figure 3. Associations between 'GDP per capita' and 'days to work at the minimum wage to cover 30 days of treatment with a bDMARD' with '% bDMARD use'.

countries are associated with clinical outcomes. Indeed we found a statistically significant relationship between a country's proportion of bDMARD-usage and DAS28 or proportion of patients in DAS28-remission (Fig 4A, 4B). DAS28 was -0.14 (-0.28; -0.0054) point lower, and 2.8% (-0.13; 5.8) more patients achieved DAS28-remission, for every 10% increase in proportion of patients using a bDMARD. However, bDMARD-usage was not associated with better functional ability [-0.024 (-0.091; 0.042) lower HAQ-score for

Prescription and reimbursement rules, access to medication and bDMARD-usage

Since bDMARD-usage is not only influenced by a country's SES, it was subsequently assessed whether the stringency of prescription- and reimbursement-rules and 'access to medication' were associated with proportion of bDMARD-usage.

We found that bDMARD-usage is less if limitative regulations are stricter: 8.5 (-2.7; 19.8) percent more bDMARD use per point increase (i.e. fewer limitations) in clinical criteria score and a trend [5.9 (-2.0; 13.8)] that better access to bDMARD-care led to more bDMARD-usage (figure 4D and 4E).

This shows that the previous relationship found between a country's SES and quality of RA care measured as a country's mean DAS28, is (among others) confounded by regulations. Relatively strict prescription- and reimbursement rules in the UK, a high SES country, result in a proportion of bDMARD-usage as low as in India and Mexico, which both have a low GDP per capita.

Finally, we calculated the quotient of a country's mean drug-price and the GDP per capita (as proxy for affordability, the lower the quotient, the less affordable the drug) and found 1) that even in countries with a same level of affordability (e.g. EU countries) significant differences in bDMARD-usage exist, apparently due to other mechanisms than drug-prices alone; and 2) that affordability of bDMARDs in some countries is so low that bDMARD-usage is virtually zero.



Figure 4. Associations between the '% of patients that ever used a bDMARD' and the 'composite score clinical criteria', 'composite score access to medication' and clinical outcomes.

DISCUSSION

Worldwide, treatment options and clinical outcomes of RA-patients have greatly improved, but not all RA-patients have benefitted similarly. We hypothesized that differences in SES have an impact on bDMARD-usage and on clinical outcomes across countries. Indeed, in this study including a large number of patients from 12 countries, among which several countries that have never been investigated before in this context, we have found substantial differences in DMARD-prices, affordability of these medications and bDMARDusage across countries. We found that in countries with a lower SES disease activity was generally higher and bDMARD-usage was lower. But a country's proportion of bDMARD- usage was also associated with restrictions through prescription and reimbursement rules, and with affordability of bDMARDs, as defined by us.

It is attractive to assume that higher country's bDMARD-usage will result in a lower country's mean DAS28, and that a lower country's GDP will hinder a sufficiently high proportion of RA-patients getting proper access to care with bDMARDs. But reality is more complicated. The effectiveness of bDMARD-usage in countries' all-day clinical practice may be overstated; previous research estimated that 'only' 7% of the effect of GDP per capita on DAS28 was mediated by the uptake of bDMARDs.[4] We found 'only' 2.8% more patients in DAS28-remission for every additional 10% patients using a bDMARD. A positive effect of bDMARDs on RA treatment effectiveness thus appears to be quite small. Vice versa, this suggests that in low-income countries other factors than 'only' access to bDMARDs determine the success of RA treatment. Nevertheless, a general trend between countries' proportions of bDMARD-usage and countries' mean-DAS28 remains obvious. Remarkably, we did not find an association between countries' SES and countries' mean HAQ-score. Here, the effect of outliers is relatively important. In particular India, the country with lowest GDP, reported a low HAQ-score compared to a high DAS28. Moreover, there may be socio-demographic and cultural differences in the way patients experience or report limitations in function.[17, 18] We could not assess the potential contribution of factors such as general access to health care and other drug and non-drug therapies, comorbidities and health barriers and support systems.[19]

Previous studies have mentioned associations between access to medication, SES and disease activity.[6, 13, 20] Such findings point to the negative effects of inequity: budget restrictions, strict regulations as well as limited access to drugs may be a hurdle for starting optimal treatment as recommended in clinical guidelines.[13, 21]

But this study also shows that several other factors play a role in determining the success of RA-treatment (here approximated by the countries' mean DAS28). We know several of these factors: countries' SES in general, the presence of a proper functioning health-care system that may assure access to care to those who are in need, DMARD prices and valid national regulations that are in place to constrain the expenses for bDMARD-usage. [4, 6-9, 13, 21] It appears obvious that the country's mean level of DAS28 is the resultant of a complicated interplay of a country's SES, drug prices and regulations. In addition, it is difficult to argue that unlimited access to expensive effective treatments makes the difference between 'good and bad care' for RA-patients, nor can we claim that countries with similar GDP per capita or similar levels of access to care have similar proportions op patients on expensive bDMARDs; there is huge variation. Nevertheless, penetration of bDMARDs in low GDP-countries stays behind and it is to be expected that this –among others- may go at the cost of effectiveness of RA-care. It is impossible to conclude from this study whether this is due to drug-prices, failing health care systems or simply worse access to optimal care. We can only conclude that there are substantial differences in

mean DAS28 (as a proxy for quality of RA-care) across countries.

This study has some strengths and many limitations. A strength of this study is that it captures real life clinical data from 12 countries world-wide with large differences in wealth, totally different (if any) health-care and health-insurance systems and many RA-patients. As such, this study can be considered a 'big-data study' allowing subtle differences across countries to be elucidated.

But the strengths of our study (real life observational, size and international diversity) also carry limitations: case-ascertainment (cases cannot be verified), completeness of data (we had to statistically impute missing data) and reliability of data-points (we had to rely on the report of the participating physicians) are among them. Other epidemiological limitations are that only few centres per country participated and we had to assume that these centres were to some extent representative of the country. In addition, we had to make certain assumptions to facilitate computations, such as declaring bDMARD-reimbursement as 'absent', if according to the rheumatologist's questionnaires less than 20% of patients in a country had health insurance coverage. Such assumptions –if flawed- may influence the reported associations. In a few cases, we relied on regional health-economical information rather than on country-specific data, in the appreciation that within a big country access to health care and regulations can be very different.

A final limitation of this database is that it will only include RA-patients that have come to the attention of the rheumatologist. If countries differ with regard to access to a rheumatologist, patients per country cannot be assumed to be comparable. Consequently, associations may be spurious. With regard to this latter argument, it can be postulated that the associations in this study are conservative and will likely be more exaggerated in real life.

Epidemiological limitations of 'big-data studies' restrict their interpretability. As such, causal interpretations will never be possible and should always be mistrusted. We have taken care to not exaggerate our conclusions that all remain at the level of associations and allow the possibility of bias and confounding as explanatory factors. Still, 'big data studies' make sense in that they can point to relevant differences between countries, that may help policymakers to guide necessary change, pharmaceutical industry to direct market access and drug-prices, and rheumatologists and health-care workers to help improving access to rheumatology care.

In conclusion, we have documented using a registry of patients with RA spanning 12 countries world-wide that mean DAS28 as well as bDMARD-usage varies across countries. While we suggest an inverse relationship between the countries' bDMARD-usage and mean DAS28, this relationship is influenced by many other factors, including countries' GDP per capita, strictness of prescription and reimbursement rules and affordability of bDMARDs. All together these findings point to the existence of worldwide inequity with regard to optimal (access to) RA health care.

REFERENCES

- Quinn MA, Conaghan PG, Emery P. The therapeutic approach of early intervention for rheumatoid arthritis: what is the evidence? *Rheumatology (Oxford, England)* 2001; 40:1211-20.
- Nam JL, Ramiro S, Gaujoux-Viala C et al. Efficacy of biological disease-modifying antirheumatic drugs: a systematic literature review informing the 2013 update of the EULAR recommendations for the management of rheumatoid arthritis. Ann Rheum Dis 2014; 73:516-28.
- Smolen JS, Breedveld FC, Burmester GR et al. Treating rheumatoid arthritis to target: 2014 update of the recommendations of an international task force. Ann Rheum Dis 2016; 75:3-15.
- 4. Putrik P, Ramiro S, Keszei AP *et al.* Lower education and living in countries with lower wealth are associated with higher disease activity in rheumatoid arthritis: results from the multinational COMORA study. *Ann Rheum Dis* 2016; 75:540-6.
- Smolen JS, Landewe R, Bijlsma J et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. Ann Rheum Dis 2017; 76:960-77.
- Putrik P, Ramiro S, Kvien TK, Sokka T, Uhlig T, Boonen A. Variations in criteria regulating treatment with reimbursed biologic DMARDs across European countries. Are differences related to country's wealth? Ann Rheum Dis 2014; 73:2010-21.
- Emery P, Van VR, Ostergaard M et al. Guidelines for initiation of anti-tumour necrosis factor therapy in rheumatoid arthritis: similarities and differences across Europe. Ann Rheum Dis 2009; 68:456-9.
- Orlewska E, Ancuta I, Anic B *et al*. Access to biologic treatment for rheumatoid arthritis in Central and Eastern European (CEE) countries. *Med Sci Monit* 2011; 17:SR1-13.
- Pease C, Pope JE, Truong D *et al*. Comparison of anti-TNF treatment initiation in rheumatoid arthritis databases demonstrates wide country variability in patient parameters at initiation of anti-TNF therapy. *Semin Arthritis Rheum* 2011; 41:81-9.
- Laires PA, Exposto F, Mesquita R, Martins AP, Cunha-Miranda L, Fonseca JE. Patients' access to biologics in rheumatoid arthritis:

a comparison between Portugal and other European countries. *The European journal of health economics : HEPAC : health economics in prevention and care* 2013; 14:875-85.

- Bergstra SA, Machado PM, van den Berg R, Landewe RB, Huizinga TW. Ten years of METEOR (an international rheumatoid arthritis registry): development, research opportunities and future perspectives. *Clin Exp Rheum* 2016; 34:S87-s90.
- Schafer JL. Analysis of Incomplete Multivariate Data.: Boca Raton, FL: Chapman & Hall/CRC; 1997.
- Putrik P, Ramiro S, Kvien TK *et al*. Inequities in access to biologic and synthetic DMARDs across 46 European countries. *Ann Rheum Dis* 2014; 73:198-206.
- Wikipedia, List of minimum wages by country, https://en.wikipedia.org/wiki/List_ of_minimum_wages_by_country, accessed 24-05-2017.
- 15. The World Bank, http://www.worldbank.org/, accessed 24-05-2017.
- OECD, OECD Better Life Index: Income, http://www.oecdbetterlifeindex.org/topics/ income/, accessed 24-05-2017.
- Hifinger M, Norton S, Ramiro S, Putrik P, Sokka-Isler T, Boonen A. Equivalence in the Health Assessment Questionnaire (HAQ) across socio-demographic determinants: Analyses within QUEST-RA. Semin Arthritis Rheu 2017.
- Putrik P, Ramiro S, Hifinger M et al. In wealthier countries, patients perceive worse impact of the disease although they have lower objectively assessed disease activity: results from the cross-sectional COMORA study. Ann Rheum Dis 2016; 75:715-20.
- van den Hoek J, Roorda LD, Boshuizen HC, Tijhuis GJ, van den Bos GA, Dekker J. Physical and Mental Functioning in Patients with Established Rheumatoid Arthritis over an 11-year Followup Period: The Role of Specific Comorbidities. J Rheumatol 2016; 43:307-14.
- Putrik P, Ramiro S, Chorus AM, Keszei AP, Boonen A. Socioeconomic inequities in perceived health among patients with musculoskeletal disorders compared with other chronic disorders: results from a cross-sectional Dutch study. *RMD open* 2015; 1:e000045.
- Guillemin F, Carruthers E, Li LC. Determinants of MSK health and disability--social determinants of inequities in MSK health. *Best Pract Res Clin Rh* 2014; 28:411-33.