

The extended Planetary Nebula Spectrograph (ePN.S) early-type galaxy survey. The kinematic diversity of stellar halos and the relation between halo transition scale and stellar mass

C. Pulsoni^{1,2}, O. Gerhard¹, M. Arnaboldi³, L. Coccato³, A. Longobardi⁵, N. R. Napolitano⁶, C. Narayan^{1,8}, V. Gupta^{1,7}, A. Burkert⁴, M. Capaccioli⁹, A. L. Chies-Santos¹⁰, A. Cortesi¹¹, K. C. Freeman¹², K. Kuijken¹³, M. R. Merrifield¹⁴, A. J. Romanowsky^{15,16}, and C. Tortora¹⁷

¹ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany

² Excellence Cluster Universe, Boltzmannstraße 2, 85748, Garching, Germany

³ European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748 Garching, Germany

⁴ University Observatory Munich, Scheinerstraße 1, 81679 Munich, Germany

⁵ Kavli Institute for Astronomy and Astrophysics, Peking University, 5 Yiheyuan Road, Haidian District, Beijing 100871, PR China

⁶ INAF - Astronomical Observatory of Capodimonte, Salita Moiariello, 16, 80131, Naples, Italy

⁷ Department of Physics, Cornell University, Ithaca, New York 14853, USA

⁸ Current address: 71, Akashganga IUCAA Post Bag 4, Ganeshkhind Pune University Campus, Pune - 411007, India

⁹ University of Naples "Federico II", C.U. Monte Sant'Angelo, via Cinthia, I-80126, Naples, Italy

¹⁰ Departamento de Astronomia, Instituto de Física, Universidade Federal do Rio Grande do Sul, Porto Alegre, R.S 90040-060, Brazil

¹¹ Departamento de Astronomia, Instituto de Astronomia, Geofísica e Ciências Atmosféricas da USP, Cidade Universitária, CEP:05508900 São Paulo, SP, Brazil

¹² Research School of Astronomy and Astrophysics, Mount Stromlo Observatory, Cotter Road, Weston Creek, ACT 2611, Australia

¹³ Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands

¹⁴ School of Physics and Astronomy, The University of Nottingham, University Park, Nottingham, NG7 2RD, UK

¹⁵ Department of Physics and Astronomy, San Jose State University, One Washington Square, San Jose, CA 95192, USA

¹⁶ University of California Observatories, 1156 High Street, Santa Cruz, CA 95064, USA

¹⁷ Kapteyn Astronomical Institute, University of Groningen, Postbus 800, NL-9700 AV Groningen, the Netherlands

November 13, 2018

ABSTRACT

Context. In the hierarchical two-phase formation scenario, the extended halos of early type galaxies (ETGs) are expected to have different physical properties from those of the galaxies' central regions.

Aims. This work aims at characterizing the kinematic properties of ETG halos using planetary nebulae (PNe) as tracers, which allow us to overcome the limitations of absorption line spectroscopy of continuum at low surface brightness.

Methods. We present two-dimensional velocity and velocity dispersion fields for 33 ETGs, including both fast and slow rotators, making this the largest kinematic survey to-date of extragalactic PNe. The survey is based on data from the custom built Planetary Nebula Spectrograph (PN.S) instrument, supplemented with PN based kinematics from counter-dispersed imaging and from high-resolution spectroscopy of narrow band-selected PN samples. The velocity fields are reconstructed from the measured PN velocities using an adaptive kernel procedure validated with simulations. They typically extend out to 6 effective radii (R_e), with a range $[3R_e - 13R_e]$ for the PN.S ETGs. We complemented the PN kinematics with absorption line data from the literature, in order to have a complete description of the kinematics, from the central regions to the outskirts.

Results. We find that ETGs typically show a kinematic transition between inner regions and halo. Estimated transition radii in units of R_e anticorrelate with stellar mass. Slow rotators have increased but still modest rotational support at large radii, while most of the fast rotators show a decrease in rotation, due to the fading of the stellar disk in the outer, more slowly rotating spheroid. 30% of the fast rotators are dominated by rotation also at large radii. Most ETGs have flat or slightly falling halo velocity dispersion profiles, but 15% of the sample have steeply falling profiles. One third of the fast rotators show kinematic twists, misalignments, or rotation along two axes, indicating that they turn from oblate near the center to triaxial in the halo.

Conclusions. ETGs have more diverse kinematic properties in their halos than in the central regions, and a significant fraction shows signatures of triaxial halos in the PNe data. The observed kinematic transition to the halo and its dependence on stellar mass is consistent with Λ CDM simulations and supports a two-phase formation scenario.

Key words. Galaxies: elliptical and lenticular, cD - Galaxies: general - Galaxies: halos - Galaxies: kinematics and dynamics - Galaxies: structure

1. Introduction

Observations and cosmological simulations suggest a two-phase formation scenario for early type galaxies (ETGs) (e.g. Trujillo

et al. 2006; Oser et al. 2010; van Dokkum et al. 2010; Rodriguez-Gomez et al. 2016; Qu et al. 2017). An initial fast assembly stage, in which the ETGs grow through rapid star formation fueled by the infall of cold gas ($z \gtrsim 1.5$) or through major merger

events (Wuyts et al. 2010; Bournaud et al. 2011; Sommer-Larsen & Toft 2010; Wellons et al. 2016), is followed by a series of merger episodes which enrich the galaxy halos of stars and make them grow efficiently in size (Oser et al. 2010; Gabor & Davé 2012; Lackner et al. 2012; Buitrago et al. 2017). The hierarchical accretion scenario finds its best evidence in the observations of a rapid growth of stellar halos at redshift $\lesssim 2$ with little or no star formation (e.g. Daddi et al. 2005; Trujillo et al. 2007; van Dokkum et al. 2010; Damjanov et al. 2011). In this context ETGs are layered structures in which the central regions are the remnant of the primordial stars formed in-situ, while the external halos are principally made of accreted material. The consequence is that the halos are expected to show significant variation with radius of properties, such as light profiles (Huang et al. 2013; D’Souza et al. 2014; Iodice et al. 2016; Spavone et al. 2017), and kinematics (Romanowsky & Fall 2012; Coccato et al. 2009; Arnold et al. 2014; Foster et al. 2016).

Long slit spectroscopic observations of ETGs (e.g. Davies et al. 1983; Franx et al. 1989; Bender et al. 1994) revealed that this apparently homogeneous class of objects actually displays a kinematic diversity which also correlates with the isophote shape (Bender 1988b; Kormendy & Bender 1996). Disky ellipticals generally rotate fast, while slowly rotating ellipticals have a rather boxy shape. A remarkable step forward in the comprehension of the nature of ETGs has been attained by the ATLAS^{3D} project (Cappellari et al. 2011), which for the first time applied integral-field spectroscopy (IFS) over a statistically-significant sample, mapping kinematics, dynamics, and stellar population properties within one effective radius (R_e). A new paradigm for ETGs was proposed, which distinguishes between fast and slow rotators according to the central projected specific angular momentum, λ_R (Emsellem et al. 2007). Fast rotators include also S0 galaxies and represent the great majority (86%) of ETGs. These are oblate systems with regular disk-like kinematics along the photometric major axis. Slow rotators, on the other hand, often display kinematic features such as counter-rotating cores or twist of the kinematic position angle. They are relatively more spheroidal systems, mildly triaxial, and tend to be massive (Cappellari et al. 2013).

The two classes have been interpreted as the result of the variety of processes shaping galaxies, leading to a sequence of baryonic angular momentum (Emsellem et al. 2011; Naab et al. 2014; Wu et al. 2014). Following this argument Emsellem et al. (2011) suggest the possibility of a continuous range of properties among fast rotators that links them to slow rotators. On-going surveys like MANGA (Bundy et al. 2015), CALIFA (Sánchez et al. 2012), SAMI (Croom et al. 2012; Bryant et al. 2015), and MASSIVE (Ma et al. 2014) are currently working on increasing the size of the sample of IFS mapped objects, and extending the study to a wider range of environment and mass.

However, a classification scheme based on the characteristics of the galaxies in the central regions (inside $\sim 1R_e$) may not be fully representative of the nature of these objects (e.g. Bellstedt et al. 2017b), raising the question of how complete our understanding is without a full knowledge of their properties at larger scales. The halos, in fact, contain half of the stars of the galaxies and most of their dynamical mass. Dark matter is known to dominate there (e.g. Mandelbaum et al. 2006; Humphrey et al. 2006; Koopmans et al. 2009; Churazov et al. 2010) and a dynamical modeling of the outskirts is essential to constrain its distribution at intermediate radii (e.g. Gerhard et al. 2001; Romanowsky et al. 2003; Thomas et al. 2009; Napolitano et al. 2011; Morganti et al. 2013). Stellar halos are predicted to host mostly accreted star material as shown by particle tagging sim-

ulations (Cooper et al. 2013) and hydro-dynamical simulations (Rodriguez-Gomez et al. 2016). In addition, these regions provide insight into the most recent dynamical phase of the galaxy. In the halo the settling times are of order 1 Gyr and so signatures of the most recent assembly events may still be apparent, providing a mine of information about the mechanisms of formation and evolution (e.g. Bullock & Johnston 2005; Tal et al. 2009; Romanowsky et al. 2012; Coccato et al. 2013; Duc et al. 2015; Longobardi et al. 2015). Thus extending investigations to the outer halos is crucial for having a complete picture of ETGs.

This issue particularly affects kinematic measurements based on stellar absorption lines for ETG halos, which generally lack cold gas (and so the 21 cm HI emission) used to probe the outer parts of spiral galaxies. Since the continuum light from the stars quickly drops with radius, this kind of observation is challenging beyond $1 - 2 R_e$, limiting the assessment of the dynamics. This limitation is particularly frustrating, because the complicated kinematics of ETGs, dominated by dispersion, necessitates a good knowledge of the higher moments of the line of sight velocity distribution (LOSVD) in order to alleviate the anisotropy-potential-degeneracy (e.g. Gerhard 1993; Rix et al. 1997; Thomas et al. 2009; de Lorenzi et al. 2009; Napolitano et al. 2009).

Kinematic studies of ETGs from integrated-light spectra out to large radii have been performed by Kelson et al. (2002); Weijmans et al. (2009); Coccato et al. (2010); Murphy et al. (2011); Barbosa et al. (2017) using long slit spectroscopy or IFS on individual objects. More recently the SLUGGS survey (Arnold et al. 2014; Foster et al. 2016; Bellstedt et al. 2017a), the MASSIVE survey (Raskutti et al. 2014; Veale et al. 2017), and Boardman et al. (2017) generated kinematic data from integral field spectrographs (IFSs) for larger samples of ETGs, but never reaching beyond $3 - 4R_e$ ¹.

The only possibility to probe the kinematics of a large sample of galaxies out to the very outskirts is through kinematic tracers that overcome the limit of the decreasing surface brightness, like globular clusters (e.g. Schuberth et al. 2010; Strader et al. 2011) or planetary nebulae.

Planetary nebulae (PNe) are established probes of the stellar population in ETG halos (e.g. Longobardi et al. 2013; Hartke et al. 2017). Their bright [OIII] line stands out against the faint galaxy background, making them relatively easy to detect. Since they are drawn from the main stellar population, their kinematics traces the bulk of the host-galaxy stars, and are directly comparable to integrated light measurements (Hui et al. 1995; Arnaboldi et al. 1996; Méndez et al. 2001; Coccato et al. 2009; Cortesi et al. 2013a). This makes the PNe the ideal kinematic probes for the halos of ETGs; globular clusters and do not follow the surface brightness distribution of the stars and do not show the same kinematics (e.g. Brodie & Strader 2006; Coccato et al. 2013; Veljanoski et al. 2014) and their color bimodality, suggesting two distinct formation mechanisms (Renaud et al. 2017, and references therein), complicates the interpretation of the kinematics when they are used as tracers. The pioneering work of Coccato et al. (2009) studied the kinematics of 16 ETGs traced with PNe out to $8R_e$, finding evidence for kinematic transitions at large radii from the trends observed in the central regions. These may be interpreted as the sought transition from in-situ to accreted components, as also shown by Arnold et al. (2014).

¹ The values of R_e used by most kinematic studies are measured in the bright central regions of galaxies, and may underestimate the half light radii (see discussion in section 8.4).

The extended Planetary Nebula Spectrograph (ePN.S) survey is based on observation mostly done with the PN.S, and consists of catalogs of PNe for 33 ETGs. This dataset is the largest survey to-date of extragalactic PNe identified in the halos of ETGs, covering from 3 to $13 R_e$, and complementing the absorption line kinematics available in the literature. The rationale of the survey, the sample definition, and the construction of the catalogs are described in detail in Arnaboldi et al. (in preparation). Section 2 is a brief description of the ePN.S sample. Section 3 describes the general procedure adopted for extracting the mean velocity fields from the measured radial velocities of PNe, and reviews the adaptive kernel smoothing technique introduced by Coccato et al. (2009). In section 4 we evaluate the systemic velocity of the galaxies. The point-symmetry of the smoothed velocity fields is studied in section 5, while the trends of the kinematic parameters, such as rotational velocity, kinematic position angle, and velocity dispersion, are derived in section 6. The results are described in section 7 with a detailed analysis of slow rotators and fast rotators. A discussion of the results is presented in section 8. In section 9 we give a summary of the work and draw our conclusions.

2. Description of the Sample - Observations - Data reduction

This work is based mostly on data collected with the Planetary Nebula Spectrograph at the William Herschel Telescope in La Palma. This is a custom-built instrument designed for counter-dispersed imaging (Douglas et al. 2002). We collected catalogs of PNe for 25 galaxies, to which we added catalogs from the literature and two unpublished catalogs for two galaxies from Counter Dispersed Imaging (CDI) observations, for a total of 33 ETGs. The ePN.S sample is magnitude limited and covers a wide range of internal parameters, such as luminosity, central velocity dispersion, ellipticity, boxy/diskyness (table 1 summarizes the properties of the sample and the origin of the catalogs). Our catalogs contain a total of 8636 PNe, with data typically extending out to $6R_e$, with a range $[3R_e-13R_e]$. This makes the ePN.S the largest kinematic survey to-date of extragalactic PNe in the outer halos of ETGs.

Arnaboldi et al. (in prep.) give a full discussion of the extraction of the catalogs; here we provide a brief description of the adopted procedures. All the datasets (the new sample of galaxies, as well as the extended PN catalogs from the literature) are uniformly validated, in order to obtain a homogeneous sample of ETGs whose properties can be consistently compared.

After the raw catalog has been obtained, it is uniformly cleaned from spurious sources and so-called "outliers", i.e., candidate sources that are classified as spurious detections. The first step for the outlier removal among the PN candidates is the removal of all the detections with signal-to-noise ratio below a given threshold. We adopted $S/N \geq 2.5$ as good compromise value between a reasonable signal-to-noise value and number of detections that satisfy this requirement. We then separate between PNe belonging to any satellites from those in the hosts. We used the probability membership method from McNeil-Moylan et al. (2012), which uses both kinematic and photometric information to assign to each star a probability of belonging to the satellite/host. The membership to the host galaxy is assigned only if the probability is greater than 90%.

The last step is the removal of the contamination from other narrow emission-line sources (i.e. background star-forming galaxies) that are not resolved and appear point-like in the

counter-dispersed images, similar to the monochromatic [OIII] $5007\text{\AA}$ emission from a PN. We identify them using a robust mean/sigma clipping procedure. The algorithm derives a robust mean velocity (v_{mean}) and velocity dispersion σ , using a running average in the phase space with a window of N data points ($15 \leq N \leq 30$, according to the number of tracers in each galaxy) and 3 data points step. An iterative procedure clips away the PN candidates whose $|v - v_{\text{mean}}| > 2\sigma$, and evaluates v_{mean} and σ until the number of clipped objects stabilizes. In each iteration σ is corrected by a factor 1.14 to account for the 2σ cut of the LOSVD tails. At last, the sample of N_{val} validated PNe is added a number of $N_{\text{val}} \times 4.5\%$ among the wings clipped objects closest to the 2σ contours, so that the final sample also includes the objects in the tails of the LOSVD.

In the case of disk galaxies, dominated by rotation, a disk/spheroid decomposition was performed following Cortesi et al. (2011): using both photometric data and kinematic information we assigned to each PN the probability of being associated with each photometric component. The tagging of the outliers from the disk and the spheroid separately allows us to account for their different kinematics when using the robust mean/sigma clipping procedure. The disk is processed first, and its flagged PNe are added to the PNe of the bulge during the last iteration; the flagged bulge PNe are then considered as outliers of the entire galaxy.

The datasets processed in this way are the Bona Fide PNe catalogs used in the following analysis.

3. Kernel smoothing method

The measured line-of-sight (LOS) velocities of the PNe are random samplings of the galaxy LOSVD function at the position of the source. Therefore each velocity measurement randomly deviates from the local mean velocity by an amount that depends on the local LOS velocity dispersion.

In order to extract the mean LOS velocity and the LOS velocity dispersion fields from this discrete velocity field, we use an adaptive kernel smoothing technique, as described in Coccato et al. (2009), that performs a local average of the measured discrete LOS velocities. In the following section we briefly review the smoothing technique, while we refer to Coccato et al. (2009) for a more detailed discussion. In appendix A we validated the adopted procedure on simulated data, in order to test the effects of different statistical realizations of a system, limited number of tracers, and different V/σ ratios on the estimate of the kinematic parameters.

3.1. Averaging the discrete velocity field with the adaptive kernel smoothing technique

The smoothing of the discrete velocity field is carried out by computing the velocity at each position (x, y) in the sky as a weighted mean $\tilde{v}(x, y)$ of all the PN LOS velocities v

$$\tilde{v}(x, y) = \frac{\sum_i v_i w_{i,p}}{\sum_i w_{i,p}} \quad (1)$$

while the velocity dispersion $\tilde{\sigma}(x, y)$ is given by the square root of the variance of varound $\tilde{v}$

$$\begin{aligned} \tilde{\sigma}(x, y) &= (\langle v^2 \rangle - \langle v \rangle^2 - \delta v^2)^{1/2} \\ &= \left[\frac{\sum_i v_i^2 w_{i,p}}{\sum_i w_{i,p}} - \tilde{v}(x, y)^2 - \delta v^2 \right]^{1/2} \end{aligned} \quad (2)$$

The weight $w_{i,p}$ of each PN is defined using a Gaussian Kernel that depends on the distance of the PN from the position (x, y) , normalized by a kernel amplitude K

$$w_{i,p} = \exp \frac{-D_i^2}{2K(x,y)^2} \quad ; \quad K(x,y) = A \sqrt{\frac{M}{\pi\rho}} + B \quad (3)$$

The latter controls the spatial scale of the region over which the smoothing is performed, and hence the spatial resolution of the kinematic study. Large values of K , in fact, lead to smoother profiles in the mean LOS velocity fields, highlighting the general trends, but also suppressing the small scale structures, while smaller values of K allow a better spatial resolution but may amplify any noise pattern. Hence the optimal K should be a compromise between spatial resolution and statistical noise smoothing. The amplitude K is defined to be linearly dependent on the distance between the position (x, y) and the M^{th} closest PN, so that K is a function of the local density of tracers $\rho(x, y)$:

This allows K to be smaller in the innermost, dense regions of galaxies, and larger in the outskirts, where their density is usually lower. The optimal kernel parameters A and B are derived as described in section 3.1.1. We chose $M = 20$, but Coccato et al. (2009) tested the procedure with $10 < M < 60$ finding no significant differences in the results.

Errors on $\tilde{v}(x, y)$ and $\tilde{\sigma}(x, y)$ are obtained using Monte Carlo simulations, as discussed in Coccato et al. (2009). For each galaxy, 100 datasets of PNe are built with simulated radial velocity at the same positions as the observed dataset. The radial velocities for each simulated object is calculated from the two-dimensional smoothed velocity field by adding a random value, extracted from a Gaussian distribution centered at 0 and with dispersion $\sigma = \sqrt{\tilde{\sigma}^2 + \delta v^2}$, where δv is the velocity error. These simulated datasets were smoothed with the same kernel K as the real sample, and the standard deviation of the simulated velocity and velocity dispersion fields give the errors on $\tilde{v}$ and $\tilde{\sigma}$.

3.1.1. Deriving the optimal A and B kernel parameters

The parameters A and B in equation (3) are chosen so that the best compromise between spatial resolution and noise smoothing is achieved. We developed an iterative procedure in order to derive the optimal kernel parameters that realize this condition.

We first estimated the velocity gradient to be resolved by the smoothing procedure by performing a preliminary averaging with a fully adaptive kernel ($A = 1$ and $B = 0$). The derived mean velocity field, $\tilde{v}_{A=1, B=0}$, is fitted with a cosine function, which, in general, approximately describes the velocity profiles of early type galaxies (see Cohen & Ryzhov 1997):

$$\tilde{v}_{A=1, B=1}(\phi) = V_{\text{max}} \cos(\phi - PA_{\text{kin}}) + \text{const} \quad (4)$$

This interpolation function provides a measure of the position angle of the kinematic major axis (PA_{kin}), along which the steepest velocity gradient $d\tilde{v}$ is expected to lie. The gradient $d\tilde{v}$ is obtained by fitting a straight line to the velocities $\tilde{v}$ of the PNe lying in a section along the PA_{kin} direction, as a function of the radius.

The best kernel parameters that allow to resolve spatial substructures with typical velocity gradient $d\tilde{v}$ are derived by building simulated sets of PNe. The stars are spatially distributed according to a given density $\bar{\rho}$, while their velocities are assigned using the derived velocity gradient and adding a dispersion equal to the standard deviation of the observed radial velocities. The artificial sets are processed using different values of the kernel K

until the simulated input velocity field is recovered: this provides the best $k(\bar{\rho})$. The procedure is repeated for different values of ρ , and the optimal A and B are the best fit values of equation (3) on the derived best K as a function of ρ .

3.2. Fitting a rotation model

The mean velocity fields, derived from smoothing the discrete velocities, are divided into radial bins of equal numbers of PNe such that they contain at least 30 stars. If a galaxy contains less than 60 tracers, we divide the sample in two bins in order to study possible radial trends.

The bins are spherical for galaxies either with small flattening, i.e. $10 \times \epsilon < 3$, or whose PN spatial distribution has a rather square shape. For all the other galaxies we use elliptical bins oriented along the photometric major axis, with a flattening equal to a characteristic ellipticity of the isophotes of the galaxy (the adopted ellipticity, ϵ , and photometric position angle PA_{phot} are given in table 1).

We found that the results of our analysis do not depend on the chosen flattening but, since the spatial distribution of the PNe follows the light, and so it may be rather flattened, elliptical bins help to sample the field homogeneously in an azimuthally-unbiased way.

We fitted the mean velocity $\tilde{v}(\phi, R)$ as a function of the position angle in each bin with a rotation model (see section 3.2.1). The position angle of the stars in the elliptical bins is defined as $\phi(x, y) = \arctan[y/((1 - \epsilon)x)]$, where ϵ is the ellipticity. R is the mean major axis distance of the PNe in each bin.

3.2.1. Point-symmetric rotation model

A velocity map $\tilde{v}(R, \phi)$ is a periodic function in ϕ , so it can be expanded in a Fourier series and approximated by a finite number of harmonics:

$$\tilde{v}(R, \phi) = a_0(R) + \sum_{n=1}^N a_n(R) \cos(n\phi) + \sum_{n=1}^N b_n(R) \sin(n\phi) \quad (5)$$

Elliptical galaxies in dynamical equilibrium are triaxial systems (e.g. Statler 1994, and references therein), so the projection on the sky of the mean velocity field should be *point-symmetric* with respect to the systemic velocity a_0 (see Krajnović et al. 2006; Coccato et al. 2013), i.e. symmetric positions have equal velocities with opposite sign ($\tilde{v}(\phi) = -\tilde{v}(\phi + \pi)$). Deviations from this behavior arise from perturbations from the equilibrium status that may be due to interaction or merger episodes. If one of these processes, which plays a role in the formation and evolution of early type galaxies, occurred relatively recently (a few Gyrs ago), it is likely that some signatures in the kinematics and orbital structure of the galaxy are still observable, especially in the halo where the dynamic time-scales are longer.

The requirement of point-symmetry on $\tilde{v}(R, \phi)$, namely $\tilde{v}(R, \phi) - a_0 = -[\tilde{v}(R, \phi + \pi) - a_0]$, allows only odd values to n .

In addition, the expansion in equation (5) can be rewritten in a more direct way, as a rotation around the kinematic axis plus higher order modes. This is achieved through a re-phasing such that

$$\begin{aligned} \tilde{v}(R, \phi) = & a_0(R) + \sum_{k=1,3,\dots} c_k(R) \cos(k\phi - k\alpha) + \\ & + \sum_{k=1,3,\dots} d_k(R) \sin(k\phi - k\alpha) \end{aligned} \quad (6)$$

Table 1. Properties of the ETG sample analyzed in this paper, and list of references

Galaxy NGC	$M_K^{(a)}$ [mag]	D ^(b) [Mpc]	class ^(c)	$R_e^{(d)}$ [arcsec]	$R_{\max}/R_e^{(e)}$	$PA_{\text{phot}}^{(f)}$ [degrees]	$\epsilon^{(g)}$	$N_{\text{PNe}}^{(h)}$	References ⁽ⁱ⁾ PN data	References ^(l) abs.line data
0584	-24.23	20.2	F	33 (1)	7.4	63	0.339	25	(7)	(21)
0821	-23.99	23.4	F	40 (2)	4.8	31.2	0.35	122	(8)	(18);(40);(41)
1023	-23.89	10.5	F	48 (2)	6.8	83.3	0.63	181	(8);(9)	(18);(30)
1316	-26.02	21.0	F	109 (3)	4.7	50	0.29*	737	(10)	(22)
1344	-24.21	20.9	F	30 (4)	7.8	167	0.333	192	(12)	(23)
1399	-25.29	20.9	S	127 (3)	4.	110	0.1*	145	(11)	(24)
2768	-24.77	22.4	F	63 (2)	6.2	91.6	0.57	312	(9)	(18);(31)
2974	-23.76	22.3	F	38 (2)	5.8	44.2	0.37	22	(7)	(18)
3115	-24.02	9.5	F	93 (6)	4.7	43.5	0.607	183	(9)	(18);(25)
3377	-22.78	11.0	F	35.5 (2)	7.7	46.3	0.33	136	(8)	(18); (33)
3379	-23.80	10.3	F	40 (2)	5.3	68.2	0.13*	189	(8)	(19);(32);(40)
3384	-23.51	11.3	F	32.5 (2)	6.8	53	0.5	85	(9)	(19)
3489	-23.04	12.0	F	22.5 (2)	4.8	70.5	0.45	57	(9)	(19)
3608	-23.69	22.8	S	29.5 (2)	8.2	82	0.2*	92	(8)	(18)
3923	-25.33	23.1	S	86.4 (1)	4.9	48	0.271	99	(15)	(26)
4278	-23.80	15.6	F	31.5 (2)	7.6	39.5	0.09*	69	(7)	(18)
4339	-22.62	17.0	F	30 (2)	3.	15.7	0.07*	44	(7)	(20);(38)
4365	-25.19	23.1	S	52.5 (2)	5.6	40.9	0.24*	227	(7)	(18)
4374	-25.12	18.5	S	52.5 (2)	5.9	128.8	0.05*	445	(8)	(18)
4472	-25.73	16.7	S	95.5 (2)	8.4	154.7	0.19*	431	(7)	(20);(37)
4473	-23.76	15.2	F	27. (2)	5.6	92.2	0.43	153	(7)	(18)
4494	-24.17	17.1	F	49 (2)	4.8	176.3	0.14*	255	(8)	(18);(36)
4552	-24.32	16.0	S	34. (2)	9.2	132	0.11*	227	(7)	(19);(38)
4564	-23.10	15.9	F	20.5 (2)	6.5	48.5	0.53	47	(8)	(18)
4594	-24.93	9.5	F	102 (5)	4.	88	0.521	258	(16)	(27)
4636	-24.35	14.3	S	89. (2)	3.	144.2	0.23*	189	(7)	(20);(39)
4649	-25.35	16.5	F	66 (2)	4.5	91.3	0.16*	281	(13)	(18);(34)
4697	-24.14	12.5	F	61.5 (1)	4.5	67.2	0.32	525	(14)	(18);(35)
4742	-22.60	15.8	F	14.4 (4)	13.1	80	0.351	64	(7)	(28)
5128	-24.16	4.1	F	162.6 (1)	11.9	30	0.069*	1222	(17)	(29)
5846	-25.04	24.6	S	59 (2)	4.3	53.3	0.08*	118	(8)	(18)
5866	-23.99	14.8	F	36 (2)	9.4	125	0.58	150	(7)	(18)
7457	-22.38	12.9	F	36 (2)	3.2	124.8	0.47	108	(9)	(18)

Notes.

^(a) M_K is the total absolute luminosity in the K band. These values are obtained from the total apparent total magnitudes K_T of the 2MASS atlas (Skrutskie et al. 2006) using the distance D, and correcting for foreground galactic extinction A_B (Schlegel et al. 1998): $M_K = K_T - 5 \log_{10} D - 25 - A_B/11.8$. We assume $A_B/A_K = 11.8$, consistently with Cappellari et al. (2011). The K_T magnitudes are from integrating the surface brightness profiles ($\propto \exp(-r/r_a)^{1/\beta}$), extrapolated from the 20 mag/arcsec² isophote to $\sim 5r_a$ (Jarrett et al. 2003)

^(b) Distances of galaxies derived from surface brightness fluctuation method. Whenever possible we adopt the distance moduli measured by Blakeslee et al. (2009, B09), otherwise we used the values from Jensen et al. (2003, J03) or from Tonry et al. (2001, T01). The distance moduli from J03 were rescaled to the same zero-point calibration of B09 by applying a shift of +0.1 mag, while the distance moduli from T01 were zero-point- and bias-corrected using the formula from Blakeslee et al. (2010), and the data quality factor Q given by T01.

^(c) The sample is divided into slow (S) and fast rotators (F), according to the definition of Emsellem et al. (2011), from the kinematics within $1R_e$.

^(d) Adopted effective radius. The index in parenthesis corresponds to the refence: (1) Ho et al. (2012), (2) Cappellari et al. (2011), (3) Caon et al. (1994), (4) Blakeslee et al. (2001), (5) Kormendy & Westpfahl (1989), (6) Capaccioli et al. (1987).

^(e) Mean radius of the last radial bin in units of effective radii.

^(f) Average value of the photometric position angle and

the ^(g) ellipticity (ϵ), from Krajnović et al. (2011) (within $2.5 - 3R_e$) and Ho et al. (2011) (in the outer regions, where they converge to a constant value). ^(*) These objects are divided into spherical radial bins ($\epsilon = 0$), see section 3.2.

^(h) Number of detected PNe.

⁽ⁱ⁾ References for the PNe datasets: (7) are new ePN.S catalogs presented in Arnaboldi et al. (in prep.), (8) Coccato et al. (2009), (9) Cortesi et al. (2013a), (10) McNeil-Moylan et al. (2012), (11) McNeil et al. (2010), (12) Teodorescu et al. (2005), (13) Teodorescu et al. (2011), (14) Méndez et al. (2009), (15) unpublished data from counter dispersed imaging (Arnaboldi et al. in prep.), (16) unpublished data from narrow band imaging and spectroscopic follow up (Arnaboldi et al. in prep.), (17) Peng et al. (2004) and Walsh et al. (2015)

^(l) References for absorption line data: kinemetry from (18) Foster et al. (2016) on SLUGGS+ATLAS^{3D} data, from (19) Krajnović et al. (2008) and from (20) Krajnović et al. (2011), major axis long slit spectroscopy from (21) Davies & Illingworth (1983), (22) Bedregal et al. (2006), (23) Teodorescu et al. (2005), (24) Saglia et al. (2000) and Scott et al. (2014), (25) Norris et al. (2006), (26) Carter et al. (1998), (27) Kormendy & Illingworth (1982), (28) Davies et al. (1983), (29) Marcelin (1983), (30) Simien & Prugniel (1997c), (31) Simien & Prugniel (1997a), (32) Statler & Smecker-Hane (1999), (33) Coccato et al. (2009), (34) De Bruyne et al. (2001), (35) de Lorenzi et al. (2008), (36) Napolitano et al. (2009), (37) Veale et al. (2017), (38) Simien & Prugniel (1997b), (39) Pu & Han (2011), (40) Weijmans et al. (2009), (41) Forestell & Gebhardt (2010).

with

$$\begin{cases} a_n = c_n \cos(n\alpha) - d_n \sin(n\alpha) \\ b_n = c_n \sin(n\alpha) + d_n \cos(n\alpha) \end{cases} \quad (7)$$

The phase α can be chosen so that the amplitude of the first order sine term is 0: $d_1 = 0$ if $\alpha = \arctan(b_1/a_1)$. This implies that

$$\begin{aligned} c_1 &= \sqrt{a_1^2 + b_1^2} \\ c_k &= \frac{b_k}{\sin(k\alpha)} - \frac{d_k}{\tan(k\alpha)} \\ d_k &= \left(\frac{b_k/a_k - \tan(k\alpha)}{1 + \tan(k\alpha)b_k/a_k} \right) c_k \end{aligned} \quad (8)$$

and

$$\begin{aligned} \tilde{v}(R, \phi) &= a_0(R) + c_1(R) \cos(\phi - \alpha(R)) + \\ &+ \sum_{k=3,5,\dots} c_k(R) \cos(k\phi - k\alpha(R)) + \sum_{k=3,5,\dots} d_k(R) \sin(k\phi - k\alpha(R)) \end{aligned} \quad (9)$$

In this notation α coincides with the position angle of the kinematic major axis, PA_{kin} , a_0 is the mean velocity of the PNe in the bin, and c_1 is the amplitude of the projected rotation, V_{rot} . The amplitudes of the higher order harmonics, c_k and d_k , are corrections that account for deviations of the galaxy motion from the simple cosine rotation.

The kinematic quantities PA_{kin} and V_{rot} obtained fitting the model in equation (9) on the smoothed velocity fields are comparable to the results from a kinemetry fit on IFS data (Krajnović et al. 2006, 2011; Foster et al. 2016). However, we do not apply kinemetry, because this would mean fitting ellipses to the PN smoothed velocity fields. Since these have been derived from small samples of discrete tracers which, by nature, have lower spatial resolution and signal-to-noise ratio, a more straightforward approach is required.

3.2.2. Errors on the fitted parameters

The errors on the fitted parameters, $a_0(R)$, $PA_{\text{kin}}(R)$, V_{rot} , $c_k(R)$, and $d_k(R)$, are evaluated via Monte Carlo simulations: the 100 simulated datasets produced for deriving the errors on $\tilde{v}$ and $\tilde{\sigma}$ (see section 3.1) are divided into radial bins and modeled with equation (9). The errors are the standard deviations on the fitted parameters.

4. Systemic velocity subtraction

A measure of the systemic velocity of the galaxies is provided by the fit of the PN smoothed velocity field in radial bins with the harmonic expansion in equation (9). The bins are built as described in section 3.2 and the adopted geometry for each galaxy (i.e. ellipticity, ϵ , and photometric position angle, PA_{phot}) is listed in table 1. The $a_0(R)$ parameter, in fact, represents the mean velocity of the tracers in the radial bin with radius R . When the galaxy does not display kinematic substructures (bulk motions), this mean velocity is an estimate of the systemic velocity of the galaxy which is constant with radius for a gravitationally bound system.

Since the PNe are not distributed uniformly on the sky, $a_0(R)$ gives actually a more precise evaluation of the systemic velocity than a straight average of the measured LOS velocities. The fit of

equation (9) removes any contribution to the mean from rotation and is not sensitive to azimuthal completeness. Hence we can perform the fit leaving the parameter $a_0(R)$ free to vary in each bin. We find that $a_0(R)$ is generally constant with radius within the errors. Therefore we adopt, for each galaxy, a mean systemic velocity, V_{sys} , defined as a mean of the $a_0(R)$ values, weighted with the errors on the fit:

$$V_{\text{sys}} = \frac{\sum_{\text{bins}} a_0(R_{\text{bin}}) / \Delta a_0^2(R_{\text{bin}})}{\sum_{\text{bins}} 1 / \Delta a_0^2(R_{\text{bin}})} \quad (10)$$

We conservatively consider as error on V_{sys} , ΔV_{sys} , the mean of the errors $\Delta a_0(R)$, since the single measures of $a_0(R)$, coming from a smoothed velocity field, are not independent quantities.

We find that $a_0(R)$ does sometimes display a trend with radius within the errors. This is due to the interplay between spatial inhomogeneities and smoothing, which may result in a slight asymmetry of one side of the galaxy with respect to the other. This effect naturally disappears as soon as the catalogs are folded by point-symmetry transformation (see section 5), but we keep track of it in the uncertainties, by adding in quadrature the scatter of the $a_0(R)$ values to the error ΔV_{sys} .

NGC 1316 and NGC 5128 are treated separately with respect to the other galaxies. Their fitted $a_0(R)$ are constant in most radial bins, but they deviate in localized bins from this constant by more than twice the errors. At these radii the galaxies display important features in their velocity fields which cause an offset of the average velocity from the systemic value. We masked the most irregular bins and use the fitted $a_0(R)$ on the others to compute the mean V_{sys} .

The measured values of V_{sys} for all the galaxies are reported in table 2. We do not observe any systematic bias in the measured values, and they all agree within twice the error on V_{sys} with the literature. Hereafter we will refer to the baricentric velocities using V , and to the smoothed baricentric velocities using $\tilde{V}$.

5. Point symmetry analysis of the sample

In this section we investigate whether the galaxies in the ePN.S sample show any deviation from point-symmetry.

We studied the point-symmetry of the velocity fields of the galaxies by comparing the velocities $\tilde{V}(R, \phi)$ with $0 \leq \phi < \pi$ with those with $\pi \leq \phi < 2\pi$, changed in sign, in each radial bin. Asymmetries in the velocity fields are visible where these quantities significantly differ from each other. Figure 1 shows a few examples of this analysis. NGC 4649 is point-symmetric, while the others show significant deviations. In the sample of 33 galaxies, 5 are found to be non-point-symmetric: NGC 1316, NGC 2768, NGC 4472, NGC 4594 and NGC 5128. The others are consistent with point-symmetry, so for these systems we used the folded catalogs to reconstruct the final velocity and velocity dispersion fields, as described in section 6.1.

Galaxies for which we find evidence for asymmetries are those with the richest PN catalogs. For these objects the kinematic details are best recovered. Since the mean velocity fields are the result of a smoothing procedure, their point symmetry does not rule out kinematic asymmetries on smaller scales.

5.1. Testing the significance of the deviations from point-symmetry

We can evaluate the significance of the observed deviations from point symmetry by using models of the galaxies, built as described in appendix A.3. These are built using the positions (x, y)




Fig. 1. Mean velocity $\bar{V}(R, \phi)$ as a function of the position angle ϕ , folded around π , for each radial bin of NGC 1316, NGC 2768, NGC 4472, NGC 4594, NGC 4649, and NGC 5128. The black solid line is the fitted point-symmetric rotation model. The red points are the PNe with position angle $0 \leq \phi < \pi$; the blue points are those located at $\pi \leq \phi < 2\pi$, with their velocity changed in sign and position angle shifted of π , i.e. $V(\phi)$ are compared with $-V(\phi + \pi)$. The overlap of PNe of opposite sides shows possible asymmetries in the velocity fields. NGC 4649 has rather symmetric PN smoothed velocity field. NGC 2768 and NGC 4594 show localized small scale deviations from point symmetry, that do not influence the kinematic analysis. NGC 1316, NGC 4472, and NGC 5128 have non-point-symmetric velocity fields.

of the PNe from the real dataset, and, by construction, have a point symmetric mean velocity field. If the local deviations from point symmetry that we observe appear also in the smoothed velocity fields of the models, then we know that they are artifacts of the smoothing over that particular spatial distribution, and not properties of the intrinsic galaxy velocity field. We considered as possible kinematic features in the velocity fields any groups of at least three tracers whose velocity $\bar{V}$ deviates more than twice the errors from the fitted point symmetric model. The fitted harmonic expansion generally provides a consistent description of the galaxy velocity field, especially where the spatial distribution of the tracers is not azimuthally complete. For these groups of stars we computed the distribution of their deviations in 100 models. These distributions will give the probability of occurrence of the features as statistical fluctuations.

We found that the features in NGC 2768 and NGC 4594 have a probability $< 1\%$ to happen in the symmetric models so they are likely real. Those in NGC 2768 may be related to the

stream/shell seen in deep optical images (e.g. Duc et al. 2015)². The features in NGC 4594 are more likely due to extinction effects from its dusty disk, which hampers the detection of a complete sample of PNe in that area.

For NGC 1316, NGC 4472, and NGC 5128 the velocity offsets and the phase-angle shifts of the $\bar{V}(R, \phi)$ in ($0 \leq \phi < \pi$, red in figure 1) with respect to ($\pi \leq \phi < 2\pi$, blue in figure 1) cannot be reproduced by the point symmetric models. These galaxies are well known recent mergers. Their halos are dominated by the recently accreted component which is not yet in a phase-mixed equilibrium with the surroundings and hence it still maintains peculiar kinematics (see also appendix B).

One may be tempted to identify the groups of PNe whose velocity significantly deviate from the model as those associated to the structure, but we need to keep in mind that their velocities are the result of an averaging procedure, and that the different kinematic components can only be separated by analyzing the full

² <http://www-astro.physics.ox.ac.uk/atlas3d/>

phase space (see e.g. the GMM modeling of Longobardi et al. 2015); such a study is beyond the scope of this paper.

6. The halo kinematics of ETGs

6.1. Velocity fields

A point symmetric system is, by definition, such that each point of the phase space (x, y, V) has a point-reflected counterpart $(-x, -y, -V)$. For the galaxies that do not show any significant deviation from point symmetry (section 5), we assume that point symmetry holds. In these cases we can double the number of data-points by adding to the observed dataset its mirror dataset, and creating in this way a folded catalog (e.g. Arnaboldi et al. 1998; Napolitano et al. 2001; Peng et al. 2004; Coccato et al. 2009). This helps in reducing the fluctuations in the recovered velocity fields. The results obtained using the doubled catalogs are consistent with those from the unfolded datasets within the errors. Therefore for the galaxies consistent with point symmetry we will use the folded catalogs to produce the final mean velocity fields; for the others (i.e. NGC 1316, NGC 2768, NGC 4472, NGC 4594 and NGC 5128) the original catalogs are used.

Figure 2 shows the result for two galaxies with a similar number of tracers, NGC 4494 and NGC 4552. Both are point symmetric, so the velocity fields in figure 2 are built using the folded catalogs. NGC 4494 is a fast rotator showing some rotation also in the halo. Its velocity dispersion field reveals that σ decreases with radius. The slow rotator NGC 4552, by contrast, displays increasing rotation velocity about two perpendicular axes, and increasing velocity dispersion with radius.

The smoothed velocity fields for all the galaxies of the ePN.S sample are shown in appendix C. For a more immediate visualization we present interpolations of the velocity fields, based on computing $\tilde{V}$ and $\tilde{\sigma}$ on a regular grid. The kinematics typically extends to $\sim 6R_e$, covering a minimum of $3R_e$ to a maximum of $13R_e$. The adopted R_e values are listed in table 1. Table 1 also shows the mean radius of the last radial bins, in which we can statistically determine $\tilde{V}$ and $\tilde{\sigma}$.

The typical errors on the mean velocities and on the velocity dispersions, evaluated with Monte Carlo simulations, range from 10 to 40 km s⁻¹, being smaller for galaxies with a larger number of tracers and higher $\tilde{V}/\tilde{\sigma}$.

These errors on the mean velocity fields, the mean errors on the radial velocity measurement, the kernel parameters A and B used in the smoothing procedure, and the systemic velocities subtracted are reported in table 2.

A visual comparison with the kinematic maps published by the ATLAS^{3D} (Krajnović et al. 2011) and SLUGGS (Arnold et al. 2014; Foster et al. 2016) surveys shows a general good agreement for all the galaxies in the regions of overlap; see the appendix B for a detailed description of the individual objects.

6.2. Kinematic parameters

We can quantify the properties of the reconstructed mean velocity fields by evaluating the amplitude of rotation, the variation of the PA_{kin} with radius, and the possible misalignments with PA_{phot} . Therefore we model the velocities in each radial bin as a function of the position angle ϕ (positive angles are from North to East, with the zero at North) with the rotation model in equation (9).

The Fourier expansion can in practice be truncated at the third order as the higher order amplitudes are generally zero




Fig. 2. Top row: smoothed velocity fields of NGC 4494 and NGC 4552; bottom row: velocity dispersion fields. The fields are built using the *folded catalogs*, but only the positions of the actual PN data points are shown. The images in the background are from the Digitized Sky Survey (DSS); north is up, east is left.

within the errors. The systemic velocity has already been subtracted, so

$$\tilde{V}(R, \phi) = V_{\text{rot}}(R) \cos(\phi - PA_{\text{kin}}(R)) + s_3(R) \sin(3\phi - 3PA_{\text{kin}}(R)) + c_3(R) \cos(3\phi - 3PA_{\text{kin}}(R)) \quad (11)$$

and the only free parameters are PA_{kin} , the projected rotation amplitude V_{rot} , and the amplitudes s_3 and c_3 .

We divided the sample of ETGs into fast and slow rotators according to the definition of Emsellem et al. (2011). In figure 3 we show separately for both families the fitted parameters V_{rot} , s_3 and c_3 , as functions of the major axis distance R in units of R_e . This is a reasonable choice in case of flattened systems rotating along the photometric major axis. In case of misalignment or twist of the PA_{kin} , R does not correspond to the position of the peak in $\tilde{V}$ but to the major axis of the elliptical bin in which the amplitude $\tilde{V}$ is calculated. Figure 4 shows the misalignment Ψ of PA_{kin} with respect to PA_{phot} , $\Psi = PA_{\text{kin}}(R) - PA_{\text{phot}}$. If the difference $PA_{\text{kin}}(\text{bin1}) - PA_{\text{phot}}$ (where $PA_{\text{kin}}(\text{bin1})$ is the value measured in the first radial bin), is greater than 90 degrees, we define Ψ as $PA_{\text{kin}}(R) - PA_{\text{phot}} - \pi$. Since PA_{phot} is a constant value for each galaxy, a variation of PA_{kin} with radius corresponds to a variation of Ψ . We do not use the definition of Franx et al. (1991), $\sin \Psi = \sin(PA_{\text{kin}}(R) - PA_{\text{phot}})$, as it does not allow the description of large position angle twists. The values and the references for the PA_{phot} used are in table 1.

Both V_{rot} and Ψ are compared with literature values in figures 3 and 4. When available, we show the profiles from the kinematic analysis of Foster et al. (2016) on the SLUGGS+ATLAS^{3D} data, or the kinematic profiles from Krajnović et al. (2008). In these cases we rescale the radii of the profiles to major axis distances using the flattening $q_{\text{kin}} = q_{\text{phot}}$ given by Foster et al.

(2016), or $\langle q_{\text{kin}} \rangle$ given by Krajnović et al. (2008). For the other galaxies we plot the corresponding quantities from the kinematics of Krajnović et al. (2011, namely k_1^{max} and PA_{kin} from their table D1), or the kinematic profiles from long slit spectroscopy similarly rescaled (references in table 1). While comparing with the literature, it is important to note the following effect. A kinematic measurement from a slit along the major axis of an edge-on fast rotating galaxy will give high velocities and low dispersions. On the other hand, the PN velocity fields are the results of a smoothing procedure, which averages together PNe belonging to the thin disk with PN belonging to the spheroid. This might result in a systematically lower rotation and higher velocity dispersion (see equation 2) in the PN velocity fields. A disk/spheroid decomposition of the PNe in some ETGs has already been performed by Cortesi et al. (2013b), and it is beyond the scope of this paper to extend this to all fast rotators. In addition, if the number of tracers or the ratio V/σ is low, our kinematic analysis provides a lower limit for the rotation velocity and an upper limit for the velocity dispersion. This issue is addressed in appendix A. In such cases, the kinematics traced by the PNe may systematically not match that from the integrated light as consequence of the discrete spatial sampling of the velocity field by the adopted tracers.

The higher order harmonics amplitudes s_3 and c_3 differ from zero whenever the smoothed velocity field deviates from simple disk-like rotation, i.e. if it is cylindrical (see for example the case of NGC 3115), or in correspondence to components rotating at different position angles (e.g. NGC 4649).

Misalignments and twists of the PA_{kin} are typically displayed by triaxial galaxies (Stark 1977; de Zeeuw & Franx 1991); see section 6.4. Figure 4 shows that both fast and slow rotators can have radial variation of the PA_{kin} or a constant non-negligible Ψ . These galaxies may have a triaxial halo. A few galaxies instead have kinematically decoupled halos with respect to the regions $\lesssim 2R_e$. Section 6.4 validates these results for each galaxy using models.

The asymmetric galaxies (i.e. NGC 1316, NGC 2768, NGC 4472, NGC 4594, and NGC 5128) are, by construction, not well represented by the point symmetric model and increasing the number of harmonics does not improve the quality of the fit. We can however still use the fitted parameters to obtain an approximate description of the shape of their velocity field.

6.3. Velocity dispersion profiles

Figure 5 shows the velocity dispersion profiles, azimuthally averaged in radial bins. These have been calculated using two different methods. In the first we used the interpolated velocity dispersion field $\tilde{\sigma}(x, y)$ in elliptical annuli of growing radius, with position angle and ellipticity as in table 1. The values shown in the plots (solid lines) are the means of the values over each ellipse, and the errors, calculated through Monte Carlo simulation (section 3.2.2), are taken conservatively as the means of the simulated errors for all points on the elliptical annulus (dotted lines). The second method is binning the measured radial velocities v of the PNe in the radial bins built as described in section 3.2. The PN catalogs are folded by point symmetry and the dispersion σ_{bin} with respect to the weighted mean velocity is computed in each bin. The weights are computed from the measurement errors. The errors on the dispersion are given by the expression: $\Delta\sigma_{\text{bin}} = \sigma_{\text{bin}} / \sqrt{2(N_{\text{bin}} - 1)}$, where N_{bin} is the number of PNe in each bin. The values and the trends given by the two methods are generally in good agreement.

The profiles obtained are compared with dispersion profiles from integrated light (red stars). For the galaxies in common with the SLUGGS survey, we show the profiles from the kinematic analysis of Foster et al. (2016) on the SLUGGS+ATLAS^{3D} data in elliptical radial bins. For NGC 3384, NGC 3489, NGC 4339, NGC 4552, and NGC 4636, we extracted the azimuthally averaged profiles from the ATLAS^{3D} data (Cappellari et al. 2011; Emsellem et al. 2011) in elliptical bins (geometry in table 1). For the other galaxies we show the velocity dispersion along the PA_{phot} from long slit spectroscopy (references in table 1).

Our dispersion profiles generally compare well with the literature in the regions of overlap (typically $R \lesssim 2R_e$). The results are described in section 7, separately for fast and slow rotators.

We tested whether it is possible that the large scale trends in the velocity dispersion profiles are the result of statistical and smoothing effects, by using 100 models of the galaxies, built as described in section A.3, created with a constant dispersion profile with radius. The velocity dispersion profiles are recovered with the same procedure as for the measured PN sample. We find that although artificial local structures may sometimes appear in the velocity dispersion maps, they are not such as to influence the trends with radius of the large scale velocity dispersion fields, and typically the error bars from Monte Carlo simulations give a good estimate of the uncertainties.

6.4. Triaxiality

Significant twists of the PA_{kin} , as well as its departures from PA_{phot} imply an intrinsic triaxial shape for the system. In an axisymmetric object both the projected photometric minor axis and the intrinsic angular momentum are aligned with the symmetry axis of the system, while in a triaxial galaxy the rotation axis can be in any direction in the plane containing both the short and long axis. This is because in a triaxial potential the main families of stable orbits are tube orbits which loop around the minor (z-tube orbits) or the major axis (x-tube orbits). The relative number of z- and x-tube orbits determines the direction of the intrinsic angular momentum. Thus, depending on the variation of this ratio with radius we can have that

- the measured PA_{kin} shows a smooth radial twist (e.g. NGC 4552 in figure 4, see section 6.4.1);
- PA_{kin} has a sudden change in direction (i.e. the galaxy has a kinematically decoupled halo, like NGC 1399, see section 6.4.2);
- PA_{kin} has a constant misalignment with respect to the PA_{phot} (e.g. NGC 3923, see section 6.4.3).

Therefore we consider as possible triaxial galaxies the objects displaying at least one of those features in their velocity fields with statistical significance determined by our MC modeling. Table 3 provides a summary of the results, which are discussed in the following sections.

6.4.1. Galaxies with radial variation in the kinematic position angle

Figure 4 shows that the fitted PA_{kin} may show a smooth variation with radius. This happens in NGC 3379, NGC 4649, NGC 5128, and NGC 5866, among the fast rotators, and in NGC 3608, NGC 4472, and NGC 4552 among the slow rotators.

We tested whether the variation with radius of the PA_{kin} for a galaxy is an artifact of the combination of a small number of tracers and the smoothing procedure. We looked at the fitted PA_{kin} in




Fig. 3. Fitted rotation velocities $V_{\text{rot}}(R)$ (full circles) and third order harmonics amplitudes, $c_3(R)$ in green and $s_3(R)$ in orange, as functions of the major axis distance for slow and fast rotators. The comparison values for V_{rot} from absorption line data from the literature are shown with colored stars. Whenever available we show kinematic profiles (light blue stars); for the other galaxies we show velocities from long slit spectroscopy along the photometric major axis (red stars) or minor axis (green stars). The references are in table 1. For NGC 4594 we show in addition the stellar kinematics from a slit along PA_{phot} , offset by 30 arcsec (purple stars). The dashed vertical lines for the fast rotators show the disk half light radius $R_{1/2}$, see section 7.2.1. The red dot-dashed vertical lines report the radial transition range $R_T \pm \Delta R_T$, see section 8.4.




Fig. 4. Misalignments $\Psi(R)$ (full circles) as functions of the major axis distance for fast and slow rotators. The horizontal solid line shows the $\Psi = 0$ axis. The light blue stars are the Ψ values calculated on PA_{kin} from the kinemetry of Foster et al. (2016), Krajnović et al. (2008), and Krajnović et al. (2011); for the other galaxies the PA_{kin} is not previously available in the literature. The red dot-dashed vertical lines report the radial transition range $R_T \pm \Delta R_T$, see section 8.4. Signatures of triaxial halos are seen in NGC 0821, NGC 1344, NGC 1399, NGC 3379, NGC 3608, NGC 3923, NGC 4365, NGC 4374, NGC 4472, NGC 4473, NGC 4552, NGC 4636, NGC 4649, NGC 4742, NGC 5128, NGC 5846, and NGC 5866.




Fig. 5. Azimuthally averaged velocity dispersion profiles as functions of the major axis distance in unit of R_e for slow and fast rotators (full circles). The gray solid lines represent the interpolated profiles and the 1σ level (dashed lines). The stars show dispersion from integrated light: when available we plot the kinematic analysis of Foster et al. (2016) on the SLUGGS+ATLAS^{3D} data in elliptical bins (light blue stars); for NGC 3384, NGC 3489, NGC 4339, NGC 4472, NGC 4552, and NGC 4636, we show azimuthally averaged profiles from ATLAS^{3D} data (Cappellari et al. 2011; Emsellem et al. 2011); for the other galaxies we show data from long slit spectroscopy along PA_{phot} (red stars, references in table 1). For NGC 4594 we also plot the stellar velocity dispersion profile in a slit parallel to the major axis but 30 arcsec offset from the center (purple stars).

100 models of each galaxy, built as described in appendix A.3, which have by construction the PA_{kin} of the mean velocity field aligned with PA_{phot} . For each radial bin we computed the probability of obtaining the observed misalignment Ψ from the distribution of the misalignments in the models.

We found that the probability of observing in models of NGC 3379 a twist of 71 ± 28 is $\sim 3\%$, and it is $\sim 2\%$ for the twist of 20 ± 13 degrees of NGC 3608. For the other galaxies, none of the 100 models produces the observed trends of PA_{kin} .

For NGC 4472 the determination of PA_{kin} is influenced by the kinematics of the in-falling satellite UGC 7636. An inspection of its smoothed velocity field suggests that the main body of the galaxy has approximately major axis rotation, once the PNe of the satellite are excluded. Nevertheless we include this galaxy in the sample of potentially triaxial galaxies, and we refer to Hartke et al. (in prep.) for a more detailed study.

NGC 5128 shows non point-symmetric kinematics, and rotation along both the photometric major and minor axes. The high number of tracers available for this galaxy (1222 PNe) makes this kinematic signature unambiguous.

NGC 4494 has a velocity field similar to NGC 3379, with the principal rotation along PA_{phot} and a fainter component rotating along the minor axis. Their superposition does not show a statistically significant twist of the PA_{kin} .

NGC 1316 shows a small but significant jump of the PA_{kin} at $\sim 200 - 250$ arcsec, that is related to the perturbed kinematics of this galaxy.

Therefore, excluding NGC 1316 that has local and irregular variations of PA_{kin} , there are 7 galaxies in the sample showing significant kinematic twist, of which 4 are fast and 3 slow rotators.

6.4.2. Galaxies with kinematically decoupled halos

Galaxies with kinematically decoupled halos are galaxies whose outskirts rotate about a different direction than their inner regions, hence PA_{kin} shows a step function with radius. NGC 1399 and NGC 4365 both show this feature beyond $2R_e$.

NGC 1399 is found to be slowly rotating around its PA_{phot} (i.e. $\Psi \sim 90^\circ$) at 31 ± 30 km s⁻¹ inside $1R_e$, in very good agreement with the integral field spectroscopic data of Scott et al. (2014).

The halo PA_{kin} , by contrast, is almost aligned with the PA_{phot} (i.e. $\Psi = 0$). We studied whether such a misalignment is an artifact of our procedure, using models that mimic the inner kinematics, i.e. with PA_{kin} aligned with the photometric minor axis. The probability of measuring a misalignment of the halo similar to the observed one is 2%.

The PA_{kin} of NGC 4365 is ill-constrained in the innermost regions where the kinematics is compatible with no rotation. At the center we do not recover the rolling about the minor axis visible in velocity fields from absorption line data (Emsellem et al. 2011; Arnold et al. 2014), because of smoothing over the inner velocity gradients. In these regions the bright background of the galaxy hampers the detection of PNe, and the resulting low number of tracers combined with the low V/σ leads to heavily smoothed velocities. We do detect a significant outer ($R \gtrsim 3R_e$) rotation of ~ 50 km s⁻¹ along PA_{phot} ($\Psi \sim 0^\circ$), misaligned with respect to the inner kinematics reported in the literature. So we built mock models as described in appendix A.3 but with PA_{kin} given by IFS data up to $1 R_e$ (references are given in table 1). We found that none of the models displays the observed step function in the PA_{kin} values.

We therefore conclude that the signature of a kinematically decoupled halo has a high probability to be real in both galaxies.

6.4.3. Galaxies with constant offset between photometric and kinematic major axis

The galaxies showing an approximately constant misalignment of the PA_{kin} with respect to the PA_{phot} are NGC 0821, NGC 1344, NGC 4473 and NGC 4742 among the fast rotators, and NGC 3923, NGC 4374, NGC 4636, and NGC 5846 among the slow rotators (see figure 4).

In these cases we can define the PA_{kin} using all the PNe, without radial binning. The derived quantities can be compared to the PA_{kin} measured in mock models built as described in appendix A.3.

For most of the listed galaxies none of the models reproduces the observed misalignments. NGC 0821 has a misalignment of 50 degrees with $\sim 4\%$ probability in the Monte Carlo models. Because of the smaller number of tracers, this probability is higher for NGC 4742 and NGC 5846 (17% and 9%, respectively). The misalignment of NGC 4473 has already been studied by Foster et al. (2013) who, using absorption line data, detected a significant rotation along the minor axis, and already proved this object to be triaxial.

Therefore in the ePN.S sample a total of 8 galaxies, 4 fast and 4 slow rotators, show a significant constant misalignment of PA_{kin} with PA_{phot} .

6.4.4. Summary

We can conclude that a total of 17 galaxies (50%) of the ePN.S sample show smoothed velocity fields that reveal their triaxial nature. 7 objects (4 fast and 3 slow rotators) have significant kinematic twists, and 8 (4 fast and 4 slow rotators) show a significant constant misalignment of PA_{kin} with PA_{phot} . In addition two slow rotators have a kinematically decoupled halo.

The observed features are more than 2 sigma significant for most of the cases (1.3 sigma for NGC 4742 and 1.7 sigma for NGC 5846), and they are not effects of the folding operation on the catalogs nor of the smoothing procedure.

All in all, we found that all the slow rotators and 8 out of 24 fast rotators show indications of intrinsic triaxial morphology in the PN data.

7. Results per family

7.1. Slow rotators

In the sample of 33 galaxies 9 are slow rotating. Figure 3 shows that they typically display some more pronounced rotation at large radii when compared to rotation in their central regions as measured from absorption line spectroscopy. The PN velocity fields show gently increasing profiles for the V_{rot} amplitude which, eventually, flatten around 50 km s⁻¹. Twists or misalignments of the PA_{kin} are commonly observed, so that all the slow rotators show signatures of a triaxial halo (see figure 4). In particular, we found that the halos of NGC 1399 and NGC 4365 are kinematically decoupled with respect to the innermost regions as mapped by Scott et al. (2014) and Arnold et al. (2014). NGC 4472 has a non point-symmetric velocity field, perturbed by a recent accretion event. The complicated kinematics of the slow rotators is also reflected in the amplitudes of the third order harmonics, which describe the presence of additional kinematic components, and twists of the PA_{kin} .

The velocity dispersion profiles, shown in figure 5, are generally flat in the halo. Some galaxies (e.g. NGC 1399, NGC 3608) reach such a constant value around $\sim 1 - 2R_e$. Others (e.g. NGC 4374, NGC 4552) flatten only beyond $\sim 4 - 5R_e$, after a small increase. NGC 4636 is the only slow rotator showing a falling profile.

7.2. Fast rotators

In the ePN.S sample 24 galaxies are classified as fast rotators (table 3). Figures 3 and 4 show that the majority of the objects have regular rotation along the photometric major axis. The comparison between inner and outer parts reveals that the rotation amplitudes V_{rot} show declining trends with more or less steep gradients for 14 out of 24 fast rotators: some galaxies show very small or no rotation in the outskirts (e.g. NGC 4278 and NGC 4473), while others reach a minimum, after which their rotation increases (e.g. NGC 4494 or NGC 4697). Among the remaining galaxies, 3 have fairly constant profiles (NGC 2768, NGC 4564, and NGC 5866), and 3 show increasing rotation (NGC 3384, NGC 4742, and NGC 7457). For NGC 0584, NGC 2974, NGC 3489 and NGC 4339, the limited number of tracers allows the estimate of a lower limit for the rotation in the halo only.

The PA_{kin} is well aligned with the photometric major axis in the majority of cases, but 8 out of 24 galaxies display a kinematic twist (NGC 3379, NGC 4649, NGC 5128, and NGC 5866) or a constant misalignment with PA_{phot} (i.e. NGC 0821, NGC 1344, NGC 4473 and NGC 4742). The smoothed velocity fields of four galaxies (NGC 3379, NGC 4649, NGC 5128, and, marginally, NGC 4494, see figures in appendix C) show hints of additional components rotating along the minor axis of the system, while NGC 1344 has prolate rotation. All these features are generally interpreted as evidences of triaxiality of the systems (see section 6.4) and a third of the fast rotating galaxies of the sample displays them in the halo also.

The azimuthally averaged velocity dispersion profiles (figure 5) are found to be either constant (e.g. NGC 3377), or decreasing with radius. Some profiles decline gently (e.g. NGC 1344), while others decrease steeply in the halo (e.g. NGC 3379). This diversity between flat and falling profiles is also reflected in the variety of the V_{rms} profiles, as already observed by Coccato et al. (2009), and are the results of differences in the mass distributions (Gerhard et al. 2001; Romanowsky et al. 2003; Deason et al. 2012; Morganti et al. 2013; Cappellari et al. 2015; Alabi et al. 2016; Veale et al. 2018) or the presence of radial anisotropy in the orbits (Gerhard et al. 1998; de Lorenzi et al. 2009; Napolitano et al. 2009), which may contribute to a lower projected velocity dispersion. For some lenticular galaxies (e.g. NGC 1023 or NGC 3115), the two-dimensional velocity dispersion maps (see figures in appendix C) reveal the presence of the colder disk along its major axis, while the dispersion is higher along the minor axes. For NGC 1023 the presence of the disk is evident also in the azimuthally averaged velocity dispersion profile, which increases with radius as the contribution of the disk to the light decreases.

Among the fast rotating galaxies there are two mergers, NGC 1316 and NGC 5128, whose velocity fields are highly disturbed by the recent accretion events and are not very well described by point-symmetric rotation models (see appendix B for more details).

The comparison of the PN kinematics with integrated light data shows a general good agreement in the overlapping regions, confirming once again that the PNe are reliable tracers of the kinematics of the parent stellar population. The tension with the

stellar kinematics in cases of high rotation and low dispersion (see e.g. NGC 1023, NGC 2768, NGC 4594) is primarily related to the presence of a known near edge-on thin disk that dominates the major axis stellar kinematics but not that of the PNe. Because our velocity fields are the results of an averaging operation that does not distinguish the PNe from the disk from those belonging to the spheroid, the velocity gradients are underestimated. Likewise the discrepancy between the integrated light data of NGC 4494 and its PNe is explained by the disk fading into the spheroid at radii beyond the coverage of the stellar kinematics, combined with the lower spatial resolution of the PN smoothed velocity fields.

In addition, if the disk is thin or obscured by dust, the fraction of observed disk PNe is reduced, causing a drop in the measured rotation and a higher dispersion. An example is the case of NGC 4594. This galaxy has a dusty thin disk which affects the detections of PNe in that region, leading to heavily absorbed light in this component and hence inhibit the PNe detections in that region where the rotation is highest. The rotational velocity and the dispersion profiles from PNe agree well with slit data along a direction slightly offset (30 arcsec) from the disk plane (purple stars in figures 3, 5 and 6), but are offset with respect to the major axis profiles. With this in mind, we shall consider the smoothed velocity fields as giving a global description of the halo kinematics, but of not their small-scale spatial structures, unless the number of tracers is very large (as, for example, in the case of NGC 1316).

7.2.1. Embedded disk in fast rotators

The observed negative gradients of $V_{\text{rot}}(R)$ observed for many fast rotators has been interpreted as signatures of a disk embedded into a dispersion dominated spheroidal component (Coccato et al. 2009; Arnold et al. 2014). Using a photometric disk-spheroid decomposition and maximum-likelihood fit, Cortesi et al. (2013b) reconstructed the kinematics of the disk and the spheroid separately in 6 of the ePN.S lenticular galaxies (NGC 1023, NGC 2768, NGC 3115, NGC 3384, NGC 3489 and NGC 7457). Arnold et al. (2014) reproduced the rotation profile of the E5-6 galaxy NGC 3377 also performing a disk-bulge decomposition. These works showed that the variation in rotation reflects the transition between disk and bulge dominance in light, and their different spatial contribution in each galaxy explains the variety in the observed λ_R profiles. The detailed study of the λ_R profiles for the ePN.S sample is the subject of a separate paper (Coccato et al. in prep.).

Here we can verify this interpretation by estimating and comparing the "size" of the disks given by the photometry, and the radius at which we observe the decrease in rotation. Figure 6 shows a subsample of the galaxies with the highest number of tracers, in order to have the best possible statistics. Most of them show a decreasing rotation amplitude. We fitted their radially binned smoothed velocity fields with a cosine, whose position angle is aligned with the photometric major axis

$$\tilde{V}(R, \phi) = V_{\text{maj.ax.}}(R) \cos(\phi - PA_{\text{phot}}) \quad (12)$$

in order to extract the velocity profiles along the major axis. The fitted amplitudes are shown in figure 6. Overlaid in red is the stellar kinematics along the photometric major axis from literature slit data (references in table 1). The orange vertical lines indicate the characteristic scale of the disks. For all the galaxies shown, the quantity plotted is the disk half light radius: $R_{1/2} \simeq 1.67R_h$. R_h is the disk scale length from an exponential fit of the disk




Fig. 6. Rotation velocity profiles for the fast rotators with highest number of tracers along the major axis. The full circles show the fitted amplitude $V_{\text{maj,ax}}$ on the PN velocity fields. The red stars are stellar kinematics along the photometric major axis from the literature (references in table 1). The PN kinematics of NGC 4594 is compared with the stellar kinematics from a slit parallel to the major axis, offset by 30 arcsec (purple stars). The orange vertical lines mark the sizes of the photometric disks (see text). NGC 4494 shows a decrease in $V_{\text{maj,ax}}$, but there is no photometric evidence for a stellar disk. When the disk fades in the slowly rotating spheroid, the rotation velocity decreases. In addition, the PN rotation velocity may be lower than the values from absorption line spectroscopy at the same radii because of the smoothing that averages disk and spheroid PNe for near edge-on disks.

component (R_h from Krajnović et al. 2013; Cortesi et al. 2013b; Beletsky et al. 2011; Burkhead 1986; Laurikainen et al. 2010; Scorza et al. 1998; Pignatelli 1999, for figures 3 and 6).

We can see that the radial distance at which the decrease in the rotation occurs is consistent with the size of the disk. Therefore we interpret this behavior as the transition between a component that rotates fast and a dispersion dominated spheroid. For NGC 4494 there is no photometric evidence for a disk component. In this case the stellar disk might be absent or very faint if the galaxy is seen nearly face on ($\epsilon \sim 0.14$). In the other fast rotators that are not shown in figure 6, but show a drop in the amplitude of rotation, the comparison between slow rotating PN system and rapid rotation in the absorption line kinematics, which is related to an inner disk, suggests similar transition at smaller radii (see figure 3).

Figure 3 shows that the three galaxies with approximately constant V_{rot} profiles (NGC 2768, NGC 4564, and NGC 5866) do actually show a small decrease in rotation in correspondence to $R_{1/2}$, when compared with the values from absorption line

spectroscopy. At larger radii their fast-rotating spheroidal halo (NGC 2768, Cortesi et al. 2013b), or alternatively an unidentified outer disk, dominates the kinematics. Among the remaining three galaxies in figure 3 with increasing V_{rot} profiles, two of them (NGC 3384 and NGC 7457) have an extended disk component, to which most of the PNe belong (Cortesi et al. 2013b). NGC 4742, by contrast, does not have any photometric evidence for a disk component, hence the high rotation at large radii is likely associated with the spheroid.

7.3. Summary

Our results show that the kinematics of ETGs at large radii can be substantially different from that in the inner regions. For the slow rotators we observe a growth in the amplitude of rotation. For the fast rotators, this variation in the kinematics manifests as a decrease in the amplitude of rotation or a twist in the PA_{kin} . We interpreted this behavior as the transition from the inner disk




Fig. 7. $V/\sigma(R_e)$ from absorption line data compared with $V/\sigma(\text{halo})$ from PN data. References for the absorption line data are in table 1; note that for NGC 1399 and NGC 3923 data are available up to $R_e/2$ and $R_e/4$, respectively. The flattening of the ellipses used to plot the galaxies correspond to the ϵ values of table 1. Fast and slow rotators are shown with different colors, as are the triaxial fast rotators. The gray open ellipses represent the galaxies (all fast rotators) with fewer tracers, for which our analysis provides a lower limit for $V/\sigma(\text{halo})$. The solid line traces equal values of $V/\sigma(R_e)$ and $V/\sigma(\text{halo})$. The orange circles show the V/σ of simulated galaxies from Wu et al. (2014), divided by a factor 0.57 to rescale their two dimensional flux-weighted measurement to one-dimensional quantities (Cappellari et al. 2007).

component into the spheroidal halo, which is dispersion dominated and might deviate from axisymmetry.

Figure 7 illustrates this conclusion by comparing the V/σ ratio in the halo and in the inner regions (at $1R_e$). The $V/\sigma(R_e)$ values are derived by interpolating the V_{rot} and σ profiles from integrated light (shown in figures 3 and 5, see references in table 1) at $R = R_e$, while $V/\sigma(\text{halo})$ is the ratio of V_{rot} and σ estimated in the outermost radial bin of the PN velocity fields (see table 1 for the outermost mean radius).

All the slow rotators fall below the 1:1 line, showing higher rotational support at large radii. The spread of the fast rotators in the diagram reflects their different intrinsic structure and kinematics. The halos of most fast rotators have V/σ ratio similar to the $V/\sigma(\text{halo})$ of slow rotators. Among these the scatter in $V/\sigma(R_e)$ is probably driven by the presence of a more or less prominent disk component seen at different inclinations, and embedded into the dispersion-dominated spheroid. The flattening of the ellipses in figure 7, in fact, shows that galaxies with higher ϵ also display higher $V/\sigma(R_e)$.

A second group of fast rotators with high rotational support in the halo populate the diagram on the right of the 1:1 line. These galaxies are either dominated by disk rotation at all radii (NGC 3384 and NGC 7457), or have a rapidly rotating spheroid (NGC 2768, Cortesi et al. 2013b), or either of these (NGC 4564, NGC 4742).

The fast rotators with triaxial halos typically show equal values of $V/\sigma(R_e)$ and $V/\sigma(\text{halo})$, spanning all values in $V/\sigma(\text{halo})$. It is possible that the group of galaxies near the 1:1 line but without signature of triaxiality in the current data could show such signatures with higher resolution in the kinematics. Therefore we cannot presently determine whether these last two subset of fast rotators are different or not.

The orange full circles in the V/σ plot show the inner-halo kinematics from Wu et al. (2014), who studied the kinematics of ETGs in cosmological zoom simulations (Oser et al. 2010; Naab et al. 2014) out to $5R_e$. Their $V/\sigma(R_e)$ and $V/\sigma(5R_e)$ were divided by a factor 0.57 to rescale their two dimensional flux-weighted measurements to one-dimensional quantities (Cappellari et al. 2007). The comparison with the observations shows that these simulations do not adequately reproduce the observed properties of fast rotating galaxies, which span a much wider spectrum of kinematic properties at large radii.

8. Discussion

8.1. Halo rotation versus central rotation

The PN velocity fields show that ETGs may have very different kinematics at large radii compared to their inner regions. We found that slow rotators typically increase their rotation in the halo, while most of the fast rotators display a decrease of V_{rot} towards large radii.

The observed variety of radial trends for the rotation amplitudes is consistent with the different shapes of the λ_R parameter profiles observed in the smaller sample of Coccato et al. (2009), and confirmed by Arnold et al. (2014) and Foster et al. (2016). The study of the λ_R profiles for the current sample galaxies is the subject of a future paper (Coccato et al. in prep.).

The existence of these radial trends was recently questioned by Raskutti et al. (2014) and Boardman et al. (2017), who found no evidence for a change in the stellar kinematics beyond R_e . The divergence from our results may arise from the different radial coverage of their kinematics, which does not reach out to the distances where the drop in rotation typically occurs, $\sim 3R_e$. An example is the lenticular galaxy NGC 1023, in common between the two samples. For this galaxy Boardman et al. (2017) report a rising λ_R profile to almost $2R_e$, in agreement with our observation of a decrease in rotation beyond $3R_e$.

The onset of rotation of the slow rotators at larger radii, often along directions that do not coincide with the major axis of the galaxy, may trace the accreted stellar component, which still maintains a memory of the orbital angular momentum of the accreted progenitors.

8.2. Fast rotators with disks embedded in slowly rotating halos

The sharp drop in angular momentum of the fast rotators has been interpreted as the fading of an embedded disk-like structure of in-situ stars into a dispersion dominated spheroidal halo (Coccato et al. 2009; Arnold et al. 2014). We qualitatively verified this scenario in section 7.2.1, where we observed that the radius at which the rotation along the major axis drops is consistent with the dimension of the disk component.

Our results show that fast rotators come with a more or less extended disk component, embedded in a more or less prominent halo with its own kinematic signature. This is reflected in the distribution of these objects in the V/σ plane (figure 7). We found that the halos of most fast rotators have rotational support

comparable to that of the halos of slow rotators. A smaller group of fast rotators shows high rotation in the halo also.

This variety of kinematic properties is consistent with the variety of physical processes that may drive the evolution of these objects (minor mergers, major mergers that lead to a spin-up of the remnant, gas accretion, interactions with the environment, secular evolution; Duc et al. 2011; Naab et al. 2014; Penoyre et al. 2017; Smethurst et al. 2018) while preserving a rotating disk structure at the center. The comparison of our observations with simulated values from Wu et al. (2014) in figure 7 indicates, however, that those simulations do not reproduce the full diversity in the variety of these processes.

8.3. Signatures of triaxial halos

In section 6.4 we studied the velocity fields and their kinematic position angle profiles PA_{kin} , and linked the observed misalignments and twists with the triaxiality of the halos. Slow rotators are known to be mildly triaxial in their central regions, while the fast rotators are predominantly oblate (Weijmans et al. 2014; Foster et al. 2017). Kinematic twists and misalignments for the fast rotators are rare and modest in the central regions. By sampling the kinematics at more than twice the distances probed by previous studies, the PNe line-of-sight velocities show that these features become more frequent and pronounced in the outskirts. Based on these signatures, the halos of a third of the fast rotator galaxies are classified as triaxial, while the remaining ones are still consistent with being axisymmetric at the resolution of the ePN.S survey. For these objects the triaxiality of the halos is not ruled out. On the other hand, it is in principle possible that axisymmetric galaxies with a recent merger might display kinematic twists and misalignments because of the contribution of the unrelaxed accreted component to the PN velocity field. However, if the size of the accreted satellite is such as to significantly contribute to the PN population of the host halo, and so as to produce features in the PN velocity field, then the effect is short lived. This makes the probability of occurrence of such circumstance low.

A previous study of the intrinsic shapes of ETG halos was performed by Foster et al. (2016) who, using the observed distribution of ellipticities and misalignments for the SLUGGS galaxies out to $\sim 2.5R_e$ were unable to rule out triaxiality in the outskirts of fast rotators. The same authors had previously found strong evidence for minor axis rotation, hence a triaxial halo, in NGC 4473 (Foster et al. 2013). The extended PN kinematics allow us to generalize this result to a significant fraction of the fast rotators.

An examination of the smoothed velocity fields in appendix C shows that for some galaxies the trends in PA_{kin} with radius reflect a two-component velocity field in which rotation along one axis dominates at intermediate radii and rotation along the other axis dominates further out. This is the case for NGC 3379 and NGC 4649 among the fast rotators, and for NGC 4552 and NGC 4636 among the slow rotators. The velocity field of NGC 1344, instead, shows that this galaxy is a prolate rotator. Up to now there are not many observations of this phenomenon (Tsatsi et al. 2017, and references therein), which is probably related to a recent merger (Ebrowá & Łokas 2017), as the shells of this galaxy attest. Our sample contains two other galaxies (NGC 3923 and NGC 4365) known to have prolate rotation at their centers, but their halos are decoupled with respect to the inner kinematics.

8.4. Kinematic transition radius

We found that most of the galaxies in the sample show a transition in their kinematics, marked by a change in the rotation velocity V_{rot} or in the kinematic position angle PA_{kin} . In the framework of the two-phase formation scenario, we could interpret such kinematic differences as tracing stellar components with different origins, such as the in- and the ex-situ components. From simulations (Rodríguez-Gomez et al. 2016, and references therein) it is predicted that the in-situ stars are concentrated in the central regions of galaxies, while the accreted stars dominate the halos, and that their relative contribution is related primarily to the total mass. Within this framework we can define a transition radius R_T as the distance at which we observe the described transition in the kinematics, and compare it with the total stellar mass.

The radial range $R_T \pm \Delta R_T$ is quantified using (A) in case of a declining V_{rot} profile, the interval between the radius at which V_{rot} is maximum and the radius at which it decreases by $\sim 50 \text{ km s}^{-1}$ (B) in case of a increasing V_{rot} profile, the radial range in which V_{rot} increases from ~ 0 to $\sim 50 \text{ km s}^{-1}$ (C) in case of a kinematic twist, the radial range in which PA_{kin} changes significantly. Table 3 lists the $R_T \pm \Delta R_T$ values measured, and specifies the criteria (A,B,C) used for deriving it. $R_T \pm \Delta R_T$ is also plotted in figures 3 and 4 with vertical lines.

For each galaxy we estimated the stellar mass using two different approaches, designated by M_{*1} and M_{*2} , to have an estimate of the systematic uncertainties. The values for the masses are listed in table 2, along with the description of the procedure used for calculating them.

Figure 8 shows the stellar mass of the galaxies versus R_T/R_e . The full circles show the same quantities in three bins of R_T/R_e ; the error bars represent the standard deviation of the mass and of the R_T/R_e ratio in the bin. A clear correlation exists between total stellar mass and R_T/R_e , in the sense that the more massive galaxies tend to have transition radii at smaller fractions of R_e . The shaded region in figure 8 shows the corresponding quantities from the N-body simulations of Cooper et al. (2013). Using their stellar mass surface density profiles in bins of dark halo virial mass ($M_{200}/M_\odot$; their figure 6), we estimated the transition radius in units of half mass radii as the distance at which the accreted component overcomes the in-situ component. The total masses M_{200} are converted to stellar masses via abundance matching, using the prescription given by Behroozi et al. (2010). Both data and simulation follow the same trend.

We also observe a higher scatter in R_T/R_e for the lower mass galaxies, in agreement with the results of Rodríguez-Gomez et al. (2016) from the Illustris simulations. They defined a three-dimensional transition radius r_T as the distance at which the accreted stellar mass fraction overcomes the in-situ, and normalized it by the stellar half mass radius $r_{\text{half},*}$. Their study shows an additional dependence of $r_T/r_{\text{half},*}$ on morphology, halo formation time, and recent merger history, and that $r_T/r_{\text{half},*}$ tightly correlates with the ex-situ stellar mass fraction.

We tested the dependence of the correlation found on the choice of the value of R_e . The used values (listed in table 1) derive mostly from a calibrated average between 2MASS and RC3 determinations (Cappellari et al. 2011). Deeper photometric data would deliver larger values for R_e for the brightest galaxies (Kormendy et al. 2009). For this reason we carried out a simple photometric analysis, to be completed in a future study, with the goal of having a homogeneous determination of R_e for all the sample




Fig. 8. Transition radius in units of R_e versus total stellar mass (open circles). The full circles show the same quantities in bins of R_T/R_e . Different colors show the results of two different procedures for calculating the total stellar mass, see table 2 for details. The shaded region shows the corresponding quantities from the simulations of Cooper et al. (2013), see text.




Fig. 9. Transition radius in units of R_{halfight} versus total stellar mass (open circles). The full circles show the same quantities in bins of R_T/R_{halfight} . Different colors show the results of two different procedures for calculating the total stellar mass, see table 2 for details.

galaxies which takes into account the shallower surface brightness profile of the brightest galaxies. Using the surface brightness profiles available in the literature (references in table 2), we fitted a Sérsic law in the outermost regions of the galaxies and extrapolated the profiles up to very large radii to include all the stellar light. We then evaluated the half-light radius R_{halfight} from the growth curves (see table 2). Figure 9 shows that with these half-light radii, the relation between R_T/R_{halfight} and total stellar mass becomes steeper and agrees even better with the predicted trend from the simulations.

Spavone et al. (2017) quantified transition radii for a subsample of the ePN.S slow rotators using a parametrization of the light profiles with a multi-components Sérsic fit. Their three-components model describes the superposition of an in-situ component dominating the central regions, a relaxed, phase-mixed accreted component, to which is associated the transition radius R_{tr1} , and an outer diffuse component of unrelaxed accreted material with transition radius R_{tr2} . The transition radius is defined there as the radius at which one component overcomes the other. The R_T derived from the kinematics generally compares with R_{tr1} for the same galaxies.

D’Souza et al. (2014), using image stacking of a large sample of galaxies, defined R_{acc} as the radius at which the outer component of a double Sérsic model begins to dominate the integrated stellar light. They found a dependence of R_{acc} on the stellar mass, and, at comparable masses, their photometric values R_{acc} [kpc]

are in the same range as our kinematic R_T [kpc]. This is interesting considering the different approaches used to derive these quantities.

8.5. The diverse halo formation histories

Cosmological simulations (Naab et al. 2014; Penoyre et al. 2017) show that fast rotators, as well as slow rotators, are the result of different formation pathways, characterized by factors like the number of mergers, the merger mass ratio, the timing of mergers, and the gas fraction. We note, however, that the adopted star formation and AGN feedback models influence the resulting morphologies and kinematic properties, so that definitive statements about the origin of fast and slow rotators cannot yet be made (see Naab & Ostriker 2017, for a recent review). In the two-phase scenario, if the formation history is dominated by dissipation, then it is likely that the system becomes a fast rotator. Dissipation is particularly important in the early stage of galaxy formation, when the in-situ component generally settles down in a rotating disk (Rodríguez-Gomez et al. 2016; Qu et al. 2017), so that before redshift ~ 1 the progenitors of the fast and slow rotators are nearly indistinguishable (Penoyre et al. 2017). Later on, the evolution of galaxies is determined by the stochasticity of merging events, which at low redshift are mostly gas-poor (e.g. Lee & Yi 2013). Major mergers contribute to the loss of spin of the remnant, while minor mergers are responsible for the growth in size and mass. The accretion of stars happens preferentially on radial orbits, and frequent mergers efficiently mix the in-situ and the accreted stellar populations while changing the intrinsic shape towards a more spheroidal form (e.g. Jesseit et al. 2005; Hilz et al. 2012, 2013; Röttgers et al. 2014). Fast rotating galaxies experience a lower number of mergers when compared to the more massive slowly rotating systems, so their in-situ component is preserved in the central regions, in an axisymmetric, rotationally supported disk (e.g. Naab et al. 2014; Penoyre et al. 2017). Their halos, which host most of the accreted component, are not necessarily axisymmetric.

9. Summary and conclusions

In this paper we reconstructed and analyzed the velocity fields of 33 ETGs into their outer halos using planetary nebulae as tracers. For the total ePN.S sample, we combined 25 galaxies from the PN.S ETG survey with 8 objects observed with counter-dispersed imaging or multi object spectroscopy from other sources (Arnaboldi et al. in prep.). In the ePN.S sample, 24 are classified as fast rotators (following the definition by Emself et al. 2011), and 9 are slow rotators. The kinematic information typically extends to 6 effective radii (R_e), from a minimum of $3R_e$ to a maximum of $13R_e$.

We derived kinematic quantities such as the amplitude of rotation, the kinematic position angle (PA_{kin}), and the velocity dispersion, using an adaptive kernel smoothing technique validated with simulated velocity fields.

In this process, we checked whether the galaxy velocity fields were point-symmetric as expected for triaxial galaxies in dynamical equilibrium, and whether the systemic velocity is unchanged in the outer halo.

We supplemented the PN kinematics with absorption line kinematic data from the literature (typically limited to the regions inside $\sim 1 - 2R_e$), in order to have a complete picture of the nature of the objects, and compared the results from different datasets in the regions of overlap. Highlights of the results are:

(1) Most fast rotators (70%) have decreasing rotation amplitudes with radius, while a minority shows constant (15%) or increasing (15%) trends. Two thirds have PA_{kin} in their halos aligned with the photometric major axis at the resolution of the ePN.S survey. The slow rotators display modest but significant rotation in their outskirts. Among the fast rotators, 10% have very weak or no rotation in their outskirts, and half of the slow rotator sample have significant rotation at large radii.

(2) Slow rotators have approximately constant velocity dispersion profiles with radius, except for NGC 4636 which shows a steep drop in dispersion beyond 200 arcsec. The fast rotators display a variety in the shape of the velocity dispersion profiles: some are approximately constant, others decrease steeply with radius (figure 5).

(3) From their velocity fields, all of the slow rotators and one third of the fast rotators show indications of a triaxial halo (figure 4). For these galaxies we observe either a twist in the PA_{kin} or a constant misalignment with the photometric major axis. In addition two slow rotators have halos that are decoupled in their PA_{kin} .

(4) Five galaxies show deviations from point symmetry. For NGC 2768 and NGC 4594 these effects are small (probably related to a small stellar stream, in the former, and to extinction effects in the latter) and do not influence our kinematic analysis. NGC 1316, NGC 4472, and NGC 5128 instead display strong features, that indicate halos not in dynamical equilibrium. NGC 5128 is a merger remnant with clear minor axis rotation, and hence it is a triaxial system. For NGC 4472 the determination of the PA_{kin} is influenced by the presence of an in-falling satellite. While we included this slow rotator among the triaxial objects, we note that the galaxy would have approximately major axis rotation once the PNe of the satellite are removed, and we refer to Hartke et al. (in prep.) for a more detailed study. NGC 1316 is a merger remnant with pronounced major axis rotation in the halo.

(5) In a $V/\sigma(R_e)$ versus $V/\sigma(\text{halo})$ diagram (figure 7), the previous results imply different locations for the slow and the fast rotators. The latter separate further between fast rotators with slowly rotating outer spheroid and fast rotators that rotate rapidly at large radii also. The high values of $V/\sigma(\text{halo})$ are either due to a dominating disk component (e.g. NGC 7457) or to a rapidly rotating outer spheroid (NGC 2768). The triaxial fast rotators show approximately equal values of $V/\sigma(R_e)$ and $V/\sigma(\text{halo})$, and span all values of $V/\sigma(\text{halo})$.

(6) In the 11 fast rotator galaxies with the largest number of PN tracers, we see a decrease in the rotation velocity profile. In 10 cases this occurs approximately at a transition radius R_T similar to the scale radius of the known stellar disk, suggesting that the transition from rapid to slow rotation is due to the radial fading of the stellar disk in a slowly rotating outer spheroid. For NGC 4494 there is no photometric evidence for a disk. In the other fast rotators with fewer tracers that show a decrease in rotation, the comparison between the slow rotation of the PN system and the rapid rotation in the absorption line kinematics, which is related to an inner disk, suggests a similar transition at smaller radii.

(7) For slow rotators, we estimate a kinematic transition radius R_T from a variation in the rotation or in the PA_{kin} profile. Combining with the transition radii of fast rotators, we find a relation between the ratio R_T/R_e and stellar mass, such that the most massive galaxies have the lowest R_T/R_e (figure 8). In the framework of the two-phase formation scenario we can interpret R_T as marking the transition between the inner regions, dominated by the in-situ stellar component, and the halo, which

is mostly accreted. The comparison with photometric studies (D'Souza et al. 2014) and cosmological simulations (Cooper et al. 2013; Rodriguez-Gomez et al. 2016) shows good agreement, in particular the prediction from simulations is quantitatively confirmed.

In the data presented here, we see

- slow rotators
- fast rotators without apparent disks (NGC 4494 and NGC 4742)
- fast rotators with only inner disks and slowly rotating spheroids (e.g. NGC 3377)
- fast rotators with dominant disks all the way to their outermost regions (e.g. NGC 7457)
- fast rotators with inner disks and rapidly rotating spheroids (e.g. NGC 2768)
- fast rotators with triaxial halos that are dominated by dispersion (e.g. NGC 4649) or that rotate rapidly (NGC 4742 and NGC 5866)

Thus we conclude that ETGs show considerably more diversity in their halos than is apparent from their central bright regions. We also see clear signatures of the two-phase formation scenario: three galaxies show out of equilibrium kinematics in their halos, and more generally the inner and outer regions of ETGs often have different kinematic properties, where the transition radius depends on the stellar mass as predicted by cosmological simulations.

Acknowledgements.

We are grateful to Nigel G. Douglas for his fundamental contribution to the foundation of the Planetary Nebula Spectrograph instrument. We thank Luis Ho and Zhao-Yu Li for providing us with the Carnegie photometric data for the ePN.S ETG sample. C.P. thanks I. Soeldner-Rembold and J. Hartke for useful comments and discussions. C.P. is also grateful to F. Hofmann for advice and suggestions on the manuscript. A.J.R. was supported by National Science Foundation grant AST-1616710, and as a Research Corporation for Science Advancement Cottrell Scholar. C.T. is supported through an NWO-VICI grant (project number 639.043.308). This research has made use of NASA's Astrophysics Data System, of the data products from the Two Micron All Sky Survey (University of Massachusetts and the IPAC/CalTech), of the NASA/ IPAC Infrared Science Archive (Jet Propulsion Laboratory, CalTech), of the NASA/IPAC Extragalactic Database (NED), and of the VizieR catalogue access tool, CDS, Strasbourg, France. We acknowledge the usage of the HyperLeda database (<http://leda.univ-lyon1.fr>) as well.

Table 2. Measured parameters and typical errors for the smoothed velocity and velocity dispersion fields

Galaxy name	A ^(a) [arcsec ⁻¹]	B ^(a)	$\delta v^{(b)}$ [km s ⁻¹]	$\langle \Delta \tilde{v} \rangle^{(c)}$ [km s ⁻¹]	$\langle \Delta \tilde{\sigma} \rangle^{(d)}$ [km s ⁻¹]	$V_{\text{sys}}^{(e)}$ [km s ⁻¹]	$R_{\text{halflight}}^{(f)}$ [arcsec]	$\log_{10} \left(\frac{M_{*1}}{M_{\odot}} \right)^{(g)}$	$\log_{10} \left(\frac{M_{*2}}{M_{\odot}} \right)^{(h)}$
NGC0584	0.37	63.2	21.	21.	15.	1901. $\pm$ 27.	30	11.07	11.04 (4)
NGC0821	0.50	9.7	21.	22.	16.	1697. $\pm$ 16.	38	10.97	10.87 (1);(2)
NGC1023	0.26	15.3	14.	19.	13.	618. $\pm$ 9.	39	10.93	11.06 (6)
NGC1316	0.46	21.3	30.	24.	21.	1749. $\pm$ 30.	115	11.84	11.64 (4)
NGC1344	0.40	10.8	10.	21.	13.	1190. $\pm$ 13.	44	11.06	11.01 (7)
NGC1399	0.84	5.2	37.	27.	18.	1401. $\pm$ 19.	122	11.53	11.51 (4)
NGC2768	0.33	17.5	20.	26.	22.	1378. $\pm$ 14.	70	11.30	10.90 (15)
NGC2974	0.57	33.6	21.	25.	21.	1803. $\pm$ 42.	43	10.87	11.06 (4)
NGC3115	0.28	19.6	20.	30.	21.	624. $\pm$ 16.	73	10.98	10.96 (4)
NGC3377	0.32	19.0	20.	12.	8.	708. $\pm$ 9.	51	10.45	10.37 (1);(2);(3)
NGC3379	0.34	16.2	20.	21.	14.	934. $\pm$ 17.	58	10.89	10.89 (8);(9)
NGC3384	0.25	18.9	20.	14.	10.	722. $\pm$ 9.	42	10.77	10.88 (16)
NGC3489	0.32	14.7	20.	14.	11.	707. $\pm$ 12.	22	10.56	10.61 (16)
NGC3608	0.37	24.6	21.	15.	11.	1235. $\pm$ 16.	108	10.84	11.18 (2);(3)
NGC3923	0.60	24.3	30.	36.	26.	1677. $\pm$ 26.	164	11.54	11.66 (4)
NGC4278	0.88	2.4	18.	18.	14.	605. $\pm$ 17.	59	10.89	10.87 (12)
NGC4339	0.43	20.3	21.	8.	6.	1300. $\pm$ 7.	27	10.38	10.18 (13)
NGC4365	0.91	0.2	21.	20.	14.	1273. $\pm$ 11.	111	11.48	11.51 (14)
NGC4374	0.86	1.6	23.	27.	19.	1050. $\pm$ 18.	122	11.45	11.56 (14)
NGC4472	0.74	11.3	20.	38.	33.	959. $\pm$ 22.	193	11.71	11.76 (14)
NGC4473	0.91	0.2	21.	18.	12.	2236. $\pm$ 13.	38	10.87	10.78 (17)
NGC4494	0.41	12.8	21.	17.	12.	1348. $\pm$ 10.	49	11.05	10.87 (10)
NGC4552	0.90	0.6	21.	23.	16.	361. $\pm$ 15.	101	11.11	11.18 (14)
NGC4564	0.24	16.5	21.	19.	12.	1169. $\pm$ 17.	24	10.59	10.94 (1)
NGC4594	0.54	19.3	30.	32.	28.	1060. $\pm$ 16.	102	11.37	11.10 (4)
NGC4636	0.68	9.5	21.	24.	16.	903. $\pm$ 19.	172	11.12	11.27 (14)
NGC4649	0.86	1.5	20.	25.	18.	1059. $\pm$ 18.	132	11.55	11.55 (14)
NGC4697	0.45	6.0	35.	23.	16.	1274. $\pm$ 11.	123	11.03	11.05 (2);(11)
NGC4742	0.20	16.9	21.	13.	10.	1305. $\pm$ 9.	13	10.38	10.37 (4)
NGC5128	0.29	25.6	4.	25.	21.	536. $\pm$ 17.	414	11.04	11.20 (4)
NGC5846	0.90	0.1	21.	25.	17.	1716. $\pm$ 19.	193	11.42	11.49 (5);(18)
NGC5866	0.34	18.7	21.	15.	10.	782. $\pm$ 10.	48	10.97	11.01 (16)
NGC7457	0.28	10.5	20.	7.	6.	843. $\pm$ 5.	62	10.28	10.63 (16)

Notes. ^(a) Kernel parameters, A and B, used in the smoothing procedure, see section 3.

^(b) Mean error on the measured radial velocities.

^(c) Mean error on the smoothed velocity field.

^(d) Mean error on the velocity dispersion field.

^(e) Subtracted systemic velocity.

^(f) Half-light radius, see text.

^(g) Logarithm of the stellar mass M_{*1} in solar units. M_{*1} are derived from the K-band luminosities M_K listed in table 1, corrected for missing flux as in Scott et al. (2013): $M_{K\text{corr}} = 1.07 \times M_K + 1.53$, and applying a mass-to-light ratio of 1 (appropriate for old stellar population with a Kroupa IMF, Forbes et al. 2016).

^(h) Logarithm of the stellar mass M_{*2} in solar units. M_{*2} are derived using total luminosities obtained by fitting the surface brightness profiles with a Sérsic law and integrating till very large radii. The apparent magnitudes obtained are converted to absolute magnitudes using the distances listed in table 1, corrected for foreground galactic extinction, and homogenized to B magnitude using color indexes from the HyperLeda^a database and from Sandage & Visvanathan (1978). We then applied a stellar mass to light ratio given by the relation from Bell et al. (2003): $\log_{10} \frac{M}{L_B} = -9.42 + 1.737 \times (B - V)$, where the $(B - V)$ color index is from HyperLeda.

References for the photometry: (1) Goudfrooij et al. (1994), (2) Lauer et al. (2005), (3) Jedrzejewski (1987), (4) Li et al. (2011), (5) Kronawitter et al. (2000), (6) Noordermeer et al. (2008), (7) Sikkema et al. (2007), (8) Capaccioli et al. (1990), (9) Gebhardt et al. (2000), (10) Napolitano et al. (2009), (11) de Lorenzi et al. (2008), (12) Peletier et al. (1990), (13) Caon et al. (1994), (14) Kormendy et al. (2009), (15) Hopkins et al. (2009), (16) Krajnović et al. (2013), (17) Caon et al. (1990), (18) Spavone et al. (2017)

^a Makarov et al. (2014), <http://leda.univ-lyon1.fr/>

Table 3. Results of the kinematic analysis

Galaxy NGC	class ^(a)	point symm. ^(b)	features in $PA_{\text{kin}}(\text{R})$ ^(c)	multiple kin. comp. ^(d)	triaxial ^(e)	R_T ^(f) arcsec	NOTES ^(g)
0584	F	y	n	n	n	—	small number of tracers
0821	F	y	y	n	y	60 ± 30	(A,C)
1023	F	y	n	n	n	160 ± 20	(A)
1316	F	n	y	y	n	75 ± 25	(A) - merger remnant
1344	F	y	y	n	y	30 ± 15	(C) - prolate rotation
1399	S	y	y	y	y	30 ± 15	(B) - kinematically decoupled halo
2768	F	n	n	n	n	145 ± 45	(A) - fast rotating halo
2974	F	y	n	n	n	—	small number of tracers
3115	F	y	n	n	n	135 ± 45	(A)
3377	F	y	n	n	n	55 ± 15	(A)
3379	F	y	y	y	y	60 ± 20	(A,C) - hints for growing rotation in the outskirts
3384	F	y	n	n	n	—	extended disk
3489	F	y	n	n	n	—	small number of tracers
3608	S	y	y	n	y	55 ± 35	(B)
3923	S	y	y	n	y	70 ± 30	(B)
4278	F	y	n	n	n	25 ± 15	(A)
4339	F	y	n	n	n	—	small number of tracers
4365	S	y	y	y	y	125 ± 25	(C) - kinematically decoupled halo
4374	S	y	y	n	y	83 ± 33	(B)
4472	S	n	y	y	y	25 ± 10	(B) - recent merger; kinematics dominated by in-falling satellite
4473	F	y	y	n	y	25 ± 13	(A) - weak halo minor axis rotation
4494	F	y	n	n	n	93 ± 43	(A) - hints for additional rotation along the photometric minor axis
4552	S	y	y	y	y	73 ± 58	(B,C)
4564	F	y	n	n	n	85 ± 45	(A) - fast rotating halo
4594	F	y	n	n	n	180 ± 90	(A) - no detections in the plane of the disk (dust absorption)
4636	S	y	y	y	y	140 ± 40	(C)
4649	F	y	y	y	y	115 ± 15	(A) - no detections on the west side of the galaxy
4697	F	y	n	n	n	85 ± 65	(A) - hints for growing rotation in the outskirts
4742	F	y	y	n	y	43 ± 32	(C)
5128	F	n	y	y	y	95 ± 55	(C) - recent merger
5846	S	y	y	n	y	100 ± 50	(B)
5866	F	y	y	n	y	75 ± 25	(A) - fast rotating halo
7457	F	y	n	n	n	—	extended disk

Notes. ^(a) The sample is divided into slow (S) and fast rotators (F), according to the definition of Emsellem et al. (2011).

^(b) Point symmetry properties of the PN smoothed velocity fields: "y" marks point symmetric galaxies while "n" marks galaxies showing asymmetries, see section 5

^(c) Significant kinematic twist or misalignment of PA_{kin} with respect to PA_{phot} : "y" marks galaxies showing at least one of these, see section 6.4

^(d) For the galaxies marked with "y", the PN smoothed velocity fields reveal rotation along two perpendicular axes or a halo which is kinematically decoupled with respect to the innermost regions, see sections 7 and 6.4.2.

^(e) Galaxies marked with "y" have been classified as triaxial in section 6.4.

^(f) $R_T \pm \Delta R_T$ is the radial range of the kinematic transition.

^(g) Criteria (A, B, C) used to define the radial range $R_T \pm \Delta R_T$ as described in the text (section 8), and notes on individual galaxies (see also the dedicated appendix B).

References

- Alabi, A. B., Forbes, D. A., Romanowsky, A. J., et al. 2016, MNRAS, 460, 3838
- Alabi, A. B., Foster, C., Forbes, D. A., et al. 2015, MNRAS, 452, 2208
- Arnaboldi, M., Freeman, K. C., Gerhard, O., et al. 1998, ApJ, 507, 759
- Arnaboldi, M., Freeman, K. C., Mendez, R. H., et al. 1996, ApJ, 472, 145
- Arnold, J. A., Romanowsky, A. J., Brodie, J. P., et al. 2011, ApJ, 736, L26
- Arnold, J. A., Romanowsky, A. J., Brodie, J. P., et al. 2014, ApJ, 791, 80
- Arrigoni Battaia, F., Gavazzi, G., Fumagalli, M., et al. 2012, A&A, 543, A112
- Barbosa, C. E., Arnaboldi, M., Coccato, L., et al. 2017, ArXiv e-prints [arXiv:1710.08941]
- Bedregal, A. G., Aragón-Salamanca, A., Merrifield, M. R., & Milvang-Jensen, B. 2006, MNRAS, 371, 1912
- Behroozi, P. S., Conroy, C., & Wechsler, R. H. 2010, ApJ, 717, 379
- Beletsky, Y., Gadotti, D. A., Moiseev, A., Alves, J., & Kniazev, A. 2011, MNRAS, 418, L6
- Bell, E. F., McIntosh, D. H., Katz, N., & Weinberg, M. D. 2003, ApJS, 149, 289
- Bellstedt, S., Forbes, D. A., Foster, C., et al. 2017a, MNRAS, 467, 4540
- Bellstedt, S., Graham, A. W., Forbes, D. A., et al. 2017b, MNRAS, 470, 1321
- Bender, R. 1988a, A&A, 202, L5
- Bender, R. 1988b, A&A, 193, L7
- Bender, R., Saglia, R. P., & Gerhard, O. E. 1994, MNRAS, 269, 785
- Blakeslee, J. P., Cantiello, M., Mei, S., et al. 2010, ApJ, 724, 657
- Blakeslee, J. P., Jordán, A., Mei, S., et al. 2009, ApJ, 694, 556
- Blakeslee, J. P., Lucey, J. R., Barris, B. J., Hudson, M. J., & Tonry, J. L. 2001, MNRAS, 327, 1004
- Blom, C., Forbes, D. A., Brodie, J. P., et al. 2012, MNRAS, 426, 1959
- Boardman, N. F., Weijmans, A.-M., van den Bosch, R., et al. 2017, MNRAS, 471, 4005
- Bois, M., Emsellem, E., Bournaud, F., et al. 2011, MNRAS, 416, 1654
- Bournaud, F., Chapon, D., Teyssier, R., et al. 2011, ApJ, 730, 4
- Bridges, T. J., Rhode, K. L., Zepf, S. E., & Freeman, K. C. 2007, ApJ, 658, 980
- Brodie, J. P. & Strader, J. 2006, ARA&A, 44, 193
- Brough, S., Forbes, D. A., Kilborn, V. A., & Couch, W. 2006, MNRAS, 370, 1223
- Bryant, J. J., Owers, M. S., Robotham, A. S. G., et al. 2015, MNRAS, 447, 2857
- Buitrago, F., Trujillo, I., Curtis-Lake, E., et al. 2017, MNRAS, 466, 4888
- Bullock, J. S. & Johnston, K. V. 2005, ApJ, 635, 931
- Bundy, K., Bershad, M. A., Law, D. R., et al. 2015, ApJ, 798, 7
- Burkhead, M. S. 1986, AJ, 91, 777
- Caon, N., Capaccioli, M., & D'Onofrio, M. 1994, A&AS, 106, 199
- Caon, N., Capaccioli, M., & Rampazzo, R. 1990, A&AS, 86, 429
- Capaccioli, M., Held, E. V., Lorenz, H., & Vietri, M. 1990, AJ, 99, 1813
- Capaccioli, M., Held, E. V., & Nieto, J.-L. 1987, AJ, 94, 1519
- Cappellari, M., Emsellem, E., Bacon, R., et al. 2007, MNRAS, 379, 418
- Cappellari, M., Emsellem, E., Krajnović, D., et al. 2011, MNRAS, 413, 813
- Cappellari, M., McDermid, R. M., Alatalo, K., et al. 2013, MNRAS, 432, 1862
- Cappellari, M., Romanowsky, A. J., Brodie, J. P., et al. 2015, ApJ, 804, L21
- Carter, D., Thomson, R. C., & Hau, G. K. T. 1998, MNRAS, 294, 182
- Churazov, E., Tremaine, S., Forman, W., et al. 2010, MNRAS, 404, 1165
- Ciardullo, R., Jacoby, G. H., & Dejonghe, H. B. 1993, ApJ, 414, 454
- Coccato, L., Arnaboldi, M., & Gerhard, O. 2013, MNRAS, 436, 1322
- Coccato, L., Arnaboldi, M., Gerhard, O., et al. 2010, A&A, 519, A95
- Coccato, L., Gerhard, O., Arnaboldi, M., et al. 2009, MNRAS, 394, 1249
- Cohen, J. G. & Ryzhov, A. 1997, ApJ, 486, 230
- Cooper, A. P., D'Souza, R., Kauffmann, G., et al. 2013, MNRAS, 434, 3348
- Cortesi, A., Arnaboldi, M., Coccato, L., et al. 2013a, A&A, 549, A115
- Cortesi, A., Merrifield, M. R., Arnaboldi, M., et al. 2011, MNRAS, 414, 642
- Cortesi, A., Merrifield, M. R., Coccato, L., et al. 2013b, MNRAS, 432, 1010
- Crnojević, D., Sand, D. J., Spekkens, K., et al. 2016, ApJ, 823, 19
- Croom, S. M., Lawrence, J. S., Bland-Hawthorn, J., et al. 2012, MNRAS, 421, 872
- Daddi, E., Renzini, A., Pirzkal, N., et al. 2005, ApJ, 626, 680
- Damjanov, I., Abraham, R. G., Glazebrook, K., et al. 2011, ApJ, 739, L44
- Davies, R. L., Efstathiou, G., Fall, S. M., Illingworth, G., & Schechter, P. L. 1983, ApJ, 266, 41
- Davies, R. L. & Illingworth, G. 1983, ApJ, 266, 516
- Davies, R. L., Kuntschner, H., Emsellem, E., et al. 2001, ApJ, 548, L33
- De Bruyne, V., Dejonghe, H., Pizzella, A., Bernardi, M., & Zeilinger, W. W. 2001, ApJ, 546, 903
- de Lorenzi, F., Gerhard, O., Coccato, L., et al. 2009, MNRAS, 395, 76
- de Lorenzi, F., Gerhard, O., Saglia, R. P., et al. 2008, MNRAS, 385, 1729
- de Zeeuw, T. & Franx, M. 1991, ARA&A, 29, 239
- Deason, A. J., Belokurov, V., Evans, N. W., & McCarthy, I. G. 2012, ApJ, 748, 2
- Debatista, V. P., Corsini, E. M., & Aguerri, J. A. L. 2002, MNRAS, 332, 65
- Douglas, N. G., Arnaboldi, M., Freeman, K. C., et al. 2002, PASP, 114, 1234
- Dowell, J. L., Rhode, K. L., Bridges, T. J., et al. 2014, AJ, 147, 150
- D'Souza, R., Kauffman, G., Wang, J., & Vegetti, S. 2014, MNRAS, 443, 1433
- Duc, P.-A., Cuillandre, J.-C., Karabal, E., et al. 2015, MNRAS, 446, 120
- Duc, P.-A., Cuillandre, J.-C., Serra, P., et al. 2011, MNRAS, 417, 863
- Ebrov, I. & Łokas, E. L. 2017, ApJ, 850, 144
- Emsellem, E., Cappellari, M., Krajnović, D., et al. 2011, MNRAS, 414, 888
- Emsellem, E., Cappellari, M., Krajnović, D., et al. 2007, MNRAS, 379, 401

- Emsellem, E., Cappellari, M., Peletier, R. F., et al. 2004, *MNRAS*, 352, 721
- Forbes, D. A., Romanowsky, A. J., Pastorello, N., et al. 2016, *MNRAS*, 457, 1242
- Forestell, A. D. & Gebhardt, K. 2010, *ApJ*, 716, 370
- Foster, C., Arnold, J. A., Forbes, D. A., et al. 2013, *MNRAS*, 435, 3587
- Foster, C., Pastorello, N., Roediger, J., et al. 2016, *MNRAS*, 457, 147
- Foster, C., Spitler, L. R., Romanowsky, A. J., et al. 2011, *MNRAS*, 415, 3393
- Foster, C., van de Sande, J., D'Eugenio, F., et al. 2017, *MNRAS*, 472, 966
- Fouque, P., Gourgoulhon, E., Chamaraux, P., & Paturel, G. 1992, *A&AS*, 93, 211
- Franx, M., Illingworth, G., & de Zeeuw, T. 1991, *ApJ*, 383, 112
- Franx, M., Illingworth, G., & Heckman, T. 1989, *ApJ*, 344, 613
- Gabor, J. M. & Davé, R. 2012, *MNRAS*, 427, 1816
- Garcia, A. M. 1993, *A&AS*, 100, 47
- Gebhardt, K., Richstone, D., Kormendy, J., et al. 2000, *AJ*, 119, 1157
- Gerhard, O., Jeske, G., Saglia, R. P., & Bender, R. 1998, *MNRAS*, 295, 197
- Gerhard, O., Kronawitter, A., Saglia, R. P., & Bender, R. 2001, *AJ*, 121, 1936
- Gerhard, O. E. 1993, *MNRAS*, 265, 213
- Goudfroot, P., Hansen, L., Jorgensen, H. E., et al. 1994, *A&AS*, 104, 179
- Graham, A. W., Colless, M. M., Busarello, G., Zaggia, S., & Longo, G. 1998, *A&AS*, 133, 325
- Halliday, C., Davies, R. L., Kuntschner, H., et al. 2001, *MNRAS*, 326, 473
- Hartke, J., Arnaboldi, M., Longobardi, A., et al. 2017, *A&A*, 603, A104
- Haynes, R. F., Cannon, R. D., & Ekers, R. D. 1983, *Proceedings of the Astronomical Society of Australia*, 5, 241
- Hilz, M., Naab, T., & Ostriker, J. P. 2013, *MNRAS*, 429, 2924
- Hilz, M., Naab, T., Ostriker, J. P., Thomas, J. A., & Burkert, A., & Jesseit, R. 2012, *MNRAS*, 425, 3119
- Ho, L. C., Li, Z.-Y., Barth, A. J., Seigar, M. S., & Peng, C. Y. 2011, *ApJS*, 197, 21
- Ho, L. C., Li, Z.-Y., Barth, A. J., Seigar, M. S., & Peng, C. Y. 2012, *VizieR Online Data Catalog*, 219
- Hopkins, P. F., Cox, T. J., Dutta, S. N., et al. 2009, *ApJS*, 181, 135
- Huang, S., Ho, L. C., Peng, C. Y., Li, Z.-Y., & Barth, A. J. 2013, *ApJ*, 768, L28
- Hui, X., Ford, H. C., Freeman, K. C., & Dopita, M. A. 1995, *ApJ*, 449, 592
- Humphrey, P. J., Buote, D. A., Gastaldello, F., et al. 2006, *ApJ*, 646, 899
- Iodice, E., Capaccioli, M., Grado, A., et al. 2016, *ApJ*, 820, 42
- Iodice, E., Spavone, M., Capaccioli, M., et al. 2017, *ApJ*, 839, 21
- Janowiecki, S., Mihos, J. C., Harding, P., et al. 2010, *ApJ*, 715, 972
- Jarrett, T. H., Chester, T., Cutri, R., Schneider, S. E., & Huchra, J. P. 2003, *AJ*, 125, 525
- Jedrzejewski, R. & Schechter, P. L. 1988, *ApJ*, 330, L87
- Jedrzejewski, R. I. 1987, *MNRAS*, 226, 747
- Jensen, J. B., Tonry, J. L., Barris, B. J., et al. 2003, *ApJ*, 583, 712
- Jesseit, R., Naab, T., & Burkert, A. 2005, *MNRAS*, 360, 1185
- Kelson, D. D., Zabludoff, A. I., Williams, K. A., et al. 2002, *ApJ*, 576, 720
- Koopmans, L. V. E., Bolton, A., Treu, T., et al. 2009, *ApJ*, 703, L51
- Kormendy, J. & Bender, R. 1996, *ApJ*, 464, L119
- Kormendy, J., Fisher, D. B., Cornell, M. E., & Bender, R. 2009, *ApJS*, 182, 216
- Kormendy, J. & Illingworth, G. 1982, *ApJ*, 256, 460
- Kormendy, J. & Westpfahl, D. J. 1989, *ApJ*, 338, 752
- Krajnović, D., Alatalo, K., Blitz, L., et al. 2013, *MNRAS*, 432, 1768
- Krajnović, D., Bacon, R., Cappellari, M., et al. 2008, *MNRAS*, 390, 93
- Krajnović, D., Cappellari, M., de Zeeuw, P. T., & Copin, Y. 2006, *MNRAS*, 366, 787
- Krajnović, D., Emsellem, E., Cappellari, M., et al. 2011, *MNRAS*, 414, 2923
- Kronawitter, A., Saglia, R. P., Gerhard, O., & Bender, R. 2000, *A&AS*, 144, 53
- Lackner, C. N., Cen, R., Ostriker, J. P., & Joung, M. R. 2012, *MNRAS*, 425, 641
- Lauer, T. R., Faber, S. M., Gebhardt, K., et al. 2005, *AJ*, 129, 2138
- Laurikainen, E., Salo, H., Buta, R., Knapen, J. H., & Comerón, S. 2010, *MNRAS*, 405, 1089
- Lee, J. & Yi, S. K. 2013, *ApJ*, 766, 38
- Li, Z.-Y., Ho, L. C., Barth, A. J., & Peng, C. Y. 2011, *ApJS*, 197, 22
- Longobardi, A., Arnaboldi, M., Gerhard, O., et al. 2013, *A&A*, 558, A42
- Longobardi, A., Arnaboldi, M., Gerhard, O., & Mihos, J. C. 2015, *A&A*, 579, L3
- Loubser, S. I., Sansom, A. E., Sánchez-Blázquez, P., Soechting, I. K., & Bromage, G. E. 2008, *MNRAS*, 391, 1009
- Ma, C.-P., Greene, J. E., McConnell, N., et al. 2014, *ApJ*, 795, 158
- Madore, B. F., Freedman, W. L., & Bothun, G. D. 2004, *ApJ*, 607, 810
- Makarov, D., Prugniel, P., Terekhova, N., Courtois, H., & Vauglin, I. 2014, *A&A*, 570, A13
- Malin, D. F. & Carter, D. 1980, *Nature*, 285, 643
- Malin, D. F., Quinn, P. J., & Graham, J. A. 1983, *ApJ*, 272, L5
- Mandelbaum, R., Seljak, U., Kauffmann, G., Hirata, C. M., & Brinkmann, J. 2006, *MNRAS*, 368, 715
- Marcelin, M. 1983, in *IAU Symposium*, Vol. 100, *Internal Kinematics and Dynamics of Galaxies*, ed. E. Athanassoula, 335
- McNeil, E. K., Arnaboldi, M., Freeman, K. C., et al. 2010, *A&A*, 518, A44
- McNeil-Moylan, E. K., Freeman, K. C., Arnaboldi, M., & Gerhard, O. E. 2012, *A&A*, 539, A11
- Méndez, R. H., Riffeser, A., Kudritzki, R.-P., et al. 2001, *ApJ*, 563, 135
- Méndez, R. H., Teodorescu, A. M., Kudritzki, R.-P., & Burkert, A. 2009, *ApJ*, 691, 228
- Michard, R. & Marchal, J. 1994, *A&AS*, 105, 481
- Morganti, L., Gerhard, O., Coccatto, L., Martínez-Va Ipuesta, I., & Arnaboldi, M. 2013, *MNRAS*, 431, 3570
- Murphy, J. D., Gebhardt, K., & Adams, J. J. 2011, *ApJ*, 729, 129
- Naab, T., Oser, L., Emsellem, E., et al. 2014, *MNRAS*, 444, 3357
- Naab, T. & Ostriker, J. P. 2017, *ARA&A*, 55, 59
- Napolitano, N. R., Arnaboldi, M., Freeman, K. C., & Capaccioli, M. 2001, *A&A*, 377, 784
- Napolitano, N. R., Romanowsky, A. J., Capaccioli, M., & Douglas, N. G., et al. 2011, *MNRAS*, 411, 2035
- Napolitano, N. R., Romanowsky, A. J., Coccatto, L., et al. 2009, *MNRAS*, 393, 329
- Neumayer, N. 2010, *PASA*, 27, 449
- Noordermeer, E., Merrifield, M. R., Coccatto, L., et al. 2008, *MNRAS*, 384, 943
- Norris, M. A., Gebhardt, K., Sharples, R. M., et al. 2012, *MNRAS*, 421, 1485
- Norris, M. A., Sharples, R. M., & Kuntschner, H. 2006, *MNRAS*, 367, 815
- Oser, L., Ostriker, J. P., Naab, T., Johansson, P. H., & Burkert, A. 2010, *ApJ*, 725, 2312
- Peletier, R. F., Davies, R. L., Illingworth, G. D., Davis, L. E., & Cawson, M. 1990, *AJ*, 100, 1091
- Peng, E. W., Ford, H. C., & Freeman, K. C. 2004, *ApJ*, 602, 685
- Peng, E. W., Ford, H. C., Freeman, K. C., & White, R. L. 2002, *AJ*, 124, 3144
- Penoyre, Z., Moster, B. P., Sijacki, D., & Genel, S. 2017, *MNRAS*, 468, 3883
- Pignatelli, E. 1999, *ArXiv Astrophysics e-prints [astro-ph/9906378]*
- Pota, V., Brodie, J. P., Bridges, T., et al. 2015, *MNRAS*, 450, 1962
- Pota, V., Forbes, D. A., Romanowsky, A. J., et al. 2013, *MNRAS*, 428, 389
- Proctor, R. N., Forbes, D. A., Romanowsky, A. J., et al. 2009, *MNRAS*, 398, 91
- Pu, S.-B. & Han, Z.-W. 2011, *Research in Astronomy and Astrophysics*, 11, 909
- Qu, Y., Helly, J. C., Bower, R. G., et al. 2017, *MNRAS*, 464, 1659
- Raskutti, S., Greene, J. E., & Murphy, J. D. 2014, *ApJ*, 786, 23
- Rejkuba, M., Minniti, D., Courbin, F., & Silva, D. R. 2002, *ApJ*, 564, 688
- Renaud, F., Agertz, O., & Gieles, M. 2017, *MNRAS*, 465, 3622
- Richtler, T., Hilker, M., Kumar, B., et al. 2014, *A&A*, 569, A41
- Rix, H.-W., de Zeeuw, P. T., Cretton, N., van der Marel, R. P., & Carollo, C. M. 1997, *ApJ*, 488, 702
- Rodríguez-Gómez, V., Pillepich, A., Sales, L. V., et al. 2016, *MNRAS*, 458, 2371
- Romanowsky, A. J., Douglas, N. G., Arnaboldi, M., et al. 2003, *Science*, 301, 1696
- Romanowsky, A. J. & Fall, S. M. 2012, *ApJS*, 203, 17
- Romanowsky, A. J., Strader, J., Brodie, J. P., et al. 2012, *ApJ*, 748, 29
- Röttgers, B., Naab, T., & Oser, L. 2014, *MNRAS*, 445, 1065
- Saglia, R. P., Kronawitter, A., Gerhard, O., & Bender, R. 2000, *AJ*, 119, 153
- Sambhus, N., Gerhard, O., & Méndez, R. H. 2006, *AJ*, 131, 837
- Sánchez, S. F., Kennicutt, R. C., Gil de Paz, A., & van de Ven, G., et al. 2012, *A&A*, 538, A8
- Sandage, A. & Visvanathan, N. 1978, *ApJ*, 223, 707
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schuberth, Y., Richtler, T., Hilker, M., et al. 2010, *A&A*, 513, A52
- Schuberth, Y., Richtler, T., Hilker, M., et al. 2012, *A&A*, 544, A115
- Schweizer, F. 1980, *ApJ*, 237, 303
- Scorza, C., Bender, R., Winkelman, C., Capaccioli, M., & Macchetto, D. F. 1998, *A&AS*, 131, 265
- Scott, N., Davies, R. L., Houghton, R. C. W., et al. 2014, *MNRAS*, 441, 274
- Scott, N., Graham, A. W., & Schombert, J. 2013, *ApJ*, 768, 76
- Sengupta, C. & Balasubramanyam, R. 2006, *MNRAS*, 369, 360
- Sikkema, G., Carter, D., Peletier, R. F., et al. 2007, *A&A*, 467, 1011
- Simien, F. & Prugniel, P. 1997a, *A&AS*, 122, 521
- Simien, F. & Prugniel, P. 1997b, *A&AS*, 126, 15
- Simien, F. & Prugniel, P. 1997c, *A&AS*, 126, 519
- Skutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Suis, A. P. N. & Williams, T. B. 2006, *AJ*, 131, 2089
- Smethurst, R. J., Masters, K. L., Lintott, C. J., et al. 2018, *MNRAS*, 473, 2679
- Sommer-Larsen, J. & Toft, S. 2010, *ApJ*, 721, 1755
- Spavone, M., Capaccioli, M., Napolitano, N. R., et al. 2017, *A&A*, 603, A38
- Spiniello, C., Napolitano, N. R., Coccatto, L., et al. 2015, *MNRAS*, 452, 99
- Stark, A. A. 1977, *ApJ*, 213, 368
- Statler, T. S. 1994, *ApJ*, 425, 500
- Statler, T. S. & Smecker-Hane, T. 1999, *AJ*, 117, 839
- Strader, J., Romanowsky, A. J., Brodie, J. P., et al. 2011, *ApJS*, 197, 33
- Tal, T., van Dokkum, P. G., Nelan, J., & Bezanson, R. 2009, *AJ*, 138, 1417
- Teodorescu, A. M., Méndez, R. H., Bernardi, F., et al. 2011, *ApJ*, 736, 65
- Teodorescu, A. M., Méndez, R. H., Saglia, R. P., et al. 2005, *ApJ*, 635, 290
- Thomas, J., Jesseit, R., Saglia, R. P., et al. 2009, *MNRAS*, 393, 641
- Tonry, J. L., Dressler, A., Blakeslee, J. P., et al. 2001, *ApJ*, 546, 681
- Trujillo, I., Conselice, C. J., Bundy, K., et al. 2007, *MNRAS*, 382, 109
- Trujillo, I., Förster Schreiber, N. M., Rudnick, G., et al. 2006, *ApJ*, 650, 18
- Tsatsi, A., Lyubenova, M., van de Ven, G., et al. 2017, *A&A*, 606, A62
- van den Bergh, S. 1961, *AJ*, 66, 562
- van der Marel, R. P., Rix, H. W., Carter, D., et al. 1994, *MNRAS*, 268, 521
- van Dokkum, P. G., Whitaker, K. E., Brammer, G., et al. 2010, *ApJ*, 709, 1018
- Veale, M., Ma, C.-P., Greene, J. E., et al. 2018, *MNRAS*, 473, 5446
- Veale, M., Ma, C.-P., Thomas, J., et al. 2017, *MNRAS*, 464, 356
- Veljanoski, J., Mackey, A. D., Ferguson, A. M. N., et al. 2014, *MNRAS*, 442, 2929
- Walsh, J. R., Rejkuba, M., & Walton, N. A. 2015, *A&A*, 574, A109
- Weijmans, A.-M., Cappellari, M., Bacon, R., et al. 2009, *MNRAS*, 398, 561
- Weijmans, A.-M., de Zeeuw, P. T., Emsellem, E., et al. 2014, *MNRAS*, 444, 3340
- Wellons, S., Torrey, P., Ma, C.-P., et al. 2016, *MNRAS*, 456, 1030
- Wu, X., Gerhard, O., Naab, T., et al. 2014, *MNRAS*, 438, 2701
- Wuyts, S., Cox, T. J., Hayward, C. C., et al. 2010, *ApJ*, 722, 1666

Appendix A: Testing the adaptive kernel smoothing procedure on simulated data

Our procedure for the measurement of the mean velocity and velocity dispersion fields from the observed LOS velocities of the PNe is based on performing a weighted local average, where the weights depend on the local density of tracers. The derived quantities depend on how well the detected PNe statistically sample the parent galaxy in the phase space. In particular the spatial distribution of the PNe plays an important role, as the smoothing averages over objects that are close together on the sky. Appendix A.1 describes what determines the observed PN spatial distribution and how this, combined with a smoothing procedure, may affect the results. Appendices A.2 and A.3 study and quantify these effects on a simulated galaxy (with variable number of tracers and V/σ ratio) and on simulated velocity fields, with the same spatial distribution as the observed galaxies. We found that the smoothing procedure does not create artifacts in the velocity fields above the 1σ level; any features above this threshold are probably real.

Appendix A.1: PN spatial distribution and completeness

The observed PN spatial distribution depends on several factors. The most evident one is the incompleteness in the central high surface brightness regions because of the difficulty in detecting faint point sources against a bright background (Arnaboldi et al. in prep., Coccato et al. 2009). This is a strong function of the radius and it is quantified by the *completeness function*, the fraction of detected objects in different radial ranges.

In addition, the PN number density is proportional to the local surface brightness of the galaxy. This means that the number of tracers decreases with radius following the steep decrease of the surface brightness, and features in the surface brightness distribution may generate under- or over-densities of PNe. Of course at our typical number of tracers (~ 200) these features in density are barely distinguishable from statistical effects.

The adopted smoothing procedure takes into account the natural non-uniformity of the PNe distribution through the use of an adaptive kernel technique, which optimizes the size of the kernel to the local density of tracers. Nevertheless a local average of the velocities over a non-uniform spatial distribution, combined with the statistics in the LOSVD sampling, may in principle generate features in the kinematic maps or biases in the estimate of the kinematic parameters. On the other hand, averaging over voids or over sparsely distributed tracers unavoidably leads to over-smoothed velocity amplitudes and to under-resolved velocity gradients. The residual velocities from the unresolved gradients may artificially boost the dispersion, creating local maxima in the velocity dispersion maps. In appendices A.2 and A.3 we assess the impact of such biases using simulated data.

Appendix A.2: Tests with data from a simulated galaxy

We used a simulated major merger remnant, sampled by 10000 particles. We reproduced a realistic observed dataset by locating the simulated galaxy at a distance of 20 Mpc, and applying to it a completeness function similar to those observed (see Arnaboldi et al. in prep.). We observed the galaxy in a field of view similar to that of the ePN.S, 800×800 arcsec² wide centered on to the galaxy. At this point, the simulated galaxy can be used as a test case for the procedure: we can explore how the results change by varying the number of tracers at our disposal, by selecting randomly typical numbers of observed PNe; different random

extractions of tracers give us different statistical realizations of the system, and different projections of the galaxy on the sky plane give us different V/σ ratios.

Figure A.1 shows an example of such an experiment. The smoothed velocity and velocity dispersion fields are plotted for the 10000 particle simulation, together with the maps for 400 stars and 200 stars. For the 200 particles case we show two typical statistical realizations (A and B) in order to illustrate the effects of limited statistics on the kinematic quantities. The panels at the bottom right show the fitted kinematic position angles, the amplitudes of the rotational velocity, and the azimuthally averaged velocity dispersion profiles in circular bins. The kinematic quantities are in general well recovered within the errors. The simulation is point symmetric in the region selected, and so are the limited statistical realizations. The statistical fluctuations that appear in the smoothed fields usually do not cause a deviation from point symmetry nor a variation of the kinematic position angle larger than the error bars. The velocity dispersion may be marginally boosted at the center of the galaxy, associated with the lack of detections, (see the case of 200 stars in figure A.1), but the dispersion profile is recovered within the errors. So we conclude that the procedure to calibrate the smoothing parameters through Monte Carlo simulations (section 3.1.1) works very well in reconstructing the mean properties of the velocity and velocity dispersion fields. Even though the smoothing procedure may create artificial peaks or deeps in the velocity maps, all these features are generally within the 1σ level. Any radial trends can therefore be considered significant only if it is outside the statistical uncertainties.

However, we can notice that as the number of tracers drops or the ratio V/σ decreases, the amplitude of rotation is systematically underestimated, while the velocity dispersion is systematically higher (see also Napolitano et al. 2001). This happens because lower numbers of stars or higher statistical noise require larger smoothing scales, which eventually become larger than the actual spatial scales of the velocity gradients. Oversmoothed velocity amplitudes, i.e. lower $\langle v \rangle$, imply higher $\tilde{\sigma}$ in equation (2). There is no way to estimate this offsets unless the mean velocity and velocity dispersion fields of the galaxy are known from independent data, since the effects of the smoothing jointly depend on the local density of tracers, on the V/σ ratio, and on the particular statistical realization (e.g. A and B in figure A.1), hence they vary from case to case. It is however safe to say that for the galaxies with limited statistics the recovered kinematics provides a lower limit to the amplitude of rotation and an upper limit for the velocity dispersion in the halos.

Appendix A.3: Statistical tests with simulated velocity fields on real galaxies

Simulated velocity fields were used to verify the results obtained for the ePN.S sample of galaxies, such as assessing the significance of deviations from point symmetry or the radial trends of the kinematic parameters. These simulated velocity fields are featureless and trend-less (fixed kinematic position angle and constant velocity dispersion) and are built for each galaxy using the positions of the observed PNe. This means that the incompleteness function is already built into the simulated catalogs and is identical to that for the real galaxy under study. In this way we can check how the smoothing procedure correlates spatially close PNe, as the degree of spatial correlation in the models will be identical to that of the real datasets.

The simulated velocity field at the positions (x_i, y_i) of the i th PNe, from the real datasets, is interpolated with the simple




Fig. A.1. Smoothed velocity and velocity dispersion fields in the ePN.S field of view for a simulated ETG, traced by 10000 particles, 400, and 200 stars after the convolution with a completeness function. For the 200 particle case, we show two typical statistical realizations (A and B) of the extracted sample, to illustrate the effects of the limited statistics on the smoothed quantities. The global features of the simulation are preserved in the different fields but, as expected, the spatial and velocity resolution of the local gradients is downgraded when the number of tracers decreases. The lower right panels display the fitted kinematic position angle PA , the rotation velocity V_{rot} , and the azimuthally averaged velocity dispersion σ in circular radial bins as functions of radius. It is evident that the radial gradients are less resolved with fewer tracers, and that the amplitude of rotation for the case with 200 stars is oversmoothed. This happens because the kernel amplitude is larger than the real spatial scale of the velocity gradients due to the sparse distribution of the tracers. The blue full circles show a simulation with higher V/σ ratio and two of its typical statistical realizations sampled by 200 objects to illustrate how of the accuracy of the reconstructed velocity amplitudes depends on the statistical noise. Errors for the kinematic profiles are determined from Monte Carlo simulations using the smoothed velocity and velocity dispersion fields as described in section 3.2.2.

function

$$V(\phi_i, R_i) = V_{\text{rot}} \cos(\phi_i - PA_{\text{phot}}) \operatorname{arctanh}(R_i/h)$$

$$(A.1) \quad \text{where } \phi_i \text{ is the position angle of the } i\text{th PN with coordinates } (x_i, y_i), R_i \text{ its distance from the center, and } h \text{ is a scale-length}$$

that defines the steepness of the central velocity radial gradient. V_{rot} is chosen to be equal to the maximum fitted rotational amplitude for the galaxy in consideration, while h has the dimension of the central void of the detections (~ 50 arcsec). We choose to align the rotation of the simulated velocity field with the real photometric position angle of the galaxy, in order to simulate the velocity field of a regular, point-symmetric disk. The model velocity field $V(x_i, y_i)$ is sampled with a constant dispersion σ , which is equal to the mean velocity dispersion of the galaxy under study. This is done by extracting random values from a Gaussian distribution centered at 0 and with dispersion σ , and adding them to $V(x_i, y_i)$. The assumption of constant dispersion is certainly unrealistic for the central regions of the galaxies, but it is reasonable at large radii. We produced 100 statistical realizations for each galaxy, and each of those is treated as a real dataset for obtaining a total of 100 smoothed velocity fields.

The smoothed velocity fields obtained in this way were used to assess whether an observed feature in the galaxy velocity or velocity dispersion field is real, by studying whether it can be produced by statistical effects in a featureless velocity field, and how typically this happens (see sections 5.1 and 6.3). Also, the 100 models can provide a statistic for the fitted kinematic parameters (equation 9) to check the probability that statistical noise combined with the smoothing procedure artificially produce effects like twists of the kinematic position angle or misalignments with the photometric axes (section 6.4).

Appendix B: Notes on single galaxies and comparison with the literature

Appendix B.1: Fast rotators

- **NGC 0584.** This is the brightest member of a small cluster containing mostly E galaxies (van den Bergh 1961). It is classified as a E4 in the NED catalog, but it seems to contain a disk structure (Michard & Marchal 1994). PN kinematic data show that the galaxy halo rotates by at least ~ 50 km s $^{-1}$ along the photometric major axis, and has a velocity dispersion of ~ 100 km s $^{-1}$.
- **NGC 0821.** This is a field E6 galaxy. The ATLAS^{3D} velocity field (Emsellem et al. 2004) maps the inner regions of the galaxy where it behaves like a regular rotator along the photometric major axis. The PNe sample, instead, probes the kinematics from 1 to $5R_e$ and show a sustained rotation along the photometric minor axis with a maximum amplitude of 52 ± 19 km s $^{-1}$, suggesting the triaxiality of the object. No rotation is detected along the major axis. The long-slit data from Proctor et al. (2009) and the integral field map of Arnold et al. (2014) agree in detecting rotation along the photometric major axis, with decreasing amplitude (80 km s $^{-1}$ at $0.3R_e$ to very faint rotation at $R < 2R_e$). They do not detect any rotation along the minor axis, but their do not extend far beyond $1.5R_e$ along that axis. Long slit data from Weijmans et al. (2009) and Forestell & Gebhardt (2010) do show some minor axis rotation at large radii. PN data are consistent with the kinematics of the blue GCs (see Pota et al. 2013), which are found to rotate at ~ 85 km s $^{-1}$ along a direction consistent with the photometric minor axis. The red GCs instead appear to faintly counter-rotate with respect to the host galaxy stars. The outer gradient of the velocity dispersion radial profile has been debated in the literature, with implications on the inferred dark matter content of the galaxy. The PN data combined with the long slit stellar kinematics show a decrease of the velocity dispersion as a function of radius, consistently
- as both red and blue GCs show, while Proctor et al. (2009) and Forestell & Gebhardt (2010) found a flat profile within 100 arcsec.
- **NGC 1023.** This is a barred lenticular galaxy (SB0), brightest member of a group of 13 galaxies (Tonry et al. 2001). The ATLAS^{3D} velocity map (Emsellem et al. 2004) covers the innermost regions of the galaxy, up to half effective radii, and displays a featureless regular rotation, with a maximum amplitude of ~ 120 km s $^{-1}$. The stellar kinematics along the major axis of Debattista et al. (2002) and Boardman et al. (2017) show that the rotation continues to increase mildly in the outer regions exceeding 200 km s $^{-1}$ beyond $2R_e$. Both the PN and SLUGGS (Arnold et al. 2014) velocity maps show decreasing rotation beyond $3R_e$, (see also Noordermeer et al. 2008; Coccato et al. 2009; Cortesi et al. 2013b). The galaxy does not show any twist of the kinematic position angle or any misalignment with the photometric axis. The non zero third order harmonics amplitudes describe a rather cylindrical velocity field. The velocity dispersion map shows minimum values ($\sigma \sim 40$ km s $^{-1}$) at the location of the disk, while the outermost (bulge) PNe are hotter ($\sigma \sim 120$ km s $^{-1}$).
- **NGC 1316.** This is a giant elliptical (also classified as a peculiar S0), member of the Fornax cluster (Fornax A). NGC 1316 is a known merger remnant. Its photometric substructures were mapped for the first time by Schweizer (1980), and were recently studied by Iodice et al. (2017). The PN velocity field is highly asymmetric, showing that the galaxy is in a non-equilibrium phase. The modeling with a point-symmetric rotation model gives an approximate description of the properties of the galaxy. The fit of the $a_0(R)$ parameter of the rotation model (see equation 11) in radial bins shows the presence of peculiar bulk motions in groups of PNe. There is no kinematics from integral field spectroscopy in literature, and the comparison with the long-slit observations from Bedregal et al. (2006) shows general good agreement with our data in the regions of overlap. The PNe in the halo of NGC 1316 (figure B.1) reveal that the galaxy rotates along the major axis with an amplitude of ~ 65 km s $^{-1}$, as already observed by Arnaboldi et al. (1998); McNeil-Moylan et al. (2012). The structures in the velocity field can be spatially associated with corresponding surface brightness features in the same location. There is an overdensity of PNe on the north-eastern part, co-located spatially with the ripple R4 (the designations of the photometric features are from Schweizer 1980, and their positions are indicated in black in figure B.1), with velocities that do not have a point symmetric counterpart. Other groups of PNe that result in non-point-symmetric velocities are those possibly related with the loop L1, on the south-west side, and to the plume P in the north-west. Figure 1 shows that the PNe of NGC 1316 in the innermost radial bin (~ 100 arcsec) have a velocity offset of $\gtrsim 150$ km s $^{-1}$ with respect to the systemic velocity. This is produced by a group of relatively slower PNe with mean velocity ~ 1325 km s $^{-1}$ and dispersion ~ 60 km s $^{-1}$ (highlighted in figure B.1 with white circles, along with the photometric feature O1 found by Richtler et al. 2014, in red). These PNe are not associated with any substructures in the light, and the fact that they are very localized in the phase space challenges (but does not rule out) the possibility that they could be contaminants. More likely this kinematic feature is the result of a combination of spatial incompleteness and differential absorption from localized dust in the galaxy, producing local lacks of detections in the phase space and so more negative




Fig. B.1. Smoothed velocity field and velocity dispersion field of NGC 1316 shown on the DSS image; north is up, east is left. The circles highlight the PNe belonging to a structure in the phase space, discussed in the text. The black contours trace features in the surface brightness distribution (designations from Schweizer 1980). In red is marked the position of the feature O1 observed by Richtler et al. (2014). The arrow at the bottom right shows the direction of the PA_{phot} .

smoothed velocities $\tilde{V}$. A more detailed study is beyond the scope of this paper.

- **NGC 1344.** This galaxy, also known as NGC 1340, is an E5 elliptical belonging to the Fornax cluster. It is characterized by the presence of internal and external concentric shells (Malin & Carter 1980), as consequences of a recent merger activity. The smoothed velocity field does not deviate from point symmetry, however it shows that the galaxy in the outskirts rolls around the photometric major axis. The velocity dispersion field shows a gently decreasing profile. Unfortunately there is no integral field stellar kinematics available for a comparison with the PNe kinematics. The long slit stellar kinematics (Teodorescu et al. 2005) show sustained rotation along the photometric major axis that reaches $\sim 100 \text{ km s}^{-1}$ at $1.3R_e$ and a decreasing velocity dispersion. The σ values from the spectroscopy are in agreement with those found with the PN data. We do not detect such a strong rotation, probably because of the different radial coverage of the PN data.
- **NGC 2768.** This S0 is a field galaxy (Madore et al. 2004) or member of a poor group (Fouque et al. 1992). NGC 2768 has a cylindrical velocity field (Emsellem et al. 2004) whose amplitude remains constant at around $\sim 130 \text{ km s}^{-1}$ from the innermost regions (Emsellem et al. 2004) to the outskirts (see Proctor et al. 2009; Pota et al. 2013). Our velocity map is in good agreement with that presented by Arnold et al. (2014) and extends it beyond $6R_e$. There is some discrepancy between the PN V_{rot} and the kinemetry fit from Foster et al. (2016) in the innermost radial bins where the disk strongly dominates. This is due to the fact that the PN smoothed velocity field average together the fast rotating PNe of the disk with those more slowly rotating from the spheroid. A kinematic study of the different components of this galaxy has been done by Cortesi et al. (2013b). At larger radii ($r \gtrsim 3R_e$)

the contribution of the disk is weaker and the PN kinematics traces the spheroid. There we found a constant sustained ($V \sim 130 \text{ km s}^{-1}$) rotation along the photometric major axis. The velocity dispersion profile is flat at $\sim 140 \text{ km s}^{-1}$. The PN smoothed velocity field of NGC 2768 shows localized, small scale deviations from point symmetry, which might be related to a small stream visible in optical light (Duc et al. 2015), while the main body of the galaxy is dynamically relaxed. These asymmetries do not influence the kinematic analysis, but we used the unfolded catalog to build the velocity fields. Our PN kinematics agrees with that of GCs. Pota et al. (2013) found significant rotation only for the red GCs along the photometric major axis and with constant amplitude. Their velocity dispersion is decreasing with radius.

- **NGC 2974.** This is an elliptical (E4) field galaxy. Its kinematics was mapped by the SAURON survey (Emsellem et al. 2004), showing that the galaxy is rotating at $\sim 120 \text{ km s}^{-1}$ along the photometric major axis, while the velocity dispersion has a steep decreasing profile. PN kinematic data show that the galaxy halo rotates at least at $\sim 70 \text{ km s}^{-1}$ along the photometric major axis, and has a velocity dispersion of 150 km s^{-1} .
- **NGC 3115.** This is an S0 associated with a loose group of galaxies (Fouque et al. 1992). The long slit spectroscopy from Norris et al. (2006) shows the rotation velocity increasing steeply in the innermost regions and flattens at 230 km s^{-1} around $\sim 1R_e$. Our data show a cylindrical velocity field, which rotates with a gently decreasing amplitude that reaches $\sim 120 \text{ km s}^{-1}$ at $\sim 3R_e$, where the spheroid dominates. No twist or misalignment with the photometric position angle are observed. The velocity dispersion field is colder ($\sigma \sim 120 \text{ km s}^{-1}$) in the region dominated by the disk, and hotter along the minor axis ($\sigma \sim 220 \text{ km s}^{-1}$). Cortesi et al. (2013b) performed a kinematic study of the different

components of the galaxy. Our kinematics is in good agreement with the stellar and globular cluster kinematics from Arnold et al. (2011).

- **NGC 3377.** It is an E6 galaxy in the Leo I group. The absorption line kinematics (Coccato et al. 2009; Emsellem et al. 2004; Foster et al. 2016) shows that the galaxy displays a disk-like rotation in the inner regions, which starts to decrease at $\sim 1.5R_e$. Our analysis of the PN kinematics shows that the decreasing trend continues out to large radii, where it gets comparable to zero at $\sim 3R_e$ and remains negligible beyond $\sim 7R_e$. The kinematic major axis position angle is poorly determined as the rotation amplitude is really low, comparable with zero within the errors, but it shows to be aligned with the photometric position angle. The velocity dispersion at large radii decreases very gently with radius, from $\sim 80 \text{ km s}^{-1}$ at $1 R_e$ and to $\lesssim 60 \text{ km s}^{-1}$ at $8 R_e$. This is consistent with the GC study of Pota et al. (2013), whose kinematics reaches $\sim 8R_e$. While the red GCs show a constant rotation of $\sim 50 \text{ km s}^{-1}$, the blue ones do not rotate significantly in agreement with the PNe. The velocity dispersion of the whole GC sample is consistent with that from PNe.
 - **NGC 3379.** This is the largest galaxy in the Leo I group, classified as an E1. The PN system of this galaxy is has been widely studied in literature (Ciardullo et al. 1993; Romanowsky et al. 2003; Sluis & Williams 2006). The velocity maps from Emsellem et al. (2004) reveal that the galaxy rotates regularly along its photometric major axis ($V_{\text{rot}}^{\text{MAX}} \sim 63 \text{ km s}^{-1}$), while the velocity dispersion has a central peak of $\sim 220 \text{ km s}^{-1}$ and it decreases to 180 km s^{-1} around $0.25R_e$. Then, the long slit data of Statler & Smecker-Hane (1999) and Weijmans et al. (2009) show that the profile gently declines to $\sim 120 \text{ km s}^{-1}$ at $\sim 2R_e$. The PN smoothed velocity field is in good agreement with the inner kinematics and extends it to $6R_e$. We find that the galaxy has a rotation along an axis that twists from the photometric major axis position angle ($\sim 70^\circ$ at $\sim 15 \text{ arcsec}$) towards higher values at large radii. We observe the onset of a rotation along the minor axis at $\sim 2.5R_e$. This is reflected in the trend of the fitted kinematic position angle and in the amplitudes of the third order harmonics. The amplitude of this additional rotation is $\sim 30 \text{ km s}^{-1}$. At the moment there are no integral field kinematic data covering these large distances from the center of the galaxy. The velocity dispersion map shows that the decline continues up to $6R_e$ where it reaches values around 70 km s^{-1} .
 - **NGC 3384.** This lenticular galaxy is a member of the Leo I group (Fouque et al. 1992). We find a rotation velocity of $160 \pm 11 \text{ km s}^{-1}$ at $\sim 7R_e$, consistent with the rotation amplitude published by Krajnović et al. (2011). We do not observe any twist of the kinematic position angle or any misalignment with the photometric one, and the motion is that of a regular rotator. The velocity dispersion map in Emsellem et al. (2004) shows that the σ has a double peak at the center ($\sigma \sim 155 \text{ km s}^{-1}$), and a declining profile that reaches $\sim 80 \text{ km s}^{-1}$ at $\sim 0.5R_e$. The PN data show a velocity dispersion of $\sim 80 \text{ km s}^{-1}$. Cortesi et al. (2013b) performed a kinematic analysis of the disk and spheroid component separately.
 - **NGC 3489.** This S0 galaxy belongs to the Leo I group. The ATLAS^{3D} kinematic map (Krajnović et al. 2011) shows a fast rotation along the photometric major axis, whose amplitude reaches $\sim 115 \text{ km s}^{-1}$ at $\sim 1R_e$. PN data show that the halo of this galaxy is also rotating at $> 40 \text{ km s}^{-1}$. The dispersion is around 80 km s^{-1} , consistent with Cappellari et al. (2011).
- We refer to Cortesi et al. (2013b) for a kinematic analysis of the the disk and spheroid component separately.
- **NGC 4278.** This a rather round elliptical galaxy (E1-2), member of the Coma I group. The ATLAS^{3D} kinematic map Krajnović et al. (2011) shows that stars are regularly rotating only within $0.5R_e$ along a direction slightly offset from the photometric major axis, while there is very weak or no rotation farther out. The kinemetry of Foster et al. (2016) shows that the galaxy does not significantly rotate beyond $1R_e$. The PN velocity field does not show any significant rotation, hence also the kinematic position angle is not very well defined. Pota et al. (2013) do not find any evidence of rotation for the metal rich GCs, while the blue metal poor GCs display a weak rotation in the outer regions, along a direction between the major and the minor axes. The velocity dispersion falls rapidly with radius from a central value of $\sim 270 \text{ km s}^{-1}$ to $\sim 120 \text{ km s}^{-1}$ at $\sim 2R_e$ (Bender et al. 1994; Emsellem et al. 2004; Foster et al. 2016). Our map shows values of σ almost constant at 130 km s^{-1} , in good agreement with the literature.
 - **NGC 4339.** This is a member of the Virgo cluster, classified as intermediate type between E0 and S0. It is regular fast rotator, showing no misalignment between the photometric and kinematic major axis (Cappellari et al. 2011). The PN motion agrees with the stellar kinematics showing a rotation of $> 20 \text{ km s}^{-1}$ and a dispersion of 30 km s^{-1} .
 - **NGC 4473.** This Virgo galaxy is an E5 disk elliptical. It is well known for its multicomponent central kinematics (Emsellem et al. 2004): it is a fast but non-regular rotator ($V_{\text{rot}}^{\text{MAX}} = 63 \text{ km s}^{-1}$, along the photometric major axis) and the velocity dispersion map displays a double peak, which have been interpreted as the presence of a counter-rotating disk (Cappellari et al. 2007), maybe formed in binary disk major mergers (Bois et al. 2011). We detect a low amplitude rotation of $\sim 30 \text{ km s}^{-1}$ along a direction tilted with respect to the photometric major axis, suggesting the triaxial shape of the system. Our velocity dispersion map does show the double peaked structure along the major axis, where $\sigma \sim 160 \text{ km s}^{-1}$. The velocity dispersion decreases to $\sim 100 \text{ km s}^{-1}$ at $10R_e$. We are consistent with the slit stellar kinematics presented in Foster et al. (2013), and the comparison with the smoothed kinematic maps from integral light (Arnold et al. 2014) and from GCs (Foster et al. 2013) confirms the presence of similar features as those observed from the PN kinematics. In particular, the analysis of Alabi et al. (2015) shows that the kinematics of the stars is in agreement with the red population of GCs.
 - **NGC 4494.** This is an E1 galaxy in the NGC 4565 group. The velocity map from Krajnović et al. (2011) shows a kinematically distinct core. The galaxy rotates almost constantly at $\sim 60 \text{ km s}^{-1}$ up to $2R_e$ (Napolitano et al. 2009; Foster et al. 2016). In the halo we find that this amplitude decreases with radius. There is evidence for an additional component rotating along the photometric minor axis at $\sim 25 \text{ km s}^{-1}$, but there is no significant trend in the measured kinematic position angle as a function of radius. The velocity dispersion field shows a peak at the center ($\sigma \sim 150 \text{ km s}^{-1}$), and then decreases to $\sim 80 \text{ km s}^{-1}$ at $R \gtrsim 5R_e$, in very good agreement with the stellar kinematics. The four-fold structure that appears in the σ map is a result from smoothing procedure, and reflects some unresolved gradient in the velocity field. The trend of the rotation amplitude from PN kinematics seems to be confirmed by the kinematic map from Arnold et al. (2014), but their data do not extend out in radii enough to

show any rotation along the minor axis. The kinematics of GCs is studied by Foster et al. (2011) out to $\sim 3.5R_e$. They found that the metal poor GCs rotate with similar amplitude as the galaxy stars, while the metal-rich GCs show marginal rotation. The velocity dispersion profile of the global GC sample is consistent with that of the stars.

- **NGC 4564.** NGC 4564 is an E4 galaxy in the Virgo cluster. It shows a prominent stellar disk along the apparent major axis, so that it has also been referred as an S0 (Michard & Marchal 1994). We agree with the stellar kinematic from Foster et al. (2016) and we extend it beyond $6R_e$, where the galaxy is still clearly rotating along its photometric major axis ($V_{\text{rot}} \sim 95 \text{ km s}^{-1}$). The velocity dispersion profile is decreasing, from the central peak of 180 km s^{-1} (Emsellem et al. 2004) to $\sim 70 \text{ km s}^{-1}$ at $6R_e$.
- **NGC 4594.** NGC 4594, also known as "the Sombrero" (M 104), is an SAa galaxy in the Virgo cluster. It has an extremely bright bulge, while its disk contributes only 10% of the light (Burkhead 1986) and has pronounced dust lanes. Kinematics from line-profile measurements show that the galaxy rotates along its major axis reaching $\sim 250 \text{ km s}^{-1}$ at $\sim 1R_e$, where the rotation curve flattens to a constant amplitude (Kormendy & Illingworth 1982; van der Marel et al. 1994). The velocity dispersion declines from the central value of $\sim 280 \text{ km s}^{-1}$ to values close to 100 km s^{-1} where the disk component dominates. The bulge, instead, has a dispersion remarkably constant at 210 km s^{-1} . Our PN data extend beyond $4R_e$, but there is a void in the detections along the major axis because of the dust lane. For this reason the PNe do not trace the fast rotating disk, and their kinematics is in very good agreement with the integrated light kinematics of Kormendy & Westpfahl (1989) from a slit parallel to the major axis, offset by 30 arcsec. Extinction effects are probably also responsible for the slight asymmetry of the smoothed velocity field, so the kinematic analysis is performed on the unfolded catalog. These asymmetries are localized and do not significantly influence the kinematic results. The only studies of the kinematics of this galaxy at large radii is from Bridges et al. (2007) and Dowell et al. (2014) using GCs. The latter sample distances out to $24'$ with 360 tracers, finding little or no evidence of rotation in the GC system as a whole, and within the red and blue subpopulations.
- **NGC 4649.** Also known as M 60, it is an E2 galaxy in the Virgo cluster. The ATLAS^{3D} velocity map shows a regular, cylindrical rotation of $\sim 95 \text{ km s}^{-1}$ (Krajnović et al. 2011), while the dispersion has a central peak of $\sim 370 \text{ km s}^{-1}$ and it decreases to $\sim 260 \text{ km s}^{-1}$ at $0.5R_e$. The kinematic maps from Arnold et al. (2014) shows that the cylindrical rotation extends up to $2R_e$, while there the dispersion is $\sim 120 \text{ km s}^{-1}$. The PN data extend out to $5R_e$. The smoothed PN velocity field shows that the amplitude of the rotation along the photometric major axis decreases with radius, and no rotation along the major axis is detected beyond $2R_e$. In the outer regions, instead, the system rotates along a nearly orthogonal direction, suggesting a triaxial halo. The velocity dispersion is roughly constant at $\sim 200 \text{ km s}^{-1}$ from 1 to $5R_e$. We see a slight bump in the dispersion profile at $3R_e$, but it may be associated to an unresolved velocity gradient. There are several studies of the GC kinematics in NGC 4649. The most recent, with the largest sample is from Pota et al. (2015). The red GCs display a nearly constant rotation with radius along the photometric major axis. Interestingly they observe a dip in rotation velocity between 100 and 200 arcsec, where we observe the inner rotation decreasing also. The blue GCs also show hints of rotation at all radii, with lower amplitude, and generally along the major axis of the galaxy. Blue GCs also appear to have a minor axis rotating component at 300 arcsec, coinciding with the one found with the PNe. The velocity dispersion of GCs is constant at $\sigma = 240 \pm 30 \text{ km s}^{-1}$.
- **NGC 4697.** This is an E6 galaxy in the Virgo southern extension, belonging to a group of 18 galaxies (Garcia 1993). This galaxy has 535 PN detections by Méndez et al. (2001). In this dataset Sambhus et al. (2006) found evidence for two separate PN populations, with different luminosity functions, spatial distributions, and radial velocities. The velocity field shows a rotation along the photometric major axis with amplitude decreasing with radius from the maximum $\sim 90 \text{ km s}^{-1}$ at $1R_e$ to $\sim 15 \text{ km s}^{-1}$ at $2.5R_e$. There is a hint for an increase in the amplitude of rotation at larger distances ($\sim 40 \text{ km s}^{-1}$ at $4.5R_e$). The kinematic position angle is constant, aligned with the photometric major axis. The velocity dispersion steady declines with radius, from $\sim 180 \text{ km s}^{-1}$ at $0.5R_e$ to $\sim 100 \text{ km s}^{-1}$ at $4.5R_e$. The kinematic maps from Krajnović et al. (2011) and Spiniello et al. (2015) show a regular disk-like rotation with a maximum amplitude of 111 km s^{-1} , confirming the trend for the rotation amplitudes that we find. Arnold et al. (2014) also observes a clear decline of the rotation and of the velocity dispersion, in agreement with PN data.
- **NGC 4742.** It is a field E4 galaxy. It is a faint object, so the number of PN detection is relatively low. We detect a rotation of at least $\sim 74 \text{ km s}^{-1}$, in agreement with the stellar kinematics of Davies et al. (1983). The direction of rotation is along a direction misaligned with respect to the photometric position angle. The dispersion decreases from 100 km s^{-1} at 1.5 arcsec (Davies et al. 1983) to below 50 km s^{-1} in the outskirts.
- **NGC 5128.** Also known as Centaurus A, it is a giant elliptical (often classified as a peculiar lenticular) in the Centaurus group. It shows clear signs of its accretion history in tidal streams and shells (Malin et al. 1983; Peng et al. 2002; Crnojević et al. 2016), evidences of major nuclear activity (see e.g. Neumayer 2010), and jet-induced star forming regions (e.g. Rejkuba et al. 2002). Its kinematics probed by PNe reflects the presence of regions which are not in equilibrium, showing a highly asymmetric and rich in subcomponents velocity field. The fit of the $a_0(R)$ parameter of the rotation model (11) in radial bins shows the presence of peculiar bulk motions in groups of PNe. The galaxy shows a rotation along the photometric major axis (NE direction) with increasing amplitude, from $\sim 50 \text{ km s}^{-1}$ at $\sim 200 \text{ arcsec}$ ($\sim 1.5R_e$) till at least $\sim 150 \text{ km s}^{-1}$ at 800 arcsec ($\sim 5R_e$). The system also displays strong rotation ($\sim 100 \text{ km s}^{-1}$) along the minor axis, revealing its triaxiality (see also Peng et al. 2004). The deviation from point-symmetry of the kinematics in almost all the radial bins is accompanied by several features in the light. In particular the outermost regions, from 700 arcsec outwards, show the most important kinematic features (Coccatto et al. 2013), in correspondence to the elongated structure (sketched with black contours in figure B.2) in light extending roughly along the radio jets (Haynes et al. 1983).
- **NGC 5866.** NGC 5866 is an SA0 galaxy belonging to a loose group (Sengupta & Balasubramanyam 2006). It has a narrow, clear-cut dark lane, running along it and making an angle with the major axis. It is a regular, disk-like rotator with a maximum amplitude of $\sim 180 \text{ km s}^{-1}$ (Krajnović et al. 2011) at $\sim 2R_e$, beyond which the rotation decreases to 110 km s^{-1} .




Fig. B.2. Smoothed velocity field and velocity dispersion field of NGC 5128 shown on the DSS image; north is up, east is left. Features in the surface brightness distribution spatially consistent with features in the PN kinematics are contoured in black. The arrow at the bottom left shows the direction of the PA_{phot} .

s^{-1} at $\sim 3 R_e$. Our velocity map shows that the galaxy keeps rotating at approximately constant speed ($\sim 85 \text{ km s}^{-1}$) at large radii (up to $9 R_e$), with a progressive twist of the position angle. The PN velocity dispersion map has constant values around 90 km s^{-1} , within the errors. The PN velocity dispersion profile is constant around $\sim 95 \text{ km s}^{-1}$ from 2 up to $9 R_e$. We refer to Cortesi et al. (in prep.) for a kinematic study of the disk and spheroid components.

- **NGC 7457** This is a field S0. The velocity maps from Emsellem et al. (2004) and Arnold et al. (2014) show that this galaxy is a regular rotator along the major axis. Our PN velocity map shows that the rotation velocity is growing with radius, reaching 100 km s^{-1} at $\sim 3.5 R_e$. No twisting of the kinematic major axis is observed. The velocity dispersion is relatively low, around $\sim 60 \text{ km s}^{-1}$ at $1 R_e$, $\sim 40 \text{ km s}^{-1}$ at $\sim 3 R_e$. There is good agreement between the PN kinematics and the kinemetry from Foster et al. (2016).

Appendix B.2: Slow rotators

- **NGC 1399.** It is a CD galaxy in the Fornax I cluster. We observe a very low amplitude ($31 \pm 30 \text{ km s}^{-1}$) rotation inside $1 R_e$ along the photometric minor axis, and a slow rotation also in the halo ($R \sim 4 R_e$), which almost counter-rotates with respect to the inner regions. The rotation velocity is also very low in the innermost regions as reported by integral light kinematics studies (e.g. Graham et al. 1998; Saglia et al. 2000; Loubser et al. 2008; Scott et al. 2014), which are in good agreement with the PN data. In particular the integral field map from Scott et al. (2014) shows that NGC 1399 does not display important ordered motions in the central regions, while at a radius of $\sim 30''$ it rotates along the minor axis ($\sim 45 \text{ km s}^{-1}$). The velocity dispersion, instead, is relatively high: it rises steeply in the inner 10 arcsec, reaching $\sim 370 \text{ km s}^{-1}$ at the center (e.g. Saglia et al. 2000), while at

large radii (from $\sim 1 R_e$) the PN σ flattens at $\lesssim 200 \text{ km s}^{-1}$. McNeil et al. (2010) showed that kinematics of the red GCs of NGC 1399 is in excellent agreement with the PNe.

- **NGC 3608.** This is an E2 galaxy in the Leo II group. It is known to have a counter-rotating core (Jedrzejewski & Schechter 1988) in the central $0.27 R_e$, aligned with the photometric major axis and rotating at $\sim 20 \text{ km s}^{-1}$ (Halliday et al. 2001; Emsellem et al. 2004). From $\sim 1 R_e$ galaxy starts to rotate at approximately along the photometric major axis (Foster et al. 2016). The PN kinematics shows that the growing trend continues at larger radii, where the amplitude of rotation reaches $\sim 65 \text{ km s}^{-1}$. The twist of the PA_{kin} reveals that the system is triaxial. The velocity dispersion decreases from the central value of $\sim 220 \text{ km s}^{-1}$ to 100 km s^{-1} at $\sim 1 R_e$ (Halliday et al. 2001). We find a constant velocity dispersion profile in the halo ($\sim 90 \text{ km s}^{-1}$).
- **NGC 3923.** This is an E4-5 galaxy in a group of 8 galaxies (Brough et al. 2006). We detect a cylindrical velocity field along an intermediate direction between the minor and the major axes, which rotate at $\sim 75 \text{ km s}^{-1}$. The velocity dispersion is constant around 235 km s^{-1} . Long-slit kinematics from Carter et al. (1998) inside $\sim 1 R_e$ shows that NGC 3923 does not rotate significantly along the photometric major axis for $R \lesssim 0.5 R_e$, but it displays a weak rotation ($\sim 20 \text{ km s}^{-1}$) along the photometric minor axis. They also found a falling dispersion profile, from the central value of $\sim 285 \text{ km s}^{-1}$ to $\sim 240 \text{ km s}^{-1}$ at $0.7 R_e$. Norris et al. (2012) studied the kinematics of 79 GC/UCD extended to more than $6 R_e$. They do not show appreciable rotation, and their velocity dispersion seem constant with radius.
- **NGC 4365.** It is an E3 galaxy, one of the brightest member of the Virgo cluster. It has a complex kinematic structure with a counter-rotating core (Bender 1988a), aligned with the photometric major axis, and a prolate rotation of $\sim 61 \text{ km s}^{-1}$ (Davies et al. 2001), which indicates a triaxial potential.




Fig. B.3. Smoothed velocity field and velocity dispersion field of NGC 4472 shown on the DSS image; north is up, east is left. The black cross marks the position of the in-falling dwarf UGC 7636. The arrow at the bottom right shows the direction of the PA_{phot} .

Our PN velocity field extends to $6R_e$. The inner kinematics ($R \lesssim 2R_e$) is compatible with no rotation. We do not detect the rolling about the minor axis that Arnold et al. (2014) report, probably because of the smoothing over the inner velocity gradients. We do measure a significant outer ($R \gtrsim 3R_e$) rotation of $\sim 50 \text{ km s}^{-1}$ along the major axis, counter-rotating with respect to the kinematically decoupled core. The velocity dispersion map has almost constant values, around $\sim 150 \text{ km s}^{-1}$, with slightly lower values in the region along the major axis ($\sim 140 \text{ km s}^{-1}$). The velocity dispersion measured by Davies et al. (2001) shows a inner value of $> 250 \text{ km s}^{-1}$ which decreases to $\sim 185 \text{ km s}^{-1}$ at $0.5R_e$. Arnold et al. (2014) measure a $\sigma \sim 150 \text{ km s}^{-1}$ at $2R_e$, consistent with the values found with PNe. Blom et al. (2012), studying the GC kinematics, found that the intermediate metallicity subpopulation (green) rotates along the photometric minor axis, as the stellar population at intermediate radii, while the red GCs rotate only at large radii along the photometric major axis in the opposite direction of the core of the galaxy, in agreement with the PN kinematics at large radii.

- **NGC 4374.** Also known as M84, it is a bright E1 galaxy in the Virgo cluster. The galaxy does not show significant rotation in the innermost $\sim 0.4R_e$, as reported by Emsellem et al. (2004) and Foster et al. (2016), while the velocity dispersion steeply decreases from 310 km s^{-1} at the center to $\sim 200 \text{ km s}^{-1}$ at $\sim 1R_e$. The PN velocity field shows evidence for rotation only at large radii ($R \gtrsim 2R_e$), with constant amplitude $\sim 60 \text{ km s}^{-1}$ and along a direction compatible with the minor axis. The velocity dispersion is rather constant around $\sim 220 \text{ km s}^{-1}$ up to $6R_e$. Our kinematics is in agreement with that from the SLUGGS velocity maps (Arnold et al. 2014; Foster et al. 2016), but their data extend only to $\sim 2R_e$. The long slit kinematics of Coccato et al. (2009) reaches almost $3R_e$ and agrees in showing no signs of rotation along the major axis, the onset of rotation along the minor, and constant velocity dispersion in the outer regions around $\sim 230 \text{ km s}^{-1}$.

- **NGC 4472.** NGC 4472 (M 49) is an E2 galaxy, the brightest object in the Virgo cluster. It is surrounded by a complex system of diffuse shells and other features in light which are evidences of its recent and still ongoing accretion activity (see Janowiecki et al. 2010; Arrigoni Battaia et al. 2012, Hartke et al. in prep.). It is classified as a non regular rotator by Krajnović et al. (2011), who also found evidence for a counter-rotating core. From $R \sim 0.2R_e$ the galaxy rotates at $\sim 60 \text{ km s}^{-1}$ (Cappellari et al. 2011; Veale et al. 2017). Beyond $\sim 3R_e$ the PN data show a complicated and out of equilibrium kinematics. The halo rotates along a direction compatible with the galaxy minor axis. This motion is neither point- nor axisymmetric, and is dominated by the kinematics of the in-falling satellite, the dwarf UGC 7636. A visual inspection of the smoothed velocity field reveals that the main body of the galaxy would show major axis rotation once the PNe belonging to the satellite are excluded (see Hartke et al. in prep. for a detailed study). While the velocity dispersion profile is flat over $4R_e$ at $\sim 250 \text{ km s}^{-1}$, we notice a high dispersion feature with $\sigma \sim 350 \text{ km s}^{-1}$ in a position corresponding to the coordinates of UGC 7636.

- **NGC 4552.** NGC 4552 (M 89) is an E0-1 galaxy in Virgo. It known to possess a kinematically distinct core, with the innermost region rotating at $\sim 30 \text{ km s}^{-1}$, and the region outside $0.3R_e$ having very weak rotation (Simien & Prugniel 1997b; Emsellem et al. 2004). Our PN smoothed velocity field shows that the galaxy start rotating beyond $3R_e$. The smoothed velocity field reveal that the rotation is about two perpendicular directions. This is reflected in the twist of the kinematic position angle with radius. The velocity dispersion map from the data is consistent with the one from the integrated light (Emsellem et al. 2004), and it shows constant values with radius, around 180 km s^{-1} .
- **NGC 4636.** This is an E0-1 galaxy in the Virgo cluster. The velocity map from Krajnović et al. (2011) does not show significant rotation in the innermost region ($R < 0.25R_e$). The


velocity dispersion, instead, has a central value of $\sim 240 \text{ km s}^{-1}$ and it decreases to 190 km s^{-1} at $0.25R_e$. The long slit kinematics from Pu & Han (2011) show that the galaxy starts rotating along the minor axis at $\sim 40 \text{ arcsec}$ ($0.2 R_e$), and along the major axis at $\sim 85 \text{ arcsec}$ ($\sim 1R_e$). The σ profile is gently decreasing along the minor axis, reaching $\sim 100 \text{ km s}^{-1}$ at $0.5 R_e$, while it is rather flat along the major, around 190 km s^{-1} , then it decreases to $\sim 150 \text{ km s}^{-1}$ where the rotation starts. We detect a weak rotation ($42 \pm 28 \text{ km s}^{-1}$) along a direction compatible with the minor axis. In addition, in the smoothed velocity map there is evidence for a rotation of comparable magnitude at larger radii ($R \sim 1R_e$) around the major axis, in very good agreement with Pu & Han (2011). The PN velocity dispersion profile decreases abruptly from $\sim 180 \text{ km s}^{-1}$ at $R < 2R_e$ to $\sim 110 \text{ km s}^{-1}$ at $3R_e$. Schuberth et al. (2012) find a strong rotation signature for the blue GCs ($V \sim 88 \text{ km s}^{-1}$ along the major axis, but in opposite direction with respect to the outermost PNe). The red GCs have a weaker rotation ($V \sim 30 \text{ km s}^{-1}$) along an axis that changes with being aligned with the minor axis to the photometric major axis, in agreement with the PN kinematics. For radii beyond $1.5R_e$ the axis of rotation remains constant along the direction of the major axis. The dispersion profile of the blue GC declines with radius, while the red GCs show values of $\sigma \sim 135 \text{ km s}^{-1}$.

- **NGC 5846.** This is an E0 galaxy, the brightest member in a group of ten. Emsellem et al. (2004) detect a low rotation of $\lesssim 30 \text{ km s}^{-1}$ and a high central dispersion of 255 km s^{-1} . PN smoothed velocity field is consistent with low amplitude rotation. We find constant values for the velocity dispersion around 180 km s^{-1} . The SLUGGS kinematic maps Arnold et al. (2014); Foster et al. (2016) are consistent with PN data. Pota et al. (2013) showed that neither the GC subpopulations show significant rotation. The velocity dispersion of the red GCs has a flat profile, around values comparable with those found for the PNe. The blue GCs have systematically higher $\sigma \sim 260 \text{ km s}^{-1}$.


Appendix C: Velocity and Velocity Dispersion Fields

The following figures show the smoothed velocity and the velocity dispersion fields for all the galaxies of the sample. The maps are interpolated on a regular grid of coordinates, as described in section 6.1. The fields are rotated such that the photometric major axis is horizontal; the orientation is specified by the direction of the arrows indicating North and East. The X and Y axes units are arcseconds.


NGC0584




NGC0821




NGC1023




NGC 1316




NGC 1344




NGC 1399




NGC2768




NGC2974




NGC3115




NGC3377




NGC3379




NGC3384




NGC3489




NGC3608




NGC3923




NGC4278




NGC4339




NGC4365




NGC4374




NGC4472




NGC4473




NGC4494




NGC4552




NGC4564




NGC4594




NGC4636




NGC4649




NGC4697




NGC4742




NGC5128



NGC5846



NGC5866



NGC7457

