

Revised SED of the triple protostellar system VLA 1623–2417

N. M. Murillo¹, D. Harsono¹, M. McClure², S.-P. Lai^{3,4}, and M. R. Hogerheijde^{1,2}

¹ Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA, Leiden, the Netherlands
e-mail: nmurillo@strw.leidenuniv.nl

² Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, The Netherlands

³ Institute of Astronomy and Department of Physics, National Tsing Hua University, 101 Section 2 Kuang Fu Road, Hsinchu 30013, Taiwan

⁴ Academia Sinica Institute of Astronomy and Astrophysics, P.O. Box 23-141, Taipei 10617, Taiwan

July 3, 2018

ABSTRACT

Context. VLA 1623–2417 is a triple protostellar system deeply embedded in Ophiuchus A. Sources A and B have a separation of 1.1", making their study difficult beyond the submillimeter regime. Lack of circumstellar gas emission suggested that VLA 1623 B has a very cold envelope and is much younger than source A, generally considered the prototypical Class 0 source.

Aims. We explore the consequences of new ALMA Band 9 data on the spectral energy distribution (SED) of VLA 1623–2417 and their inferred nature.

Methods. Using dust continuum observations spanning from centimeter to near-infrared wavelengths, the SED of each component in VLA 1623–2417 is constructed and analysed.

Results. The ALMA Band 9 data presented here show that the SED of VLA 1623–2417 B does not peak at 850 μm as previously expected, but instead presents the same shape as VLA 1623–2417 A at wavelengths shorter than 450 μm .

Conclusions. The results presented here indicate that the previous assumption that the flux in *Herschel* and *Spitzer* observations is solely dominated by VLA 1623–2417 A is not valid, and instead, VLA 1623–2417 B most likely contributes a significant fraction of the flux at $\lambda < 450 \mu\text{m}$. These results, however, do not explain the lack of circumstellar gas emission and puzzling nature of VLA 1623–2417 B.

Key words. stars: formation - stars: low-mass - ISM: individual objects: VLA 1623–2417 - methods: observational - techniques: interferometric

1. Introduction

VLA 1623–2417 (hereafter VLA 1623) is a deeply embedded protostellar system in Ophiuchus A (L1688), and first observed by André et al. (1990). Recent observations place L1688 and VLA 1623 at a distance of 137.3 ± 1.2 pc (Ortiz-León et al. 2017). VLA 1623 is generally considered to be the prototypical Class 0 source (André et al. 1993). Observations at centimeter wavelengths with the Very Large Array (VLA) revealed several continuum peaks (Bontemps & André 1997), which were interpreted as the central object of VLA 1623 and the jet knots A, B and C. However, observations of the outflow suggested more than one component is present in VLA 1623 (Dent et al. 1995; Yu & Chernin 1997; Caratti o Garatti et al. 2006). Interferometric observations in submillimeter revealed that VLA 1623 is actually a triple protostellar system (Looney et al. 2000; Murillo & Lai 2013), with each component driving an outflow of its own (Murillo & Lai 2013; Santangelo et al. 2015). The VLA interferometric centimeter peaks A and B matched with sources B and W, respectively (Ward-Thompson et al. 2011; Murillo & Lai 2013). No submillimeter counterpart was observed for the centimeter peak C detected with the VLA.

VLA 1623 A is brightest in the submillimeter wavelength range, presenting a centrally peaked source with extended dust emission coinciding with the Keplerian disk traced in C^{18}O (Murillo et al. 2013). VLA 1623 B is separated from source A by 1.1" to the west, and presents a strongly collimated outflow (Santangelo et al. 2015). Interferometric observations do

not detect warm (C^{18}O , $c\text{-C}_3\text{H}_2$) or cold (DCO^+ , N_2D^+ , N_2H^+) gas emission associated with VLA 1623 B (Murillo et al. 2013, 2016, 2018). Submillimeter Array (SMA, Ho et al. 2004) observations detected SO peaking between sources A and B (Murillo & Lai 2013). Even more interesting is the detection of a variable H_2O maser centered on VLA 1623 B (Furuya et al. 2003), detected between 8 to 13 km s^{-1} . Suggestions have been made that VLA 1623 B could be a background object, such as a galaxy. The velocity shift of the H_2O maser would suggest a redshift of $3\text{--}4 \times 10^{-5}$. Thus, if the maser is associated with VLA 1623 B, then this source is located within our galaxy. VLA 1623 W, separated by 10" from VLA 1623 A, presents hints of a rotating envelope or disk-like structure, and a slightly different systemic velocity ($\sim 1 \text{ km s}^{-1}$) from its companions VLA 1623 A and B ($\sim 3.5 \text{ km s}^{-1}$) (Murillo et al. 2013).

In this paper, we present Atacama Large Millimeter/submillimeter Array (ALMA) Cycle 2 Band 9 dust continuum observations of VLA 1623. These data, combined with previous ALMA Band 6 and 7 observations, continuum fluxes found in literature, and fluxes from *Spitzer* and *Herschel* observations, are used to construct and analyze the spectral energy distribution (SED) of each component in VLA 1623. In addition, newly reduced *Spitzer* IRS spectra for VLA 1623 is also presented here.

2. Observations

Table 1 details the ALMA observations presented here, including the field of view, angular resolution and typical noise level.

Cycle 2 observations in Band 9 (Project code: 2012.1.00707.S; PI: D. Salter) have phase center $\alpha_{J2000} = 16:26:26.42$; $\delta_{J2000} = -24:24:30.0$. However the observations required rephasing, and the phase center was shifted to the position of VLA 1623 A, $\alpha_{J2000} = 16:26:26.39$; $\delta_{J2000} = -24:24:30.688$, consistent with the Cycle 2 Band 6 and 7 observations (see below). The spectral set-up was configured to observe mainly CO, HCO⁺ and continuum. J1924–2914 was used for bandpass calibration. Phase calibration was done using the check source J1700–2610, with a time bin of 24 seconds, instead of 6 seconds. Flux calibration was done with J1733–130. Ceres was observed, but it was resolved on some baselines, and thus did not provide a reliable flux calibration. Observations toward VLA 1623 did not have any system temperature measurements, so the system temperature of the phase calibrators was used.

Cycle 2 observations in Bands 6 and 7 (Project code: 2013.0.01004; PI: S.-P. Lai) have a phase center of $\alpha_{J2000} = 16:26:26.390$; $\delta_{J2000} = -24:24:30.688$, and targeted several molecular lines (e.g., DCO⁺, CO isotopologues and c-C₃H₂ in Band 6, DCO⁺, N₂H⁺ and H₂D⁺ in Band 7) along with continuum. The molecular line observations are detailed in Santangelo et al. (2015); Murillo et al. (2015, 2018). ALMA Cycle 0 Band 6 observations (Project code: 2011.0.00902.S; PI: N. Murillo) of VLA 1623, with phase center $\alpha_{J2000} = 16:26:26.419$; $\delta_{J2000} = -24:24:29.988$, slightly offset from the position of VLA 1623 A, were previously reported in Murillo et al. (2013).

The Spitzer IRS spectra are new reductions of serendipitous observations of VLA 1623. The VLA 1623 W and blended VLA 1623 A & B components were aligned along the LL slit of the second target in the clustered AOR 12697600 (PID 2) and are visible in the second nod. The basic-calibrated data were downloaded from the Spitzer Heritage Archive and reduced by hand using the AdOpt optimal point source extraction routines from the CASSIS low-resolution pipeline update, version 7 (Lebouteiller et al. 2010) provided in SMART (Higdon et al. 2004). The VLA 1623 W and blended VLA 1623 A & B components were fit simultaneously, along with a low-order polynomial sky background, at each wavelength. The resulting spectra are complete in LL (14–35 microns) for source W, but the flux of the blended sources A & B is only detected longwards of 19 microns.

3. Results

Measured fluxes from ALMA observations are listed in Table 2. VLA 1623 A presents a central bright component surrounded by extended emission in all bands (Fig. 1). In the Band 9 observations, source A does not show further multiplicity, and the clumpiness of the extended dust emission is most likely due to the difficulty of recovering extended structure with high spatial resolution observations. The extended dust emission from Cycle 2 Band 6 data was found to have a radius of ~ 100 AU (Persson et al. 2016), smaller than that of the Keplerian disk (150 AU) traced with C¹⁸O (Murillo et al. 2013). In contrast, VLA 1623 B is observed to be compact, almost point-like in all Bands. The extended emission in the Band 9 image is most likely not real, and product of the difficulty of recovering structure with high spatial resolution observations. Similarly, VLA 1623 W is also




Fig. 1. VLA 1623 observed with ALMA. The phase center for all observations is similar and located around the position of VLA 1623 A. Contours are in steps of 3, 8, 15, 20, 50 σ , with σ listed in Table 1. The green ellipses show the beamsize of the observations. The green box in the middle panel outlines the area of the Band 9 observations presented in the bottom panel.

point-like in Bands 6 and 7, but with a continuum flux 3 to 5 times weaker than VLA 1623 B (Table 2). Fluxes for each component are measured using a region which is defined by the 3 σ contour of the continuum emission. For Band 9 continuum, we use the same region used for Band 6. For VLA 1623 A, the extended dust disk component is included in the region used for flux measurement. Measured fluxes are listed in Table 2. VLA 1623 A is brighter than VLA 1623 B by a factor up to 2, although this difference in fluxes might be caused by the inclusion of the dust disk in the flux of VLA 1623 A, and assuming that VLA 1623 B has no disk. VLA 1623 W is dimmer than VLA 1623 A and B by about a factor of 5 to 12.

Table 1. ALMA observations

Project ID	Band	Wavelength	UV-baseline range k λ	Largest scale "	Field of view "	Beam "	P.A. °	Noise mJy beam ⁻¹
2011.0.00902.S	6	1.3 – 1.4 mm	25 ~ 310	2.5	28.1	0.79'' \times 0.54''	-85.9	3.5
2013.0.01004.S	6	1.3 – 1.4 mm	18 ~ 791	2.0	28.8	0.45'' \times 0.27''	78.4	12
2013.0.01004.S	7	828 – 832 μ m	18 ~ 420	3.1	17.2	0.86'' \times 0.53''	-87.1	4.2
2012.1.00707.S	9	421 – 429 μ m	30 ~ 2280	1.6	9.0	0.25'' \times 0.14'' ^a	-85.1	23.9 ^a
2012.1.00707.S	9	421 – 429 μ m	30 ~ 2280	1.6	9.0	0.17'' \times 0.1'' ^b	85.6	14.3 ^b

Notes. ^(a) Continuum image generated with natural weighting. ^(b) Continuum image generated with robust (Briggs) weighting.

4. Analysis

In order to construct the SEDs of VLA 1623, the fluxes presented in this work are used, together with those found in literature. To constrain the peak of the SEDs, archival *Herschel Space Observatory* (Pilbratt et al. 2010) PACS photometric maps for Ophiuchus from the Gould Belt Survey (André et al. 2010; Pezzuto et al. 2012) and from the program of PI: P. Ábrahám (Alves de Oliveira et al. 2013) were obtained from the *Herschel Science Archive* (Observation IDs: 134222148 and 1342238816). The maps produced with the JScanmap method were used. The 160 and 100 μ m maps show the emission toward VLA 1623 to be concentrated mainly on the position of VLA 1623 A and B. A distinct VLA 1623 W emission is not resolved at either wavelength, most likely due to the resolution of the *Herschel* observations. At 70 μ m, two distinct continuum emission components are detected in the maps, one centered on VLA 1623 A & B, and the other on VLA 1623 W. Both components show extended emission, most likely arising from their outflow cavities. Due to the nature of the sources, PSF photometry is performed on the PACS maps to extract fluxes using *StarFinder* (Diolaiti et al. 2000). The PSF for each wavelength is extracted from the corresponding PACS maps using sources that appear as point sources and have little to no extended emission around them. An aperture mask is applied to the PSF, thus minimizing the extended emission that can be included during photometry. Due to the aperture mask, an aperture correction factor obtained from Balog et al. (2014) is applied to the fluxes. At 160 and 100 μ m, deblending of the detected sources was done in order to obtain fluxes for VLA 1623 W. The method is further detailed in Murillo et al. (2016) and Diolaiti et al. (2000).

Table 2 lists the fluxes and Fig. 2 shows the SED of each component. At wavelengths smaller than 450 μ m, VLA 1623 A and B are spatially unresolved. Observations with SMA at 1.3 mm and BIMA at 2.7 mm (Looney et al. 2000; Murillo & Lai 2013) detect all three components. In both datasets, VLA 1623 A is the brightest of the three sources, while VLA 1623 B and W are point-like. The extended emission around VLA 1623 A is resolved with the SMA (synthesized beam: 0.61'' \times 0.43'', 10°) but not in the BIMA observations (synthesized beam: 0.95'' \times 0.39'', 18°). Spitzer observations detect VLA 1623 W in all IRAC and MIPS channels, while unresolved VLA 1623 A and B are detected only in MIPS1 (Murillo & Lai 2013), indicating that both sources are deeply embedded. IRS spectra for VLA 1623 W, and unresolved VLA 1623 A and B are included in the SEDs.

The SEDs are fit using the Scipy method `curve_fit`¹ with a single-temperature graybody, and with optical depth $\tau \propto \nu^\beta$, where β is the dust emissivity index. The fit is done in the wavelength range from 7 mm to 22 μ m. The best fit curves are shown

in Fig. 2. The graybody fits are practically the same with or without the *Herschel* PACS fluxes, which is to be expected since the *Herschel* fluxes are picking up the envelope material as well. In the submillimeter regime, the SEDs of VLA 1623 A and B are quite similar, as shown in Fig. 2 with open and filled symbols. For sources A and B, the unresolved emission at $\lambda < 450 \mu\text{m}$ is fitted in three ways. First, assuming all the infrared emission belongs to one of the sources (Fig. 2 solid red line). Second, the infrared emission is evenly distributed between both sources (Fig. 2 dashed red line). Third, VLA 1623 B only contributes one-tenth of the infrared emission (Fig. 2 dotted red line). The resulting dust temperatures T_{dust} are similar for all three fits, and for both systems, between 40 to 50 K. However, it is not certain if the emission is evenly distributed at all wavelengths or varies with wavelength. Thus, there are large uncertainties in the SED fit of VLA 1623 A and B. However, the derived dust temperature for VLA 1623 B (38 to 49 K) is larger by an order of magnitude than previously expected ($T_{\text{dust}} = 5$ K, Murillo & Lai 2013). The derived dust temperature for VLA 1623 W (66 K) is also higher than previously reported ($T_{\text{dust}} = 41$ K, Murillo & Lai 2013). The higher dust temperature is most likely due to the addition of the IRS spectra to the SED, which was not included in previous studies.

The bolometric luminosity is calculated based on the curve of the fits to the constructed SEDs using the trapezoid method. Considering the three cases used for SED fitting, the bolometric luminosity of VLA 1623 A is between $\sim 0.4 - 0.5 L_\odot$, while for VLA 1623 B it is between $\sim 0.2 - 0.3 L_\odot$. It must be stressed that due to the SED being unresolved at wavelengths shorter than 450 μ m for VLA 1623 A and B, parameters derived from the SED are uncertain, and these bolometric luminosities are rough estimates. In addition, source A appears brighter since the emission from the dust disk is being included. For VLA 1623 W, the derived bolometric luminosity is on the order of $0.13 \pm 0.01 L_\odot$.

To calculate the envelope mass, the formula from Jørgensen et al. (2009), which takes into account that brighter sources have somewhat higher dust temperatures, is used and expressed as:

$$M_{\text{env}} = 0.44 M_\odot \left(\frac{L_{\text{bol}}}{1 L_\odot} \right)^{-0.36} \left(\frac{S_{850\mu\text{m}}}{1 \text{ Jy beam}^{-1}} \right)^{1.2} \left(\frac{d}{125 \text{ pc}} \right)^{1.2}. \quad (1)$$

For this calculation, the 850 μ m fluxes obtained with ALMA Cycle 2 Band 7 observations are used, in conjunction with the bolometric luminosities derived from the SEDs and described above. This method yields a mass of ~ 0.8 and $0.2 M_\odot$ for VLA 1623 A and B, respectively, for a distance of 137.3 pc. For VLA 1623 W, the mass is $0.04 \pm 0.005 M_\odot$. If using the previous distance to VLA 1623 of 120 pc (Loinard et al. 2008), the masses are decreased by 15%. Several caveats need to be highlighted here. The 850 μ m flux from ALMA Cycle 2 observations only recovers emission from the disk and the inner envelope (up to 3'', ~ 400

¹ <https://www.scipy.org/>




Fig. 2. SED of VLA 1623 including the Spitzer IRS spectra. The red curves show the best fit graybody to each SED, with the dust temperature T_{dust} and dust emissivity index β indicated in each panel. For VLA 1623 A (top) and B (middle), the dashed curves show the best fit assuming that the emission at $\lambda < 450 \mu\text{m}$ is evenly distributed among the two sources. The dotted curve assumes VLA 1623 B only contributes one-tenth of the total emission for $\lambda < 450 \mu\text{m}$. The open symbols in the top and middle panel are the fluxes for VLA 1623 B and VLA 1623 A, respectively, and are included for ease of comparison.

AU), but not from the large scale gas (> 400 AU). Thus the mass estimate reported here is not representative of the whole envelope mass. Furthermore, if VLA 1623 B is observed through the envelope of A, then the mass estimate for B is in fact not indicative of its actual mass.

5. Summary

In this paper we present new ALMA Band 9 continuum data toward VLA 1623, resolving the components A and B. PSF photometry of *Herschel* PACS maps provides unresolved fluxes between 70 to 160 μm , which trace the peak of the SED. The updated SEDs show that VLA 1623 B does not peak at 1000 μm , as suggested by Murillo & Lai (2013) based on SMA 850 μm observations, but might instead peak at around 100 μm , similar to VLA 1623 A. The best fit using a graybody indicates that the dust temperature of both sources is in the 40 to 50 K range. There are large uncertainties, however, due to the unresolved fluxes of VLA 1623 A and B at wavelengths shorter than 450 μm , and thus the derived dust temperature should be considered with caution. With these results, the previous assumption that the continuum flux at shorter wavelengths is dominated by VLA 1623 A is not well based, and instead VLA 1623 B could be the dominant source in the infrared regime.

Deriving source parameters or evolutionary stage from unresolved SEDs is non-trivial, and can lead to under- or over-estimation of the parameters (see e.g., Murillo et al. 2016). Keeping this caveat in mind, the bolometric luminosity derived for VLA 1623 A and B from the SEDs presented in this work, together with the rough estimate of the disk and inner envelope mass on scales up to 400 AU, provide constraints on the source parameters for each component, albeit with large uncertainties. The bolometric luminosity for VLA 1623 A and B varies by a factor of ~ 2 , and the mass of the disk plus inner envelope varies by a factor of ~ 4 since it is dependent on the luminosity as well as the 850 μm flux. All parameters derived from the SED indicate that both sources are Class 0. In the scenario where source B would dominate the infrared emission, its bolometric luminosity would rise and the mass would decrease. For VLA 1623 W, the luminosity is lower by a factor of ~ 4 compared to source A, while the envelope mass is lower by a factor of 20 compared to that of source A, consistent with its classification as a Class I source.

Along with separating the fluxes of components in a multiple protostellar system, the variability of the sources must also be taken into account when understanding the nature of these systems. Recent studies show variation in the brightness of protostellar systems, possibly due to episodic accretion processes (e.g., Herczeg et al. 2017; Johnstone et al. 2018). For the case of VLA 1623, however, all three components have not shown significant variations in brightness in the submillimeter range. This can be noted in the 1.3 mm fluxes (Table 2) which have been obtained with the SMA (2007 and 2009), and with ALMA in Cycle 0 (2011) and Cycle 2 (2013). The relative similarity of the fluxes at 1.3 mm suggests that there has been no large variations in brightness. There could be variations on the order of 10% at 1.3 mm, but given the uncertainties of the flux calibration, such variations would not be detected. In addition to variability in brightness, chemical signatures in the envelope gas can be used to probe episodic accretion (Visser et al. 2015; Jørgensen et al. 2015; Frimann et al. 2017; Hsieh et al. 2018). Considering the characterized gas structure of VLA 1623 (Murillo et al. 2013, 2015, 2018), no chemical signatures of luminosity variations toward the components of VLA 1623 are detected.

The presented results contribute to the puzzle of VLA 1623 B. The derived dust temperature from the SED fitting (40 to 50 K) suggests there should be warm gas emission from this source. However, no circumstellar gas emission in either cold or warm molecules (Murillo et al. 2013, 2015, 2018) has been detected to be associated with VLA 1623 B. In contrast, VLA 1623 B drives

Table 2. SED for each component in VLA1623

	VLA 1623 A		VLA 1623 B		VLA 1623 W			
RA	16:26:26.390		16:26:26.309		16:26:25.636			
Dec	−24:24:30.688		−24:24:30.588		−24:24:29.488			
Wavelength μm	S_{int} mJy	error mJy	S_{int} mJy	error mJy	S_{int} mJy	error mJy	Ref.	Telescope
3.6					0.74	0.12	1	Spitzer/IRAC1
4.5					2.35	0.19	1	Spitzer/IRAC2
5.8					3.26	0.33	1	Spitzer/IRAC3
8.0					2.60	0.26	1	Spitzer/IRAC4
24		40.4 ± 4.2^a			164.26	4.54	2,1	Spitzer/MIPS1
70 ^b		17424.5 ± 3.2^a			5790.8	3.88	3	Herschel/PACS
100 ^b		55092.3 ± 8.7^a			10284.9 ^c	8.6	3	Herschel/PACS
160 ^b		81340.6 ± 10.3^a			21457.2 ^c	11.5	3	Herschel/PACS
450	3820.0	4.2	1723.0	5.7			3	ALMA/Band 9 Natural weighting
450	2910.0	6.4	1653.0	6.8			3	ALMA/Band 9 Briggs weighting
850	1139.5	3.4	373.0	4.6	69.0	7.0	3	ALMA/Band 7
1300	184.0	13.8	104.0	15.3	38.0	3.8	3	ALMA/Band 6 Cycle 0
1300	178.0	5.1	106.0	5.3	30.0	3.0	3	ALMA/Band 6 Cycle 2
1300	152.0	8.0	106.0	8.0	25.7	8.0	4	SMA
2700	25.5	6.3	25.8	3.5	3.1	1.5	5	BIMA
7000	1.2	0.2	2	0.2			6	VLA
36000	0.13	0.02	0.4	0.02	0.2	0.02	6,7	VLA
44000	<0.2		<0.2				6	MERLIN

Notes. ^(a) Unresolved fluxes. ^(b) Fluxes obtained with *StarFinder* PSF photometry using an aperture mask of 11.2", 12.8" and 19.2" for 70, 100 and 160 μm , respectively. The aperture correction factors are 0.791, 0.795 and 0.789 for 70, 100 and 160 μm , respectively (Balog et al. 2014). ^(c) VLA 1623 W fluxes obtained using the deblend option in *StarFinder*, since no distinct peak for VLA 1623 W is observed.

References. (1) Gutermuth et al. (2009); (2) c2d catalog; (3) this work; (4) Murillo & Lai (2013); (5) Looney et al. (2000); (6) Ward-Thompson et al. (2011); (7) Bontemps & Andre (1997)

a fast and collimated jet and outflow (Santangelo et al. 2015), and presents a variable H₂O maser (Furuya et al. 2003). The lack of detected circumstellar gas emission around VLA 1623 B is difficult to explain. One possibility could be gas stripping due to interaction with the outflow from VLA 1623 A or VLA 1623 W (e.g. B1-b: Hirano & Liu 2014), but this would not explain why the dust was not stripped or presence of the outflow. Another explanation would be that VLA 1623 B is more evolved than VLA 1623 A. However this would not fully explain the complete lack of gas in submillimeter when VLA 1623 W does present circumstellar gas (Murillo et al. 2013). A final possibility would be that VLA 1623 B is being obscured, either by the envelope of VLA 1623 A, or circumstellar matter with high optical depth. This would also introduce further uncertainties to the calculated envelope mass. Observations with subarcsecond resolution in the infrared are required to determine the nature of this object, for which JWST will be ideal.

Acknowledgements. This paper made use of the following ALMA data: ADS/JAO.ALMA 2011.0.00902.S, 2012.1.00707.S, and 2013.1.01004.S. ALMA is a partnership of ESO (representing its member states), NSF (USA), and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO, and NAOJ. The authors acknowledge assistance from Allegro, the European ALMA Regional Center node in the Netherlands. Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. PACS has been developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany); INAF-IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); IAC (Spain). This development has been supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy), and CICYT/MCYT (Spain). This work is based in part on observations made with the Spitzer Space Telescope,

which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. NMM and DH are supported by the European Union A-ERC grant 291141 CHEMPLAN, by the Netherlands Research School for Astronomy (NOVA), by a Royal Netherlands Academy of Arts and Sciences (KNAW) professor prize. MM receives funding through a Marie Skłodowska-Curie action in the Horizon 2020 program. SPL acknowledges support from the Ministry of Science and Technology of Taiwan with Grant MOST 106-2119-M-007-021-MY3. MRH is support by grant 614.001.352 from the Netherlands Organisation for Scientific Research (NWO). The authors wish to thank Doug Johnstone for the helpful and insightful comments that greatly improved this work.

References

- Alves de Oliveira, C., Ábrahám, P., Marton, G., et al. 2013, *A&A*, 559, A126
 André, P., Martin-Pintado, J., Despois, D., & Montmerle, T. 1990, *A&A*, 236, 180
 André, P., Men'shchikov, A., Bontemps, S., et al. 2010, *A&A*, 518, L102
 André, P., Ward-Thompson, D., & Barsony, M. 1993, *ApJ*, 406, 122
 Balog, Z., Müller, T., Nielbock, M., et al. 2014, *Experimental Astronomy*, 37, 129
 Bontemps, S. & Andre, P. 1997, in *IAU Symposium*, Vol. 182, Herbig-Haro Flows and the Birth of Stars, ed. B. Reipurth & C. Bertout, 63
 Caratti o Garatti, A., Giannini, T., Nisini, B., & Lorenzetti, D. 2006, *A&A*, 449, 1077
 Dent, W. R. F., Matthews, H. E., & Walther, D. M. 1995, *MNRAS*, 277, 193
 Diolaiti, E., Bendinelli, O., Bonaccini, D., et al. 2000, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 4007, *Adaptive Optical Systems Technology*, ed. P. L. Wizinowich, 879–888
 Frimann, S., Jørgensen, J. K., Dunham, M. M., et al. 2017, *A&A*, 602, A120
 Furuya, R. S., Kitamura, Y., Wootten, A., Claussen, M. J., & Kawabe, R. 2003, *ApJS*, 144, 71
 Gutermuth, R. A., Megeath, S. T., Myers, P. C., et al. 2009, *ApJS*, 184, 18
 Herczeg, G. J., Johnstone, D., Mairs, S., et al. 2017, *ApJ*, 849, 43
 Higdon, S. J. U., Devost, D., Higdon, J. L., et al. 2004, *PASP*, 116, 975
 Hirano, N. & Liu, F.-c. 2014, *ApJ*, 789, 50

- Ho, P. T. P., Moran, J. M., & Lo, K. Y. 2004, ApJ, 616, L1
- Hsieh, T.-H., Murillo, N. M., Belloche, A., et al. 2018, ApJ, 854, 15
- Johnstone, D., Herczeg, G. J., Mairs, S., et al. 2018, ApJ, 854, 31
- Jørgensen, J. K., van Dishoeck, E. F., Visser, R., et al. 2009, A&A, 507, 861
- Jørgensen, J. K., Visser, R., Williams, J. P., & Bergin, E. A. 2015, A&A, 579, A23
- Lebouteiller, V., Bernard-Salas, J., Sloan, G. C., & Barry, D. J. 2010, PASP, 122, 231
- Loinard, L., Torres, R. M., Mioduszewski, A. J., & Rodríguez, L. F. 2008, ApJ, 675, L29
- Looney, L. W., Mundy, L. G., & Welch, W. J. 2000, ApJ, 529, 477
- Murillo, N. M., Bruderer, S., van Dishoeck, E. F., et al. 2015, A&A, 579, A114
- Murillo, N. M. & Lai, S.-P. 2013, ApJ, 764, L15
- Murillo, N. M., Lai, S.-P., Bruderer, S., Harsono, D., & van Dishoeck, E. F. 2013, A&A, 560, A103
- Murillo, N. M., van Dishoeck, E. F., Tobin, J. J., & Fedele, D. 2016, A&A, 592, A56
- Murillo, N. M., van Dishoeck, E. F., van der Wiel, M. H. D., et al. 2018, ArXiv e-prints [arXiv:1805.05205]
- Ortiz-León, G. N., Loinard, L., Kounkel, M. A., et al. 2017, ApJ, 834, 141
- Persson, M. V., Harsono, D., Tobin, J. J., et al. 2016, A&A, 590, A33
- Pezzuto, S., Elia, D., Schisano, E., et al. 2012, A&A, 547, A54
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, A&A, 518, L1
- Santangelo, G., Murillo, N. M., Nisini, B., et al. 2015, A&A, 581, A91
- Visser, R., Bergin, E. A., & Jørgensen, J. K. 2015, A&A, 577, A102
- Ward-Thompson, D., Kirk, J. M., Greaves, J. S., & André, P. 2011, MNRAS, 415, 2812
- Yu, T. & Chernin, L. M. 1997, ApJ, 479, L63