



Universiteit
Leiden
The Netherlands

Role of environment and gas temperature in the formation of multiple protostellar systems: molecular tracers

Murillo Mejias, N.M.; Dishoeck, E.F. van; Tobin, J.J.; Mottram, J.C.; Karska, A.

Citation

Murillo Mejias, N. M., Dishoeck, E. F. van, Tobin, J. J., Mottram, J. C., & Karska, A. (2018). Role of environment and gas temperature in the formation of multiple protostellar systems: molecular tracers. *Astronomy And Astrophysics*, 620, A30.
doi:10.1051/0004-6361/201832954

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/71681>

Note: To cite this publication please use the final published version (if applicable).

The role of environment and gas temperature in the formation of multiple protostellar systems: molecular tracers

N. M. Murillo¹, E. F. van Dishoeck^{1,2}, J. J. Tobin³, J. C. Mottram⁴, and A. Karska⁵

¹ Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA, Leiden, the Netherlands
e-mail: nmurillo@strw.leidenuniv.nl

² Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße 1, 85748, Garching bei München, Germany

³ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 W. Brooks Street, Norman, Oklahoma 73019, USA

⁴ Max Planck Institute for Astronomy, Königstuhl 17, 69117, Heidelberg, Germany

⁵ Centre for Astronomy, Nicolaus Copernicus University, Faculty of Physics, Astronomy and Informatics, Grudziadzka 5, 87100, Torun, Poland

September 14, 2018

ABSTRACT

Context. Simulations suggest that gas heating due to radiative feedback is a key factor in whether or not multiple protostellar systems will form. Chemistry is a good tracer of the physical structure of a protostellar system, since it depends on the temperature structure.

Aims. To study the relationship between envelope gas temperature and protostellar multiplicity.

Methods. Single dish observations of various molecules that trace the cold, warm and UV-irradiated gas are used to probe the temperature structure of multiple and single protostellar systems on 7000 AU scales.

Results. Single, close binary and wide multiples present similar current envelope gas temperatures, as estimated from H₂CO and DCO⁺ line ratios. The temperature of the outflow cavity, traced by c-C₃H₂, on the other hand, shows a relation with bolometric luminosity and an anti-correlation with envelope mass. Although the envelope gas temperatures are similar for all objects surveyed, wide multiples tend to exhibit a more massive reservoir of cold gas compared to close binary and single protostars.

Conclusions. Although the sample of protostellar systems is small, the results suggest that gas temperature may not have a strong impact on fragmentation. We propose that mass, and density, may instead be key factors in fragmentation.

Key words. astrochemistry - stars: formation - stars: low-mass - ISM: molecules - methods: observational

1. Introduction

Multiple protostellar systems are widely thought to be formed through fragmentation of the cloud core and/or disk within which they form. This process is expected to either be induced by turbulence (e.g., Offner et al. 2010) or through instabilities in the disk that can lead to fragmentation of the disk material (e.g., Stamatellos & Whitworth 2009; Kratter et al. 2010). Each mechanism is proposed to produce multiple protostellar systems, but since they operate on different spatial scales (disks: ~100 AU, vs. cloud core: ~1000 AU) they result in different separations between the sources. Turbulent fragmentation predicts the initial formation of wide companions, whereas disk fragmentation can produce close companions, on scales of the disk radius. The time when these processes occur, upon initial collapse or after one source has formed, can also alter the resulting multiple protostellar system and its evolution. Observational studies of massive star formation provide conflicting evidence regarding the role of turbulence in core fragmentation (e.g., Wang et al. 2014; Palau et al. 2015; Beuther et al. 2018). Another potentially relevant factor in regulating fragmentation may be magnetic support. This mechanism suggests that magnetic fields reduce the number of fragments formed in a cloud core (Commerçon et al. 2010; Hennebelle et al. 2011). Observations of fragmentation in massive dense cloud cores (Tan et al. 2013; Fontani et al. 2018), where both high- and low-mass protostars can form, suggest that

magnetic fields can shape the fragment mass and distribution. The role that magnetic support plays in low-mass multiple star formation is not yet entirely understood.

The factors that enhance fragmentation of cloud cores need to be studied in order to understand how multiple protostellar systems form. Radiative feedback and gas heating have been raised as key factors in the fragmentation of protostellar cores (e.g., Krumholz 2006; Bate 2012; Krumholz et al. 2014). Simulations and models show that fragmentation is suppressed by heated gas due to the increase in the thermal Jeans mass needed for collapse. An accreting protostar heats up its surrounding gas, even as early as the first collapse of the core (Boss et al. 2000; Whitehouse & Bate 2006). Thus it is expected that fragmentation can be considerably suppressed even as the protostellar object forms. Numerical simulations show that as stars begin to form they can heat surrounding gas out to a few thousand AU, with the gas being continuously heated out to larger expanses as more objects form (Bate 2012). This is expected to considerably reduce fragmentation, and consequently the formation of multiple protostellar systems on envelope scales (few thousand AU). Models considering the effect of accretion luminosity on the temperature structure of cloud cores suggest that cores can be heated above 100 K out to a few hundred AU, and 30 K out to a few thousand AU (Krumholz et al. 2014). This is thought to significantly hinder cloud core fragmentation and the consequent formation of multiple stellar systems (Krumholz 2006).

Observations of young low-mass embedded protostellar systems, however, seem to show a different picture. Construction of the SEDs of all known embedded protostellar systems in the Perseus star forming region ($d \sim 235$ pc, Hirota et al. 2011) found, for separations larger than $7''$, that higher order multiples have a tendency for one of the sources to be at a different evolutionary stage than the rest, i.e. non-coeval systems (Murillo et al. 2016). For these non-coeval systems to occur, one of the sources was most likely formed after the other protostars were formed. During the star formation process, episodic accretion bursts produce quiescent phases, lasting on the order of 10^3 – 10^4 yr (Scholz et al. 2013; Visser et al. 2015), which allow enough time for the envelope to cool and thus be more conducive to fragmentation and collapse. However, objects undergoing episodic accretion do not always show multiplicity (e.g., very low luminosity objects, VeLLOs; Hsieh et al. 2018), and not all multiples present signatures of accretion bursts (e.g., Frimann et al. 2017). Furthermore, the recently fragmented circumbinary disk of the deeply embedded protostellar system L1448 N (Tobin et al. 2016a) suggests that instabilities could overcome heating-suppressed fragmentation, since this disk is most certainly heated by both the central binary and through accretion.

Observational evidence for gas and dust heating on scales of a few thousand AU by UV radiation escaping through outflow cavities comes from several lines of evidence. Multi-wavelength observations of dust emission around low-mass protostars suggest indeed elevated temperatures out to such distances (e.g., Hatchell et al. 2013; Sicilia-Aguilar et al. 2013). More relevant are measurements of the gas temperature, since at low cloud densities (10^4 cm $^{-3}$) gas and dust temperatures may be decoupled (Evans et al. 2001; Galli et al. 2002), and it is the gas temperature that enters the formulation for suppressing fragmentation (Offner et al. 2010). The best diagnostics are the ^{13}CO mid-J lines, especially ^{13}CO J = 6–5, first demonstrated by Spaans et al. (1995) and modeled in detail by Visser et al. (2012). The use of ^{13}CO 6–5 as a temperature probe on extended scales has been quantified observationally by van Kempen et al. (2009) and Yıldız et al. (2013, 2015), showing temperatures of 30–50 K on scales of a few thousand AU.

Relating the temperature structure and multiplicity of a protostellar system can provide constraints on the temperature-fragmentation relation. Simulations including radiative feedback (e.g., Krumholz 2006; Bate 2012) would lead us to expect that non-coeval multiple protostellar systems may form in much colder cloud cores than single and coeval binary protostellar systems, in order to have further fragmentation. Thus, the temperature structure of protostellar systems needs to be characterized in order to test these models. Because heating is time-dependent and we cannot observe the temporal history of protostellar envelope heating, protostellar objects at different evolutionary stages and having recently undergone processes such as accretion bursts and fragmentation need to be studied and compared.

Molecular excitation and chemistry provide an excellent tool to probe the temperature structure of a protostellar system. Using selected molecules that trace the cold and warm envelope gas (e.g., Murillo et al. 2015, 2018), it can be established how the gas heating is being distributed throughout the cloud core. In addition, we use new ^{13}CO 6–5 data combined with existing ^{13}CO 3–2 spectra (Mottram et al. 2017) and ^{13}CO 10–9 spectra from the WILL survey (Mottram et al. 2017), observed with the James Clerk Maxwell Telescope (JCMT) and *Herschel Space Observatory* (Pilbratt et al. 2010), respectively. Relating this to

the multiplicity and coevality¹ can then provide information on how temperature affects fragmentation.

This work presents single-dish observations of embedded multiple and single protostellar systems aiming to address the relation between temperature and fragmentation at the envelope scale (~ 7000 AU). The sample selection criteria for the protostellar systems studied in this work are described in Section 2. Section 3 describes the Atacama Pathfinder EXperiment (APEX; Güsten et al. 2006) observations. Results and analysis of the data obtained from the observations are given in Sections 4 and 5, respectively. Comparison between single and multiple protostellar systems is made in Section 6, considering evolutionary stage and whether they are located in a crowded or isolated environment. The conclusions of this work are given in Section 7, along with the resulting insight on the temperature-fragmentation relation.

2. System sample

The Perseus molecular cloud ($d = 235$ pc) large-scale structure has been well studied in continuum (e.g. Hatchell et al. 2005; Enoch et al. 2006; Chen et al. 2016; Pokhrel et al. 2018) and molecular line emission (e.g. Arce et al. 2010; Curtis et al. 2010a,b; Curtis & Richer 2011; Walker-Smith et al. 2014; Hacar et al. 2017). The small scale structure of the region has been studied through the characterization of individual systems (e.g. Kwon et al. 2006; Enoch et al. 2009; Mottram et al. 2013; Hirano & Liu 2014; Ching et al. 2016). The evolutionary classification of the protostars in the region has been carefully studied through molecular line and continuum observations (Enoch et al. 2009; Carney et al. 2016; Mottram et al. 2017). Recently, Tobin et al. (2016b) conducted an unbiased 8 mm survey of all protostars in Perseus down to 15 AU separation with the Karl G. Jansky Very Large Array (VLA), thus characterizing the multiplicity of the star forming region.

The evolutionary stages of each source in embedded wide multiple protostellar systems in Perseus have been characterized through the construction of spectral energy distributions (SEDs) and the parameters derived from the SEDs including *Herschel* Space Observatory PACS maps (Murillo et al. 2016). This provides information on the coevality of wide multiple protostellar systems and can help to understand how these systems are formed (Murillo et al. 2016). Further examination of coevality, core structure and protostar distribution was done by Sadavoy & Stahler (2017), studying possible formation mechanisms. Dust emission observed with the VLA toward several disk-candidate embedded protostellar systems was examined by Segura-Cox et al. (2016), and the dust continuum was found to present disk-shaped structures. In addition, envelopes and outflows driven by protostars in Perseus have been studied both at scales larger than 4000 AU (Davis et al. 2008; Curtis et al. 2010a,b; Arce et al. 2010; Mottram et al. 2013; Karska et al. 2014; Yıldız et al. 2015; Mottram et al. 2017) and below 2000 AU (Persson et al. 2012; Plunkett et al. 2013; Maret et al. 2014; Lee et al. 2016). Together, previous work provides an extensive database of information about the protostars in the Perseus molecular cloud.

For this study, a sample of 12 low-mass protostellar systems in Perseus were selected from the work of Tobin et al. (2016b) and Murillo et al. (2016). The sample is listed in Table 1, along with coordinates, source separations, type of region where they are located and the bolometric luminosity L_{bol} calculated from

¹ Coevality is here used as defined in Murillo et al. (2016), which is the relative evolutionary classes of sources in a multiple protostellar system.

Table 1. Sample of protostellar systems

System	Source	RA	Dec	Separation (")	Class	Region ^b	L_{bol} (L_{\odot})
Wide multiples							
L1448N	A	03:25:36.53	+30:45:21.35	...	I	non-clustered	5.88 ± 0.93
	B	03:25:36.34	+30:45:14.94	7.3	0		2.15 ± 0.33
	C	03:25:35.53	+30:45:34.20	16.3	0		1.22 ± 0.19
NGC1333 SVS13	A	03:29:03.75	+31:16:03.76	...	I	clustered	119.28 ± 18.31
	B	03:29:03.07	+31:15:52.02	14.9	0		10.26 ± 1.57
	C	03:29:01.96	+31:15:38.26	34.7	0		2.22 ± 0.34
NGC1333 IRAS5	Per63	03:28:43.28	+31:17:32.9	...	I	clustered	1.38 ± 0.21
	Per52	03:28:39.72	+31:17:31.9	45.7	I		0.12 ± 0.02
NGC1333 IRAS7	Per18	03:29:11.26	+31:18:31.08	...	0	clustered	4.77 ± 0.73
	Per21	03:29:10.67	+31:18:20.18	13.3	0		3.50 ± 0.54
	Per49	03:29:12.96	+31:18:14.31	27.5	I		0.65 ± 0.10
B1-b	S	03:33:21.30	+31:07:27.40	...	0	non-clustered	0.32 ± 0.05
	N	03:33:21.20	+31:07:44.20	17.4	0		0.16 ± 0.05
	W	03:33:20.30	+31:07:21.29	13.9	I		0.10 ± 0.02
IC348 Per8+Per55 ^a	Per8	03:44:43.94	+32:01:36.09	...	0	non-clustered	1.96 ± 0.30
	Per55	03:44:43.33	+32:01:31.41	9.6	I		1.58 ± 0.26
Close binaries							
NGC1333 IRAS1		03:28:37.00	+31:13:27.5	1.908	I	clustered	11.00 ± 1.78
IRAS 03282+3035		03:31:21.00	+30:45:30.0	0.098	0	non-clustered	1.49 ± 0.23
IRAS 03292+3039 ^a		03:32:17.00	+30:49:47.0	0.085	0	non-clustered	0.89 ± 0.14
Single systems							
IRAS 03271+3013		03:30:15.00	+30:23:49.0	...	I	non-clustered	1.62 ± 0.26
L1455-Per25		03:26:37.46	+30:15:28.01	...	0	non-clustered	1.09 ± 0.17
NGC1333 SK1		03:29:00.00	+31:12:00.7	...	0	clustered	0.71 ± 0.11

Notes. ^(a) Only observed in 13 CO with APEX. ^(b) Clustered regions are defined to have 34 YSO pc⁻¹, while non-clustered regions present 6 YSO pc⁻¹, (Plunkett et al. 2013).

the SEDs. Figure 1 shows *Hersechel* PACS thermal continuum mini-maps of the sample obtained from the Gould Belt Survey (André et al. 2010), along with the constructed SEDs from Murillo et al. (2016). The selected systems are young embedded protostars in the Class 0 and I evolutionary stages. Both single and multiple (i.e., binary and higher order multiples) protostellar systems are included, with the multiple systems spanning a range of separations from $\sim 0.1''$ to $46''$ (~ 23.5 AU to 11000 AU). Thus, both close and wide multiple protostellar systems are considered in this study. Finally, the systems are located in both clustered (NGC1333) and non-clustered regions (L1448, L1455 and B1). Selecting systems from both clustered and non-clustered regions (34 and 6 YSO pc⁻², respectively; Plunkett et al. 2013) allows the impact of external heating on the measured gas temperatures to be assessed.

This sample then allows the evolutionary stage, multiplicity, and region to be compared with temperature, both from the UV-heated gas and the envelope gas temperature. Since the timescale for protostellar evolution (of order a few $\times 10^5$ yrs, Evans et al. 2009; Mottram et al. 2011; Heiderman & Evans 2015; Carney et al. 2016) is considerably longer than can be observed in human lifespan, the evolution of the protostellar temperature structure cannot be directly observed. Hence, systems in the Class 0 and I evolutionary stages need to be compared, as well as single and multiple protostellar systems. Evolutionary classes are determined based on the shape of the SED, derived parameters such as bolometric temperature T_{bol} , and the structure of the system (e.g. envelope, outflow opening angle). Thus, the temperature-multiplicity-age relation can be studied, which can in turn provide constraints for hydrodynamical models with radiative feedback.

3. Observations

3.1. Single pointing

APEX observations of 10 out of 12 systems in our sample were carried out with the Swedish Heterodyne Facility Instrument (SHeFI; Nyström et al. 2009) in position switching mode. The APEX-1 band was used for observations on 1 December 2016 with a precipitable water vapor (PWV) ~ 1.6 mm (O-098.F-9320B.2016, NL GTO time) using one spectral setting with the central frequency set to 217.11258 GHz and a bandwidth of 4 GHz. This setting targeted the molecules DCO⁺, DCN, c-C₃H₂ and H₂CO. In addition, transitions of SO and CH₃OH were detected. Typical noise levels for the observations ranged between 15 to 70 mK for a channel width of 0.4 km s⁻¹ and a HPBW of 28.7''. The beam efficiency η_{mb} for observations at 230 GHz is 0.75.

APEX-2 band observations were carried out from 7 to 12 July 2017 with PWV between 0.37 and 1.5 mm (M-099.F-9516C-2017) using two spectral settings with central frequencies of 350.33746 and 361.16978 GHz, and bandwidth of 4 GHz. The molecules targeted were C₂H, DCO⁺, DCN and H₂CO. HNC was also detected. Typical noise levels range from 20 to 100 mK for a channel width of 0.4 km s⁻¹, a HPBW of 18'' and $\eta_{\text{mb}} = 0.73$. For both bands, calibration uncertainties are on the order of 20%. The molecules targeted in these observations probe the cold (DCO⁺, H₂CO 218.222 GHz) and warm (DCN, c-C₃H₂, C₂H, H₂CO) gas of the envelope at scales of 7000 AU and a gas temperature range of 10 to 100 K (Table 2).




Fig. 1. *Herschel* PACS maps of the systems sampled in this work together with their respective SEDs. The 70 and 160 μm emission are shown in color-scale and contours, respectively. Each stamp spans a region of $80'' \times 80''$ and is centered on the position of the OTF maps, except for NGC1333 IRAS5 whose sources have a separation of $45.7''$. Red symbols represent the sources of a system and the positions of the APEX single-pointing observations. Circles denote systems with additional unresolved multiplicity. Diamonds indicate single sources within multiple protostellar systems. Crosses indicate single protostars. The SEDs are overlaid on the average SEDs from Enoch et al. (2009) for reference (dashed lines), with early Class 0 (red), late Class 0, early Class I, late Class I, and Class II (gray).

3.2. OTF maps

On-the-fly (OTF) maps of all 12 systems were obtained with two instruments: CHAMP+ (Kasemann et al. 2006), and SEPIA B9 (Baryshev et al. 2015; Belitsky et al. 2018), in order to observe ^{13}CO 6–5. ^{13}CO 6–5 is a particularly useful tracer of UV heated gas, in contrast to ^{12}CO 6–5 (Yıldız et al. 2012; van Kempen et al. 2010).

CHAMP+ was used to observe three systems: NGC1333 IRAS7, IC348 Per8+Per55 and IRAS 03292+3029, with a spec-

tral set-up targeting ^{13}CO 6–5 (661.06728 GHz) and a beam of $9.4''$ (HPBW). Maps of $45'' \times 45''$ were centered on the target system in the case of IC348 Per8+Per55 and IRAS 03292+3029. In the case of NGC1333 IRAS7, the maps were centered at a position equidistant from all sources. Observations took place in two epochs, from 26 August to 19 September 2014 (M-094.F-0006.2014), and 2 to 13 of August 2015 (M-095.F-0023.2014).

SEPIA B9 maps ($45'' \times 45''$) were made for the remaining 9 systems in our sample, with the spectral set-up targeting ^{13}CO 6–5 (661.06728 GHz) and a beam of $9.4''$ (HPBW).

Table 2. Molecular species in this work

Molecule	Transition	Frequency GHz	E_{up} K	$\log_{10} A_{ij}$
Single pointing				
SO	5 ₅ -4 ₄	215.22065	44.10	-3.92
DCO ⁺	3-2	216.11258	20.74	-2.62
c-C ₃ H ₂	3 _{3,0} -2 _{2,1}	216.27876	19.47	-3.33
DCN	3-2	217.23863	20.85	-4.24
c-C ₃ H ₂ ^a	6-5	217.82215	38.61	-3.23
c-C ₃ H ₂	5 _{1,4} -4 _{2,3}	217.94005	35.42	-3.35
p-H ₂ CO	3 _{0,3} -2 _{0,2}	218.22219	20.96	-3.55
CH ₃ OH	4 _{2,2} -3 _{1,2}	218.44005	45.45	-4.33
p-H ₂ CO	3 _{2,2} -2 _{2,1}	218.47563	68.09	-3.80
p-H ₂ CO	3 _{2,1} -2 _{2,0}	218.76007	68.11	-3.80
C ₂ H	4-3 J=9/2-7/2 F=5-4	349.33771	41.91	-3.88
C ₂ H	4-3 J=9/2-7/2 F=4-3	349.33899	41.91	-3.89
C ₂ H	4-3 J=7/2-5/2 F=4-3	349.39927	41.93	-3.90
C ₂ H	4-3 J=7/2-5/2 F=3-2	349.40067	41.93	-3.92
o-H ₂ CO	5 _{1,5} -4 _{1,4}	351.76864	62.45	-2.92
DCO ⁺	5-4	360.16978	51.86	-2.42
DCN	5-4	362.04575	52.12	-3.13
HNC	4-3	362.63030	43.51	-2.64
p-H ₂ CO	5 _{0,5} -4 _{0,4}	362.73602	52.31	-2.86
APEX OTF maps				
¹³ CO	4-3	440.76517	52.9	-5.27
¹³ CO	6-5	661.06728	111.1	-4.73
JCMT and Herschel HIFI				
¹³ CO	3-2	330.58797	31.7	-5.66
¹³ CO	10-9	1101.34960	290.8	-4.05

Notes. ^(a) Contains both ortho- and para forms.

References. All rest frequencies were taken from the Cologne Database for Molecular Spectroscopy (CDMS; Endres et al. 2016). The SO entry is based on Clark & De Lucia (1976). The DCO⁺ entries are based on Caselli & Dore (2005). The c-C₃H₂ entry was based on Bogey et al. (1987) with transition frequencies important for our survey from Bogey et al. (1986) and from Spezzano et al. (2012). The DCN entries are based on Okabayashi & Tanimoto (1993) and Brünken et al. (2004). The entry for CH₃OH is based on the line list from Xu & Lovas (1997). The H₂CO entries are based on experimental data from Bocquet et al. (1996). The C₂H entry is based on Padovani et al. (2009) with additional important data from Müller et al. (2000) and Sastry et al. (1981), and on Cazzoli et al. (2012), respectively. The entry for HNC is based on Okabayashi & Tanimoto (1993). The ¹³CO entries are based on Klapper et al. (2000).

Two maps were made for each of the systems NGC1333 SVS13 and NGC1333 IRAS5, given the separation between the sources. For NGC1333 SVS13, the maps were centered on the A and C sources. Observations took place from 13 August to 25 November 2016 (O-098.F-9320A.2016), with an additional science verification observation of B1-B on 29 July 2016 (E-097.F-9810A.2016).

The systems IRAS 03282+3035 and IRAS 03292+3029 were further observed using FLASH (Heyminck et al. 2006) with a spectral set-up targeting ¹³CO 4-3 (440.76517 GHz), since observations of ¹³CO 3-2 were not available with JCMT for these two systems. Observations were carried out on 26 August 2014 (M-094.F-0006.2014).

In order to compare the observations from FLASH, CHAMP+ and SEPIA B9 with the Herschel HIFI ¹³CO 10-9 observations (RMS noise: 0.03 K; Mottram et al. 2017), the data were averaged within a box of approximately 19.3'', the HPBW of the HIFI observations, centered on the position of the HIFI observations. The FLASH observations provide RMS noise of 0.44 and 0.75 K for IRAS 03292+3029 and IRAS 03282+3035, respectively, for a channel width of 0.4 km s⁻¹. For the CHAMP+ observations, typical RMS noise is between 0.03 to 0.08 K for a channel width of 0.4 km s⁻¹. For the SEPIA B9 observations the

typical RMS noise level is between 0.1 to 0.2 K for a channel width of 0.4 km s⁻¹. The main beam efficiencies η_{mb} being used are 0.60 for 440 GHz, and 0.56 for 660 GHz. Typical calibration uncertainties are about 10 to 20%.

4. Results

4.1. Cold and warm gas

The observed molecular lines trace the cold and warmer envelope gas of the systems in our sample. For the multiple protostellar systems, single pointing observations for each source were taken. However, the beams of the observations (28.7'' and 18'') are comparable to the source separations, and the spectra of the individual sources in a wide multiple system are similar. As an example, the spectra of the individual sources for NGC1333 IRAS7 are shown in Fig. A.1. Thus the spectra of all three sources in the wide multiple systems are averaged together, except for NGC1333 IRAS5 which has a separation of 45.7'' and is treated in this work as two separate single protostars Per63 and Per52. The results of the averaged spectra are discussed and analyzed in this work. Figures 2 and 3 show the resulting spectra of the observed sample. Observed line peak temperatures, RMS




Fig. 2. APEX single pointing observations of DCO^+ , DCN , SO , CH_3OH and HNC . Except for DCO^+ 3–2 and HNC , the spectra are multiplied by a factor of 3 to make the features more prominent. The spectra for the wide multiple systems are averaged for all the sources, except for NGC1333 IRAS5. The vertical blue line marks the systemic velocity.

noise, line widths and integrated fluxes are listed in Tables B.1 to B.6.

HNC was detected toward all the systems in the sample, however, the other observed molecules were not detected in the envelope of every system. Three out of the five wide multiple protostellar systems show detections with signal-to-noise (S/N) above 3 in 0.4 km s^{-1} channels for all targeted molecular lines. B1-b, also a wide multiple system, shows above 3σ detections in all lines except $\text{H}_2\text{CO } 3_{2,2}-2_{1,2}$, $\text{DCN } 5-4$ and $\text{C}_2\text{H } 4-3 \text{ J}=7/2-5/2$. SO and methanol (CH_3OH) were detected toward four wide multiple systems. IRAS5 Per63 has detections above 3σ only in DCO^+ 3–2, $\text{H}_2\text{CO } 3_{0,3}-2_{0,2}$ and $\text{H}_2\text{CO } 5_{0,5}-4_{0,4}$. In contrast, IRAS5 Per52 presents detections in DCO^+ 3–2, $c\text{-C}_3\text{H}_2$ 3–2 and 6–5, and $\text{H}_2\text{CO } 3_{0,3}-2_{0,2}$.

The close binary system NGC1333 IRAS 1 presents emission with $S/N > 3$ in SO , DCO^+ 3–2, all transitions of the warm molecules $c\text{-C}_3\text{H}_2$ and C_2H , as well as $\text{H}_2\text{CO } 3_{0,3}-2_{0,2}$. In con-

trast, IRAS 03282+3035 presents emission only in both transitions of DCO^+ , $\text{DCN } 3-2$ and $\text{H}_2\text{CO } 3_{0,3}-2_{0,2}$.

The single protostellar systems show little to no emission. NGC1333 SK1 presents emission above 3σ in DCO^+ 3–2, $\text{H}_2\text{CO } 3_{0,3}-2_{0,2}$ and $\text{H}_2\text{CO } 5_{0,5}-4_{0,4}$; while L1455-Per25 shows only DCO^+ 3–2 and HNC emission. The other single protostar, IRAS 03271+3013, presented no detection beyond HNC . This might be due to the protostar being a Class I protostar, however, NGC1333 IRAS 1 and both sources of NGC1333 IRAS5 are also Class I objects but show more line detections than IRAS 03271+3013. In summary, wide multiple systems generally present more line emission than single protostars, most likely related to the higher envelope mass, and hence larger molecular column density.

DCO^+ presents strong emission towards most systems. Since DCO^+ is formed from the reaction of $\text{H}_2\text{D}^+ + \text{CO}$, with H_2D^+ greatly enhanced at low temperatures due to the freeze-out of CO ($< 30 \text{ K}$), DCO^+ has been found to be a good tracer of cold




Fig. 3. Same as Fig. 2 but for $c\text{-C}_3\text{H}_2$, C_2H and H_2CO .

gas (Jørgensen et al. 2005; Mathews et al. 2013; Murillo et al. 2015). It should be noted, however, that CO freezes out onto the dust grains at densities above $10^4 \sim 10^5 \text{ cm}^{-3}$ (e.g., Caselli et al. 1999; Bergin et al. 2002; Jørgensen et al. 2005), thus at these densities DCO^+ may be dependent on density as well as temperature. The low-lying transition of H_2CO $3_{0,3}-2_{0,2}$ is the strongest among the five transitions, with peaks a factor of ~ 10 higher than the higher-lying H_2CO transitions. The peaks of the H_2CO 5–4 transitions appear stronger than the higher-lying 3–2 transitions, this is due to the smaller HPBW of the APEX-2 observations ($18''$). DCN is a warm gas tracer, which can be formed and fractionated through a higher temperature route starting with CH_2D^+ (e.g., Favre et al. 2015). Similarly to DCN, the higher-lying transitions of H_2CO also trace warm gas. The weak emission of DCN and the higher-lying transitions of H_2CO would suggest that the envelope gas is relatively cold and currently not being strongly heated. The molecules $c\text{-C}_3\text{H}_2$ and C_2H trace the warm UV-irradiated gas (e.g., Nagy et al. 2015; Guzmán et al. 2015), most likely located along the outflow cavity (e.g., Fontani et al. 2012; Jørgensen et al. 2013; Murillo et al. 2018). The peak intensities of $c\text{-C}_3\text{H}_2$ vary by less than a factor of 3 among all three observed transitions. C_2H is detected in both spin doubling transitions with each transition showing a characteristic double hyperfine structure pattern. Methanol (CH_3OH) and SO are mainly formed on grain surfaces, and either are sputtered off the grains by shocks (e.g., Buckle & Fuller 2002; Burkhardt et al. 2016) or sublimated into the gas phase in hot, dense regions of the outflow (e.g., van der Tak et al. 2000; Palau et al. 2017).

The different systemic velocities of each region within Perseus are reflected in the observed spectra (Fig. 2 and 3). For the systems located in NGC1333, there is also a slight difference in the systemic velocity between the systems located closer to the cluster center (NGC1333 SVS13, NGC1333 IRAS7 and NGC1333 IRAS5; $v_{\text{LSR}} = 8.0\text{--}8.5 \text{ km s}^{-1}$) and those located in the outer part of the cluster (NGC1333 IRAS1 and NGC1333 SK1; $v_{\text{LSR}} \sim 7.3 \text{ km s}^{-1}$).

4.2. ^{13}CO maps

The results from the ^{13}CO maps are described in this section. The spectra extracted from the maps are shown in Fig. B.1, along with the spectra from JCMT and *Herschel* HIFI observations, for comparison. Spectra from the JCMT and APEX observations are smoothed to the beam of the HIFI observations. Observed line peak temperatures, RMS noise, line widths and integrated fluxes are listed in Table B.7.

^{13}CO 4–3 was observed toward the close binary systems IRAS 03282+3035 and IRAS 03292+3039, with emission being detected only toward IRAS 03292+3039 ($S/N \sim 7\sigma$). IRAS 03282+3035 does not show emission at the natural resolution of the FLASH observations ($\sigma \sim 0.5 \text{ K}$), nor smoothed to the *Herschel* HIFI observations HPBW = $19.25''$ ($\sigma \sim 1 \text{ K}$).

^{13}CO 6–5 was mainly detected toward the wide multiple systems and one close binary system (Fig. B.1). NGC1333 IRAS7 and IC348 Per8+Per55, observed with CHAMP+, present strong emission. For NGC1333 IRAS7, the bulk of the emission

(S/N = 24σ) is located between Per18 and Per21, with little emission toward Per49, while IC348 Per8+Per55 shows centrally concentrated emission with S/N = 42σ . IRAS 02393+3039 did not show any significant line emission in the CHAMP+ observations with a noise of 0.12 K.

Observations with SEPIA B9 detected ^{13}CO 6–5 toward L1448N, NGC1333 SVS13 and NGC1333 IRAS1. From these systems, NGC1333 SVS13 presents the strongest emission, peaking at ~ 4.7 K (average of the central position of the maps centered on sources A and C). L1448N presents weak emission, despite presenting relatively strong line detections in all the other molecules observed with APEX-1 and APEX-2 (Fig. 2 and 3). However, the weak ^{13}CO 6–5 emission is consistent with the observations of ^{13}CO using the JCMT and *Herschel* HIFI (Table B.7 and Fig. B.1).

The SEPIA B9 observations show considerably more noise by a factor of 3 higher than those of the CHAMP+ observations. Considering the signal-to-noise ratio of the detected emission, however, the non-detections are not due to the higher noise level of the SEPIA B9 observations, but most likely from the compact ^{13}CO 6–5 emission toward these systems, which is diluted in the larger beam.

5. Analysis

5.1. Line emission and system parameters

The observed molecular line emission is compared to system luminosity and envelope mass in this section. Bolometric luminosity L_{bol} is obtained from the SEDs of the observed systems, with L_{bol} for the wide multiple systems derived from the combined SEDs (Murillo et al. 2016). Envelope mass M_{env} , listed in Table 3, is calculated from the $850\ \mu\text{m}$ peak intensity $S_{850\mu\text{m}}$, L_{bol} and distance d using the formula from Jørgensen et al. (2009) expressed as:

$$M_{\text{env}} = 0.44M_{\odot} \left(\frac{L_{\text{bol}}}{1L_{\odot}} \right)^{-0.36} \left(\frac{S_{850\mu\text{m}}}{1\text{ Jy beam}^{-1}} \right)^{1.2} \left(\frac{d}{125\text{ pc}} \right)^{1.2} \quad (1)$$

which takes into account that more luminous systems have somewhat higher dust temperatures. The relation assumes optically thin emission and typical dust-to-gas ratio of 1:100, and is derived from the power-law relations that arise between envelope mass obtained from radiative transfer models, and the observed peak flux, and luminosity. The power-law relations are $M_{\text{env}} \propto S_{850}^{1.2}$ and $M_{\text{env}}/S_{850} \propto L_{\text{bol}}^{0.36}$.

The $850\ \mu\text{m}$ peak intensity used in this work is measured from the COMPLETE survey map of Perseus taken with SCUBA on the JCMT (Kirk et al. 2006), which has a beam of $15''$. The peak intensity was measured in a circular region of $28.7''$ (HPBW of the APEX-1 observations) centered on each system. For the wide multiple systems, the total flux of all sources is used. In addition, the observed line emission is compared to the ratio of envelope mass to bolometric luminosity, ($M_{\text{env}}/L_{\text{bol}}$), listed in Table 3. This ratio gives insight into the amount of mass heated by the luminosity of the protostellar sources within each system. Because younger systems are expected to have more mass in their envelopes, higher ratios are expected to correspond to younger sources (Bontemps et al. 1996). This holds true for the close binary and single systems in our sample. But it is more difficult to disentangle for the wide multiple systems, since the individual sources present different evolutionary stages (Fig. 1 and Table 1).

System type (wide multiple, close binary and single protostar) is indicated in the plots with different symbols in order

to determine if there is any relation with respect to line emission or system parameters. For systems with molecular line non-detections, the upper limits in the plots are placed at 3σ . A linear fit to the data is used to identify trends between the observed line emission peaks and system parameters in cases where there is a significant correlation. Further statistical analysis is treated in Sec. 5.3.

Peak antenna temperatures are compared with the envelope mass in Fig. 4. The peak intensities of H_2CO $3_{0,3}-2_{0,2}$, and DCO^+ 3–2 increase with envelope mass. Wide multiple systems have larger envelope masses than the close binaries and single protostars, with the exception of NGC1333 IRAS5, where Per63 and Per52 have envelope masses comparable to single protostars. This can be interpreted as wide multiple systems having more massive reservoirs of cold gas compared to close binaries and single protostellar systems. On the other hand, the warm gas being traced by DCN and the two higher-lying transitions of H_2CO 3–2 do not show a dependency on the envelope mass, degree of multiplicity or region type. Instead the line peaks are practically constant with envelope mass. Methanol, SO and the three transitions of $\text{c-C}_3\text{H}_2$ do not show particular dependency on envelope mass either. For ^{13}CO 3–2, there appears to be a slight correlation between envelope mass and peak antenna temperature, but there are not enough data points to be certain. The 6–5 and 10–9 transitions also do not show a correlation to envelope mass.

Figure 5 shows the observed line peak antenna temperatures compared with L_{bol} . The warm molecule C_2H appears to be associated with L_{bol} , as well as the $\text{c-C}_3\text{H}_2$ 6–5 and 5–4 transitions. Since these molecules are generally formed in irradiated regions (Fontani et al. 2012; Jørgensen et al. 2013; Nagy et al. 2015; Guzmán et al. 2015), the correlation between C_2H and $\text{c-C}_3\text{H}_2$, and L_{bol} can be explained by the outflow cavity being irradiated by the central protostar, as was also found for O[*I*] and H_2O (e.g. Mottram et al. 2017). Thus, the more luminous the central protostar, the deeper the outflow cavity is irradiated and more $\text{c-C}_3\text{H}_2$ and C_2H is produced. The two transitions of H_2CO 5–4 show a correlation to bolometric luminosity, whereas the higher lying transitions of H_2CO 3–2 do not, despite the similar upper energy levels ($E_{\text{up}} = 52\text{--}68$ K). The reason for this discrepancy could be the difference in beam size from the observations. The H_2CO 3–2 transition was observed with a beam of $28.7''$, compared to $18''$ for the 5–4 transition. This suggests that the H_2CO 5–4 transition is picking up emission from material at smaller scales, closer to the protostellar source(s), and thus related to the luminosity of the protostellar source(s). HNC, and all three transitions of ^{13}CO also show relation to L_{bol} , as previously found for a larger sample by e.g., San José-García et al. (2013) and Yıldız et al. (2013). The other molecules do not present any correlation to bolometric luminosity, not even SO and CH_3OH which are expected to trace shocks, or the higher transitions of H_2CO 3–2 which trace warmer gas. The lack of correlation could be due to the low number of detections, and are further explored through statistical analysis in Section 5.3.

The peak antenna temperatures compared with $M_{\text{env}}/L_{\text{bol}}$ are presented in Fig. B.2. There appears to be no relation between any of the observed molecular line peaks and $M_{\text{env}}/L_{\text{bol}}$. This indicates that when the amount of mass being illuminated, and thus heated, by the central protostellar system is taken into account, the observed protostellar systems present similar peak antenna temperatures.

These results can be summarized as follows. The bulk of the cold envelope gas is traced by DCO^+ and H_2CO $3_{0,3}-2_{0,2}$ while the warm gas is traced by DCN and the two higher-lying transi-

Table 3. Source parameters

System	850 μ m peak intensity Jy beam ⁻¹	M_{env} M_{\odot}	L_{bol} L_{\odot}	$M_{\text{env}} / L_{\text{bol}}$ M_{\odot} / L_{\odot}
Wide multiples				
L1448N	5.69	2.920	13.08	0.22
SVS13	4.63	1.023	121.07	0.01
Per63	0.33	0.215	1.38	0.16
Per52	0.04	0.041	0.12	0.34
IRAS7	1.76	0.820	8.92	0.09
B1-b	2.33	3.052	0.59	5.17
Per8+Per55	0.88	0.506	3.38	0.15
Close binaries				
IRAS1	0.86	0.322	11	0.03
IRAS 03282	1.17	0.957	1.49	0.64
IRAS 03292	2.43	2.768	0.89	3.11
Singles				
IRAS03271	0.24	0.139	1.62	0.09
Per25	0.35	0.252	1.09	0.23
SK1	0.33	0.274	0.71	0.39

tions of H₂CO. c-C₃H₂ and C₂H trace the warm irradiated outflow cavity, rather than envelope material.

5.2. Line ratios and implied physical conditions

The gas temperature structure can be probed with the several transitions of c-C₃H₂, H₂CO and DCO⁺ that were observed. In addition, the ratio of DCN/DCO⁺ can be used to compare the warm (DCN) and cold gas (DCO⁺) from the envelope. Since the two transitions of DCO⁺ were observed with a different beam (3–2: 28.7''; 5–4: 18''), a beam dilution correction factor of 0.39 is applied to the 5–4 transition. Not considering the different beam sizes of the observations, would result in the DCO⁺ ratio being overestimated by a factor of 2 to 3. The line ratios of c-C₃H₂, H₂CO, DCO⁺ and DCN/DCO⁺ are listed in Table 4. While two transitions of DCN were observed, ratios can only be obtained for half the systems due to low detection rates, of which three are upper limits. These ratios are thus also listed for each system where both lines were detected in Table 4 but not discussed further.

High- J CO lines ($J_u > 3$) can be used as diagnostics of temperature and density, as well as UV photon-heated gas (e.g., Yıldız et al. 2012, 2015). Hence the ¹³CO 4–3 and 6–5 observations presented in this work are compared with peak antenna temperatures from the JCMT and *Herschel* HIFI observations. For this purpose, the spectra from the JCMT and APEX data were smoothed to the beam of the *Herschel* HIFI observations (19.25''). Only part of our sample has data from the JCMT and/or *Herschel* HIFI (Table B.7). Thus, only those systems are considered for ¹³CO ratios (Table 5).

Using RADEX (van der Tak et al. 2007), non-LTE excitation and radiative transfer calculations were performed to study the variation of the c-C₃H₂, H₂CO, DCO⁺ and ¹³CO ratios with H₂ density and temperature (Fig. 6). The molecular data files for the RADEX calculations were obtained from the Leiden Atomic and Molecular Database (LAMDA; Schöier et al. 2005). The collisional rate coefficients for DCO⁺ are based on the results of Botschwina et al. (1993) and Flower (1999). For c-C₃H₂, H₂CO and ¹³CO, the collisional rate coefficients are based on Chandra & Kegel (2000), Wiesenfeld & Faure (2013) and Yang et al. (2010), respectively.

Ratios are not calculated for the systems with non-detections in both transitions used in the ratio (Tables B.1, B.3, B.5 and B.7) and are thus not considered in the following analysis (Table 4 and 5). Upper limits are given when one of the transitions is a non-detection (Table 4), and are not considered for the linear fits but are shown in the figures for reference.

Kinetic gas temperatures are derived from the c-C₃H₂, H₂CO, DCO⁺ and ¹³CO ratios by averaging over a range of H₂ densities, and are listed in Table 6. An H₂ density n_{H_2} range between 10⁵ to 10⁶ cm⁻³, typical in the envelopes of embedded protostellar objects on the scales of the beam, is assumed for the c-C₃H₂, H₂CO, and DCO⁺ calculations. A column density of 10¹² cm⁻² is adopted for the DCO⁺, c-C₃H₂ and H₂CO line ratio calculations, ensuring that the lines are optically thin. For ¹³CO, a column density of 10¹⁶ cm⁻² is adopted for a line width of 2 km s⁻¹, which produces optically thin emission. Yıldız et al. (2012) found a column density of 10¹⁷ cm⁻² for a line width of 10 km s⁻¹ toward NGC1333 IRAS4A and IRAS4B. Thus our adopted ¹³CO column density is reasonable. Adopting a column density of 10¹⁷ cm⁻² for a line width of 2 km s⁻¹ would provide optically thick emission.

Figure 7 compares the calculated ratios with envelope mass M_{env} , bolometric luminosity L_{bol} , and $M_{\text{env}}/L_{\text{bol}}$. Since molecular line ratios are related to gas temperature (Fig. 6), the comparison serves to determine relations between temperature and system parameters.

With respect to L_{bol} , both ratios of c-C₃H₂ show good correlation with luminosity. On the other hand, c-C₃H₂ shows an anti-correlation with $M_{\text{env}}/L_{\text{bol}}$, to be expected based on the correlation with L_{bol} . Both c-C₃H₂ ratios point to gas temperatures between 10 and 60 K, with the ratios of NGC1333 IRAS7 and NGC1333 IRAS1 pointing to higher temperatures at n_{H_2} of a few 10⁵ cm⁻³. These results suggest that c-C₃H₂ is dependent on the protostellar system luminosity. This makes sense if it is considered that c-C₃H₂ traces the outflow cavity, which is irradiated by the central protostar, and thus the L_{bol} . A higher luminosity will lead to higher temperatures traced by c-C₃H₂. If the envelope mass is large, however, then more material needs to be heated by the protostar and the gas temperature traced by c-C₃H₂ will be lower. Since the density in the outflow cavity is expected to decrease and the outflow to become hotter as protostars evolve




Fig. 4. Peak intensities of the observed molecular lines compared to the envelope mass (M_{\odot}) of each systems. The dashed lines are linear fits to the data for the cases where a correlation is found. Circles, diamonds and stars show single, close binary and wide multiple protostellar systems, respectively. Note that the more massive envelopes show an increase in the peak intensities of DCO^+ and the low-lying transition of H_2CO , which trace cold gas in the envelope. Molecules tracing warm gas have similar peak intensities regardless of envelope mass.

from Class 0 to I (e.g., Nisini et al. 2015; Mottram et al. 2017), the results of $c\text{-C}_3\text{H}_2$ toward our sample may indicate an evolutionary effect. The separation of the sources in multiple systems or their covality does not show any effect on the temperature traced by the $c\text{-C}_3\text{H}_2$ ratio (Fig. 7). While there is no difference in temperature between close binaries and wide multiples, the effect of multiplicity, on the other hand, cannot be fully determined, since none of the single protostars in our sample present $c\text{-C}_3\text{H}_2$ detections.

The ratios of H_2CO , DCO^+ and ^{13}CO are quite constant in relation to all system parameters, and suggest overall cooler temperatures. System type, region and evolutionary stage do not present any correlation either. The ratios of H_2CO and DCO^+ indicate temperature between 10 and 60 K. Considering higher n_{H_2} , alters the temperature by only a few degrees, otherwise the temperatures stay mainly constant. Thus, all embedded protostars appear to have envelope gas with similar, and relatively cold, temperatures regardless of their multiplicity (Fig. 7).

The ratios obtained from ^{13}CO vary little among the systems, regardless of system parameters, multiplicity and evolutionary stage. The low ratios suggest temperatures typically below 60 K, with only L1448 N showing temperatures closer to 100 K at n_{H_2} of a few 10^5 cm^{-3} . This is consistent with the work of Yıldız et al. (2015), which found typical gas temperatures of 30 to 50 K

toward protostellar systems. Given that high- J ^{13}CO lines trace UV photon-heated gas, the low ratios and derived temperatures would indicate that the envelope is not being heated out to large radii. Furthermore, the lack of line detection toward some of the systems would suggest that ^{13}CO emission is concentrated closer to the source of heating, and thus is being diluted in the APEX beam, and even more so when smoothed to the *Herschel* HIFI beam.

In order to determine whether our sample of protostellar systems is particularly cold, we construct a histogram of ^{13}CO ratios from Appendix C of Yıldız et al. (2013) and those derived from the observations in this work (Fig. 8). In total the sample includes 33 protostellar systems, 26 from the work of Yıldız et al. (2013), and 7 from this work. The sample from Yıldız et al. (2013) includes protostellar systems from different star forming regions, but does not overlap with the sample in this work. The histogram is then fit with a Gaussian distribution to find the mean ratio and standard deviation. For the ^{13}CO 6–5/3–2 ratio, the mean ratio is 0.74 with a standard deviation of 0.46, and the ratios indicating a range of temperatures. For 10–9/3–2 and 10–9/6–5, the ratios are below 0.3 and indicate cool temperatures in general. For the ^{13}CO 10–9/3–2 ratio, the mean ratio is 0.05 with a standard deviation of 0.04, whereas for 10–9/6–5, the mean ratio and standard deviation are 0.08 and 0.07, respectively. The




Fig. 5. Peak intensities of the observed molecular lines compared to the bolometric luminosity L_{bol} (L_{\odot}) of each system. The dashed lines are linear fits to the data for the cases where a correlation is found. Circles, diamonds and stars show single, close binary and wide multiple protostellar systems, respectively. C_2H and $\text{c-C}_3\text{H}_2$ show somewhat higher peak intensities in systems with relatively higher luminosities.

upper limits would decrease the mean ratio for the ^{13}CO 10–9/3–2 and 10–9/6–5 ratios, but not so much for the 6–5/3–2 ratio.

In summary, the cool envelope temperatures found in this work are not from particularly cold protostellar systems, nor is the Perseus molecular cloud producing uncommon protostars. It would seem, instead, that the central protostar does not extensively heat the envelope to high temperatures, and that single protostars do not heat the envelope differently than multiple protostellar systems.

5.3. Statistical analysis

In order to determine quantitatively if there is a relation between the observed line peaks and derived quantities, and the system parameters, the Generalized Kendall’s rank correlation is used (Isobe et al. 1986). This method measures the degree of association between two quantities which contain upper limits (censored data), with the null hypothesis being that the values are uncorrelated. Thus, if the significance level $p > 0.05$, the probability of the values being correlated is less than 3σ , while $p < 0.05$ indicates a correlation at better than 3σ significance. The standard normal score z and the significance level p of the Generalized Kendall’s rank correlation are listed in Table C.1, with values indicating a correlation highlighted in bold text. The significance level is calculated from the standard normal score z by

the relation

$$p = 1 - 0.5 * (1 + \text{erf}\left(\frac{|z|}{\sqrt{2}}\right)), \quad (2)$$

where $\text{erf}(x)$ is the error function.

The results confirm the correlations seen by eye (Sec. 5.1). The peak antenna temperatures of the molecules tracing cold gas, namely DCO^+ and H_2CO are associated with system envelope mass, but not with luminosity. On the other hand, the peak antenna temperatures of $\text{c-C}_3\text{H}_2$ and C_2H , both tracers of warm gas, are correlated with luminosity but not system envelope mass. The three transitions of ^{13}CO present correlation with luminosity, while the 6–5 transition also presents a correlation with the mass to luminosity ratio. The 6–5 / $3_{3,0}-2_{2,1}$ ratio of $\text{c-C}_3\text{H}_2$ shows correlations with luminosity and the mass to luminosity ratio, but not envelope mass. The DCN/DCO^+ and ^{13}CO 10–9/3–2 ratios present correlation with envelope mass, but not the other system parameters.

6. Discussion

6.1. Observed line detections

The observations appear to show a tendency for wide multiple protostellar systems (separations $> 7''$) to have more molecular




Fig. 6. Calculated line ratios for $c\text{-C}_3\text{H}_2$, H_2CO , DCO^+ and ^{13}CO . The black lines show the modelled ratios assuming a column density of 10^{12} cm^{-2} for $c\text{-C}_3\text{H}_2$, H_2CO and DCO^+ , and 10^{16} cm^{-2} for ^{13}CO . For the ^{13}CO 6-5/3-2 and 6-5/4-3 panels, the two lower lines show the 0.02 and 0.04 ratios. The shaded areas show the results for the gas temperature calculations and adopted H_2 density range for individual systems: L1448N (blue), NGC1333 IRAS7 (light blue), NGC1333 IRAS1 or IRAS 03292+3039 (gray) and NGC1333 IRAS5 Per52 or NGC1333 SK1 (red).

Table 4. Peak main beam temperature line ratios

System	$c\text{-C}_3\text{H}_2$		H_2CO	DCO^+	DCN	DCN/DCO^+
	6-5 / 3 _{3,0} -2 _{2,1}	5 _{1,4} -4 _{2,3} / 3 _{3,0} -2 _{2,1}	3 _{2,2} -2 _{2,1} / 3 _{0,3} -2 _{0,2}	5-4/3-2	5-4/3-2	3-2
Wide multiples						
L1448N	0.77 ± 0.05	0.54 ± 0.04	0.09 ± 0.02	0.1 ± 0.01	0.26 ± 0.09	0.08 ± 0.01
SVS13	1.15 ± 0.07	0.67 ± 0.06	0.1 ± 0.01	0.15 ± 0.03	0.35 ± 0.10	0.24 ± 0.02
Per63	<0.47	<0.19	...	<0.36
Per52	0.38 ± 0.12	<0.33	<0.23	<0.11	...	<0.21
IRAS7	0.8 ± 0.12	0.75 ± 0.1	0.06 ± 0.01	0.08 ± 0.01	<0.41	0.12 ± 0.01
B1-b	0.44 ± 0.06	0.3 ± 0.06	<0.07	0.07 ± 0.01	<0.35	0.1 ± 0.01
Close binaries						
IRAS1	0.92 ± 0.09	0.74 ± 0.07	<0.24	0.12 ± 0.1	...	<0.15
IRAS 03282	<0.69	<0.12	<0.76	0.1 ± 0.03
Singles						
IRAS 03271
Per25	<0.13	...	<0.52
SK1	<0.12	<0.05	...	<0.1




Fig. 7. Calculated molecular line ratios of $c\text{-C}_3\text{H}_2$ (top row), H_2CO (second row), DCO^+ (third row) and ^{13}CO (bottom row) compared to the system parameters bolometric luminosity L_{bol} (left column), envelope mass M_{env} (middle column) and the mass to luminosity ratio $M_{\text{env}}/L_{\text{bol}}$ (right column). The black and red dashed lines in the top row are linear fits to the data for the $6\text{-}5/3_{3,0}\text{-}2_{2,1}$ and $5_{1,4}\text{-}4_{2,3}/3_{3,0}\text{-}2_{2,1}$ ratios, respectively. Circles, diamonds and stars show single, close binary and wide multiple protostellar systems, respectively. The $c\text{-C}_3\text{H}_2$ ratios appear to be somewhat related to luminosity, whereas the ratios from H_2CO and DCO^+ are constant regardless of system parameter.

line detections with strong peak intensities than close binaries (Fig. 9). In contrast, single protostars present very weak molecular line emission. There seems to be a relation, however, between the envelope mass and the number of molecular line detections. As noted in Sect. 5.1, the ratio $M_{\text{env}}/L_{\text{bol}}$ correlates with evolutionary stage for the close binary and single systems. However, the ratio, and thus the evolutionary stage, does not seem to have relation to the number of line detections (Fig. 9d). These relations appear to not be affected by clustering either. However, the envelope mass is larger for wide multiple protostellar systems in contrast to that of close binary and single protostars. Core mass would then be more related to formation of non-coeval wide multiple protostellar systems. There is no apparent rela-

tion between the bolometric luminosity L_{bol} and the number of line detections (Fig. 9c) nor their strength (Fig. 5). The systems L1448 N, NGC1333 IRAS7 and NGC1333 SVS13, which have combined bolometric luminosities above $8 L_{\odot}$, show strong detections of all the molecular species. Clustering does not seem to particularly enhance the line strength or number of line detections in the envelope (Fig. 9). L1448 N and B1-b are both located in non-clustered environments and present the same chemical richness as the multiple protostars located in NGC1333, which is a clustered region. Both Class 0 and I systems are present in wide multiples, close binaries and single protostellar systems, but no effect is seen on the line detections. It must be highlighted, however, that the sample size presented in this work is small and




Fig. 8. ^{13}CO ratio stacked histogram including JCMT and *Herschel* data from Yıldız et al. (2013) (top row) and this work (bottom row). Upper limits are shown in gray. Ratios from Yıldız et al. (2013) are for the systems in the Water In Star-forming regions with *Herschel* (WISH) key project (van Dishoeck et al. 2011), which do not overlap with the sample studied in this work. The black dashed line shows a Gaussian distribution fit to the total sample of ^{13}CO ratios.

Table 5. ^{13}CO peak main beam temperature line ratios

System	6-5/3-2	6-5/4-3	10-9/3-2	10-9/4-3	10-9/6-5
Wide multiples					
L1448N	0.26 ± 0.07		0.04 ± 0.004		0.15 ± 0.04
IRAS7 ^a	0.48 ± 0.02		0.02 ± 0.004		0.03 ± 0.01
PER8+PER55	0.31 ± 0.01		0.02 ± 0.003		0.07 ± 0.01
Close binaries					
IRAS1	0.29 ± 0.06		0.02 ± 0.01		0.07 ± 0.02
IRAS 03282	
IRAS 03292		<0.11		<0.03	...
Singles					
IRAS 03271	...		<0.02		...
Per25					...
SK1	<0.19		<0.02		...

Notes. ^(a) Includes only Per18 and Per21.

more protostellar systems would be needed to determine if the trends found in this work are real or product of the small sample size.

6.2. Envelope gas temperature and multiplicity

The H_2CO and DCO^+ ratios point to cool envelopes ($T_{\text{gas}} < 60$ K at scales of 7000 AU) for all the systems in our sample, regardless of multiplicity, clustering and evolutionary stage. Only $\text{c-C}_3\text{H}_2$ presents a correlation with bolometric luminosity and an anti-correlation with $M_{\text{env}}/L_{\text{bol}}$. Since a formation path of $\text{c-C}_3\text{H}_2$ is through the breakdown of large hydrocarbons by UV photons, these correlations could be related to the amount of material that is irradiated by the central protostar, with more luminous protostars irradiating more material. Alternatively, it could be an evolutionary effect where the jet becomes hotter while the envelope and outflow cavity mass decreases, resulting in more material being irradiated by UV photons (e.g., Nisini et al. 2015; Mottram et al. 2017). The low ^{13}CO ratios, and derived temperature ($T_{\text{gas}} < 100$ K at scales of ~ 4500 AU), suggest that the

envelope and outflow cavity are not being UV photon-heated out to large extents.

The results of Maret et al. (2004) and Koumpia et al. (2016) support the possibility that the envelope gas is not being heated out to large extents. Maret et al. (2004) observed L1448 N, NGC1333 IRAS4A and IRAS4B using the single-dish IRAM 30-meter telescope and the JCMT, tracing scales of 2500 to 4000 AU (HPBW = 11'' to 17''), while Koumpia et al. (2016) observed NGC1333 IRAS4 with the JCMT (HPBW = 15'', ~ 3500 AU). Using LVG modelling under non-LTE conditions, Maret et al. (2004) derive temperatures of 50 K for NGC1333 IRAS4A, 80 K for NGC1333 IRAS4B and 90 K for L1448 N. The results presented here for L1448 N find gas temperature of 30 K at 7000 AU scales and 65 K at 4500 AU scales (Table 6). The peak gas temperatures found by Koumpia et al. (2016) for NGC1333 IRAS4 are in agreement with the results of Maret et al. (2004). However the gas temperature distribution from Koumpia et al. (2016) shows that the gas temperature drops to 40 K at ~ 2000 AU away from NGC1333 IRAS4B, and that the 60 K gas is located along the outflow cavity of NGC1333 IRAS4A, consistent

Table 6. Derived T_{kin} from line ratios averaged over H_2 density range

System	$\text{c-C}_3\text{H}_2^a$	$\text{c-C}_3\text{H}_2^a$	H_2CO^a	DCO^{+a}	$^{13}\text{CO}^a$	$^{13}\text{CO}^a$	$^{13}\text{CO}^a$
	6-5 / 3 _{3,0} -2 _{2,1}	5 _{1,4} -4 _{2,3} / 3 _{3,0} -2 _{2,1}	3 _{2,2} -2 _{2,1} / 3 _{0,3} -2 _{0,2}	5-4 / 3-2	6-5/3-2	10-9/3-2	10-9/6-5
	T_{kin} (K)	T_{kin} (K)	T_{kin} (K)	T_{kin} (K)	T_{kin} (K)	T_{kin} (K)	T_{kin} (K)
Wide multiples							
L1448N	18 ± 2	20 ± 3	30 ± 2	19 ± 3	30 ± 3	58 ± 7	86 ± 16
SVS13	30 ± 6	22 ± 4	31 ± 1	26 ± 8
Per63	<180	<29
Per52	11 ± 1	<10	<61	<21
IRAS7	18 ± 2	54 ± 30	24 ± 1	17 ± 3	42 ± 3	51 ± 6	51 ± 7
B1-b	11 ± 1	10 ± 7	<25	21 ± 5
Per8+Per55	33 ± 2	50 ± 6	66 ± 10
Close binaries							
IRAS1	22 ± 3	81 ± 49	<64	25 ± 8	33 ± 3	50 ± 6	65 ± 10
IRAS 03282	<180	<26
IRAS 03292	<20 ^b	<57 ^b	...
Singles							
IRAS 03271	<51	...
Per25	<23
SK1	<35	<13	<27	<51	...

Notes. ^(a) Assuming a H_2 density range of 10^5 to 10^6 cm^{-3} ^(b) Kinetic temperatures are for the 6-5/4-3 and 10-9/4-3 ratios.

with the findings of Yıldız et al. (2012), with the gas temperature dropping to 40 K at a distance of ~ 5000 AU. These results thus indicate that while the embedded protostar can heat up the surrounding gas to temperatures of 50 to 100 K, it can only do so out to about 1000 AU or along low density structures such as the outflow cavity. In contrast wide multiple protostellar systems are observed to have much larger separations (Table 1), with sources not likely to be located within the outflow cavity.

Non-detection within the APEX beam of molecules such as $\text{c-C}_3\text{H}_2$ and H_2CO , which trace the outflow cavity and envelope gas, respectively, would point to cooler envelopes and outflow cavities at scales of 7000 AU. In addition, DCO^+ , a cold gas tracer, is detected toward all but one system (IRAS 03271+3013, single). Non-detections of warm gas tracers in addition to the presence of DCO^+ toward single and close binary protostellar systems could be explained by these systems being somewhat cooler than wide multiple protostellar systems. Another possibility could be the evolutionary dispersal of envelope material which would cause the warm gas to be located closer to the protostellar source(s) and thus diluted in the observing beam. Interestingly enough, H_2CO 3_{0,3}-2_{0,2} and DCO^+ appear to increase with an increase in envelope mass. In other words, H_2CO 3_{0,3}-2_{0,2} and DCO^+ present stronger emission in wide multiple protostars. This could indicate the presence of massive reservoirs of cold gas in these systems.

External heating does not seem to play a significant role, as evidenced by a lack of difference in all aspects between systems in clustered and non-clustered regions. However it is not clear why close binaries present less molecular line emission than wide multiple protostars, given that some sources of the latter are often close binaries themselves (Fig. 1). A possible explanation would be the sensitivity and beam filling factor of the observations. However, this would not explain why NGC1333 IRAS1 has more line detections than NGC1333 IRAS5 Per63 when both have sensitivities of ~ 30 mK. The results presented in this work indicate that there is no significant difference in envelope temperature between single, close binary and wide multiple protostellar systems. The only difference temperature-wise

arises from $\text{c-C}_3\text{H}_2$, which traces the outflow cavity. This is consistent with previous studies that find the heating by protostellar source(s) is mainly channeled through the outflow cavity (Yıldız et al. 2012, 2015).

The envelope gas temperatures measured in our sample coincide well with the temperature expectations from Krumholz (2006), 30 K at a few thousand AU, while the results from Koumpia et al. (2016) and Maret et al. (2004) are consistent with the inner regions closer to the source being heated to 100 K. However, no extensive heating of the envelope out to thousand AU scales (Bate 2012) is found in our sample, with gas temperatures above 30 K occurring only in the outflow cavities. Based on the work presented here, the prediction that heating of the envelope should hinder fragmentation is not consistent with the results and analysis of the sample. Wide multiple systems tend to be non-coeval and are thus produced by continued fragmentation of the cloud core (Murillo et al. 2016; Sadavoy & Stahler 2017), but they do not show colder envelope gas temperatures with respect to coeval binary and single protostellar systems. Thus, if the envelope gas temperature is similar for all surveyed systems, then it is not clear why only some cores have continued fragmentation. Cloud core mass could provide a possible explanation, since wide multiple systems present higher envelope masses than coeval binary and single protostellar systems, however this would require further studies.

An interesting case is that of L1448 N, which has been found to have undergone recent fragmentation in the disk of one source (Tobin et al. 2016a), that is fragmentation at 10 to 100 AU scales. Although the recent fragmentation could have heated the envelope of this system as suggested by simulations (Whitehouse & Bate 2006; Boss et al. 2000), increasing the envelope temperature, it does not seem to have generated a considerable effect on thousand AU scales. Furthermore, the disk is certainly heated by the existing protostars but fragmentation of the disk was not hindered.




Fig. 9. Number of line detections (a), envelope mass M_{env} (b), bolometric luminosity L_{bol} (c), and envelope mass to luminosity ratio $M_{\text{env}} / L_{\text{bol}}$ (d) of each system in our sample compared to the region, clustered (circles) or non-clustered (diamonds), and system type. NGC1333 IRAS5 is considered here as two single protostellar systems, Per63 and Per52.

6.3. Accretion bursts

Accretion of material onto the protostar is not a continuous process, but has been found from observations to be episodic (e.g., Visser et al. 2015; Safron et al. 2015; Frimann et al. 2017; Hsieh et al. 2018). An accretion burst causes an increase in luminosity, and consequently heating of the envelope gas. The chemical composition of the envelope gas is altered, since cold chemistry molecules (e.g. DCO^+) move further out and molecules such as CO are evaporated off the grains. While the luminosity of the central protostar will decrease after the accretion burst has passed, it takes about 10^4 years for the gas to refreeze onto the dust grains (Johnstone et al. 2013; Jørgensen et al. 2015).

Frimann et al. (2017) studied the presence of accretion bursts toward embedded protostars in Perseus. Their sample includes four of the systems studied in this work (L1448 N, NGC1333 SVS13, NGC1333 IRAS1 and IRAS 03282), as well as two sources of NGC1333 IRAS7, Per18 and Per21. Half of these systems (Per21, IRAS1 and IRAS 03282) present indications of past accretion burst activity, while the others (L1448 N, SVS13 and Per18) do not. The results presented here do not show any difference among the systems that have evidence of having undergone an accretion burst, and those that have not. NGC1333 IRAS1 and

IRAS 03282, both close binary protostars, are suggested to have undergone an accretion burst. The difference found in this work is mainly that IRAS1 presents emission from warm molecules, while IRAS 03282 only shows emission in cold molecules. For L1448N and SVS13, both wide multiple protostars, there is also no significant difference, since both systems present the same number of line detections, with similar peak intensities except for DCO^+ which is stronger in L1448N. Analyzing NGC1333 IRAS7 Per18 and Per21 (Appendix A) separately also points to a lack of significant difference among both sources. However, it must be noted that given their separation of $13''$, the spectra of each source is contaminated by emission from the other.

7. Conclusions

Single-dish observations of a sample of 12 embedded protostellar systems in Perseus are presented here. The observations targeted molecular line emission that trace the cold, warm and UV-heated gas of the observed protostellar systems. The sample included wide multiple protostellar systems (separation $\geq 7''$, or ≥ 1600 AU), close binary protostars (separation $< 2''$, or < 470 AU) and single protostars, located in clustered and non-clustered environments, spanning Class 0 and I objects, and containing coeval and non-coeval systems. The results presented in this work examine the relationship between fragmentation and temperature, since heated gas is expected to suppress fragmentation based on simulations including radiative feedback.

Although the sample presented here is small and there are several upper limits, the envelope gas temperature is found to be similar among multiple and single protostars, regardless of evolutionary stage, coevality or clustering. These results suggest that gas temperature may not have as strong a role in suppressing fragmentation as expected from models, a result that was also stated in Offner et al. (2010), who found that temperature does not suppress turbulent fragmentation. Instead, wide multiple protostellar systems present larger envelope masses and massive cold gas reservoirs in comparison to close binary and single protostars. It seems, then, that mass, along with other factors such as turbulence, density profile and magnetic fields, rather than envelope gas temperature plays a fundamental role in fragmentation. Larger, more massive cores could then lead to further fragmentation that forms non-coeval wide multiple systems.

Further interferometric observations of the sample used in this work using the same molecular lines treated here would lend insight into the spatial distribution of the cold, warm and UV heated gas. The gas density and temperature distribution can then be compared to the multiplicity of the protostellar system. An additional topic of further research would then be to determine what causes some cores to become more massive than others.

Acknowledgements. Astrochemistry in Leiden is supported by the European Union A-ERC grant 291141 CHEMPLAN, by the Netherlands Research School for Astronomy (NOVA), by a Royal Netherlands Academy of Arts and Sciences (KNAW) professor prize. JCM acknowledges support from the European Research Council under the European Community's Horizon 2020 framework program (2014-2020) via the ERC Consolidator grant 'From Cloud to Star Formation (CSF)' (project number 648505). AK acknowledges support from the Polish National Science Center grant nr 2016/21/D/ST9/01098. We are grateful to the APEX staff for support with these observations. Observing time for the APEX data was obtained via Max Planck Institute for Radio Astronomy, Onsala Space Observatory and European Southern Observatory.

References

André, P., Men'shchikov, A., Bontemps, S., et al. 2010, A&A, 518, L102

- Arce, H. G., Borkin, M. A., Goodman, A. A., Pineda, J. E., & Halle, M. W. 2010, *ApJ*, 715, 1170
- Baryshev, A. M., Hesper, R., Mena, F. P., et al. 2015, *A&A*, 577, A129
- Bate, M. R. 2012, *MNRAS*, 419, 3115
- Belitsky, V., Lapkin, I., Fredrixon, M., et al. 2018, *A&A*, 612, A23
- Bergin, E. A., Alves, J., Huard, T., & Lada, C. J. 2002, *ApJ*, 570, L101
- Beuther, H., Mottram, J. C., Ahmadi, A., et al. 2018, *ArXiv e-prints* [arXiv:1805.01191]
- Bocquet, R., Demaison, J., Poteau, L., et al. 1996, *Journal of Molecular Spectroscopy*, 177, 154
- Bogey, M., Demuyneck, C., & Destombes, J. L. 1986, *Chem. Phys. Lett.*, 125, 383
- Bogey, M., Demuyneck, C., Destombes, J. L., & Dubus, H. 1987, *J. Mol. Spectrosc.*, 122, 313
- Bontemps, S., Andre, P., Terebey, S., & Cabrit, S. 1996, *A&A*, 311, 858
- Boss, A. P., Fisher, R. T., Klein, R. I., & McKee, C. F. 2000, *ApJ*, 528, 325
- Botschwina, P., Horn, M., Flugge, J., & Seeger, S. 1993, *J. Chem. Soc., Faraday Trans.*, 89, 2219
- Brünken, S., Fuchs, U., Lewen, F., et al. 2004, *Journal of Molecular Spectroscopy*, 225, 152
- Buckle, J. V. & Fuller, G. A. 2002, *A&A*, 381, 77
- Burkhardt, A. M., Dollhopf, N. M., Corby, J. F., et al. 2016, *ApJ*, 827, 21
- Carney, M. T., Yıldız, U. A., Mottram, J. C., et al. 2016, *A&A*, 586, A44
- Caselli, P. & Dore, L. 2005, *A&A*, 433, 1145
- Caselli, P., Walmsley, C. M., Tafalla, M., Dore, L., & Myers, P. C. 1999, *ApJ*, 523, L165
- Cazzoli, G., Cludi, L., Buffa, G., & Pazzarini, C. 2012, *ApJS*, 203, 11
- Chandra, S. & Kegel, W. H. 2000, *A&AS*, 142, 113
- Chen, M. C.-Y., Di Francesco, J., Johnstone, D., et al. 2016, *ApJ*, 826, 95
- Ching, T.-C., Lai, S.-P., Zhang, Q., et al. 2016, *ApJ*, 819, 159
- Clark, W. W. & De Lucia, F. C. 1976, *Journal of Molecular Spectroscopy*, 60, 332
- Commerçon, B., Hennebelle, P., Audit, E., Chabrier, G., & Teyssier, R. 2010, *A&A*, 510, L3
- Curtis, E. I. & Richer, J. S. 2011, *MNRAS*, 410, 75
- Curtis, E. I., Richer, J. S., & Buckle, J. V. 2010a, *MNRAS*, 401, 455
- Curtis, E. I., Richer, J. S., Swift, J. J., & Williams, J. P. 2010b, *MNRAS*, 408, 1516
- Davis, C. J., Scholz, P., Lucas, P., Smith, M. D., & Adamson, A. 2008, *MNRAS*, 387, 954
- Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., & Müller, H. S. P. 2016, *J. Mol. Spectrosc.*, 327, 95
- Enoch, M. L., Evans, II, N. J., Sargent, A. I., & Glenn, J. 2009, *ApJ*, 692, 973
- Enoch, M. L., Young, K. E., Glenn, J., et al. 2006, *ApJ*, 638, 293
- Evans, II, N. J., Dunham, M. M., Jørgensen, J. K., et al. 2009, *ApJS*, 181, 321
- Evans, II, N. J., Rawlings, J. M. C., Shirley, Y. L., & Mundy, L. G. 2001, *ApJ*, 557, 193
- Favre, C., Bergin, E. A., Cleaves, L. I., et al. 2015, *ApJ*, 802, L23
- Flower, D. R. 1999, *MNRAS*, 305, 651
- Fontani, F., Commerçon, B., Giannetti, A., et al. 2018, *A&A*, 615, A94
- Fontani, F., Palau, A., Busquet, G., et al. 2012, *MNRAS*, 423, 1691
- Frimann, S., Jørgensen, J. K., Dunham, M. M., et al. 2017, *A&A*, 602, A120
- Galli, D., Walmsley, M., & Gonçalves, J. 2002, *A&A*, 394, 275
- Güsten, R., Nyman, L. Å., Schilke, P., et al. 2006, *A&A*, 454, L13
- Guzmán, V. V., Pety, J., Goicoechea, J. R., et al. 2015, *ApJ*, 800, L33
- Hacar, A., Tafalla, M., & Alves, J. 2017, *A&A*, 606, A123
- Hatchell, J., Richer, J. S., Fuller, G. A., et al. 2005, *A&A*, 440, 151
- Hatchell, J., Wilson, T., Drabek, E., et al. 2013, *MNRAS*, 429, L10
- Heiderman, A. & Evans, II, N. J. 2015, *ApJ*, 806, 231
- Hennebelle, P., Commerçon, B., Joos, M., et al. 2011, *A&A*, 528, A72
- Heyminck, S., Kasemann, C., Güsten, R., de Lange, G., & Graf, U. U. 2006, *A&A*, 454, L21
- Hirano, N. & Liu, F.-c. 2014, *ApJ*, 789, 50
- Hirota, T., Honma, M., Imai, H., et al. 2011, *PASJ*, 63, 1
- Hsieh, T.-H., Murillo, N. M., Belloche, A., et al. 2018, *ApJ*, 854, 15
- Isobe, T., Feigelson, E. D., & Nelson, P. I. 1986, *ApJ*, 306, 490
- Johnstone, D., Hendricks, B., Herczeg, G. J., & Bruderer, S. 2013, *ApJ*, 765, 133
- Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2005, *A&A*, 435, 177
- Jørgensen, J. K., van Dishoeck, E. F., Visser, R., et al. 2009, *A&A*, 507, 861
- Jørgensen, J. K., Visser, R., Sakai, N., et al. 2013, *ApJ*, 779, L22
- Jørgensen, J. K., Visser, R., Williams, J. P., & Bergin, E. A. 2015, *A&A*, 579, A23
- Karska, A., Kristensen, L. E., van Dishoeck, E. F., et al. 2014, *A&A*, 572, A9
- Kasemann, C., Güsten, R., Heyminck, S., et al. 2006, in *Proc. SPIE, Vol. 6275, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 62750N
- Kirk, H., Johnstone, D., & Di Francesco, J. 2006, *ApJ*, 646, 1009
- Klapper, G., Lewen, F., Gendriesch, R., Belov, S. P., & Winnewisser, G. 2000, *Journal of Molecular Spectroscopy*, 201, 124
- Koumpia, E., van der Tak, F. F. S., Kwon, W., et al. 2016, *A&A*, 595, A51
- Kratter, K. M., Matzner, C. D., Krumholz, M. R., & Klein, R. I. 2010, *ApJ*, 708, 1585
- Krumholz, M. R. 2006, *ApJ*, 641, L45
- Krumholz, M. R., Bate, M. R., Arce, H. G., et al. 2014, *Protostars and Planets VI*, 243
- Kwon, W., Looney, L. W., Crutcher, R. M., & Kirk, J. M. 2006, *ApJ*, 653, 1358
- Lee, K. I., Dunham, M. M., Myers, P. C., et al. 2016, *ApJ*, 820, L2
- Maret, S., Belloche, A., Maury, A. J., et al. 2014, *A&A*, 563, L1
- Maret, S., Ceccarelli, C., Caux, E., et al. 2004, *A&A*, 416, 577
- Mathews, G. S., Klaassen, P. D., Juhász, A., et al. 2013, *A&A*, 557, A132
- Mottram, J. C., Hoare, M. G., Davies, B., et al. 2011, *ApJ*, 730, L33
- Mottram, J. C., van Dishoeck, E. F., Kristensen, L. E., et al. 2017, *A&A*, 600, A99
- Mottram, J. C., van Dishoeck, E. F., Schmalzl, M., et al. 2013, *A&A*, 558, A126
- Müller, H. S. P., Klaus, T., & Winnewisser, G. 2000, *A&A*, 357, L65
- Murillo, N. M., Bruderer, S., van Dishoeck, E. F., et al. 2015, *A&A*, 579, A114
- Murillo, N. M., van Dishoeck, E. F., Tobin, J. J., & Fedele, D. 2016, *A&A*, 592, A56
- Murillo, N. M., van Dishoeck, E. F., van der Wiel, M. H. D., et al. 2018, *ArXiv e-prints* [arXiv:1805.05205]
- Nagy, Z., Ossenkopf, V., Van der Tak, F. F. S., et al. 2015, *A&A*, 578, A124
- Nisini, B., Santangelo, G., Giannini, T., et al. 2015, *ApJ*, 801, 121
- Nyström, O., Lapkin, I., Desmaris, V., et al. 2009, *Journal of Infrared, Millimeter, and Terahertz Waves*, Volume 30, Issue 7, pp.746-761, 30, 746
- Offner, S. S. R., Kratter, K. M., Matzner, C. D., Krumholz, M. R., & Klein, R. I. 2010, *ApJ*, 725, 1485
- Okabayashi, T. & Tanimoto, M. 1993, *J. Chem. Phys.*, 99, 3268
- Padovani, M., Walmsley, C. M., Tafalla, M., Galli, D., & Müller, H. S. P. 2009, *A&A*, 505, 1199
- Palau, A., Ballesteros-Paredes, J., Vázquez-Semadeni, E., et al. 2015, *MNRAS*, 453, 3785
- Palau, A., Walsh, C., Sánchez-Monge, Á., et al. 2017, *MNRAS*, 467, 2723
- Persson, M. V., Jørgensen, J. K., & van Dishoeck, E. F. 2012, *A&A*, 541, A39
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, *A&A*, 518, L1
- Plunkett, A. L., Arce, H. G., Corder, S. A., et al. 2013, *ApJ*, 774, 22
- Pokhrel, R., Myers, P. C., Dunham, M. M., et al. 2018, *ApJ*, 853, 5
- Sadavoy, S. I. & Stahler, S. W. 2017, *MNRAS*, 469, 3881
- Safron, E. J., Fischer, W. J., Megeath, S. T., et al. 2015, *ApJ*, 800, L5
- San José-García, I., Mottram, J. C., Kristensen, L. E., et al. 2013, *A&A*, 553, A125
- Sastry, K. V. L. N., Helminger, P., Charo, A., Herbst, E., & De Lucia, F. C. 1981, *ApJ*, 251, L119
- Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, *A&A*, 432, 369
- Scholz, A., Froebrich, D., & Wood, K. 2013, *MNRAS*, 430, 2910
- Segura-Cox, D. M., Harris, R. J., Tobin, J. J., et al. 2016, *ApJ*, 817, L14
- Sicilia-Aguilar, A., Henning, T., Linz, H., et al. 2013, *A&A*, 551, A34
- Spaans, M., Hogerheijde, M. R., Mundy, L. G., & van Dishoeck, E. F. 1995, *ApJ*, 455, L167
- Spezzano, S., Tamassia, F., Thorwirth, S., et al. 2012, *ApJS*, 200, 1
- Stamatellos, D. & Whitworth, A. P. 2009, *MNRAS*, 392, 413
- Tan, J. C., Kong, S., Butler, M. J., Caselli, P., & Fontani, F. 2013, *ApJ*, 779, 96
- Tobin, J. J., Kratter, K. M., Persson, M. V., et al. 2016a, *Nature*, 538, 483
- Tobin, J. J., Looney, L. W., Li, Z.-Y., et al. 2016b, *ApJ*, 818, 73
- van der Tak, F. F. S., Black, J. H., Schöier, F. L., Jansen, D. J., & van Dishoeck, E. F. 2007, *A&A*, 468, 627
- van der Tak, F. F. S., van Dishoeck, E. F., Evans, II, N. J., & Blake, G. A. 2000, *ApJ*, 537, 283
- van Dishoeck, E. F., Kristensen, L. E., Benz, A. O., et al. 2011, *PASP*, 123, 138
- van Kempen, T. A., Kristensen, L. E., Herczeg, G. J., et al. 2010, *A&A*, 518, L121
- van Kempen, T. A., van Dishoeck, E. F., Güsten, R., et al. 2009, *A&A*, 501, 633
- Visser, R., Bergin, E. A., & Jørgensen, J. K. 2015, *A&A*, 577, A102
- Visser, R., Kristensen, L. E., Bruderer, S., et al. 2012, *A&A*, 537, A55
- Walker-Smith, S. L., Richer, J. S., Buckle, J. V., Hatchell, J., & Drabek-Mauder, E. 2014, *MNRAS*, 440, 3568
- Wang, K., Zhang, Q., Testi, L., et al. 2014, *MNRAS*, 439, 3275
- Whitehouse, S. C. & Bate, M. R. 2006, *MNRAS*, 367, 32
- Wiesenfeld, L. & Faure, A. 2013, *MNRAS*, 432, 2573
- Xu, L.-H. & Lovas, F. J. 1997, *Journal of Physical and Chemical Reference Data*, 26, 17
- Yang, B., Stancil, P. C., Balakrishnan, N., & Forrey, R. C. 2010, *ApJ*, 718, 1062
- Yıldız, U. A., Kristensen, L. E., van Dishoeck, E. F., et al. 2012, *A&A*, 542, A86
- Yıldız, U. A., Kristensen, L. E., van Dishoeck, E. F., et al. 2015, *A&A*, 576, A109
- Yıldız, U. A., Kristensen, L. E., van Dishoeck, E. F., et al. 2013, *A&A*, 556, A89

Appendix A: Single-pointing observations: NGC1333 IRAS7

APEX single pointing observations with the heterodyne instruments APEX-1 and APEX-2 were made toward 5 wide (separation $>7''$) multiple protostellar systems in Perseus. These systems are referred to as wide multiple protostars since the sources span separations greater than $7''$ (which can be resolved with *Herschel Space Observatory* PACS photometric maps, Murillo et al. 2016). One pointing per source in a wide multiple system was observed. In the main text are the parameters of the lines obtained by averaging the spectra of the individual sources for the corresponding system. This is because in all cases except NGC1333 IRAS5, the beam of APEX observations partially overlaps another source in the system.

Here the spectra for the individual sources of the system NGC1333 IRAS7 are presented (Fig. A.1). This provides an example of the similarity of the spectra of the individual sources. The spectra of Per21, which is expected to have undergone a recent accretion burst, and Per18, which has no evidence of episodic accretion, can also be compared. It is also interesting to note that Per49 presents much weaker emission than Per18 and Per21. The evolutionary stage, multiplicity and bolometric luminosity for each source in the system are listed in Table A.1 for reference.

The peak brightness temperature ratios for $c\text{-C}_3\text{H}_2$, H_2CO , DCO^+ and DCN are calculated for each source (Table A.2). A beam dilution factor of 0.39 is applied to the 5–4 transitions of DCO^+ and DCN , since the beam of the observations of the 5–4 transition is smaller than the beam for the 3–2 transition (see Sect. 3). Since the beams of the APEX-1 observations overlap for Per18 and Per21, we note that the ratios for these two sources may be contaminated with emission from each other. Table A.2 also lists the line ratios for the whole system for the purpose of comparison.

The line ratios for each source in NGC1333 IRAS7 are similar to those derived for the whole system (Table A.2), and are compared with the ratio models shown in Fig. 6. The DCN ratios are upper limits due to the non-detection of the 5–4 transition, and instead three times the noise level is used for the ratio. The DCN to DCO^+ 3–2 ratio is used to look at the ratio of warm to cold gas in the envelope, with all the ratios well below unity, indicating a larger amount of cold gas.

Gas temperatures were derived from the line ratios assuming an H_2 density range of 10^5 to 10^6 cm^{-3} for $c\text{-C}_3\text{H}_2$, H_2CO and DCO^+ . The average derived gas temperatures are listed in Table A.3. For the envelope gas, H_2CO indicates temperatures of around 30 K, while DCO^+ ratios point to gas temperatures of about 21 K. For $c\text{-C}_3\text{H}_2$, the line ratios indicate higher temperatures, between 20 and 50 K, which is to be expected for material along the outflow cavity. The gas temperatures for each source in the NGC1333 IRAS7 system are similar to the system as a whole (Table A.3).

The derived temperatures do not show relation with bolometric luminosity or envelope mass. The overall trend is consistent with that of the other systems studied, namely that the envelope gas temperature is similar for all systems, regardless of multiplicity. The source Per21 is thought to have undergone a recent accretion burst (Frimann et al. 2017), however no significant difference in the derived ratios and gas temperatures is found. It must be stressed, however, that the values toward Per21 are very likely contaminated by Per18, and viceversa, due to the beam of the APEX-1 observations.

Appendix B: Observed molecular line emission

In this appendix the peak antenna temperatures, RMS noise, measured line widths and integrated fluxes are listed for the detected molecular lines from the APEX observations. These values were obtained with a simple Gaussian fit of the detected emission line, with no fixed parameters, and binning 4 channels to obtain a velocity resolution of 0.4 km s^{-1} .

Figure B.1 shows the spectra of ^{13}CO in different transitions. The 3–2 transition was observed with the JCMT. Both the 4–3 and 6–5 transitions were observed with APEX using FLASH, CHAMP+ and SEPIA B9. The 10–9 transition was observed with *Herschel* HIFI in the WISH program. All spectra were smoothed to an angular resolution of $19.25''$, and the spectra of JCMT and APEX observations were taken with a $19.25''$ box.

Figure B.2 plots the peak antenna temperature from the APEX observations versus the mass to bolometric luminosity ratio $M_{\text{env}}/L_{\text{bol}}$. No correlation is found between the peak antenna temperatures and $M_{\text{env}}/L_{\text{bol}}$. This lack of correlation is further confirmed by the Generalized Kendall's rank correlation (See Sec. 5.3 and Appendix C), which only finds correlations between Methanol (CH_3OH) and $M_{\text{env}}/L_{\text{bol}}$.

Appendix C: Statistical analysis

The relation between the results of the observations and the system parameters is determined by the Generalized Kendall's rank (Isobe et al. 1986). The table in this appendix lists the significance level p and standard normal score z of the rank correlation test. The numbers in bold indicate real correlations ($p < 0.05$).




Fig. A.1. Spectra for the individual sources of NGC1333 IRAS7. Note that some spectra are multiplied by a factor of 3 in order to enhance the line emission features. The spectra of Per21 and Per49 are offset by 1.3 and 2.6 K from that of Per18 for clarity.

Table A.1. Source parameters for NGC1333 IRAS7

	Class	Multiplicity	Accretion burst?	850 μm peak (Jy beam^{-1})	L_{bol} (L_{\odot})	M_{env} (M_{\odot})	$M_{\odot} / L_{\text{bol}}$ (M_{\odot} / L_{\odot})
Per18	0	Binary	No	1.76	4.8	1.0	0.21
Per21	0	Single	Possible	1.76	3.5	1.1	0.33
Per49	I	Binary		1.75	0.7	2.0	2.50

Table A.2. Peak main beam temperature line ratios for NGC1333 IRAS7

	$\text{c-C}_3\text{H}_2$ 6-5 / 3 _{3,0} -2 _{2,1}	$\text{c-C}_3\text{H}_2$ 5 _{1,4} -4 _{2,3} / 3 _{3,0} -2 _{2,1}	H_2CO 3 _{2,2} -2 _{2,1} / 3 _{0,3} -2 _{0,2}	DCO^+ 5-4/3-2	DCN 5-4/3-2	DCN/DCO^+ 3-2
Per18	1.07 ± 0.15	0.63 ± 0.10	0.08 ± 0.02	0.08 ± 0.01	< 0.24	0.12 ± 0.02
Per21	0.65 ± 0.19	0.90 ± 0.22	0.08 ± 0.03	0.16 ± 0.01	< 0.18	0.17 ± 0.02
Per49	1.10 ± 0.45	< 0.90	0.10 ± 0.03	0.05 ± 0.01	< 0.50	0.08 ± 0.02
IRAS7 ^a	0.8 ± 0.12	0.75 ± 0.1	0.06 ± 0.01	0.08 ± 0.01	< 0.41	0.12 ± 0.01

Notes. ^(a) Values for the spectra of all three sources averaged together.

Table A.3. Derived T_{kin} from line ratios averaged over H_2 for NGC1333 IRAS7

	$\text{c-C}_3\text{H}_2$ 6-5 / $3_{3,0}-2_{2,1}$ T_{kin} (K)	$\text{c-C}_3\text{H}_2$ 5 _{1,4} -4 _{2,3} / $3_{3,0}-2_{2,1}$ T_{kin} (K)	H_2CO 3 _{2,2} -2 _{2,1} / $3_{0,3}-2_{0,2}$ T_{kin} (K)	DCO^+ 5-4/3-2 T_{kin} (K)
Per18	23 ± 3	21 ± 4	28 ± 3	20 ± 5
Per21	14 ± 1	67 ± 41	29 ± 2	26 ± 8
Per49	28 ± 7	< 71	32 ± 3	16 ± 4
IRAS7	18 ± 2	54 ± 30	24 ± 1	17 ± 3




Fig. B.1. ^{13}CO spectra observed with JCMT (3–2), APEX (6–5) and *Herschel* HIFI (10–9). The spectra of JCMT and APEX are smoothed to a resolution of $19.25''$, the HPBW of the HIFI observations. The short blue lines at the top of each plot indicate the systemic velocity of the system. Note that the ^{13}CO 4–3 and 10–9 spectra are multiplied by a factor of 3 and 15, respectively. The 6–5 spectra for NGC1333 IRAS1 is also multiplied by a factor of 3.

Table B.1. Peak main beam temperatures for DCO⁺ and DCN

System	DCO ⁺ 3-2			DCO ⁺ 5-4			DCN 3-2			DCN 5-4						
	T_{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹	T_{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹	T_{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹				
L1448N	3160	31	1.13	3786.7	811	33	1.04	895	267	25	1.63	453.3	69	23	1.43	105
SVS13	1213	23	1	1306.7	461	31	1.03	505	293	25	1.18	373.3	103	28	1.82	199
Per63	333	40	0.64	226.7	162	54	-	-	...	40	36	-	-
Per52	560	37	0.66	386.7	161	54	-	-	...	39	47	-	-
IRAS7	2267	27	0.76	1866.7	468	33	0.56	281	267	24	0.8	266.7	...	37	-	-
B1-b	2627	21	1.32	3680.0	446	30	0.92	438	253	19	1.05	280.0	...	30	-	-
Per8+Per55	-	-	-	-
Wide multiples																
Close binaries																
IRAS1	800	43	0.68	573.3	243	81	-	-	...	40	64	-	-
IRAS 03282	1933	73	0.6	1200.0	567	56	0.57	342	200	67	0.82	173.3	...	50	-	-
IRAS 03292	-	-	-	-
Singles																
IRAS 03271	...	71	43	-	-	...	69	41	-	-
Per25	480	67	0.4	200.0	156	52	-	-	...	83	40	-	-
SK1	1173	37	0.5	666.7	160	53	-	-	...	40	40	-	-

Notes. ^(a) In this and subsequent tables, the noise refers to a 0.4 km s⁻¹ velocity bin.

Table B.2. Peak main beam temperatures for SO, CH₃OH and HNC

System	SO 5 ₅ – 4 ₄			CH ₃ OH 4 _{2,2} –3 _{1,2}			HNC 4-3					
	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹
Wide multiples												
L1448N	280	27	0.99	30.7	187	23	1.02	200.0	1219	24	2.00	2592
SVS13	253	23	1.5	413.3	120	21	1.1	133.3	2070	26	1.32	2898
Per63	...	35	40	256	46	0.78	212
Per52	...	45	39	667	40	0.81	573
IRAS7	600	23	0.74	480.0	187	23	0.82	160.0	1120	30	0.90	1068
B1-b	627	21	0.85	573.3	187	23	0.8	160.0	490	24	1.34	699
Per8+Per55	–	–
Close binaries												
IRAS1	147	39	1.4	266.7	...	39	1284	58	0.89	1212
IRAS 03282	...	67	71	1030	44	0.84	917
IRAS 03292	–	–
Singles												
IRAS 03271	...	77	60	324	47	1.03	357
Per25	...	75	65	370	38	1.07	422
SK1	...	39	40	611	46	0.70	457

Table B.3. Peak main beam temperatures for c-C₃H₂

System	c-C ₃ H ₂ 3 _{3,0} -2 _{2,1}			c-C ₃ H ₂ 6-5			c-C ₃ H ₂ 5 _{1,4} -4 _{2,3}					
	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹
	Wide multiples											
L1448N	693	25	1.56	1146.7	533	25	1.65	946.7	373	23	1.68	666.7
SVS13	440	23	0.99	453.3	507	19	1.1	586.7	293	23	1.3	400.0
IRAS5 Per63	...	47	35	40
IRAS5 Per52	320	40	0.5	173.3	121	36	0.94	121.3	...	35
IRAS7	267	24	0.7	200.0	213	24	0.97	226.7	200	20	0.59	121.3
B1-b	360	23	1.33	506.7	160	19	0.99	173.3	107	19	1.19	13.3
Per8+Per55
	Close binaries											
IRAS1	667	44	0.66	466.7	613	40	0.75	480.0	493	33	0.62	333.3
IRAS 03282	...	68	61	55
IRAS 03292
	Singles											
IRAS 03271	...	97	61	56
Per25	...	87	69	59
SK1	...	35	47	40

Table B.4. Peak main beam temperatures for C₂H

System	C ₂ H 4-3 J=9/2 - 7/2 f=5-4			C ₂ H 4-3 J=9/2 - 7/2 f=4-3			C ₂ H 4-3 J=7/2 - 5/2 f=4-3			C ₂ H 4-3 J=7/2 - 5/2 f=3-2						
	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹	T _{mb} mK	Noise mK	Width km s ⁻¹	Integrated mK km s ⁻¹
Wide multiples																
L1448N	412	53	2.14	937	263	53	2.47	693	254	46	2.16	584	229	46	2.61	637
SVS13	742	35	1.18	928	608	35	1.06	688	594	29	1.11	703	431	29	1.31	602
Per63	0	49	-	-	0	49	-	-	0	50	-	-	0	50	-	-
Per52	0	56	-	-	0	56	-	-	0	53	-	-	0	53	-	-
IRAS7	335	24	0.61	216	276	24	1.21	356	208	29	0.65	143	221	29	0.99	232
B1-b	126	24	0.80	108	84	24	0.04	840	0	25	-	-	0	25	-	-
Per8+Per55	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Close binaries																
IRAS1	337	39	0.86	308	258	39	0.81	222	215	37	0.97	221	138	37	0.92	135
IRAS 03282	0	78	-	-	0	78	-	-	0	81	-	-	0	81	-	-
IRAS 03292	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Singles																
IRAS 03271	0	92	-	-	0	92	-	-	0	89	-	-	0	89	-	-
Per25	0	97	-	-	0	97	-	-	0	88	-	-	0	88	-	-
SK1	0	60	-	-	0	60	-	-	0	63	-	-	0	63	-	-

Table B.5. Peak main beam temperatures for H₂CO 3–2

System	H ₂ CO 3 _{0,3} –2 _{0,2}			H ₂ CO 3 _{2,2} –2 _{2,1}			H ₂ CO 3 _{2,1} –2 _{2,0}					
	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹	Peak mK	Noise mK	Width km s ⁻¹	Integrated K km s ⁻¹
	Wide multiples											
L1448N	1520	23	1.68	666.7	132	24	2.2	306.7	109	23	1.67	200.0
SVS13	1520	27	1.45	2333.3	147	20	1.5	240.0	132	21	2	293.3
IRAS5 Per63	253	37	1.32	360.0	...	40	32
IRAS5 Per52	533	40	0.76	400.0	...	40	37
IRAS7	1587	27	1.16	1946.7	97	21	1.3	133.3	80	23	1.3	116.0
B1-b	1040	19	1.3	1453.3	...	24	84	19	0.83	73.3
Per8+Per55
	Close binaries											
IRAS1	573	36	0.86	520.0	...	45	41
IRAS 03282	320	60	0.97	333.3	...	73	61
IRAS 03292
	Singles											
IRAS 03271	...	69	64	107
Per25	...	73	69	73
SK1	880	43	0.95	89.3	...	35	41

Table B.6. Peak main beam temperatures for H₂CO 5–4

System	T_{mb} mK	Noise mK	$5_{1,5-4_{1,4}}$		T_{mb} mK	Noise mK	$5_{0,5-4_{0,4}}$	
			Width km s ⁻¹	Integrated mK km s ⁻¹			Width km s ⁻¹	Integrated mK km s ⁻¹
Wide multiples								
L1448N	595	21	2.00	1265	758	40	2.03	1639
SVS13	594	23	1.80	1136	1130	27	1.38	1660
Per63	...	55	82	29	3.49	303
Per52	...	40	48
IRAS7	374	33	0.98	388	568	32	1.22	736
B1-b	127	19	1.92	260	252	25	2.52	677
Per8+Per55
Close binaries								
IRAS1	230	57	2.19	535	423	41	1.08	489
IRAS 03282	...	42	0.00	0	...	92
IRAS 03292
Singles								
IRAS 03271	...	36	90
Per25	...	28	69
SK1	161	40	0.92	158	...	62



Fig. B.2. Peak intensities of the observed molecular lines compared to the envelope mass to luminosity ratio of each system. Circles, diamonds and stars show single, close binary and wide multiple protostellar systems, respectively.

Table B.7. Peak main beam temperatures for ^{13}CO

System	3-2 ^a			4-3 ^b			6-5 ^b			10-9 ^c		
	T_{mb} K	Noise K	Width km s ⁻¹	Integrated K km s ⁻¹	T_{mb} K	Noise K	Width km s ⁻¹	Integrated K km s ⁻¹	T_{mb} K	Noise K	Width km s ⁻¹	Integrated K km s ⁻¹
L1448N	7.51	0.12	2.15	17.22	1.95	0.49	2.03	4.22	0.30	0.03		1.0034
SVS13-A					4.83	0.56	1.71	8.78				
SVS13-C					4.66	0.50	1.57	7.78				
IRAS7 ^d	7.20	0.13	2.93	22.46	3.44	0.14	1.41	5.16	0.11	0.03		0.17
B1-B					...	0.19				
PER8+PER55	8.05	0.12	2.48	21.23	2.51	0.06	2.07	5.54	0.18	0.03		0.6882
Wide multiples												
Close binaries												
IRAS1	5.47	0.13	1.40	8.16	1.57	0.33	1.10	1.84	0.11	0.03		0.2244
IRAS 03282					1.99	0.75	0.62	1.32	...	0.40
IRAS 03292					3.26	0.44	0.84	2.91	...	0.12
Singles												
IRAS 03271	3.85	0.13	1.06	4.35	...	0.37	0.09	0.03
Per25					...	0.42	0.03
SK1	4.47	0.13	2.26	10.73	...	0.29	0.03

Notes. Blank spaces indicate that there are no observations for the respective system and transition; three dots (...) indicate non-detection. ^(a) Peak antenna temperatures from JCMT observations, smoothed to a beam of 19.25'' ^(b) Peak antenna temperatures and noise from APEX observations, averaged over a box of 19.25''. ^(c) Peak antenna temperatures from *Herschel* HIFI observations with a beam of 19.25'' (Motttram et al. 2017). ^(d) IRAS 7 here includes only Per18 and Per21

Table C.1. Results of the Generalized Kendall Correlation

Molecule	Transition	Mass		L_{bol}		Mass / L_{bol}	
		z	p	z	p	z	p
Peak antenna temperatures							
SO	5 ₅ -4 ₄	2.34	0.01	0.94	0.17	-0.23	0.41
DCO ⁺	3-2	3.04	0.00	0.54	0.29	1.09	0.14
c-C ₃ H ₂	3 _{3,0} -2 _{2,1}	0.94	0.17	1.56	0.06	-1.02	0.15
DCN	3-2	1.85	0.03	1.85	0.03	-0.81	0.21
c-C ₃ H ₂	6-5	1.49	0.07	2.58	0.00	-1.57	0.06
c-C ₃ H ₂	5 _{1,4} -4 _{2,3}	1.1	0.14	3.3	0.00	-2.44	0.01
H ₂ CO	3 _{0,3} -2 _{0,2}	2.03	0.02	0.94	0.17	-0.23	0.41
CH ₃ OH	4 _{2,2} -3 _{1,2}	0.93	0.18	-0.08	0.47	1.27	0.10
H ₂ CO	3 _{2,2} -2 _{2,1}	-0.55	0.29	1.02	0.15	-0.63	0.26
H ₂ CO	3 _{2,1} -2 _{2,0}	-1.1	0.14	0.63	0.26	-0.39	0.35
C ₂ H	4-3 J=9/2-7/2 F=5-4	1.17	0.12	3.35	0.00	-2.34	0.01
C ₂ H	4-3 J=9/2-7/2 F=4-3	0.23	0.41	2.11	0.02	-1.96	0.02
C ₂ H	4-3 J=7/2-5/2 F=4-3	0.47	0.32	2.34	0.01	-1.8	0.04
C ₂ H	4-3 J=7/2-5/2 F=3-2	-0.08	0.47	1.48	0.07	-0.94	0.17
H ₂ CO	5 _{1,5} -4 _{1,4}	1.74	0.04	2.22	0.01	-1.51	0.07
DCO ⁺	5-4	2.76	0.00	1.34	0.09	0.16	0.44
DCN	5-4	-1.67	0.05	-0.56	0.29	0.16	0.44
HNC	4-3	1.63	0.05	1.95	0.03	-1.25	0.11
H ₂ CO	5 _{0,5} -4 _{0,4}	2.03	0.02	3.12	0.00	-1.49	0.07
¹³ CO	3-2 & 4-3	0.08	0.47	1.76	0.04	-0.64	0.26
¹³ CO	6-5	0.39	0.35	2.81	0.00	-2.5	0.01
¹³ CO	10-9	0.55	0.29	1.92	0.03	-0.64	0.26
Ratios							
DCN/DCO ⁺	3-2	-1.82	0.03	-0.18	0.43	-1.27	0.10
DCO ⁺	5-4/3-2	-0.9	0.18	0.9	0.18	-1.44	0.08
c-C ₃ H ₂	6-5 / 3 _{3,0} -2 _{2,1}	-0.19	0.43	2.07	0.02	-2.44	0.01
c-C ₃ H ₂	5 _{1,4} -4 _{2,3} / 3 _{3,0} -2 _{2,1}	-0.94	0.17	0.56	0.29	-1.69	0.05
H ₂ CO	3 _{2,2} -2 _{2,1} / 3 _{0,3} -2 _{0,2}	-1.46	0.07	-0.21	0.42	0.21	0.42
¹³ CO	6-5/3-2	0.51	0.30	0.94	0.17	-0.94	0.17
¹³ CO	10-9/3-2	2.03	0.02	0.23	0.41	1.6	0.05
¹³ CO	10-9/6-5	0.36	0.36	1.53	0.06	1.08	0.14