

To cite this article:

Sleuwaegen, E., Claes, L., Luyckx, K., Wilderjans, T. F., Berens, A., & Sabbe, B. (in press). Do treatment outcomes differ after 3 months DBT inpatient treatment based on borderline personality disorder subtypes. *Personality and Mental Health*. doi:10.1002/pmh.1430

**Do treatment outcomes differ after 3 months DBT inpatient treatment  
based on borderline personality disorder subtypes?**

Ellen Sleuwaegen<sup>1,2</sup>, Laurence Claes<sup>2,3</sup>, Koen Luyckx<sup>3</sup>, Tom Wilderjans<sup>3,4</sup>, Ann Berens<sup>1</sup>, &  
Bernard Sabbe<sup>1,2</sup>

<sup>1</sup>*University Department of Psychiatry, Campus Psychiatric Hospital Duffel, Duffel, Belgium*

<sup>2</sup>*Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium*

<sup>3</sup>*Faculty of Psychology and Educational Sciences, KU Leuven, Leuven, Belgium*

<sup>4</sup>*Research Group of Methodology and Statistics, Institute of Psychology, Leiden University, Leiden, The Netherlands*

*Address of correspondence*

Ellen Sleuwaegen (dra.)

University Department of Psychiatry, Campus Psychiatric Hospital Duffel, Belgium

[Ellen.sleuwaegen@uantwerpen.be](mailto:Ellen.sleuwaegen@uantwerpen.be)

Acknowledgement

This study was partly supported by the Research Foundation – Flanders (FWO), thanks to a Special PhD fellowship [1901317N].

## Abstract

Heterogeneity in Borderline Personality Disorder (BPD) drives the search for BPD subtypes to optimize the assessment and treatment of these patients. Therefore, the aim of the present study was: (1) to replicate previously identified BPD subtypes based on reactive and regulative temperament; (2) to compare them on symptomatology and coping; and (3) to investigate whether these subtypes show different treatment responses after 3 months of inpatient Dialectical Behavior Therapy (DBT). A total of 145 BPD inpatients were assessed by means of measures of temperament, symptomatology and coping. Through model-based clustering on the Behavioral Inhibition and Behavioral Activation Scales (BISBAS) and Effortful Control Scale (EC), we identified three BPD subtypes: an Emotional/Disinhibited subtype (15%, high BAS and low EC); a Low Anxiety subtype (41%, low BIS); and an Inhibited subtype (44%, low BAS). After 3 months of DBT, 75 patients completed the measures for a second time. Repeated measure ANOVA's demonstrated a general improvement on all symptoms and coping strategies. In addition, the BPD subtypes showed trajectory differences in clinical and borderline specific symptomatology and dissociation. These findings indicate that BPD subtypes based on temperament demonstrate different treatment responses, which can contribute to the search of more BPD subtype tailored treatment interventions.

## Highlights

- Three BPD subtypes were identified based on reactive and regulative temperament: the Emotional/Disinhibited, Inhibited, and Low Anxiety subtype.
- All BPD patients improved in symptomatology and coping after 3 months of DBT.
- The three BPD subtypes differed in symptom trajectory after 3 months of DBT.
- In contrast to the other two subtypes, the Low Anxiety subtype did not show a significant decrease in symptomatology.

**Keywords:** borderline personality disorder (BPD), BPD subtypes, temperament, treatment outcomes, dialectical behaviour therapy (DBT)

## 1 Introduction

Given the heterogeneity of borderline personality disorder patients (BPD), recent studies investigated the existence of personality subtypes in BPD patients (Frias et al., 2017; Sleuwaegen et al., 2017; Smits et al., 2017). A person-centered approach in a theoretical framework can explain different phenotypes of BPD. Frias et al. (2017) identified four subtypes based on personality traits, and Smits et al. (2017) distinguished three subtypes based on dimensional PD features. Since temperament is a promising transdiagnostic process that underlies psychopathology, different temperament dimensions such as reactive and regulative temperament can define subtypes. Previous studies have displayed different subtypes based on temperament dimensions and their differential associations with psychopathology within various clinical samples (e.g. Muller et al., 2014). Subtypes based on temperament dimensions, characterized by differential psychiatric comorbidity, may demonstrate different treatment responses. In a previous study, Sleuwaegen et al. (2017) distinguished four BPD subtypes based on reactive (Behavioural Activation System reactivity; BAS and Behavioural Inhibition System reactivity; BIS) and regulative temperament (effortful control): an Emotional/Disinhibited subtype (low effortful control), characterized by the highest level of symptomatology and comorbid histrionic PD; an Inhibited subtype (low BAS), demonstrating low expression of emotions and comorbid cluster C personality disorder (PD); a Low Anxiety subtype (low BIS) characterised by low anxiety, high expression of emotions and comorbid antisocial PD; and a High Self-control subtype (high effortful control) showing the highest level of functioning.

The identification of BPD subtypes may be important for enhancing the effectiveness of assessment and treatment of BPD patients (Kopala-Sibley et al., 2012), however the clinical utility of using BPD subtypes has not yet been proven. Insight in treatment trajectories of the

different BPD subtypes may support the development of more tailored treatment interventions and may guide treatment choices to optimize treatment outcome.

Several effectivity studies of DBT inpatient treatment showed significant improvement in ratings of non-suicidal self-injury (NSSI), general clinical symptomatology, borderline-related symptoms and dissociation after a 3-month DBT program in BPD samples (Bohus et al., 2004; Kröger et al., 2013). However, a third of the patients did not benefit from treatment (Kröger et al., 2013). In addition, in DBT the dropout rate was estimated around 27% and till now few predictors of dropout have been identified (e.g. Kliem et al., 2010). It is hypothesized that different BPD subtypes might show differences in treatment trajectories (e.g., Oldham, 2006), explaining the mixed treatment responses of BPD patients. To our knowledge, only one study so far, has evaluated differences in treatment responses between different BPD subtypes. Digre et al. (2009) assessed 74 BPD inpatients and identified three BPD subtypes focusing on attribution style. After six months of treatment, which consisted of some DBT skills training aspects, a significant improvement in depression, dissociation and less suicide attempts, but no significant change in NSSI was demonstrated. However, they did not find significant trajectory differences between the different BPD subtypes for each of these symptoms. BPD subtypes based on reactive and regulative temperament might demonstrate differences in treatment trajectories, since evidence from an earlier study found that clinical subtypes based on temperament dimensions were connected with clinical outcomes (Karalunas et al., 2014).

The aim of this study was threefold: (1) to replicate the prior BPD subtypes based on reactive and regulative temperament dimensions (see Sleuwaegen et al., 2017) with model-based clustering, since this method only generates multiple clusters if the data provides evidence for them; (2) to investigate differences in symptomatology and coping between the BPD subtypes at

baseline; and (3) to investigate the (potential differential) dropout rate and differences in symptomatology and coping of these BPD subtypes after 3 months of BPD inpatient treatment (Linehan, 1993). Although this study was exploratory in nature, several hypotheses were formulated based on prior research. First, we expected to find four BPD subtypes based on temperament as in a previous study (Sleuwaegen et al., 2017). Second, we expected intergroup differences in symptomatology and coping at baseline. The subtype with the highest level of BIS and BAS, and the lowest level of effortful control (Emotional/disinhibited cluster) would show the highest level of symptomatology and the worst coping compared to the other subtypes (Sleuwaegen et al., 2017). Third, it was expected that the total BPD sample would show a good treatment response (main effect of time) on general clinical and borderline specific symptoms, dissociation, NSSI and coping styles (e.g., Kröger et al., 2013). Since the exploration of trajectory differences among the BPD subtypes based on temperament dimensions is new, we did not formulate hypotheses concerning potential differences in symptomatology and coping after three months of DBT inpatient treatment. However, we hypothesized a higher dropout rate in the subtype with the lowest BIS level, consistent with past research in samples with anti-social personality characteristics (Daughters et al., 2008).

## 2 Methods

### 2.1 Participants and procedure

Our participants consisted of 145 BPD inpatients (88.3% females) hospitalized in a treatment unit for BPD patients, based on the principles of DBT (Linehan, 1993). Participants were included between May 2014 and April 2017. Patients were included if they fulfilled the criteria of BPD and they were excluded if they suffered from a current psychotic disorder or a current

substance use disorder. In case of current addiction, patients were referred to a specialized unit for detoxification and substance use disorder. Patients were allowed to be under pharmacological treatment. During the first week of admission, all patients were informed about the aims of the study and invited to participate. After giving written informed consent, patients were asked to fill out the questionnaires on a computer. All 145 patients (see Table 1 for more detailed information) met criteria for BPD diagnosis as assessed by means of the self-report Assessment of DSM-IV Personality Disorders - Borderline scale (ADP-IV; Schotte, 1994), and for 80 patients this BPD diagnosis was validated by the Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II; First et al., 1997; Dutch translation by Weertman et al., 2000). Assessment after 3 months (13 to 16 weeks) was carried out as an element of routine outcome monitoring which is integrated into the regular treatment routine to evaluate treatment response. After 3 months of DBT treatment, 75 inpatients provided data concerning clinical symptomatology and coping. Table 1 displays the sample characteristics for the sample at T1 and the sample at T2. The study was developed in accordance with the Declaration of Helsinki and approved by the local research and participating hospitals ethics committees. Participants did not receive any remuneration.

## 2.2 Measures

The diagnosis of BPD was assessed by means of the Assessment of DSM-IV Personality Disorders (ADP-IV; Schotte and De Doncker, 1994), a 94-item Dutch self-report questionnaire used to assess the presence and severity of the personality disorders as defined in the DSM-IV-TR (APA, 2000) and the Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II; First et al., 1997; Dutch translation by Weertman et al., 2000). Schotte et al. (2004)



indicated that the ADP-IV-BPD scale showed acceptable concordance with the SCID-II-BPD section ( $\kappa = 0.54$ ).

To replicate the BPD subtypes based on reactive/regulative temperament, we used the Behavioural Inhibition System/Behavioural Activation System Scales (BIS/BAS; Carver & White, 1994; Dutch translation: Franken et al., 2005) and the Effortful Control Scale (ECS) of the Adult Temperament Questionnaire-Short Form (ATQ-SF; Evans & Rothbart, 2007). The BISBAS scales consist of 24-items, rated on a 4-point Likert scale, of which seven items assess BIS reactivity ( $\alpha = .72$  in the present study), reflecting sensitivity to punishment, and 13 items assess BAS reactivity ( $\alpha = .86$ ), reflecting sensitivity to potentially rewarding outcome. Regulative temperament was assessed by means of the ECS scale with 19 items rated on a seven-point Likert scale. The internal consistency of the ECS scale was  $\alpha = .77$  in the present sample.

To validate the BPD subtypes and to investigate treatment response, several self-report measures were administered assessing clinical/BPD symptomatology and coping:

General clinical symptomatology was assessed by means of the Brief Symptom Inventory (BSI; Derogatis, 1993; translated into Dutch by de Beurs et al., 2005) consisting of 53 items, rated on a 4-point Likert scale. In the present study, the alpha coefficient for the total BSI-score was .95.

Borderline specific symptoms were measured by means of the Borderline Symptom List-23 (BSL-23; Bohus et al., 2009), in which patients rated 23 items on a 5-point Likert scale. Cronbach's alpha of the total BSL-score in the present study was .95.

Dissociation was assessed by means of the Dissociation Questionnaire (DIS-Q; Vanderlinden et al., 1993), consisting of 63-items to be rated on a 5-Likert scale. In this study the internal consistency coefficient of the total DIS-Q-score was excellent ( $\alpha = .96$ ).

Non-suicidal self-injury was assessed by means of the Self-Injury Questionnaire-Treatment Related (SIQ-TR; Claes & Vandereycken 2007). Participants had to answer in a yes/no question format whether they had ever intentionally engaged in self-injury without the intention to die (i.e., scratching, cutting, hitting, burning, picking or head banging) (life-time NSSI) and whether they were still engaging in NSSI at the current moment (current NSSI) (Claes & Vandereycken 2007).

Coping strategies were assessed by means of the Utrecht Coping List (UCL; Schreurs, et al., 1993), consisting of 47 items, to be rated on a 5-point Likert scale. Two coping styles can be distinguished: an active coping style (based on the subscales ‘active problem solving’ and ‘self-soothing thoughts’) and a passive coping style (based on the subscales ‘avoidance’, ‘depressive reactions’ and ‘expression of emotions’; Schreurs et al., 1993). In this study the alpha coefficient for active coping style was .84 and for passive coping style .69.

### 2.3 Statistical analysis

Statistical analyses were performed with SPSS version 24 and R version 3.4.3. There were no missing data given that the assessment on the computer did not allow for missing values.

To identify different BPD subtypes based on reactive and regulative temperament dimensions, we performed a model-based cluster analysis (Fraley & Raftery, 1998) on the standardized BIS/BAS and EC scale scores (i.e., *z*-scores) using the *mclust* package version 5.4 (Scrucca et al., 2016). Compared to standard clustering techniques such as hierarchical clustering and *k*-means, Model-Based Clustering (MBC) allows for more flexible cluster distributions in terms of their volume, shape and orientation. As such, MBC, in general, yields cluster solutions of a higher quality. In particular, MBC fits a mixture of Gaussian distributions to the data and

searches for the best model across a predefined set of models that differ in the volume, shape and orientation of their cluster distributions (see Scrucca et al., 2016). In our analysis, ten different models were tested with the number of clusters ranging from one to six (60 models in total). To overcome convergence problems when estimating the parameters of some of these models, we applied a Bayesian regularisation as recommended in Fraley and Raftery (2003). To choose the optimal model, we mainly relied on the Bayesian information criterion (BIC) but also took interpretability of the solution into account; BIC balances model fit with model complexity (i.e., number of clusters and complexity of the cluster distributions). We additionally compared the best MBC cluster solution with solutions obtained from the hierarchical cluster analysis and k-means clustering procedure.

To compare the different BPD-subtypes with respect to general clinical symptomatology, BPD specific symptomatology, dissociation and coping styles, we performed a MANOVA with BPD subtypes as independent variable and clinical, BPD symptomatology, dissociation and coping as dependent variables. The association between categorical variables (presence/absence of NSSI) and BPD subtype at baseline were analysed using the Chi-square test statistic.

To investigate the association between the presence/absence of dropout and BPD subtypes we used the Chi-square test statistic. In addition, to evaluate differences in the level of general clinical symptomatology, BPD specific symptomatology, dissociation and two coping styles at baseline and after 3 month of treatment in function of BPD subtype, we performed five repeated measures ANOVA's with time (baseline and 3-months) as within-subject factor, BPD subtype as between-subject factor, and general clinical symptomatology, BPD symptomatology, dissociation and two coping styles as dependent variables.

All significant group differences were evaluated using the post hoc Bonferroni test.  $P < .05$  was considered statistically significant. Partial eta-squared value was reported as a measure of effect size (Cohen, 1988).

To calculate the association between BPD subtype and NSSI throughout time, we used the Chi-Square test statistic. NSSI was coded as follows: 0 = no NSSI baseline - no NSSI 3m; 1 = NSSI baseline - no NSSI 3m; 2 = NSSI baseline - NSSI 3m).

### 3 Results

#### 3.1 BPD subtype clusters

BIS/BAS and EC scores were analyzed by means of a model-based cluster analysis to identify BPD subtypes. The three best fitting models using BIC were: a VVI (diagonal, varying volume and shape) model with 2 components,  $BIC = -1202.079$ ; a VVV (ellipsoidal, varying volume, shape, and orientation) model with 2 components,  $BIC = -1205.934$  and a VVI model with 3 components  $BIC = -1216.226$ . The other models all had a worse BIC value ( $BIC < -1231.189$ ). The two best fitting models (VVI and VVV with two clusters) gave very similar results and both clustered the patients into two groups that only differ in BIS level. As these clusters are not very heterogeneous in terms of temperamental dimensions, which was the goal of our study, we elaborated further on the third best fitting model, the three cluster model (VVI-3). This model resulted in clusters with a more distinguished profile regarding the three temperamental dimensions (BIS/BAS/EC). As, to our opinion, this solution is readily interpretable and clinically more relevant than the best fitting solution, we decided to prefer the VVI model with three clusters over the best fitting VVI2 model. Note that the difference in BIC

between both models is rather small. Moreover, the additional analyses with hierarchical and k-means clustering also favored a three-cluster solution.

The three BPD clusters showed substantively different means on the BIS/BAS and EC scale (see Figure 1). The first cluster was identified as the ‘Emotional/Disinhibited’ subtype ( $n = 22, 15\%$ ), characterized by moderately high scores on BIS, high scores on BAS and low scores on EC. The second cluster was named the ‘Low Anxiety subtype’ ( $n = 59, 41\%$ ), characterized by very low BIS, average low BAS, and neutral EC scores. The third and last cluster was called the ‘Inhibited subtype’ ( $n = 64, 44\%$ ), characterized by a high score on BIS, a low score on BAS and moderate score on EC. Note that the two clusters from the best fitting model (VV12) are closely related to the ‘Low Anxiety subtype’ and the ‘Inhibited subtype’, with the former subtype having a low BIS and the latter one a high BIS score (and subtypes not differing on BAS and EC scores). As such, the adopted three-cluster solution identified a new subtype -the ‘Emotional/Disinhibited’ subtype- with a clearly different reactive/regulative temperament, which justifies adopting the three-cluster solution instead of the slightly better fitting two-cluster solution. The patients in the three clusters did significantly differ with regard to gender [ $\chi^2(2) = 10.22, p = .006$ ] with the Low Anxiety subtype containing relatively more males (22% male and 78% female) than the ‘Emotional/Disinhibited’ (4.5% males) and the Inhibited subtype (4.7% males). There were no significant difference between the three subtypes for age [ $F(3,142) = .60, p = .55$ ], education [ $\chi^2(6) = 2.96, p = .81$ ], marital status [ $\chi^2(4) = .81, p = .94$ ] and overall medication use [yes/no:  $\chi^2(2) = 1.16, p = .56$ ].

### 3.2 Differences in symptomatology and coping of the three BPD subtypes at baseline

Table 2 lists the means and standard deviations of the different measures for each of the three BPD subtypes. The MANOVA showed only significant differences among the three BPD subtypes with respect to dissociation (total DIS-Q). The Emotional/Disinhibited subtype scored significantly higher on the DIS-Q compared to the Inhibited subtype ( $p = .002$ ), with a moderate effect size (partial  $\eta^2 = .089$ ).

There was no significant difference between the three BPD subtypes for presence of lifetime NSSI [Emotional/Disinhibited subtype: 81.8%, Low Anxiety: 84.7%, Inhibited: 79.7%;  $\chi^2(2) = .53, p = .765$ ], nor for presence of current NSSI [resp. 51.6%, 61.0%, 36.4%;  $\chi^2(2) = 4.02, p = .134$ ].

Finally, the three BPD subtypes showed no significant difference on the UCL active and passive coping styles (resp.  $p = .72$  and  $p = .20$ ).

### 3.3 Difference in dropout and treatment response after three months in the three BPD-subtypes

After three months of DBT inpatient treatment, 34.5% ( $n = 50$ ) of all patients dropped out. There was no significant difference between the three BPD subtypes and the presence/absence of dropout [Emotional/Disinhibited subtype: 27.3% ( $n = 6$ ), Low Anxiety subtype: 39% ( $n = 23$ ), and Inhibited subtype: 32.8% ( $n = 21$ );  $\chi^2(2) = 1.11, p = .573$ ].

Almost 80% of the 95 patients who were still in treatment completed the second measurement ( $n=75$ ). Table 3 displays the means and standard deviations of the different self-report measures at baseline and after three months of DBT treatment for each of the three BPD subtypes; and

Table 4 lists the results of each of the repeated measures ANOVA's.

We found significant main effects of time for all the self-report measures. The results clearly showed a decrease in general clinical symptomatology (BSI), borderline specific symptomatology (BSL), dissociation (DIS-Q) and passive coping style (UCL-PASS); and a significant increase in active coping style (UCL-ACT) from baseline till three month after DBT treatment with all large effect sizes (ranging from partial  $\eta^2 = .210$  to  $.335$ ).

Additionally, a significant interaction effect of time by BPD subtype was observed for BSI ( $p = .010$ ) and BSL ( $p = .024$ ) with large effect sizes (resp. partial  $\eta^2 = .123$  and  $.107$ ), and DIS-Q ( $p = .036$ ) with a moderate effect size (partial  $\eta^2 = .088$ ), see Table 4. Figures 2, 3 and 4 respectively illustrate the progress of general clinical symptoms, borderline specific symptoms and dissociation in the three BPD subtypes after three months of DBT treatment. It should be noted that patients of the 'Emotional/Disinhibited' subtype shows a larger progress on these measures, indicating that it is justified to consider this patient cluster as a separate subtype.

Since the interaction effect of time by subtype was significant for BSI (clinical symptoms), BSL (borderline symptoms) and DIS-Q (dissociation), we investigated the time effect (baseline - 3 months) on the different symptom measures within each subtype using paired samples t-tests. For the Emotional/Disinhibited subtype, a significant decrease was found for BSI [ $t(10) = 5.97, p < .001$ ], BSL [ $t(10) = 4.35, p = .002$ ] and DIS-Q [ $t(10) = 2.75, p = .021$ ]. For the Low Anxiety subtype, no significant difference was found in symptom measurements between baseline and 3 months (BSI [ $t(26) = 1.46, p = .156$ ], BSL [ $t(26) = 1.15, p = .260$ ], DIS-Q [ $t(26) = 1.77, p = .088$ ]). For the Inhibited subtype, a significant decrease was found for BSI [ $t(36) = 4.36, p < .001$ ], BSL [ $t(36) = 2.38, p = .023$ ] and DIS-Q [ $t(36) = 3.41, p = .002$ ].

In addition, no significant interaction effects of “time by subtype” were demonstrated for UCL active and passive coping style (resp.  $p = .35$  and  $p = .22$ ).

Finally, of all BPD patients who engaged in NSSI at baseline, 60.0% of the Emotional/Disinhibited subtype, 60.9% of the Low Anxiety and 62.5% of the Inhibited reported the absence of current NSSI at 3 months follow-up. Analysing the trajectory on current NSSI from baseline to 3 months after treatment for the three subtypes, revealed no difference in NSSI trajectory between the BPD subtypes [ $\chi^2(4) = 4.97, p = .290$ ], indicating that the different subtypes showed the same improvement for NSSI.

#### 4 Discussion

The aim of the present study was to confirm the existence of BPD subtypes based on temperament, to validate them in terms of symptomatology and coping, and to investigate their differences in treatment response after 3 months of inpatient DBT. To replicate previously identified BPD subtypes, we performed a model-based cluster analysis on reactive and regulative temperament dimensions in a sample of 145 BPD patients who were treated at a DBT unit. We identified three subtypes, the ‘Emotional/Disinhibited’, ‘Low Anxiety’, and ‘Inhibited’ subtypes, demonstrating the same characteristics as the three dominant subtypes found in a previous study using a hierarchical and k-means clustering procedure (see Sleuwaegen et al., 2017). The fourth and smallest cluster found in the previous study, namely the ‘High Self-control subtype’ (characterized by high scores on EC and moderately high scores on BIS and BAS) was not found in this sample. In this sample, the Inhibited subtype is also characterized by low levels of BAS reactivity (low reward sensitive) and moderate effortful control, however a higher level of BIS was demonstrated. The Low Anxiety subtype is characterized by very low levels of BIS reactivity



(low punishment sensitive) and low levels of BAS and a mean level of effortful control. The Emotional/Disinhibited subtype is characterized by a high level of BAS, moderate BIS and low effortful control. Patients of the Emotional/Disinhibited subtype reported a higher level of dissociation compared to the other two BPD subtypes. There was no difference in lifetime and current NSSI between the different BPD-subtypes.

The third, and most clinical relevant, aim of this study was to investigate differences in dropout and treatment response of these three BPD subtypes after 3 months of inpatient DBT treatment. Dropout rate in the total sample was 34.5%, however there was no significant difference in amount of dropout between the three BPD subtypes. Trajectory measurement on data of 75 patients who filled out the questionnaires at baseline and after 3 months DBT inpatient treatment, demonstrated a significant reduction in general clinical symptoms, borderline specific and dissociative symptoms with large effect sizes in all BPD subtypes. Moreover, there was a significant interaction effect (time by BPD subtype) on general clinical and borderline specific symptoms and dissociation, explaining a different evolution between the three subtypes on the level of symptomatology. The results demonstrated that, in contrast to the Emotional/Disinhibited subtype and Inhibited subtype, the Low Anxiety subtype did not improve significantly in general clinical and borderline specific symptoms and dissociation. These aforementioned results are in line with a previous study of Kröger et al. (2013), who also reported a reduction in clinical symptoms in almost all patients after a 3-month DBT program, but for borderline-specific symptoms only for half of them.

In the present study, almost 80% of the patients reported to engage in NSSI at baseline and 60% of these patients did not engage in NSSI anymore after 3 months of DBT inpatient treatment. However, the reduction of NSSI over time was not significant and there was no difference in NSSI trajectory between the three BPD subtypes, which is in line with the findings

of Digre et al. (2009). Finally, concerning coping style, all BPD patients showed an increase in active coping and a decrease in passive coping after three month of treatment. However, there was no evidence for significant differences in coping progress between the three BPD subtypes.

However, our study is not without limitations which need to be taken into account in future studies. The first limitation refers to the short follow-up period of three month of treatment and high drop-out rate. Therefore, future studies should try to include more patients followed during a longer time period, to assess the treatment trajectories of different BPD subtypes. A second limitation is the fact that the data were collected in a specialized DBT unit, over-represented by female and medicated patients, which makes the sample not representative for the general BPD population. As a third limitation, all baseline/outcome assessments were solely based on self-report measures, with the known limitations (Mc Donald, 2008). In addition, since we have only patients with BPD in the sample, the associations between the temperamental dimensions may not be representative for other samples.

In conclusion, the present study confirms the existence of BPD subtypes based on reactive and regulative temperament. Furthermore, in addition to the finding that all subtypes show significant improvement in level of symptomatology and coping after three months of DBT inpatient treatment, this study provides evidence for trajectory differences on general clinical and borderline specific symptoms and dissociation between the three subtypes. The Low Anxiety subtype does not demonstrate a significant reduction in symptomatology compared to the two other subtypes. This finding is an incentive for future research on the trajectory of BPD subtypes based on temperament, in larger samples and with longer follow-up periods. The conclusions may guide treatment choices and/or the development of subtype-tailored treatment in the future, which would be of clinical and social benefit.

## References


- American Psychiatric Association (2000). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed., text revision). Washington, DC: APA.
- Claes, L., & Vandereycken, W. (2007). The self-injury questionnaire-treatment related (SIQ-TR): Construction, reliability, and validity in a sample of female eating disorder patients. In P. M. Goldfarb (Ed.), *Psychological tests and testing research trends* (pp. 111–139). New York, NY: Nova Science Publishers.
- Bohus, M., Haaf, B., Simms, T., Limberger, M. F., Schmahl, C., Unckel, C., et al. (2004). Effectiveness of inpatient dialectical behavioral therapy for borderline personality disorder: A controlled trial. *Behaviour Research and Therapy*, *42*, 487–499.
- Bohus, M., Kleindienst, N., Limberger, M. F., Stieglitz, R. D., Domsalla, M., Chapman, A., et al. (2009) The short version of the borderline symptom list (BSL-23): Development and initial data on psychometric properties. *Psychopathology*, *42*, 32-39.
- Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: the BIS/BAS scales. *Journal of Personality and Social Psychology*, *67*, 319–333.
- Cohen, J. (1988). *Statistical power analysis for the behavioural sciences*. (2nd ed.). Hillsdale, NJ: L. Erlbaum associates.
- Daughters, S. B., Stipelman, B. A., Sargeant, M. N., Schuster, R., Bornovalova, M. A., & Lejuez, C., W. (2008). The interactive effects of antisocial personality disorder and court-mandated status on substance abuse treatment dropout. *Journal of Substance Abuse Treatment*, *34*, 157-164.

- De Beurs, E. (2005). *De Brief Symptom Inventory*; Handleiding [The Brief Symptom Inventory; Manual]. Leiden: Pits Publishers.
- Derogatis, L. R. (1993). *Brief Symptoms Inventory (BSI): Administration, scoring and procedures manual* (3rd ed.). Minneapolis: NCS Pearson, Inc.
- Digre, E. I., Reece, J., Johnson, A. L., & Thomas, R. A. (2009). Treatment response in subtypes of borderline personality disorder. *Personality and Mental Health, 3*, 56–67.
- Evans, D. E., & Rothbart, M. K. (2007). Developing a model for adult temperament. *Journal of Research in Personality, 41*, 868–888.
- First, M. B., Gibbon, M., Spitzer, R. L., Williams, J. B. W., & Benjamin, L. S. (1997). *Structured clinical interview for DSM-IV Axis II disorders*. Washington, DC: American Psychiatric Press.
- Fraley, C., & Raftery, A. E. (1998). How many clusters? Which clustering method? Answer via model-based cluster analysis. *The Computer Journal, 41*, 578-588.
- Fraley, C., & Raftery, A. E. (2003). Enhanced software for model-based clustering, discriminant analysis, and density estimation: MCLUST. *Journal of Classification, 20*, 263-286.
- Franken, I. H. A., Muris P., & Rassin E. (2005). Psychometric properties of the Dutch BIS/BAS scales. *Journal of Psychopathology and Behavioral Assessment, 27*, 25–30
- Frias, A., Navarro, S., Palma, C., Farraiols, N., Alagia, F., Salvador, A., et al. (2017). Defining subtypes of borderline personality disorder based on underlying dimensional personality traits. *Personality and Mental Health, 27*, 25–30
- Karalunas, S. L., Fair, D., Musser, E. D., Aykes, K., Iyer, S. P., & Nigg, J. T. (2014). Subtyping attention-deficit/hyperactivity disorder using temperament dimensions: toward biologically based nosologic criteria. *JAMA Psychiatry, 71*, 1015-1024.

- Kliem, S., Kroger, C., Kosfelder, J. (2010). Dialectical behavior therapy for borderline personality disorder: a meta-analysis using mixed-effects modeling. *Journal of Consulting and Clinical Psychology, 78*, 936-951.
- Kopala-Sibley, D. C., Zuroff, D. C., Russell, J. J., Moskowitz, D. S., & Paris, J. (2012). Understanding Heterogeneity in Borderline Personality Disorder: Differences in Affective Reactivity Explained by the Traits of Dependency and Self-Criticism. *Journal of Abnormal Psychology, 121*, 680–691.
- Kröger, C., Harbeck, S., Ambrust, W., & Kliem, S. (2013). Effectiveness, response and dropout of dialectical behavior therapy for borderline personality disorder in an inpatient setting. *Behaviour Research and Therapy, 51*, 411-416.
- Linehan (1993) Cognitive-behavioral treatment of Borderline Personality Disorder. New York, NY: The Guilford Press
- McDonald, J. D. (2008). Measuring Personality Constructs: The advantages and disadvantages of self-reports, informant reports and behavioural assessments. *Enquire, 1*, 75–94.
- Müller, A., Claes, L., Wilderjans, T. & de Zwaan, M. (2014). Temperament Subtypes in Treatment Seeking Obese Individuals: A Latent Profile Analysis. *European Eating Disorders Review, 22*, 260-266.
- Oldham, J. M. (2006). Borderline personality disorder and suicidality. *American Journal of Psychiatry, 163*, 20-26.
- Schotte, C. K. W., & De Doncker, D. (1994). *ADP-IV vragenlijst*. Antwerpen: Universitair Ziekenhuis Antwerpen.

- Schotte, C. K. W., De Doncker, D., Dmitruk, D., Van Mulders, I., D'Haenen, H., & Cosyns, P. (2004). The ADP-IV questionnaire: Differential validity and concordance with semistructured interview. *Journal of Personality Disorders, 18*, 405–419
- Schreurs, P. J. G., van de Willige, G., Brosschot, J. F., Tellegen, B., & Graus, G. M. H. (1993). *De Utrechtse Coping Lijst: UCL* [The Utrecht Coping List: UCL]. Lisse: Swets & Zeitlinger.
- Sleuwaegen, E., Claes, L., Luyckx, K., Berens, A., Vogels, C., & Sabbe, B. (2017). Subtypes in borderline patients based on reactive and regulative temperament. *Personality and Individual Differences, 108*, 14–19.
- Smits, M. L., Feenstra, D. J., Bales, D. L., de Vos, J., Lucas, Z., Verheul, R. & Luyten, P. (2017). Subtypes of borderline personality disorder patients: a cluster-analytic approach. *Borderline Personality Disorder and Emotion Dysregulation, 4:16*, 1–15.
- Vanderlinden, J., Van Dyck, R., Vandereycken, W., Vertommen, H., & Verkes, R., J. (1993). The dissociation questionnaire (DIS-Q): development and characteristics of a new self-report questionnaire. *Clinical Psychology and Psychotherapy, 1*, 21-27.
- Weertman, A., Arntz, A., & Kerkhofs, M. L. M. (2000). *Gestructureerd diagnostisch interview voor DSM-IV persoonlijkheidsstoornissen (SCID-II)* [Structured Clinical Interview for DSM-IV Personality Disorders (SCID-II)]. Lisse: Swets Test.


Figure 1. BPD subtypes based on reactive and regulative temperament dimensions in 145 BPD patients



Note: BIS = Behavioural Inhibition System; BAS = Behavioural Activation System; EC = Effortful Control

BPD Subtypes : Emo/Dis = Emotional/Disinhibited, Low Anx = Low Anxiety, Inh = Inhibited


Figure 2. Reduction of general clinical symptoms after 3 months DBT treatment for the three BPD subtypes



*Note:* BSI = Brief Symptom Inventory, T1 = baseline; T2 : after 3 months DBT inpatient treatment  
BPD Subtypes : Emo/Dis = Emotional/Disinhibited, Low Anx = Low Anxiety, Inh = Inhibited




Figure 3. Reduction of borderline specific symptoms after 3 months DBT treatment for the three BPD subtypes



Note: BSL = Borderline Symptom List, T1 = baseline; T2 : after 3 months DBT inpatient treatment  
BPD Subtypes : Emo/Dis = Emotional/Disinhibited, Low Anx = Low Anxiety, Inh = Inhibited

Figure 4. Reduction of dissociation symptoms after 3 months DBT treatment for the three BPD subtypes



Note: DIS-Q = Dissociation Questionnaire, T1 = baseline; T2 : after 3 months DBT inpatient treatment  
BPD Subtypes : Emo/Dis = Emotional/Disinhibited, Low Anx = Low Anxiety, Inh = Inhibited

Table 1. Sample characteristics for the total sample of BPD patients at baseline and for the subsample at 3 months assessment

	Total sample at T1	Sample at T2
	n=145	n=75
Gender – Female (%)	88.3	91.9
Age ( <i>M, SD</i> )	29.72 (9.32)	28.75 (8.83)
Education (%)		
Lower secondary	19.7	18.9
Higher secondary	57.8	55.4
High School	17.6	18.9
University	4.7	6.8
Marital status (%)		
Single	71.4	71.6
Living together/Married	19.7	18.9
Divorced	8.8	9.5
Medication (%)	84.8	85.1
Antidepressants	71.7	70.3
Antipsychotics	48.3	44.6
Anxiolytics	6.2	6.8
Mood Stabilizers	11.7	10.8
(Note. Antipsychotics provided as mood regulator)		

Table 2. Means (standard deviations) of the self-report scales at baseline for the three BPD subtypes.

	Emotional/Disinhibited subtype (n=22)	Low Anxiety subtypes (n=59)	Inhibited subtype (n=64)			
	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>F</i>	partial $\eta^2$	Post-hoc comparisons
BSI	123.59 (26.77)	108.78 (38.71)	122.02 (32.3)	2.80	.04	
BSL	50.55 (23.49)	48.78 (21.92)	55.02 (19.09)	1.39	.02	
DIS-Q	199.32 (25.87)	163.97 (42.70)	165.17 (42.32)	6.89*	.09	1<2,3
UCL-ACT	23.23 (7.02)	22.34 (6.19)	22.06 (5.01)	.33	.05	
UCL-PASS	49.23 (5.73)	46.58 (6.74)	46.56 (6.28)	1.61	.02	

*Note.* BSI = Brief Symptom Inventory – total score, BSL = Borderline Symptom List – total score; DIS-Q = Dissociation Questionnaire, UCL-ACT = Utrecht Coping List – Active coping style score, UCL-PASS = Utrecht Coping List – Passive coping style score

\*  $p < 0.05$ .

Table 3. Means (standard deviations) of the self-report scales at baseline (T1) and after 3 months (T2) for the three BPD subtypes.

	<u>Emotional/Disinhibited subtype</u>		<u>Low Anxiety subtype</u>		<u>Inhibited Subtype</u>	
	<u>(n=11)</u>		<u>(n=27)</u>		<u>(n=37)</u>	
	<u>T1</u>	<u>T2</u>	<u>T1</u>	<u>T2</u>	<u>T1</u>	<u>T2</u>
	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>
BSI	134.91 (17.77)	93.73 (18.69)	99.37 (33.12)	93.20 (42.70)	125.05 (30.90)	98.81 (41.64)
BSL	63.18 (17.71)	41.33 (14.54)	47.35 (20.28)	42.96 (23.65)	55.35 (19.03)	45.97 (24.14)
DIS-Q	207.82 (29.02)	164.91 (36.70)	154.30 (40.71)	143.59 (38.48)	162.97 (44.70)	146.00 (48.61)
UCL-ACT	24.64 (7.23)	29.64 (7.70)	21.26 (5.75)	23.56 (5.50)	22.81 (5.46)	26.81 (5.85)
UCL-PASS	50.18 (5.31)	43.91 (6.09)	45.26 (6.02)	43.19 (6.94)	45.27 (6.68)	42.08 (7.02)

*Note.* BSI = Brief Symptom Inventory – total score, BSL = Borderline Symptom List – total score; DIS-Q = Dissociation Questionnaire, UCL-ACT = Utrecht Coping List – Active coping style score, UCL-PASS = Utrecht Coping List – Passive coping style score

Table 4. Results for the Repeated Measurement ANOVA with time as within-subject factor, BPD subtype as between-subject factor, and clinical, BPD symptomatology and coping as dependent variables.

	Main effects			Main effects			Interaction effect		
	Time			Subtype			Time * Subtype		
	<i>F</i> (1,144)	<i>p</i>	partial $\eta^2$	<i>F</i> (2,143)	<i>p</i>	partial $\eta^2$	<i>F</i> (2,143)	<i>p</i>	partial $\eta^2$
BSI	35.78	<.001	.335	2.91	.130	.056	4.96	.010	.123
BSL	21.19	<.001	.243	1.29	.280	.038	3.96	.024	.107
DIS-Q	27.07	<.001	.273	3.79	.027	.088	3.49	.036	.088
UCL-ACT	24.18	<.001	.251	3.65	.031	.092	1.07	.348	.029
UCL-PASS	19.12	<.001	.210	1.51	.227	.040	1.53	.222	.041

*Note.* BSI = Brief Symptom Inventory – total score, BSL = Borderline Symptom List – total score; DIS-Q = Dissociation Questionnaire, UCL-ACT = Utrecht Coping List – Active coping style score, UCL-PASS = Utrecht Coping List – Passive coping style score