



Universiteit
Leiden

The Netherlands

**Human-wildlife interactions in the Western Terai of Nepal.
An analysis of factors influencing conflicts between
sympatric tigers (*Panthera tigris tigris*) and leopards
(*Panthera pardus fusca*) and local communities around
Bardia National Park, Nepal**

Upadhyaya, S.K.

Citation

Upadhyaya, S. K. (2019, April 16). *Human-wildlife interactions in the Western Terai of Nepal. An analysis of factors influencing conflicts between sympatric tigers (*Panthera tigris tigris*) and leopards (*Panthera pardus fusca*) and local communities around Bardia National Park, Nepal*. Retrieved from <https://hdl.handle.net/1887/71374>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/71374>

Note: To cite this publication please use the final published version (if applicable).

References

- Acharya, K. P., Paudel, P. K., Neupane, P. R., & Köhl, M. (2016). Human-Wildlife Conflicts in Nepal: Patterns of Human Fatalities and Injuries Caused by Large Mammals. *PLOS ONE*, *11*(9), e0161717.
- Acharya, K. P., Paudel, P. K., Jnawali, S. R., Neupane, P. R., & Köhl, M. (2017). Can forest fragmentation and configuration work as indicators of human-wildlife conflict? Evidences from human death and injury by wildlife attacks in Nepal. *Ecological Indicators*, *80*, 74–83.
- Allendorf, T., Swe, K. K., Oo, T., Htut, Y., Aung, M., Aung, M., ... Wemmer, C. (2006). Community attitudes toward three protected areas in Upper Myanmar (Burma). *Environmental Conservation*, *33*(4), 344–352.
- Allendorf, T. D., Smith, J. L. D., & Anderson, D. H. (2007). Residents' perceptions of Royal Bardia National Park, Nepal. *Landscape and Urban Planning*, *82*(1–2), 33–40.
- Allendorf, T. D. (2010). A framework for the park-people relationship: Insights from protected areas in Nepal and Myanmar. *International Journal of Sustainable Development and World Ecology*, *17*(5), 417–422.
- Amarasekare, P. (2008). Coexistence of intraguild predators and prey in resource-rich environments. *Ecology*, *89*(10), 2786–2797.
- Andheria, A. P., Karanth, K. U., & Kumar, N. S. (2007). Diet and prey profiles of three sympatric large carnivores in Bandipur Tiger Reserve, India. *Journal of Zoology*, *273*(2), 169–175.
- Athreya, V., Odden, M., Linnell, J. D. C., Krishnaswamy, J., & Karanth, U. (2013). Big Cats in Our Backyards: Persistence of Large Carnivores in a Human Dominated Landscape in India. *PLoS ONE*, *8*(3).
- Azlan, J. M., & Sharma, D. S. K. (2006). The diversity and activity patterns of wild felids in a secondary forest in Peninsular Malaysia. *Oryx*, *40*(1), 36–41.
- Bagchi, S., Goyal, S. P., & Sankar, K. (2003). Prey abundance and prey selection by tigers (*Panthera tigris*) in a semi-arid, dry deciduous forest in western India. *Journal of Zoology*, *260*(3), 285–290.
- Bahuguna, A., Sahajpal, V., Goyal, S.P., Mukherjee, S.K., & Thakur, V. (2010). *Species identification from guard hair of selected Indian Mammals: A reference Guide*. Wildlife Institute of India, Dehradun, India.
- Bailey, T. N. (1993). *The African leopard: ecology and behavior of a solitary felid* (Vol. 430). New York: Columbia University Press.

- Balme, G., Hunter, L., & Slotow, R. (2007). Feeding habitat selection by hunting leopards *Panthera pardus* in a woodland savanna: prey catchability versus abundance. *Animal Behaviour*, *74*(3), 589–598.
- Baral, N., & Heinen, J. T. (2007). Resources use, conservation attitudes, management intervention and park-people relations in the Western Terai landscape of Nepal. *Environmental Conservation*, *34*(01), 64–72.
- Basak, K., Mandal, D., Babu, S., Kaul, R., Ashraf, N. V. K., Singh, A., & Mondal, K. (2018). Prey Animals of Tiger (*Panthera tigris tigris*) in Dudhwa Landscape, Terai Region, North India. *Proceedings of the Zoological Society*, *71*(1), 92–98.
- Bauer, H., & Iongh, H. H. (2005). Lion (*Panthera leo*) home ranges and livestock conflicts in Waza National Park, Cameroon. *African Journal of Ecology*, *43*(3), 208–214.
- Bhagavatula, J., & Singh, L. (2006). Genotyping faecal samples of Bengal tiger *Panthera tigris tigris* for population estimation: A pilot study. *BMC Genetics*, *7*, 1–12.
- Bhattarai, B. R. (2009). *Human-tiger (Panthera tigris tigris) conflict in Bardia National Park, Nepal*. Ernst Moritz Arndt University of Greifswald.
- Bhattarai, B. P., & Kindlmann, P. (2012). Interactions between Bengal tiger (*Panthera tigris*) and leopard (*Panthera pardus*): Implications for their conservation. *Biodiversity and Conservation*, *21*(8), 2075–2094.
- Bhattarai, B. R., & Fischer, K. (2014). Human–tiger *Panthera tigris* conflict and its perception in Bardia National Park, Nepal. *Oryx*, *48*(04), 522–528.
- Bhattarai, B., Wright, W., & Khatiwada, A. (2016). Illegal Hunting of Prey Species in the Northern Section of Bardia National Park, Nepal: Implications for Carnivore Conservation. *Environments*, *3*(4), 32.
- Biswas, S., & Sankar, K. (2002). Prey abundance and food habit of tigers (*Panthera tigris tigris*) in Pench National Park, Madhya Pradesh, India. *Journal of Zoology*, *256*(3), 411–420.
- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., & White, J. S. S. (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution*, *24*(3), 127–135.
- Borthakur, U., Barman, R. D., Das, C., Basumatary, A., Talukdar, A., Ahmed, M. F., ... Bharali, R. (2011). Noninvasive genetic monitoring of tiger (*Panthera tigris tigris*) population of Orang National Park in the Brahmaputra floodplain, Assam, India. *European Journal of Wildlife Research*, *57*(3), 603–613.
- Bothma, J., & Le Riche, E. N. (1984). Aspects of the ecology and the behaviour of the leopard *Panthera pardus* in the Kalahari Desert. *Koedoe: African Protected Area Conservation and Science*, *27*(2), 259–279.
- Breuer, T. (2005). Diet choice of large carnivores in northern Cameroon Diet choice of large carnivores in northern Cameroon. *African Journal of Ecology*, *43*, 97–106.

- Brown, K. (1998). The political ecology of biodiversity, conservation and development in Nepal's Terai: Confused meanings, means and ends. *Ecological Economics*, 24(1), 73–87.
- Budhathoki, P. (2003). A Category V Protected Landscape approach to buffer zone management in Nepal. *Parks*, 13(2), 22–30.
- Budhathoki, P. (2004). Linking communities with conservation in developing countries: buffer zone management initiatives in Nepal. *Oryx*, 38(03), 334–341.
- Carbone, C., Teacher, A., & Rowcliffe, J. M. (2007). The costs of carnivory. *PLoS Biology*, 5(2), 0363–0368.
- Carter, N. H., Shrestha, B. K., Karki, J. B., Pradhan, N. M. B., & Liu, J. (2012). Coexistence between wildlife and humans at fine spatial scales. *Proceedings of the National Academy of Sciences*, 109(38), 15360–15365.
- Carter, N. H., Riley, S. J., Shortridge, A., Shrestha, B. K., & Liu, J. (2014). Spatial assessment of attitudes toward tigers in Nepal. *Ambio*, 43(2), 125–137.
- Carter, N., Jasny, M., Gurung, B., & Liu, J. (2015). Impacts of people and tigers on leopard spatiotemporal activity patterns in a global biodiversity hotspot. *Global Ecology and Conservation*, 3, 149–162.
- Carter, N. H., & Linnell, J. D. C. (2016). Co-Adaptation Is Key to Coexisting with Large Carnivores. *Trends in Ecology and Evolution*, 31(8), 575–578.
- Caughley, G., & Sinclair, A. R. E. (1994). *Wildlife ecology and management*. Blackwell Science.
- Chakrabarti, S., Jhala, Y. V., Dutta, S., Qureshi, Q., Kadivar, R. F., & Rana, V. J. (2016). Adding constraints to predation through allometric relation of scats to consumption. *Journal of Animal Ecology*, 85(3), 660–670.
- Chanchani, P., Lamichhane, B. R., Malla, S., Maurya, K., Bista, A., Warriar, R., ... & Dhakal, M. (2014). *Tigers of the Transboundary Terai Arc Landscape: Status, distribution and movement in the Terai of India and Nepal*. National Tiger Conservation Authority, Government of India, and Department of National Park and Wildlife Conservation, Government of Nepal NTNC/DNPWC.
- Chapron, G., Miquelle, D. G., Lambert, A., Goodrich, J. M., Legendre, S., & Clobert, J. (2008). The impact on tigers of poaching versus prey depletion. *Journal of Applied Ecology*, 45(6), 1667–1674.
- Chauhan, N. P. S., Barwal, K. S., & Kumar, – Devender. (2009). Human–Wild Pig Conflict in Selected States in India and Mitigation Strategies. *Acta Silv. Lign. Hung. Acta Silv. Lign. Hung*, 5(5), 189–197.
- Croes, B. M., Funston, P. J., Rasmussen, G., Buij, R., Saleh, A., Tumenta, P. N., & de Iongh, H. H. (2011). The impact of trophy hunting on lions (*Panthera leo*) and other large carnivores in the Bénoué Complex, northern Cameroon. *Biological Conservation*, 144(12), 3064–3072.

- Damania, R., Stringer, R., Karanth, K. U., & Stith, B. (2002). The Economics of Protecting Tiger Populations: Linking Household Behaviour to Poaching and Prey Depletion. *SSRN Electronic Journal*, 79(2), 198–216.
- Dar, N. I., Minhas, R. A., Zaman, Q., & Linkie, M. (2009). Predicting the patterns, perceptions and causes of human-carnivore conflict in and around Machiara National Park, Pakistan. *Biological Conservation*, 142(10), 2076–2082.
- Davies, T. E., Wilson, S., Hazarika, N., Chakrabarty, J., Das, D., Hodgson, D. J., & Zimmermann, A. (2011). Effectiveness of intervention methods against crop-raiding elephants. *Conservation Letters*, 4(5), 346–354.
- Dhakal, M., Karki, M., Jnawali, S. R., Subedi, N., Pradhan, N. M. B., Malla, S., ... Oglethorpe, J. (2014). *Status of tigers and prey in Nepal*. Department of National Park and Wildlife Conservation, Kathmandu, Nepal.
- Dhungana, R., Savini, T., Karki, J. B., Dhakal, M., Lamichhane, B. R., & Bumrungsri, S. (2018). Living with tigers *Panthera tigris*: Patterns, correlates, and contexts of human-tiger conflict in Chitwan National Park, Nepal. *Oryx*, 52(1), 55–65.
- Dickman, A. J. (2010). Complexities of conflict: The importance of considering social factors for effectively resolving human-wildlife conflict. *Animal Conservation*, 13(5), 458–466.
- Dinerstein, E. (1979). An ecological survey of the Royal Karnali Bardia Wildlife Reserve, Nepal. Part II: Habitat and animal interactions. *Biological Conservation*, 16(4), 265–300.
- Dinerstein, E., Loucks, C., Wikramanayake, E., Ginsberg, J., Sanderson, E., Seidensticker, J., ... Songer, M. (2007). The Fate of Wild Tigers. *BioScience*, 57(6), 508–514.
- DNPWC | Department of National Parks and Wildlife Conservation. (2018). Retrieved August 13, 2018, from <http://www.dnpwc.gov.np/>
- Dublin, H. T., & Hoare, R. E. (2004). Searching for solutions: The evolution of an integrated approach to understanding and mitigating human–elephant conflict in africa. *Human Dimensions of Wildlife*, 9(4), 271–278.
- Dutta, T., Sharma, S., Maldonado, J. E., Wood, T. C., Panwar, H. S., & Seidensticker, J. (2013). Gene flow and demographic history of leopards (*Panthera pardus*) in the central Indian highlands. *Evolutionary Applications*, 6(6), 949–959.
- Edgaonkar, A., & Chellam, R. (2002). in the Sanjay Gandhi National Park , Maharashtra , India. *Mammalia*, 66, 353–360.
- Farrell, L. E., Roman, J., & Sunquist, M. E. (2000). Dietary separation of sympatric carnivores identified by molecular analysis of scats.pdf. *Molecular Ecology*, 9, 1583–1590.
- Fisher, M. (2016). Whose conflict is it anyway? Mobilizing research to save lives. *Oryx*, 50(03), 377–378.

- Fuller, T., & Sievert, P. (2001). Carnivore demography and the consequences of changes in prey availability. *Carnivore Conservation*, (December), 163–179.
- Gilbert, M., Soutyrina, S. V., Seryodkin, I. V., Sulikhan, N., Uphyrkina, O. V., Goncharuk, M., ... Miquelle, D. G. (2015). Canine distemper virus as a threat to wild tigers in Russia and across their range. *Integrative Zoology*, 10(4), 329–343.
- Global Tiger Initiative. (2011). *Global Tiger Recovery Program 2010–2022*. Global Tiger Initiative.
- Goodrich, J. M., Kerley, L. L., Smirnov, E. N., Miquelle, D. G., McDonald, L., Quigley, H. B., ... McDonald, T. (2008). Survival rates and causes of mortality of Amur tigers on and near the Sikhote-Alin Biosphere Zapovednik. *Journal of Zoology*, 276(4), 323–329.
- Goodrich, J. M., Miquelle, D. G., Smirnov, E. N., Kerley, L. L., & Howard, B. (2010). Spatial structure of Amur (Siberian) tigers (*Panthera tigris altaica*) on Sikh ... *Journal of Mammalogy*, 91(3), 737–748.
- Gross, E. M., Drouet-Hoguet, N., Subedi, N., & Gross, J. (2017). The potential of medicinal and aromatic plants (MAPs) to reduce crop damages by Asian Elephants (*Elephas maximus*). *Crop Protection*, 100, 29–37.
- Gunn, J., Hawkins, D., Barnes, R. F. W., Mofulu, F., Grant, R. A., & Norton, G. W. (2014). The influence of lunar cycles on crop-raiding elephants; evidence for risk avoidance. *African Journal of Ecology*, 52(2), 129–137.
- Gurung, B., Smith, J. L. D., McDougal, C., Karki, J. B., & Barlow, A. (2008). Factors associated with human-killing tigers in Chitwan National Park, Nepal. *Biological Conservation*, 141(12), 3069–3078.
- Harihar, A., Pandav, B., & Goyal, S. P. (2009). Density of leopards (*Panthera pardus*) in the Chilla Range of Rajaji National Park, Uttarakhand, India. *Mammalia*, 73(1), 68–71.
- Harihar, A., Pandav, B., & Goyal, S. P. (2011). Responses of leopard *Panthera pardus* to the recovery of a tiger *Panthera tigris* population. *Journal of Applied Ecology*, 48(3), 806–814.
- Harmsen, B. J., Foster, R. J., Silver, S. C., Ostro, L. E. T., & Doncaster, C. P. (2009). Spatial and Temporal Interactions of Sympatric Jaguars (*Panthera onca*) and Pumas (*Puma concolor*) in a Neotropical Forest. *Journal of Mammalogy*, 90(3), 612–620.
- Hayward, M. W., Henschel, P., O'Brien, J., Hofmeyr, M., Balme, G., & Kerley, G. I. H. (2006). Prey preferences of the leopard (*Panthera pardus*). *Journal of Zoology*, 270(2), 298–313.
- Hayward, M. W., O'Brien, J., & Kerley, G. I. H. (2007). Carrying capacity of large African predators: Predictions and tests. *Biological Conservation*, 139(1–2), 219–229.
- Hayward, M. W., & Slotow, R. (2009). Temporal Partitioning of Activity in Large African Carnivores: Tests of Multiple Hypotheses. *South African Journal of Wildlife Research*, 39(2), 109–125.

- Hayward, M. W., Jedrzejewski, W., & Jedrzejewska, B. (2012). Prey preferences of the tiger *Panthera tigris*. *Journal of Zoology*, 286(3), 221–231.
- Heinen, J. T., & Mehta, J. N. (2000). Emerging Issues in Legal and Procedural Aspects of Buffer Zone Management with Case Studies from Nepal. *The Journal of Environment & Development*, 9(1), 45–67.
- Hoffmann, M., Hilton-Taylor, C., Angulo, A., Böhm, M., Brooks, T. M., Butchart, S. H. M., ... Stuart, S. N. (2010). The impact of conservation on the status of the world's vertebrates. *Science*, 330(6010), 1503–1509.
- Inskip, C., & Zimmermann, A. (2009). Human-felid conflict: A review of patterns and priorities worldwide. *Oryx*, 43(1), 18–34.
- Inskip, C., Fahad, Z., Tully, R., Roberts, T., & MacMillan, D. (2014). Understanding carnivore killing behaviour: Exploring the motivations for tiger killing in the Sundarbans, Bangladesh. *Biological Conservation*, 180, 42–50.
- Jacobs, J. (1974). Quantitative measurement of food selection. *Oecologia*, 14, 413–417.
- Jacobson, A. P., Gerngross, P., Lemeris Jr., J. R., Schoonover, R. F., Anco, C., Breitenmoser-Würsten, C., ... Dollar, L. (2016). Leopard (*Panthera pardus*) status, distribution, and the research efforts across its range. *PeerJ*, 4, e1974.
- Jiang, G., Wang, G., Holyoak, M., Yu, Q., Jia, X., Guan, Y., ... Ma, J. (2017). Land sharing and land sparing reveal social and ecological synergy in big cat conservation. *Biological Conservation*, 211, 142–149.
- Johnsingh, A. J. T. (1992). Prey selection in three large sympatric carnivores in Bandipur insects or lion hunting buffalo, depends on the quality and quantity of its diet time tested strategies against several ecological constraints such as prey size. *Mammalia*, 56(4), 517–527.
- Kabir, M., Ghoddousi, A., Awan, M. S., & Awan, M. N. (2014). Assessment of human-leopard conflict in Machiara National Park, Azad Jammu and Kashmir, Pakistan. *European Journal of Wildlife Research*, 60(2), 291–296.
- Kapfer, P. M., Streby, H. M., Gurung, B., Simcharoen, A., McDougal, C. C., & Smith, J. L. D. (2011). Fine-scale spatio-temporal variation in tiger *Panthera tigris* diet: effect of study duration and extent on estimates of tiger diet in Chitwan National Park, Nepal. *Wildlife Biology*, 17(3), 277–285.
- Kansky, R., & Knight, A. T. (2014). Key factors driving attitudes towards large mammals in conflict with humans. *Biological Conservation*, 179, 93–105.
- Karanth, K. K., & Nepal, S. K. (2012). Local residents perception of benefits and losses from protected areas in India and Nepal. *Environmental Management*, 49(2), 372–386.
- Karanth, K. U., & Sunquist, M. E. (1992). Population Structure, Density And Biomass Of Large Herbivores In The Tropical Forests Of Nagarhole, India. *Journal of Tropical Ecology*, 8(1), 21–35.

- Karanth, K. U., & Sunquist, M. E. (1995). Prey Selection by Tiger, Leopard and Dhole in Tropical Forests. *The Journal of Animal Ecology*, 64(4), 439.
- Karanth, K. U., & Sunquist, M. E. (2000). Behavioural correlates of predation by tiger (*Panthera tigris*), leopard (*Panthera pardus*) and dhole (*Cuon alpinus*) in Nagarahole, India. *Journal of Zoology*, 250, 255–265.
- Karanth, K. U. (2003). Tiger Ecology and Conservation in the Indian Subcontinent. *Journal of the Bombay Natural History Society*, 100(2&3), 169–189.
- Karanth, K. U., Nichols, J. D., Kumar, N. S., Link, W. A., & Hines, J. E. (2004). Tigers and their prey: Predicting carnivore densities from prey abundance. *Proceedings of the National Academy of Sciences*, 101(14), 4854–4858.
- Karanth, K. U., & Gopal, R. (2005). An ecology-based policy framework for human-tiger coexistence in India. *Conservation Biology Series, Cambridge*, 9, 381.
- Karanth, K. U., & Chellam, R. (2009). Carnivore conservation at the crossroads. *Oryx*, 43(1), 1–2.
- Karki, J., Jhala, Y., Pandav, B., Jnawali, S., Shrestha, R., Thapa, K., ... Barber-Meyer, S. M. (2016). Estimating tiger and its prey abundance in Bardia National Park, Nepal. *Banko Janakari*, 26(1), 60–69.
- Karmacharya, D., Sherchan, A. M., Dulal, S., Manandhar, P., Manandhar, S., Joshi, J., ... Hughes, J. (2018). Species, sex and geo-location identification of seized tiger (*Panthera tigris tigris*) parts in Nepal—A molecular forensic approach. *PLOS ONE*, 13(8), e0201639.
- Kawanishi, K., & Sunquist, M. E. (2004). Conservation status of tigers in a primary rainforest of Peninsular Malaysia. *Biological Conservation*, 120(3), 333–348.
- Kerley, L. L. (2010). Using dogs for tiger conservation and research. *Integrative Zoology*, 5(4), 390–396.
- King, L. E., Lawrence, A., Douglas-Hamilton, I., & Vollrath, F. (2009). Beehive fence deters crop-raiding elephants. *African Journal of Ecology*, 47(2), 131–137.
- Kolipaka, S. S., Person, G. A., De Iongh, H. H., & Srivastava, D. P. (2015). The Influence of People's Practices and Beliefs on Conservation: A Case Study on Human-Carnivore Relationships from the Multiple Use Buffer Zone of the Panna Tiger Reserve, India. *Journal of Human Ecology*, 52(3), 192–207.
- Kolipaka, S. S., Tamis, W. L. M., Van't Zelfde, M., Person, G. A., & De Iongh, H. H. (2017). Wild versus domestic prey in the diet of reintroduced tigers (*Panthera tigris*) in the livestock-dominated multiple-use forests of Panna Tiger Reserve, India. *PLoS ONE*, 12(4), 1–15.
- Lambin, E. F., & Meyfroidt, P. (2011). Global land use change, economic globalization, and the looming land scarcity. *Proc Natl Acad Sci U S A*, 108(9), 3465–3472.
- Lamichhane, B. R., Person, G. A., Leirs, H., Poudel, S., Subedi, N., Pokheral, C. P., ... de Iongh, H. H. (2018a). Spatio-temporal patterns of attacks on human and

- economic losses from wildlife in Chitwan National Park, Nepal. *PLoS ONE*, 13(4), 1–18.
- Lamichhane, B. R., Pokheral, C. P., Poudel, S., Adhikari, D., Giri, S. R., Bhattarai, S., ... Subedi, N. (2018b). Rapid recovery of tigers *Panthera tigris* in Parsa Wildlife Reserve, Nepal. *Oryx*, 52(1), 16–24.
- Lamichhane, B. R., Subedi, N., Pokheral, C. P., Dhakal, M., Acharya, K. P., Pradhan, N. M. B., ... Yackulic, C. B. (2018). Using interviews and biological sign surveys to infer seasonal use of forested and agricultural portions of a human-dominated landscape by Asian elephants in Nepal. *Ethology Ecology & Evolution*, 30(4), 331–347.
- Linkie, M., & Ridout, M. S. (2011). Assessing tiger-prey interactions in Sumatran rainforests. *Journal of Zoology*, 284(3), 224–229.
- Linnell, J. D. C., & Strand, O. (2000). Interference interactions, co-existence and conservation of mammalian carnivores. *Biodiversity Research*, 6, 169–176.
- Lovari, S., Pokheral, C. P., Jnawali, S. R., Fusani, L., & Ferretti, F. (2015). Coexistence of the tiger and the common leopard in a prey-rich area: The role of prey partitioning. *Journal of Zoology*, 295(2), 122–131.
- Loveridge, A. J., Wang, S. W., Frank, L. G., & Seidensticker, J. (2010). People and wild felids: conservation of cats and management of conflicts. *Biology and Conservation of Wild Felids*, (January), 161–195.
- Lyngdoh, S., Shrotriya, S., Goyal, S. P., Clements, H., Hayward, M. W., & Habib, B. (2014). Prey preferences of the snow leopard (*Panthera uncia*): Regional diet specificity holds global significance for conservation. *PLoS ONE*, 9(2), 1–11.
- MacArthur, R. H., & Pianka, E. R. (1966). On optimal use of a patchy Environment. *The American Naturalist*, 100(916), 603–609.
- Madden, F. (2004). Creating coexistence between humans and wildlife: Global perspectives on local efforts to address Human–Wildlife conflict. *Human Dimensions of Wildlife*, 9(4), 247–257.
- Madhusudan, M. D. (2003). Living amidst large wildlife: Livestock and crop depredation by large mammals in the interior villages of Bhadra Tiger Reserve, south India. *Environmental Management*, 31(4), 466–475.
- Manipady, S., Menezes, R. G., & Bastia, B. K. (2006). Death by attack from a wild boar. *Journal of Clinical Forensic Medicine*, 13(2), 89–91.
- Manral, U., Sengupta, S., Hussain, S. A., Rana, S., & Badola, R. (2016). Human Wildlife Conflict in India: A Review of Economic Implication of Loss and Preventive Measures. *Indian Forester*, 142(10), 928–940.
- Mazák, V. (1981). *Panthera tigris*. *Mammalian Species*, 152, 1–8.
- McDougal, C. (1988). Leopard and tiger interactions at Royal Chitwan National park, Nepal. *Journal Bombay Natural History Society*, 85(3), 609–610.

- Mehta, J. N., & Heinen, J. T. (2001). Does community-based conservation shape favorable attitudes among locals? An empirical study from Nepal. *Environmental management*, 28(2), 165-177.
- Meredith, M., & Ridout, M. (2018). Overview of the overlap package. Retrieved from <http://cran.radicaldevelop.com/web/packages/overlap/vignettes/overlap.pdf>
- Messmer, T. (2009). Human-wildlife conflicts: emerging challenges and opportunities. *Human-Wildlife Conflicts*, 3(1), 10–17.
- Miller, J. R. B., Jhala, Y. V., Jena, J., & Schmitz, O. J. (2015). Landscape-scale accessibility of livestock to tigers: Implications of spatial grain for modeling predation risk to mitigate human-carnivore conflict. *Ecology and Evolution*, 5(6), 1354–1367.
- Mills, L. S., & Allendorf, F. W. (1996). The one migrant per generation rule in conservation and management. *Conservation Biology*, 10(6), 1509–1518.
- Miththapala, S., Seidensticker, J., & O'Brien, S. J. (1996). Phylogeographic subspecies recognition in leopards (*Panthera pardus*): molecular genetic variation. *Conservation Biology*, 10(4), 1115-1132.
- Moe, S. R., & Wegge, P. (1994). Spacing behaviour and habitat use of axis deer (*Axis axis*) in lowland Nepal. *Canadian Journal of Zoology*, 72(10), 1735-1744.
- Ministry of Federal Affairs and Local Development (MoFALD) (2017). *Gaunpalika and Nagarpalika of Nepal*. Government of Nepal, Kathmandu, Nepal.
- Mondal, K., Gupta, S., Bhattacharjee, S., Qureshi, Q., & Sankar, K. (2012a). Prey selection, food habits and dietary overlap between leopard *Panthera pardus* (Mammalia: Carnivora) and re-introduced tiger *Panthera tigris* (Mammalia: Carnivora) in a semi-arid forest of Sariska Tiger Reserve, Western India. *Italian Journal of Zoology*, 79(4), 607–616.
- Mondal, K., Gupta, S., Bhattacharjee, S., Qureshi, Q., & Sankar, K. (2012b). Response of leopards to re-introduced tigers in Sariska Tiger Reserve, Western India. *International Journal of Biodiversity and Conservation*, 4(5), 228–236.
- Mondol, S., R, N., Athreya, V., Sunagar, K., Selvaraj, V. M., & Ramakrishnan, U. (2009). A panel of microsatellites to individually identify leopards and its application to leopard monitoring in human dominated landscapes. *BMC Genetics*, 10.
- Morrison, J. C., Sechrest, W., Dinerstein, E., Wilcove, D. S., & Lamoreux, J. F. (2007). Persistence of Large Mammal Faunas as Indicators of Global Human Impacts. *Journal of Mammalogy*, 88(6), 1363–1380.
- Mukherjee, S., Goyal, S. P., & Chellam, R. (1994). Standardization of scats analysis technique for leopard (*Panthera pardus fusca*) in Gir National Park, western India. *Mammalia*, 58, 139–143.
- Nepal, S. K. (2002). Linking parks and people: Nepal's experience in resolving conflicts in parks and protected areas. *International Journal of Sustainable Development and World Ecology*, 9(1), 75–90.

- Nepal, S. K., & Weber, K. E. (1995). Managing resources and resolving conflicts: National parks and local people. *International Journal of Sustainable Development and World Ecology*, 2(1), 11–25.
- Nowell, K., & Jackson, P. (1996). Status Survey and Conservation Action Plan: Wild Cats. (J. P. Kristin Nowell, Ed.), *IUCN*. Gland Switzerland: IUCN.
- Nyhus, P. J. (2016). Human–Wildlife Conflict and Coexistence. *Annual Review of Environment and Resources*, 41(1), 143–171.
- Odden, M., & Wegge, P. (2005). Spacing and activity patterns of leopards *Panthera pardus* in the Royal Bardia National Park, Nepal. *Wildlife Biology*, 11, 145–152.
- Odden, M. (2007). *Tigers, leopards and their prey in Bardia National Park, Nepal*. Department of Ecology and Natural Resource Management, Norwegian University of Life Sciences. PhD thesis.
- Odden, M., & Wegge, P. (2009). Kill rates and food consumption of leopards in Bardia National Park, Nepal. *Acta Theriologica*, 54(1), 23–30.
- Odden, M., Wegge, P., & Fredriksen, T. (2010). Do tigers displace leopards? If so, why? *Ecological Research*, 25(4), 875–881.
- Ogra, M. V. (2008). Human-wildlife conflict and gender in protected area borderlands: A case study of costs, perceptions, and vulnerabilities from Uttarakhand (Uttaranchal), India. *Geoforum*, 39(3), 1408–1422.
- Ogra, M. (2009). Attitudes toward resolution of human - Wildlife conflict among forest-dependent agriculturalists near Rajaji National Park, India. *Human Ecology*, 37(2), 161–177.
- Oli, M. K., Taylor, I. R., & Rogers, M. E. (1994). Snow leopard *Panthera uncia* predation of livestock—an assessment of local perceptions in the Annapurna Conservation area. *Nepal. Biological Conservation*, 68, 63–68.
- Oliver, J. S. (1994). Estimates of hominid and carnivore involvement in the FLK Zinjanthropus fossil assemblage: some socioecological implications. *Journal of Human Evolution*, 27(1–3), 267–294.
- Packer, C., Swanson, A., Ikanda, D., & Kushnir, H. (2011). Fear of darkness, the full moon and the nocturnal ecology of African lions. *PLoS ONE*, 6(7), 4–7.
- Palomares, F., & Caro, T. M. (1999). Interspecific Killing among Mammalian Carnivores. *The American Naturalist*, 153(5), 492–508.
- Pant, G., Dhakal, M., Pradhan, N. M. B., Leverington, F., & Hockings, M. (2016). Nature and extent of human-elephant *Elephas maximus* conflict in central Nepal. *Oryx*, 50(4), 724–731.
- Parker, G. E., & Osborn, F. V. (2006). Investigating the potential for chilli *Capsicum* spp. to reduce human-wildlife conflict in Zimbabwe. *Oryx*, 40(03), 343–346.
- Perez, I., Geffen, E., & Mokady, O. (2006). Critically Endangered Arabian leopards *Panthera pardus nimr* in Israel: Estimating population parameters using molecular scatology. *Oryx*, 40(3), 295–301.

- Pereira, H. M., Navarro, L. M., & Martins, I. S. (2012). Global biodiversity change: the bad, the good, and the unknown. *Annual Review of Environment and Resources*, 37, 25-50.
- Peterson, M. N., Birkhead, J. L., Leong, K., Peterson, M. J., & Peterson, T. R. (2010). Rearticulating the myth of human-wildlife conflict. *Conservation Letters*, 3(2), 74–82.
- Pilgrim, K. L., Mckelvey, K. S., Riddle, A. E., & Schwartz, M. K. (2005). Felid sex identification based on noninvasive genetic samples. *Molecular Ecology Notes*, 5(1), 60–61.
- Pradhan, N. M. B., & Wegge, P. (2007). Dry season habitat selection by a recolonizing population of Asian elephants *Elephas maximus* in lowland Nepal. *Acta Theriologica*, 52(2), 205–214.
- Rajaratnam, R., Sunquist, M., Rajaratnam, L., & Ambu, L. (2007). Diet and habitat selection of the leopard cat (*Prionailurus bengalensis borneoensis*) in an agricultural landscape in Sabah, Malaysian Borneo. *Journal of Tropical Ecology*, 23(2), 209–217.
- Ramakrishnan, U., Coss, R. G., & Pelkey, N. W. (1999). Tiger decline caused by the reduction of large ungulate prey: Evidence from a study of leopard diets in southern India. *Biological Conservation*, 89(2), 113–120.
- Ramesh, T., Kalle, R., Sankar, K., & Qureshi, Q. (2012). Spatio-temporal partitioning among large carnivores in relation to major prey species in Western Ghats. *Journal of Zoology*, 287(4), 269–275.
- Ramesh, T., Kalle, R., & Downs, C. T. (2017). Staying safe from top predators: patterns of co-occurrence and inter-predator interactions. *Behavioral Ecology and Sociobiology*, 71(2), 41.
- Ray, J. ., Hunter, L., & Zigouris, J. (2005). *Setting Conservation and Research Priorities for Larger African Carnivores*. Wildlife Conversation Society, New York. No. 24 (Vol. 267).
- Rayan, D. M., & Linkie, M. (2016). Managing conservation flagship species in competition: Tiger, leopard and dhole in Malaysia. *Biological Conservation*, 204, 360–366.
- R Core Team. (2018). R: The R Project for Statistical Computing. Retrieved June 21, 2018, from <https://www.r-project.org/>
- Redpath, S. M., Bhatia, S., & Young, J. (2015). Tilting at wildlife: Reconsidering human-wildlife conflict. *Oryx*, 49(2), 222–225.
- Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, 14(3), 322–337.

- Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M., ... Wirsing, A. J. (2014). Status and Ecological Effects of the World's Largest Carnivores. *Science*, 343(6167), 1241484-1241484.
- Romañach, S. S., Lindsey, P. A., & Woodroffe, R. (2007). Determinants of attitudes towards predators in central Kenya and suggestions for increasing tolerance in livestock dominated landscapes. *Oryx*, 41(2), 185-195.
- Røskaft, E., Händel, B., Bjerke, T., & Kaltenborn, B. P. (2007). Human attitudes towards large carnivores in Norway. *Wildlife Biology*, 13(2), 172-185.
- Saberwal, V. K., Gibbs, J. P., Chellam, R., & Johnsingh, A. J. T. (1994). Lion-Human Conflict in the Gir Forest, India. *Conservation Biology*, 8(2), 501-507.
- Sangay, T., & Vernes, K. (2008). Human-wildlife conflict in the Kingdom of Bhutan: Patterns of livestock predation by large mammalian carnivores. *Biological Conservation*, 141(5), 1272-1282.
- Sankar, K., Johnsingh, A. J. T. (2002). Food habits of tiger (*Panthera tigris*) and leopard (*Panthera pardus*) in Sariska Tiger Reserve, Rajasthan, India, as shown by scat analysis. *Mammalia*, 66(2), 285-289.
- Sankar, K., Qureshi, Q., Nigam, P., Malik, P. K., Sinha, P. R., Mehrotra, R. N., ... Gupta, S. (2010). Monitoring of Reintroduced Tigers in Sariska Tiger Reserve, Western India: Preliminary Findings on Home Range, Prey Selection and Food Habits. *Tropical Conservation Science*, 3(3), 301-318.
- Sapkota, S., Aryal, A., Baral, S. R., Hayward, M. W., & Raubenheimer, D. (2014). Economic Analysis of Electric Fencing for Mitigating Human-wildlife Conflict in Nepal. *Journal of Resources and Ecology*, 55(533), 237-243.
- Sarker, A. H. M. R., & Røskaft, E. (2011). Human attitudes towards the conservation of protected areas: a case study from four protected areas in Bangladesh. *Oryx*, 45(03), 391-400.
- Seidensticker, J. (1976). On the Ecological Separation between Tigers and Leopards. *Biotropica*, 8(4), 225-234.
- Seidensticker, J., Sunquist, M.E., McDougal, C. (1990). *Leopards living at the edge of the Royal Chitwan National Park Nepal*. Conservation in Developing Countries: Problems and Prospects, Bombay Oxford University Press, 415-423.
- Seidensticker, J., & McDougal, C. (1993). Tiger predatory behaviour, ecology and conservation. *Symposium of the Zoological Society of London*, (65), 105-125.
- Seidensticker, J. (2010). Saving wild tigers: A case study in biodiversity loss and challenges to be met for recovery beyond 2010. *Integrative Zoology*, 5(4), 285-299.
- Seidensticker, J., Dinerstein, E., Goyal, S. P., Harihar, A., Johnsingh, A. J. T., Manandhar, A., ... Wikramanayake, E. (2015). Tiger range collapse at the base of the Himalayas : a case study Tiger range collapse and recovery at the base of the Himalayas. *The Biology and Conservation of Wild Felids*, (September), 784.

- Shehzad, W., Nawaz, M. A., Pompanon, F., Coissac, E., Riaz, T., Shah, S. A., & Taberlet, P. (2015). Forest without prey: livestock sustain a leopard *Panthera pardus* population in Pakistan. *Oryx*, *49*(2), 248–253.
- Shrestha M. K. (2004). *Relative ungulate abundance in a fragmented landscape: Implications for tiger conservation*. University of Minnesota, PhD thesis.
- Silwal, T., Kolejka, J., Bhatta, B. P., Rayamajhi, S., Sharma, R. P., & Poudel, B. S. (2017). When, where and whom: Assessing wildlife attacks on people in Chitwan National Park, Nepal. *Oryx*, *51*(2), 370–377.
- Simcharoen, S., Barlow, A. C. D., Simcharoen, A., & Smith, J. L. D. (2008). Home range size and daytime habitat selection of leopards in Huai Kha Khaeng Wildlife Sanctuary, Thailand. *Biological Conservation*, *141*(9), 2242–2250.
- Simcharoen, A., Savini, T., Gale, G. A., Simcharoen, S., Duangchantrasiri, S., Pakpien, S., & Smith, J. L. D. (2014). Female tiger *Panthera tigris* home range size and prey abundance: Important metrics for management. *Oryx*, *48*(3), 370–377.
- Smith, J. L. D., Sunquist, M. E., Tamang, K. M., & Rai, P. B. (1983). A Technique for Capturing and Immobilizing Tigers. *The Journal of Wildlife Management*, *47*(1), 255–259.
- Smith, J. L. D., McDougal, C., & Miquelle, D. G. (1989). Scent marking in free-ranging tigers, *Panthera tigris*. *Animal Behaviour*, *37*(1), 11–27.
- Smith, J. L. D., & McDougal, C. (1991). The Contribution of Variance in Lifetime Reproduction to Effective Population Size in Tigers. *Conservation Biology*, *5*(4), 484–490.
- Smith, J. L. D. (1993). The role of dispersal in structuring the Chitwan tiger population. *Behaviour*, *124*(3), 165–195.
- Smith, J. L. D., Ahearn, S. C., & McDougal, C. (1998). Landscape Analysis of Tiger Distribution and Habitat Quality in Nepal. *Conservation Biology*, *12*(6), 1338–1346.
- Sogbohossou, E. A., De longh, H. H., Sinsin, B., De Snoo, G. R., & Funston, P. J. (2011). Human-carnivore conflict around pendjari biosphere reserve, northern benin. *Oryx*, *45*(4), 569–578.
- Steinmetz, R., Seuaturien, N., & Chutipong, W. (2013). Tigers, leopards, and dholes in a half-empty forest: Assessing species interactions in a guild of threatened carnivores. *Biological Conservation*, *163*, 68–78.
- Støen, O. G., & Wegge, P. (1996). Prey selection and prey removal by tiger (*Panthera tigris*) during the dry season in lowland Nepal. *Mammalia*, *60*(3), 363–374.
- Studsrod, J. E., & Wegge, P. (1995). Park-People Relationships: The Case of Damage Caused by Park Animals Around the Royal Bardia National Park, Nepal. *Environmental Conservation*, *22*(2), 133–142.
- Sukumar, R. (1991). The management of large mammals in relation to male strategies and conflict with people. *Biological Conservation*, *55*(1), 93–102.

- Sunarto, S., Kelly, M. J., Parakkasi, K., & Hutajulu, M. B. (2015). Cat coexistence in central Sumatra: Ecological characteristics, spatial and temporal overlap, and implications for management. *Journal of Zoology*, 296(2), 104–115.
- Sunquist, M. E. (1981). The Social Organization of Tigers (*Panthera Tigris*) in Royal Chitawan National Park, Nepal. In *Smithsonian Contributions to Zoology* 1–98.
- Sunquist, M., & Sunquist, F. (2002). *Wild Cats of the World*. University of Chicago Press.
- Swanson, A., Arnold, T., Kosmala, M., Forester, J., & Packer, C. (2016). In the absence of a “landscape of fear”: How lions, hyenas, and cheetahs coexist. *Ecology and Evolution*, 6(23), 8534–8545.
- Tamang, B., & Baral, N. (2008). Livestock depredation by large cats in Bardia National Park, Nepal: Implications for improving park-people relations. *International Journal of Biodiversity Science and Management*, 4(1), 44–53.
- Thapa, S., & Chapman, D. S. (2010). Impacts of resource extraction on forest structure and diversity in Bardia National Park, Nepal. *Forest Ecology and Management*, 259(3), 641–649.
- Thapa, S. (2010). Effectiveness of crop protection methods against wildlife damage: A case study of two villages at Bardia National Park, Nepal. *Crop Protection*, 29(11), 1297–1304.
- Thapa, K., Wikramanayake, E., Malla, S., Acharya, K. P., Lamichhane, B. R., Subedi, N., ... Vattakaven, J. (2017). Tigers in the terai: Strong evidence for meta-population dynamics contributing to tiger recovery and conservation in the terai arc landscape. *PLoS ONE*, 12(6), 1–15.
- Thinley, P., Rajaratnam, R., Lassoie, J. P., Morreale, S. J., Curtis, P. D., Vernes, K., ... Dorji, P. (2018). The ecological benefit of tigers (*Panthera tigris*) to farmers in reducing crop and livestock losses in the eastern Himalayas: Implications for conservation of large apex predators. *Biological Conservation*, 219(February), 119–125.
- Trall, L. W., Martin, J., & Owen-Smith, N. (2016). Lion proximity, not moon phase, affects the nocturnal movement behaviour of zebra and wildebeest. *Journal of Zoology*, 299(3), 221–227.
- Treves, A., & Naughton-Treves, L. (1999). Risk and opportunity for humans coexisting with large carnivores. *Journal of Human Evolution* 36(3),275–282.
- Treves, A., & Karanth, K. U. (2003). Human-Carnivore Conflict and Perspectives on Carnivore Management Worldwide. *Conservation Biology*, 17(6), 1491–1499.
- Treves, A., Wallace, R. B., Naughton-Treves, L., & Morales, A. (2006). Co-managing human–wildlife conflicts: A review. *Human Dimensions of Wildlife*, 11(6), 383–396.
- Tumenta, P. N. (2012). *A Lion Population under Threat*. Leiden University , PhD thesis.

- Tumenta, P. N., de Iongh, H. H., Funston, P. J., & Udo de Haes, H. A. (2013). Livestock depredation and mitigation methods practised by resident and nomadic pastoralists around Waza National Park, Cameroon. *Oryx*, 47(02), 237–242.
- Turner, A. (1987). *Megantereon cultridens* (Cuvier) (Mammalia, Felidae, Machairodontinae) from Plio-Pleistocene deposits in Africa and Eurasia, with comments on dispersal and the possibility of a New World origin. *Journal of Paleontology*, 61(6), 1256–1268.
- Upadhyaya, S. K., Musters, C. J. M., Lamichhane, B. R., de Snoo, G. R., Thapa, P., Dhakal, M., Karmacharya, D., Shrestha, P. M., & de Iongh, H. H. (2018). An Insight Into the Diet and Prey Preference of Tigers in Bardia National Park, Nepal. *Tropical Conservation Science*, 11, 194008291879947.
- Upadhyaya, S. K., Musters, C. J. M., Lamichhane, B. R., de Snoo, G. R., Dhakal, M., & de Iongh, H. H. (2018). Defining the risks of attacks by predators around protected areas – the case of Bardia National Park, Nepal. *Oryx* (Accepted).
- Uphyrkina, O., Johnson, W. E., Quigley, H., Miquelle, D., Marker, L., Bush, M., & O'Brien, S. J. (2001). Phylogenetics, genome diversity and origin of modern leopard, *Panthera pardus*. *Molecular ecology*, 10(11), 2617–2633.
- Van Bommel, L., Bij De Vaate, M. D., De Boer, W. E., & De Iongh, H. H. (2007). Factors affecting livestock predation by lions in Cameroon. *African Journal of Ecology*, 45(4), 490–498.
- Walston, J., Robinson, J. G., Bennett, E. L., Breitenmoser, U., da Fonseca, G. A. B., Goodrich, J., ... Wibisono, H. (2010). Bringing the tiger back from the brink—the six percent solution. *PLoS Biology*, 8(9), 6–9.
- Wang, S. W., & Macdonald, D. W. (2006). Livestock predation by carnivores in Jigme Singye Wangchuck National Park, Bhutan. *Biological Conservation*, 129(4), 558–565.
- Wang, S. W., & Macdonald, D. W. (2009). The use of camera traps for estimating tiger and leopard populations in the high altitude mountains of Bhutan. *Biological Conservation*, 142(3), 606–613.
- Wang, S. W., & Macdonald, D. W. (2009). Feeding habits and niche partitioning in a predator guild composed of tigers, leopards and dholes in a temperate ecosystem in central Bhutan. *Journal of Zoology*, 277(4), 275–283.
- Wasser, S. K., Smith, H., Madden, L., Marks, N., & Vynne, C. (2009). Scent-Matching Dogs Determine Number of Unique Individuals From Scat. *Journal of Wildlife Management*, 73(7), 1233–1240.
- Wegge, P., Odden, M., Pokharel, C. P., & Storaas, T. (2009). Predator-prey relationships and responses of ungulates and their predators to the establishment of protected areas: A case study of tigers, leopards and their prey in Bardia National Park, Nepal. *Biological Conservation*, 142(1), 189–202.

- Wegge, P., & Storaas, T. (2009). Sampling tiger ungulate prey by the distance method: Lessons learned in Bardia National Park, Nepal. *Animal Conservation*, 12(1), 78–84.
- Wegge, P., Yadav, S. K., & Lamichhane, B. R. (2018). Are corridors good for tigers *Panthera tigris* but bad for people? An assessment of the Khata corridor in lowland Nepal. *Oryx*, 52(01), 35–45.
- Wikramanayake, E. D., Dinerstein, E., Robinson, J. G., Karanth, U., Rabinowitz, A., Olson, D., ... Bolze, D. (1998). An Ecology-Based Method for Defining Priorities for Large Mammal Conservation: The Tiger as Case Study. *Conservation Biology*, 12(4), 865–878.
- Wikramanayake, E., McKnight, M., Dinerstein, E., Joshi, A., Gurung, B., & Smith, D. (2004). Designing a conservation landscape for tigers in human-dominated environments. *Conservation Biology*, 18(3), 839–844.
- Wilson, S., Davies, T. E., Hazarika, N., & Zimmermann, A. (2015). Understanding spatial and temporal patterns of human-elephant conflict in Assam, India. *Oryx*, 49(1), 140–149.
- Wilting, A., Courtiol, A., Christiansen, P., Niedballa, J., Scharf, A. K., Orlando, L., ... Kitchener, A. C. (2015). Planning tiger recovery: Understanding intraspecific variation for effective conservation. *Science Advances*, 1(5) e1400175.
- Woodroffe, R., & Ginsberg, J. R. (1998). Edge effects and the extinction of populations inside protected areas. *Science*, 280, 2126–2128.
- Woodroffe, R., Lindsey, P., Romañach, S., Stein, A., & Ole Ranah, S. M. K. (2005). Livestock predation by endangered African wild dogs (*Lycaon pictus*) in northern Kenya. *Biological Conservation*, 124(2), 225–234.
- Woodroffe, R., Frank, L. G., Lindsey, P. A., Ole Ranah, S. M. K., & Romañach, S. (2007). Livestock husbandry as a tool for carnivore conservation in Africa's community rangelands: A case-control study. *Biodiversity and Conservation*, 16(4), 1245–1260.
- Wulsch, C., Waits, L. P., & Kelly, M. J. (2014). Noninvasive individual and species identification of jaguars (*Panthera onca*), pumas (*Puma concolor*) and ocelots (*Leopardus pardalis*) in Belize, Central America using cross-species microsatellites and faecal DNA. *Molecular Ecology Resources*, 14(6), 1171–1182.

Summary

Large carnivores and humans, along with their livestock, have co-existed for thousands of years. However, human population growth and an increase in economic activities are modifying the landscape for large carnivores and their prey. Large cats, which are regarded as keystone species for maintaining a healthy ecosystem through the balance between predator and prey, are now often locally on the verge of extinction. Large predators are sensitive to human activities and have been actively persecuted from all over the world, because of their conflicts with local people. The number of tigers (*Panthera tigris*) has declined rapidly in the past century and now they are confined to 7 % of their historical range. Similarly the range of leopards (*Panthera pardus*) has also declined significantly, although they are more widespread and adaptive than tigers. Since these carnivores require large habitats, degradation and fragmentation of their habitat over past decades are regarded as the major threats for their survival. The other factors responsible for their decline include poaching for illegal trade of body parts, loss of prey species and conflicts with local communities. Local farmers often face negative interactions with wild predators thereby experiencing economic loss by the killing of livestock. These conflicts in combination with a poor understanding of the ecological role which predators play may lead to local resentment against predators and negative attitude towards predator conservation projects.

Nepal is one of the 13 countries to ratify the Global Tiger Recovery Plan at the meeting of world leaders held in St. Petersburg in 2010, aims at doubling the number of wild tigers by 2022. In order to achieve this goal the Government of Nepal has focused on better conservation practices in and around tiger conservation landscape. Tigers are found in five National Parks of the lowland plain (Terai). This area is a part of the greater Terai Arc Landscape which is located in India and Nepal. There has been an increase in the tiger population after the end of civil war in Nepal, as well as a decrease in poaching, due to new government policy which included strict protection measures in national parks and the involvement of youths in anti-poaching campaign. As a result of a significant increase in the population of tigers inside Bardia National Park they are now showing more interaction with leopards. As tigers and leopards are elusive species it is difficult to study their interaction in the field. Therefore, using camera trap data obtained during 2013

and 2016, I studied the interaction between tigers and leopards in Bardia National Park. I found that there is a significant level of spatial avoidance of leopards towards tigers in the core zone of the park whereas in the boundary zone the spatial avoidance is not significant. I found that the temporal activities of leopards in grids of the core zone where tigers and leopards both are present are significantly different compared to grids where tigers are absent. The temporal activities of tiger are not significantly different in grids with leopards present in the core zone. However, the temporal activity of tigers is significantly different in the boundary zone when leopards were present, whereas the activity of leopard does not change. This can probably be explained by the suggestion that leopards approach human settlements more easily, while tigers show strong avoidance of human settlements. This also suggests that habitat partitioning and space use may play an important role in the co-existence of tigers and leopards in Bardia National Park. In general leopards show more avoidance to tigers than the other way round.

I also compared the diet and prey preference of tigers and leopards using their scats. However, due to lower number of leopard scats, I could not process it further and had to limit my study to the diet and prey preference of tigers only. I was able to identify male and female tigers based on DNA analysis of scat. I found that both male and female tigers were relying on medium sized prey (10 kg to 40 kg) namely chital (*Axis axis*) and wild pig (*Sus scrofa*) which was the most abundant prey in Bardia. However the Jacobs index for prey preference of male tiger was for sambar (*Cervus unicolor*) and wild pig whereas for female tiger the preference was for chital and wild pig, however the results were not statistically significant. The diet of tigers showed mostly the presence of wild prey species which suggests that they are generally not involved in the killing of livestock. For the long term conservation of the tiger in Bardia National Park we need to focus on increasing the population of larger prey species in the park, so that the available prey species are energetically beneficial for tigers and there is less competition with leopards.

I also looked into the spatiotemporal pattern of the conflict incidence in Bardia NP for a period of five years, on the basis of compensation paid to the villagers for the damage caused by wildlife. I found that four animals were mainly involved in causing damage to local communities, namely elephant (*Elephas maximus*), leopard, tiger and wild boar. The elephant was responsible for most of the damage. During the five years elephants killed 14 people and wild boar killed one person. Elephants also caused serious damage to the crops, destroyed houses and raided store houses for grains. Leopards and

tigers particularly caused depredation of livestock. Elephants caused more damage during the autumn season when the crops (mainly paddy) had matured. Leopards killed more livestock during the rainy season compared to dry season, while tigers did not showed seasonal pattern of predation. Tigers and leopards were involved in significantly more killing of livestock during the new moon phase compared to the full moon phase. The conflict incidence for elephants, wild boar and leopards was higher in the southern and the western part of the buffer zone, whereas tigers had a similar level of conflict all over the park.

To know more about the attitude and knowledge of local people living in the buffer zone towards conservation, I conducted a questionnaire survey by dividing the park into four different regions. I found that the loss of livestock was significantly related to number of livestock owned, the distance to the park boundary and the ethnicity of the respondent. The attitude of the respondent towards wildlife conservation was dependent on i) knowledge about the species, ii) (higher) educational level, iii) gender (male/female) and iv) level of self-sufficiency of the respondents, with a positive attitude related to more knowledge, male gender, higher education and higher self-sufficiency level of respondents. The sharing of conservation benefits also resulted in a positive attitude among the people living in the buffer zone.

In conclusion, it can be said that, due to the increase in number of tigers inside the park leopards may be pushed out of the park, where they become involved in conflicts while there may also be a spill-over of young tigers moving to the buffer zone. For future conservation of these large cats, additional habitats need to be sought and the community forest of the buffer zone can provide such habitat. Similarly, Banke National Park which has been established to provide a sink for the Bardia source population, which can help in accommodating the dispersed population of Bardia NP. Therefore, for future conservation programs we need to focus on the conservation of tiger and leopard in human dominated landscape. Due to interaction with tigers, the leopards are living in close proximity with people and thereby get more involved in conflicts with local communities. Tigers enjoy a religious and cultural tolerance among the people living in the buffer zone whereas leopards lack such religious and cultural tolerance.



Samenvatting

Grote carnivoren en mensen, samen met hun vee, leven al duizenden jaren samen in dezelfde gebieden. Bevolkingsgroei en economische ontwikkeling veranderen het landschap van deze gebieden voor carnivoren en hun prooi dieren. Grote katten, die worden beschouwd als sleutel soorten voor gezonde ecosystemen door de wisselwerking predator-prooi, staan nu vaak lokaal op het punt om uit te sterven. Grote carnivoren zijn kwetsbaar voor menselijke activiteiten en worden wereldwijd actief vervolgd, vanwege conflicten met mensen. Het aantal tijgers (*Panthera tigris*) is de afgelopen eeuw sterk afgenomen en tijgers bevinden zich nu slechts in 7% van hun historische leefgebied. Ook het leefgebied van de luipaard (*Panthera pardus*) is sterk afgenomen, alhoewel deze soort meer algemeen en adaptief is dan de tijger. Omdat carnivoren grote leefgebieden nodig hebben, wordt de degradatie van hun habitat over de laatste tientallen jaren gezien als de belangrijkste bedreiging. Andere factoren die van invloed zijn op de afname van carnivoren betreffen stroperij voor de illegale handel in lichaamsdelen van carnivoren, de afname van prooidier populaties en conflicten met lokale gemeenschappen. Lokale boeren ondervinden vaak de negatieve gevolgen van wilde carnivoren, waarbij ze economische schade ondervinden door het doden van vee. Deze conflicten worden verergerd door een beperkt begrip van de ecologische rol van carnivoren en dit kan resulteren in een negatieve houding ten opzichte van carnivoren en gerelateerde projecten voor natuurbescherming.

Nepal is een van de dertien landen die het Global Tiger Recovery Plan hebben getekend op de bijeenkomst van wereldleiders in St Petersburg in 2010, die als doel heeft het aantal wilde tijgers te verdubbelen in 2022. Om dit doel te behalen heeft de regering van Nepal haar aandacht gegeven op verbeterde activiteiten gericht op natuurbescherming in en rond de landschappen waar tijgers voorkomen. Tijgers worden in Nepal nog aangetroffen in vijf nationale parken. In de laagvlakte (Terai). Dit gebied is een onderdeel van het grotere 'Terai Arc' landschap dat zich bevindt in India en Nepal.. De tijger populatie in Nepal is toegenomen sinds het einde van de burgeroorlog, terwijl de stroperij is afgenomen. Dit is een gevolg van het overheidsbeleid, dat strikte natuurbeschermings-activiteiten in nationale parken omvatten e o.m. de deelname van jongeren in anti-stroperij campagnes. Dit heeft geresulteerd in een sterke groei van de tijger populatie in Bardia nationaal park, waar tijgers nu meer interac-

ties hebben met luipaarden dan vroeger. Omdat tijgers en luipaarden schuwe dieren zijn is het moeilijk hun interactie in het veld te onderzoeken. Camera val surveys zijn een goede methode om de aanwezigheid en afwezigheid van tijgers en luipaarden te onderzoeken. Daarom heb ik met behulp van camera val data uit 2013 en 2016 de interacties tussen tijgers en luipaarden bestudeerd in Bardia nationaal park

Ik vond daarbij een significante ruimtelijke ontwijking van luipaarden t.o.v. tijgers in de centrale zone van het park, terwijl in de bufferzone en geen significant patroon van ontwijking was. Ik vond ook dat de activiteiten in de tijd van luipaarden in grids in de centrale zone, waar zowel tijgers als luipaarden aanwezig waren, significant verschilden met grids waar tijgers niet aanwezig waren, maar luipaarden wel. De activiteiten in de tijd van tijgers in grids waar luipaarden ook aanwezig waren in de centrale zone verschillen niet van grids zonder luipaarden. Ik vond echter in de bufferzone een significant verschil wat betreft de activiteiten van tijgers in de tijd in grids met luipaarden vergeleken met grids zonder luipaarden, terwijl de activiteiten van luipaarden in de tijd niet verschilden in grids met en zonder tijgers. Dit kan waarschijnlijk worden verklaard door de suggestie dat luipaarden dichterbij menselijke nederzettingen komen, terwijl tijgers menselijke nederzettingen vermijden. Dit geeft ook aan dat het gemeenschappelijk gebruik van habitat en ruimte een belangrijke rol speelt in het samenleven van tijgers en luipaarden in Bardia nationaal park. In het algemeen ontwijken luipaarden meer de tijgers dan andersom. Ik heb ook de diëten en prooidier voorkeuren van tijgers en luipaarden willen vergelijken op basis van uitwerpselen. Echter, vanwege een onvoldoende grote monster-grootte van luipaard uitwerpselen, kon ik deze niet verder gebruiken en heb ik mij beperkt tot het analyseren van de uitwerpselen en prooidier voorkeuren van tijgers. Gebaseerd op het DNA in de uitwerpselen kon ik een onderscheid maken tussen vrouwelijke en mannelijke tijgers. Ik vond dat zowel mannelijke als vrouwelijke tijgers vooral prooidieren eten van medium gewicht (10 tot 40 kg), namelijk Chital (*Axis axis*) en wild zwijn (*Sus Scrofa*), die de meest algemene prooidieren zijn in Bardia nationaal park. Echter de Jacobs index voor prooidier preferentie van mannelijke tijgers gaf een voorkeur aan voor Sambar hert (*Cervus unicolor*) en wild zwijn, terwijl de Jacobs index voor vrouwelijke tijgers een voorkeur aangaf voor Chital en wild zwijn.

Deze resultaten waren echter statistisch gezien niet significant. Het dieet van tijgers bestond voornamelijk uit wilde prooidier soorten, hetgeen suggereert dat tijgers over het algemeen geen vee consumeren. Voor de lange termijn bescherming van de tijger in Bardia nationaal park moet meer aandacht gegeven

worden aan de groei van prooidier populaties in het park, zodat de beschikbare prooidier populaties energetische voordelen bieden voor tijgers en er minder competitie is met luipaarden. Ik heb ook gekeken naar het verloop van conflicten in Bardia nationaal park in ruimte en tijd, over een periode van vijf jaar, op basis van de statistieken van compensatie die is betaald aan dorpelingen voor de schade veroorzaakt door wilde dieren.

Ik vond dat voornamelijk vier soorten dieren betrokken waren bij schade gevallen gemeld door de lokale bevolking; olifant, luipaard, tijger en wild zwijn. Gedurende die vijf jaar werden 14 personen gedood door olifanten en één persoon door een wild zwijn. Olifanten veroorzaakten ook ernstige schade aan gewassen, vernielden huizen en vielen opslag schuren voor graan aan. Luipaarden en tijgers veroorzaakten vooral schade aan vee. Olifanten veroorzaakten meer schade gedurende de herfst wanneer gewassen (vooral rijst) aan het rijpen waren. Luipaarden veroorzaakten meer schade aan vee gedurende het regenseizoen in vergelijking met het droge seizoen, terwijl de schade toegebracht door tijgers geen verschillende vertoonden tussen de seizoenen.. Tijgers en luipaarden veroorzaakten meer schade aan vee gedurende de donkere maan periode vergeleken met de volle maan periode. De frequentie van conflicten met luipaarden was hoger in het zuidelijke en westelijke deel van de bufferzone voor olifanten en wilde zwijnen, terwijl de frequentie van conflicten met tijgers gelijk was over het hele gebied. Om meer te weten te komen over de attitudes en kennis van lokale bewoners die in de bufferzone leven, van belang voor natuurbescherming, heb ik interviews gehouden. Dit heb ik gedaan door het park te verdelen in vier verschillende regio's. Ik vond dat de schade aan vee een significante relatie had met i) de aantallen vee in bezit van bewoners, ii) de afstand tot de park grens en ii) de etnische afkomst van de respondent. De attitude van de respondenten ten opzichte van de bescherming van wilde dieren was gerelateerd aan de volgende factoren; i) kennis over de soort ii) opleidings nivo respondent iii) gender van de respondent (man/vrouw) en iv) nivo van onafhankelijkheid van de respondent. Een positieve attitude was gerelateerd aan meer kennis over de soort, een hoog opleidings nivo, en een hoog nivo van onafhankelijkheid. Het delen van voordelen en inkomsten uit natuurbescherming resulteert ook in een positieve attitude onder de bevolking in de bufferzone. Concluderend kan gezegd worden dat als gevolg van een toename van het aantal tijgers in het park luipaarden waarschijnlijk uitwijken naar de bufferzone, waar ze in conflict komen met bewoners, terwijl ook jonge tijgers het park verlaten. Voor de toekomstige bescherming van deze grote katten is het van belang dat nieuwe habitats worden gezocht en de gemeenschapsbossen in de bufferzone zouden hier aan bij kunnen dragen. Ook kan het nabijgelegen

Banke nationaal park een *sink* vormen voor de bron-populatie in Bardia nationaal park. Dit kan de verdere verspreiding van de tijger populatie in Bardia faciliteren. Voor toekomstige natuurbeschermings programma's is het van belang zich te concentreren op de bescherming van tijger en luipaard in een door mensen gedomineerd landschap. Vanwege de interactie met tijgers leven luipaarden dichterbij mensen en komen ze zo ook vaker in conflict met lokale bewoners. Tijgers genieten een culturele en religieuze tolerantie onder de bevolking die leeft in de bufferzone terwijl luipaarden deze tolerantie niet genieten.

Acknowledgements

I would like to dedicate this work to my parents Puskar Nath Upadhyaya Gyawali and Tamkala Gyawali, who constantly inspired me during my PhD. I also want to thank my wife Hemlata, daughters Simran and Subhani and son Somin for their constant encouragement for my work and to bear with me during the period of my absence from home. I would also like to thank my brother Shyam and sisters Bina, Sarita and Parvata as well as other family members for their support.

I would like to thank my promotor Professor Geert de Snoo and co-promoters Professor Hans H de Iongh and Dr. Kees Musters from Leiden University for their constant supervision, guidance and encouragement. I also would like to thank my co-supervisor at Kathmandu University, Prof. Dr. Panna Thapa for his constant guidance and encouragement. My sincere gratitude go to Prof. Dr. Ram Kantha Makaju Shrestha (Vice Chancellor), Prof. Dr. Subodh Sharma (Registrar), Prof. Dr. Bhola Thapa (Former Registrar) and Prof. Dr. Kanhaiya Jha (Dean of School of Science) of Kathmandu University. I would also like to thank colleagues from my department at Kathmandu University for their support. I would also like to thank the Himalayan Tiger Foundation, especially Cas de Stoppelaar and his team for their support. I would like to thank Dr. Dibesh Karmacharya and his team from the Center for Molecular Dynamics Nepal for helping in carrying out the laboratory work for DNA analysis. My sincere thanks also go to Ms Patricia Chettri, (fellowship officer) of The Netherlands Consulate, Kathmandu for her help in visa processing. I want to thank Nuffic for providing funding for this project through a NFP fellowship. I would also like to thank Cindy Scotte, Marielle van Es and other employees of the department of Student and Educational Affairs (SEA) who helped me with the allotment of funds.

I would also like to thank the Director, and former and present ecologist of the Department of National Parks and Wildlife Conservation (DNPWC), namely Dr. Maheshwar Dhakal, Laxman Poudyal, Rishi Ranabhat and other staff of DNPWC for providing the permit for my research work. I also want to thank former and present senior warden of Bardia National Park, namely Ram Chandra Kandel, Ramesh Thapa, Ashok Bhandari and other staffs for their help. I would like to extend my sincere thanks to Mrs. Sarita Jnawali,

Dr. Naresh Subedi and other members of National Trust for Nature Conservation (NTNC) for their support and encouragement. Thanks also go to Rabin Kadariya, Ambika Prasad Khatiwada, Shyam Thapa, Shreeram Ghimire, Shailendra Yadav, Ramji Thapa and other staff of the Bardia Conservation Program of NTNC for providing technical and logistic support during my field work.

I want to thank from my heart Babu Ram Lamichhane for his help and support during the course of my PhD. I also acknowledge the support of friends in Leiden who made my stay comfortable namely Praduman Khayargoli, Prof. Jan Houben, Dr. Saraju Rath, Dr. Bal Gopal Shrestha, Sri Laxmi Shrestha, Dipen Lama, Ranjita Lama and Rogier van de Vorst.

I also want to thank the Scientific Director of the Institute of Environmental Sciences Leiden (CML) Prof. Dr. Arnold Tukker and Head of Environmental Biology department Prof. Dr. Peter van Bodegom. I also want to thank the Institute Manager Paul de Hoog as well as secretaries namely Jory Sjardijn and Susanna van den Oever for their help provided to me during my stay in CML. I am indebted to all staff of CML and my fellow PhD for making my stay at Leiden memorable.

I am grateful to my PhD committee members, Prof. Dr. Arnold Tukker, Prof. Dr. Peter van Bodegom, Prof. Dr. Herwig Leirs and Prof. Dr. Gerard Persoon for evaluation and providing constructive comments and suggestions on an earlier version of the thesis draft. I want to thank Professor Hans de Iongh for the Dutch translation of the summary. I would like to express my gratitude to my friend Sijan Baral for initial correction of the thesis and to Barbara Croes for the final language corrections. I also want to thank Sjoukje Rienks for the layout work. Finally, I am grateful to Sagar Giri for providing the cover photographs and photographs at the beginning of chapter two, four and six.

Curriculum Vitae



Subodh Kumar Upadhyaya was born on the 8th of October 1965 in Khanchikot, Arghakhanchi district of Nepal. He followed his primary and secondary school from 1972 to 1984 in Kendriya Vidyalaya, Shillong, India and attended there the St. Edmund's College affiliated to North Eastern hill University in Shillong, India, where he obtained a Bachelor's degree with Zoology Honours in 1987. He obtained his first Masters degree in Zoology from Tribhuvan University, Kathmandu, Nepal in 1989. and a second Masters degree in Biotechnology from Birla Institute of Technology and Science, Pilani,

India in 2004. Since 1998 he is working as a staff member in the Department of Biotechnology, School of Science, Kathmandu University, Nepal.

Subodh started his PhD at the Institute of Environmental Sciences Leiden (CML) in 2014, when he received a fellowship of the Netherlands Organization for International Cooperation in Higher Education, the Netherlands fellowship program (Nuffic-NFP). His PhD research was on the interaction between tigers and leopards in relation to human-wildlife conflict in Bardia National Park, Nepal. This PhD is part of a collaborative agreement between Leiden University, the Netherlands and Kathmandu University, Nepal and between Leiden University and the National Trust for Nature Conservation Nepal (NTNC).

