



Universiteit
Leiden
The Netherlands

Visualizing strongly-correlated electrons with a novel scanning tunneling microscope

Battisti, I.

Citation

Battisti, I. (2019, May 8). *Visualizing strongly-correlated electrons with a novel scanning tunneling microscope*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/72410>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/72410>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/72410> holds various files of this Leiden University dissertation.

Author: Battisti, I.

Title: Visualizing strongly-correlated electrons with a novel scanning tunneling microscope

Issue Date: 2019-05-08

VISUALIZING STRONGLY-CORRELATED ELECTRONS WITH A NOVEL SCANNING TUNNELING MICROSCOPE

PROEFSCHRIFT

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 8 mei 2019
klokke 11:15 uur

door

IRENE BATTISTI

geboren te Padova, Italië
in 1990

Promotor:	Prof. dr. J. Aarts	Universiteit Leiden
Co-promotor:	Dr. M.P. Allan	Universiteit Leiden
Promotiecommissie:	Prof. dr. P. Wahl	University of St. Andrews
	Prof. dr. C. Morais-Smith	Universiteit Utrecht
	Prof. dr. K. Schalm	Universiteit Leiden
	Prof. dr. E.R. Eliel	Universiteit Leiden

Casimir PhD series, Delft-Leiden 2019-09.
ISBN 978-90-8593-391-5

An electronic version of this thesis can be found at:
<https://openaccess.leidenuniv.nl>

This work is supported by the Netherlands Organization for Scientific Research (NWO/OCW) as part of the Frontiers of Nanoscience (NanoFront) program and the VIDI talent scheme (Project No. 680-47-536).

The cover shows a (Fourier-filtered) atomically-resolved STM topographic image measured by the author on Sr_2RhO_4 . Few atomic defects act as scatterers for quasiparticles, resulting in the formation of standing waves that interfere with each other. The width of the author name corresponds to a scalebar of 2 nm.

Contents

1	Introduction	1
1.1	Studying correlated electrons with a scanning tunneling microscope . . .	1
1.2	From Mott insulators to high- T_c superconductors	3
1.3	Outline of this thesis	7
2	The experimental technique: spectroscopic-imaging scanning tunneling microscopy	9
2.1	Scanning tunneling microscopy	10
2.1.1	STM as a probe of the local density of states	12
2.1.2	Tunneling into many-body systems	14
2.2	Spectroscopic-imaging STM	15
2.3	Probing momentum space: quasiparticle interference	17
2.3.1	Comparing STM and photoemission	18
2.4	STM on materials with poor electronic screening	19
2.5	Energy resolution in STM	21

3	Dome: design and construction of an ultra-stable scanning tunneling microscope	23
3.1	Introduction	25
3.1.1	The technical challenge of building an STM	25
3.1.2	Reducing vibrations in STM	27
3.2	The STM head	29
3.2.1	STM head design	30
3.2.2	Improving the geometry with finite element analysis	32
3.2.3	Measurement of resonant frequencies	35
3.3	Cryogenic insert	37
3.4	Vibration isolation table	39
3.5	UHV chamber	39
3.6	Performance	40
3.6.1	Current and height noise at the tunneling junction	40
3.6.2	Quasiparticle interference on Sr_2RhO_4	42
3.7	Conclusions	44
4	Universality of pseudogap and emergent order in lightly doped Mott insulators	45
4.1	Introduction	47
4.2	The electron-doped iridate $(\text{Sr}_{1-x}\text{La}_x)_2\text{IrO}_4$	47
4.3	Sample characterization	49
4.4	Doping level determination with STM	51
4.5	Low doping levels: frozen Mott state	52
4.6	High doping levels: pseudogap and electronic order	54
4.6.1	Phase separation	54
4.6.2	Mapping pseudogap and Mott gap	55
4.6.3	Emergent order	58
4.7	Doping evolution: a sharp transition	60
4.8	Discussion and conclusions	63

5	Poor electronic screening in lightly doped Mott insulators	65
5.1	Introduction	66
5.2	Poor electronic screening and tip-induced band bending	66
5.3	Influence of poor screening on the energy gap of $(\text{Sr}_{1-x}\text{La}_x)_2\text{IrO}_4$. . .	67
5.3.1	Calculation of the band-bending potential	69
5.3.2	Algorithm to retrieve the real energy scales in the LDOS	72
5.4	Bubbles in the conductance layers of $(\text{Sr}_{1-x}\text{La}_x)_2\text{IrO}_4$	74
5.5	Conclusions	78
6	Quasiparticle interference in the correlated metal Sr_2RhO_4	81
6.1	Introduction	82
6.2	The correlated metal Sr_2RhO_4	83
6.3	SI-STM quasiparticle interference measurements	86
6.4	Identification of scattering vectors: comparison with simulations	89
6.5	Extraction of dispersions and Fermi vectors	92
6.6	Comparison with ARPES	94
6.7	Conclusions	96
7	Conclusions and outlook	99
	Bibliography	103
	Samenvatting	115
	Publications	119
	Curriculum Vitae	121
	Acknowledgements	123

