



Universiteit
Leiden
The Netherlands

Cardiovascular effects of thoracic epidural anaesthesia

Wink, J.

Citation

Wink, J. (2019, May 9). *Cardiovascular effects of thoracic epidural anaesthesia*. Retrieved from <https://hdl.handle.net/1887/72411>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/72411>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/72411> holds various files of this Leiden University dissertation.

Author: Wink, J.

Title: Cardiovascular effects of thoracic epidural anaesthesia

Issue Date: 2019-05-09

Chapter 4

Effects of thoracic epidural anaesthesia on neuronal cardiac regulation and cardiac function

Jeroen Wink, Bernadette Th Veering,
Leon PHJ Aarts, Patrick F Wouters

Anesthesiology 2019;130:472-491

Introduction

Thoracic epidural anaesthesia (TEA) is widely applied in thoracic and abdominal surgical procedures, as it provides excellent analgesia and decreases postoperative pulmonary complications^{1,2}. Epidural anaesthesia with local anesthetics produces sensory and motor blockade but also affects the autonomic nervous system. The resultant effects on the cardiovascular system vary with the level and the extend of sympathetic blockade. Involvement of the lower thoracic region (T6-L1) by TEA is associated with increased venous capacitance and redistribution of blood to the dilated splanchnic veins. This results in decreased venous return to the heart and a reduction of cardiac preload³. Arterial vasodilation in blocked segments is counteracted by compensatory vasoconstriction in unblocked segments and the effect on cardiac afterload depends on the balance between blocked and unblocked segments⁴. Direct effects of cardiac sympatholysis have not been the subject of detailed investigation. Many studies quantified the cardiac effects of TEA using load dependent indices of contractile performance which does not allow differentiation between direct and indirect effects. Regardless, the effects of TEA have generally been considered beneficial to the cardiovascular system and protective against surgical stress⁵. Interestingly, recent systematic reviews have not been able to confirm improved cardiac outcome in surgical patients treated with TEA⁶. In contrast, some evidence was found for increased cardiovascular problems in high-risk patients receiving neuraxial block^{7, 8}. In light of these concerns a reappraisal of the cardiovascular effects associated with the use of TEA seems appropriate. Since the last published review on this subject⁹, new data have emerged from experimental and clinical studies addressing previously unexplored domains such as the effects of cardiac sympathectomy on right ventricular function, an important determinant of outcome in surgery¹⁰⁻¹².

We conducted the present review to update the knowledge in this field, with focus on the effects associated with high TEA and cardiac sympathectomy in the normal and diseased cardiovascular system.

Materials and methods

The databases PubMed, Embase and Cochrane were searched by the author and by an independent expert librarian to identify studies in which the cardiac sympathetic nerves (T1-T5) are involved in neural blockade by thoracic epidural anaesthesia. The search strategy consisted of the following thesaurus terms and text words: (“Anesthesia, Epidural”[majr] OR “Epidural Anesthesia”[ti] OR “Epidural Anaesthesia”[ti] OR “Epidural Analgesia”[ti] OR “Analgesia, Epidural”[majr]) AND (“thoracic”[ti] OR “cervicothoracic”[ti]) AND (“sympathectomy”[MeSH] OR “sympathectomy”[tw] OR “sympathicolysis”[tw] OR (“Heart”[tw] OR “Heart”[Mesh] OR “cardiac”) AND (“adrenergic activation”[tw] OR “sympathetic innervation”[tw] OR “autonomic innervation”[tw])) OR “Cardiac”[tw] OR “cardiovascular”[tw] OR “blood circulation”[MeSH] OR “circulation”[tw] OR “circulatory”[tw] OR “cardiopulmonary”[tw] OR “heart rate”[MeSH] OR “heart rate”[tw] OR “baroreceptor”[tw] OR “haemodynamic”[tw] OR “hemodynamic”[tw]

OR “hemodynamics”[MeSH] OR “hemodynamics”[tw] OR “haemodynamics”[tw] OR “Echocardiography”[Mesh] OR “Echocardiography”[tw] OR “echocardiographic”[tw] OR “Coronary Vessels”[Mesh] OR “coronary”[tw] OR “ventricular”[tw] OR “Heart Ventricles”[Mesh] OR ventricle*[tw] OR “Ventricular Function”[Mesh] OR “Diastole”[Mesh] OR “Systole”[Mesh] OR “systolic”[tw] OR “diastolic”[tw] OR “myocardial”[tw] OR “Myocardial Contraction”[Mesh] OR “Contraction”[tw] OR “contractility”[tw] OR “contractile”[tw] OR “stress”[tw] OR “exercise”[MeSH] OR “exercise”[tw] OR “ischemic heart disease”[tw] OR “ischaemic heart disease”[tw] OR “Myocardial Ischemia”[Mesh] OR “Coronary Artery Disease”[Mesh] OR “Myocardial Infarction”[Mesh] OR “Hypertension, Pulmonary”[Mesh] OR “pulmonary hypertension”[tw]). This initial search strategy yielded 1189 references.

An additional 20 references were obtained through subsequent hand search for relevant articles and authoritative texts in cited references. Only articles published in English were included. Articles were considered relevant if cardiac sympathetic blockade by TEA was demonstrated. If assessment of neural blockade was not reported, articles were excluded unless epidural puncture was mentioned to be at the cervical or high thoracic level or when the combination of puncture level and dose of local anaesthetics were shown earlier to induce cardiac sympathetic blockade. Editorials, letters to the editor, case reports, abstracts only, studies assessing pain or studies targeting children as the study population were excluded.

The first author evaluated titles and abstracts and selected articles according to relevance and to the inclusion and exclusion criteria. The remaining articles were reviewed full-text and screened for eligibility according to the inclusion and exclusion criteria. See **Figure 1**, presenting a flow diagram of literature search. The author’s main objective of this search was to compose an updated narrative review regarding the specific effects of TEA on cardiac function.

Search Results

The search results and study selection flowchart are presented in **Figure 1**. From the initial 1209 records identified through database searching and cited reference searching, 577 were duplicates or not in the English language and 560 records were excluded because the studies did not meet our inclusion criteria.




Figure 1. Flow diagram of literature search

Cardiac neurophysiology

The central nervous system exerts a beat-to-beat control on cardiac function. Specific areas in the brain involved in emotional behaviour, stress responses and homeostatic reflexes affect cardiac function¹³. These brain areas give excitatory input to the preganglionic sympathetic fibers originating from the intermediolateral cell column of the spinal cord. Preganglionic sympathetic neurons synapse on postganglionic noradrenergic cardiac nerves in the paravertebral ganglia (**Figure 2**). It is generally assumed that the cardiac sympathetic outflow emerges from spinal levels T1 to T5, with a main supply to the ventricles from T1 to T4¹⁴.

Preganglionic parasympathetic fibers originate predominantly in the nucleus ambiguus and also in the dorsal motor nucleus of the vagal nerve. Upon entering the heart, the postganglionic sympathetic and parasympathetic nerves converge into the cardiac plexus. Electrical or chemical stimulation of neuronal tissue within the cardiac sympathetic nervous system, usually the right and/or left stellate ganglion, has yielded relevant information regarding the interplay between the cardiac autonomic nervous system and cardiac function. In general, cardiac sympathetic stimulation in animals¹⁵⁻²¹ and humans²² increases inotropy, dromotropy and chronotropy of the heart. Increases in peak systolic pressure of the left ventricle LV (20-167 %) and maximum positive rate of pressure change (dP/dt max) (20-213 %) after unilateral or bilateral stellate ganglion stimulation indicate substantial increases in contractility of the LV in animals^{15, 16, 19, 21} and humans²². LV relaxation (lusitropy) was also shown to improve substantially after cardiac sympathetic stimulation^{16, 23-25}.




Figure 2. Overview of cardiac innervation

Schematic drawing of the cardiac visceral innervation system. Cardiac innervation starts with a signal from the heart or baroreceptors (e.g., on the aorta), relayed by sensory nerves (blue) giving feedback on, for instance, the levels of oxygen, carbon dioxide and blood pressure. The brain will give a signal to parasympathetic or sympathetic nerves to either relax or stimulate the heart. Parasympathetic innervation is achieved mainly via the vagal nerve (green) that will synapse in cardiac ganglia from where postganglionic nerves innervate the SA node, AV node and ventricular myocytes. Sympathetic neurons (red) start in the grey matter of the spinal cord, where interneurons (orange) from the brain project to the sympathetic neurons. Via the ventral root of the spinal cord, sympathetic nerves synapse in the sympathetic chain, from where postganglionic nerves will enter the heart. Modified with permission from Vegh et al. (Vegh et al: Part and parcel of the cardiac autonomic nerve system: unravelling its cellular building blocks during development. *J Cardiovasc Dev Dis* 2016, 3:28).

Sympathetic and parasympathetic activity depends on chemo-and/or mechanosensory input from multiple cardiac regions, the coronary vasculature and from major intrathoracic- and cervical vessels. The nucleus of the solitary tract receives input from these chemo-and /or mechanosensory receptors via the glossopharyngeal and vagal nerve and is the first relay for several cardiac and cardiovascular reflexes (Figure 3).

The baroreceptor reflex

High-pressure stretch receptors in the aortic arch and the carotid sinus are triggered when mechanical deformation of the vessel wall occurs. Increased blood pressure activates these mechanoreceptors resulting in inhibition of sympathetic outflow, and a subsequent decrease in total peripheral resistance, heart rate and myocardial contractility²⁶. Besides this sympatho-inhibitory pathway there is a cardio-inhibitory pathway, which upon excitation of the cardiovagal neurons of the nucleus ambiguus, results in a decrease in heart rate²⁶.

Cardiac reflexes

Atrial stretch reflex. Distension of low pressure receptors at the (pulmonary) vein-atrial junctions is signalled via vagal afferents to increase sympathetic activity and reduce vagal tone to the sinoatrial node. This results in increased heart rate without increased myocardial contractility (Bainbridge reflex)²⁷. Conversely, low atrial pressure causes bradycardia. This positive feedback reflex, creating a direct relationship between filling pressures and heart rate, is rarely observed in clinical practice because it is weaker than and inferior to the baroreflex, a negative feedback system. It is typically observed in neonates and infants however, where the baroreflex is not yet fully developed. In theory, the Bainbridge reflex can become more prominent in clinical conditions associated with impaired baroreflex function²⁸.

Ventricular reflex. Both ventricles contain mechanosensors and chemosensors most of which can sense mechanical and chemical changes simultaneously²⁹. Either chemical stimulation, potentially elicited by a host of chemicals (e.g. after myocardial infarction), and possibly mechanical stimulation by decreased end-systolic volumes^{30, 31} activates these receptors to decrease sympathetic outflow and increase parasympathetic tone. This results in bradycardia, vasodilatation and hypotension, a response known as the Bezold-Jarisch reflex. This cardio-inhibitory reflex was suggested to play a cardioprotective role but this has never been confirmed³².

Coronary baroreflex. Coronary arteries contain arterial baroreceptors that buffer pressure changes. However, these coronary artery receptors operate at much lower pressures than aortic and carotid baroreceptors³³. Increases in coronary artery pressure cause limited reflex activity but coronary arterial hypotension induces a powerful systemic vasoconstrictor response. The coronary baroreflex is considered a defense mechanism against myocardial hypoperfusion³³.




Figure 3. Cardiac and cardiovascular reflexes

Cardiovascular effects of TEA

The focus of this section is primarily on TEA studies where the cardiac sympathetic nerves (T1-T5) are involved in neural blockade. This definitely includes high TEA but may also apply to mid-thoracic epidural analgesia with cranial spread of anaesthetic blockade.

Effect on cardiovascular reflexes

Baroreceptor control of heart rate depends on an integrative role of the parasympathetic and sympathetic nervous system. This balance is affected when cardiac sympathetic innervation is blocked by TEA. Multiple studies have demonstrated that baroreflex sensitivity is altered by cardiac sympathectomy during cervicothoracic epidural anesthesia³⁴⁻³⁹. However, in some studies TEA attenuated the reduction in HR after blood pressure increase (pressure test) without changing the cardiac acceleration in response to blood pressure decrease (depressor test)³⁴⁻³⁶ whereas others demonstrated the opposite^{37, 38}. Yet another study reports that cervical but not lumbar epidural significantly depresses both “up-and down-sequence” baroreflex sensitivities³⁹. However in this study spontaneously occurring fluctuations in arterial pressure and heart period were used as indices of baroreflex function, a method that has been criticized⁴⁰. The contrasting results in above mentioned studies may relate to heterogeneous study design, differences in the management of TEA-induced preload changes and the use of general anesthesia. Finally, age differences between study populations may also have contributed to the differential effects of TEA on baroreflex control.

Sympathetic control of heart rate can operate indirectly by influencing vagal activity as well as directly by acting as a cardiac accelerator. Both mechanisms could be involved in the effects of TEA on baroreceptor control of heart rate. Regardless, it appears that cardiac sympathetic blockade by TEA at least partially suppresses the baroreceptor reflex. There are no studies addressing the effect of cardiac sympathectomy on other cardiovascular reflexes. It has been suggested that life threatening paradoxical bradycardia in hypotensive patients undergoing spinal and epidural anesthesia is due to attenuation of baroreflex control with subsequent unmasking of the reversed Bainbridge and/or Bezold-Jarish reflex⁴¹⁻⁴⁴. Similarly during severe hemorrhage the Bezold-Jarish reflex may predominate resulting in bradycardia and hypotension⁴³ (**Figure 3**).

Summary:

- Cardiac sympathetic blockade by TEA at least partially suppresses the baroreceptor reflex.
- Extensive neural blockade by TEA with reduction of preload to the heart may evoke hypotension and bradycardia. This has been attributed to impairment of the baroreflex and unmasking of a reversed Bainbridge reflex.

Effect on heart rate

Chronotropic control of the heart is mediated by the balance between sympathetic and parasympathetic tone and is dominated by parasympathetic tone at⁴⁵. The effects of TEA on heart rate depend on the prevailing sympathetic tone, the extent of neural blockade with its proportional impact on pre- and afterload, and the in- or exclusion of cardiac sympathetic nerves. Indeed, TEA effects on heart rate are not solely related to blockade of preganglionic cardiac accelerator nerves but also reflect TEA induced changes in preload and afterload (**Figure 3**) as described in great detail by Veering and Cousins⁴⁶. Clinical studies report no change⁴⁷⁻⁵⁰ or minor reductions in HR^{34, 39, 51-56} after TEA including cardiac sympathetic nerves. Age might affect HR response to TEA since ageing is accompanied by an increased sympathetic nervous system activity at rest⁵⁷. Two studies assessed cardiovascular effects of TEA in different age groups, however results are conflicting. Holman and co-workers⁵⁸ showed that HR reductions after TEA were most pronounced in the elderly group whereas we reported HR reductions only in the younger age group with no changes in the middle or older age groups⁵⁹. Interestingly beta blockers, the chronotropic effects of which might be comparable to those of TEA, were found to result in a more pronounced reduction in HR in young as compared to older healthy volunteers⁶⁰.

Summary:

- The reported effects of TEA on heart rate are mild and not uniform. Changes result from the complex interaction between direct cardiac sympathetic blockade and cardiovascular reflexes which occur secondary to altered preload and afterload.
- Current studies do not indicate a consistent effect of age on HR response to TEA.

Effect on ventricular contractility (Table 1-3)

The majority of TEA studies use load dependent indicators of global left ventricular (LV) performance indicators, such as ejection fraction, fractional area change, fractional shortening, cardiac output (CO), stroke volume (SV) for the assessment of cardiac function⁶¹. Load independent assessment of cardiac performance requires advanced and often invasive technology such as pressure-volume catheters . Newer echocardiographic techniques allowing calculation of the slope of the end-systolic pressure-length relationship, or indices of myocardial velocity and deformation may offer a valid noninvasive alternative for this purpose⁶²⁻⁶⁴.

Effects on LV contractile performance

Several studies have shown a reduction in inotropic state (intrinsic function) after blockade of cardiac sympathetic innervation by TEA. In anaesthetised dogs, the maximal rate of ventricular pressure increase (dp/dt max) of the LV decreased after induction of TEA but not after induction of lumbar epidural anaesthesia (LEA)^{65, 66}. These results suggest a reduction in LV contractility due to blockade of cardiac sympathetic innervation. This is supported by results obtained in pigs, where load independent indices of contractility based on pressure-volume loop analysis decreased after TEA but not after LEA^{49, 50}. In spite of the diminished contractility of the LV after TEA there was no change in global ventricular performance, due to a concomitant reduction of afterload^{49, 50, 66}. Echocardiographic studies in awake and healthy volunteers compared the cardiac effects of TEA versus LEA and also found that only TEA, but not LEA, decreased ejection fraction, fractional area change or fractional shortening and increased left ventricular end-diastolic volume and/or left ventricular end-systolic volume. They also suggest that the reduction in LV cardiac function is due to cardiac sympathetic denervation^{52, 67, 68}. Our group recently evaluated cardiac performance in awake resting patients scheduled for lung surgery using tissue Doppler based measurement of myocardial velocities. We found no effect of TEA on LV systolic pump performance, however, results may have been confounded by our study design which included pre-interventional volume loading. CO increased after TEA, presumably due to the combination of increased external volume loading and TEA-induced afterload reduction. This study in different age groups showed no effects of age on cardiovascular response to TEA⁵⁹. Using the slope of the end-systolic pressure length relationship as a load independent measure of left ventricular contractility, Goertz and colleagues demonstrated in patients under general anesthesia that TEA but not LEA decreases LV contractility by 50% (**Figure 4**)⁴⁷. The results of above mentioned studies suggest that there is a clinically relevant influence of the cardiac sympathetic nervous system on baseline LV function. Blockade of cardiac sympathetic nerves by TEA is associated with a reduction in LV contractility – the magnitude of which is likely related to the level of sympathetic tone. In cardiovascular healthy patients the cardiodepressant effects of TEA seem to be well tolerated with preservation of CO. Use of TEA in these patients is safe. The impact of cardiac sympathectomy in patients with limited cardiac reserve has not been studied specifically.

Summary:

- Load-independent indices demonstrate a 40-50% reduction in LV contractility following cardiac sympathetic blockade by TEA.
- TEA can be safely applied in patients with normal cardiovascular function.
- There are no studies addressing the clinical impact of TEA-induced cardiac sympathectomy in patients with limited cardiac reserve.

Table 1. Effects of TEA on systolic and/or diastolic cardiac function in animals

Author and Year of Publication	Data Acquisition	Human or Animal	Condition	TEA Level	Level of Anesthesia	MAP	HR	SV	CO/CI	LV Systolic Function	RV Systolic Function	LV Diastolic Function	RV Diastolic Function
Holveld <i>et al.</i> ⁸⁵ 1984	LV pressure transducer catheter	Animal N = 7	General anesthesia and β -blocker	T3–4 or T4–5	ND	↓	↓	ND	ND	dP/dt _{max} decreased	ND	ND	ND
Hirabayashi <i>et al.</i> ⁸⁶ 1996	LV pressure transducer catheter and flow probe around ascending aorta	Animal N = 16	General anesthesia TEA	T7/T8	C3/C7–T6/T9 (by ink)	↓	↓	↑	■	dP/dt _{max} decreased LVEDP increased	ND	ND	ND
Rex <i>et al.</i> ⁴⁹ 2007	LV and RV pressure–volume catheters	Animal N = 14	General anesthesia Control compared with TEA General anesthesia and pulmonary hypertension Control compared with TEA	T4/T5 (tip catheter T2)	ND	↓	■	■	■	Ees, Mw and dP/dt _{max} decreased	Ees, Mw, dP/dt _{max} and V25 unchanged	■ ↓	RV: τ increases, $\tau\%RR_{\text{interval}}$ and dP/dt _{min} = ~
Missant <i>et al.</i> ⁵⁰ 2010		Animal N = 18	General anesthesia Control baseline compared with TEA baseline and LEA baseline General anesthesia and pulmonary hypertension Control compared with TEA and LEA	T2 L2 T2	C7–T6 T13–L6 C7–T6	↓ ↓ ↓	■ ↑ ↑	ND ND ND	■ ■ ↓	Ees, Mw and dP/dt _{max} decreased, V100 increased Ees, Mw decreased and V25 increased Ees, Mw decreased	↓ ↓ ↓	ND ND ND	↓ τ increased, $\tau\%RR_{\text{interval}}$ = and dP/dt _{min} decreased ND ND

CI, cardiac input; CO, cardiac output; dP/dt_{max}, peak rate of ventricular pressure decrease; Ees, slope of the end-systolic pressure–volume relationship; HR, heart rate; LEA, lumbar epidural anesthesia; LV, left ventricle; LVEDV, left ventricular end-diastolic volume; LVEDP, left ventricle end-diastolic pressure; LVEDV, left ventricle end-diastolic volume; MAP, mean arterial pressure; Mw, slope of the preload-recruitable stroke work relationship; ND, not determined; RV, right ventricle; SV, stroke volume; τ , time constant of ventricular relaxation; $\tau\%RR_{\text{interval}}$, corrected for heart rate by normalizing to the RR interval; TEA, thoracic epidural anesthesia; V₂₅ and V_{100}}, volume intercept of end-systolic pressure–volume relation, quantified at pressure 25 and 100 mm Hg, respectively.
 ↑, increased; ↓, decreased; ■ or =, no effect

Table 2. Effects of TEA on systolic and/or diastolic ventricular function in humans.

Author and Year of Publication	Data Acquisition	Human or Animal	Condition	TEA Level	Level of Analgesia	MAP	HR	SV	CO/CI	LV Systolic Function	RV Systolic Function	LV Diastolic Function	RV Diastolic Function
Wathvil <i>et al.</i> ⁵² 1985	Echo (TTE) and systolic time intervals; PEP/LVET ratio	Human N = 9	Awake Rest Awake Exercise	T4	C8/T1–T5/T7	■ ↓	↓ ↓	↓ ND	↓ ND	PEP/LVET ratio increased EF decreased LVEDd = and LVESd increased ND	ND ND	ND ND	ND ND
Goertz <i>et al.</i> ⁴⁷ 1993	Echo (TTE); end-systolic pressure–volume relationship (ESPR)	Human N = 36	General anesthesia TEA compared with control group or LEA group (L2–L5)	T8–T11	C6/T1–T11/L4	■	■	ND	ND	E _{max} ESPVR decreased LVEDV, LVESV and fractional area change remained unchanged	ND	ND	ND
Nimi <i>et al.</i> ⁶⁷ 1997	Echo (TTE)	Human N = 24	Awake	T4–T6 T10–T12	T1–T10 T6–L2	ND ND	↓ ■	■ ■	↓ ■	FAC decreased EF tends to decrease and LVEDV and LVESV increased ■ FAC, EF, LVEDV and LVESV remained unchanged	ND ND	■ MV E, MVA and MV E/A unchanged MV DT increased ■ MV E, MV A, MV E/A and MV DT unchanged	ND ND
Shiga ⁶⁸ 1998	Echo (TEE)	Human N = 16	General anesthesia Control compared to TEA	T4	At least T1–T5	↓	↓	ND	ND	FS decreased LVEDV = LVESV increased	ND	■ MV E and MV DT unchanged MVA decreased and MV E/A increased	ND
Wink <i>et al.</i> ⁵⁹ 2014	Echo (TTE); Tissue Doppler imaging	Human N = 31	Awake	T3–T4	C4/6–L1/4	↓	■	↑	↑	MV S' and MPI = EF increased ■ or ↑	~ TAPSE and TV S' increased RV MPI = IVA decreased	↑ MV E' increased	↑ TV E' increased
Wink <i>et al.</i> ⁷¹ 2016	RV conduction catheter Atrial pacing	Human N = 11	General anesthesia Control compared with TEA GA and PHT Control compared with TEA	T3–T4	ND	■ ■	■ ■	■ ■	■ ■	ND ND	↓ Ees, stroke work, dP/d _{max} decreased and V25 increased ■	ND ND	↓ dP/dt _{max} decreased and nonsignificant increase in τ ■

CI, cardiac input; CO, cardiac output; dP/dt_{max}, peak rate of RV pressure increase; dP/dt_{min}, peak rate of ventricular pressure decrease; DT, deceleration time, time interval required for the E velocity to decline from its peak to the baseline; E/A, ratio of E to A; Ees, the slope of the end-systolic pressure–volume relationship; EF, ejection fraction; E_{max}, maximal elastance; FAC, fractional area change; FS, fractional shortening; GA, general anesthesia; HR, heart rate; LEA, lumbar epidural anesthesia; LV, left ventricle; LVEDd, left ventricular end-diastolic diameter; LVEDV, left ventricle end-diastolic volume; LVESd, left ventricular end-systolic diameter; LVESV, left ventricle end-systolic volume; LVET, left ventricular ejection time; MAP, mean arterial pressure; MPI, myocardial performance index or TEI index; MV A, peak velocity during atrial contraction phase; MV E', early diastolic velocity of the mitral annulus; MV S', systolic velocity of the mitral annulus; ND, not determined; PEP, pre-ejection period; PHT, pulmonary hypertension; RV, right ventricle; SV, stroke volume; τ, time constant of ventricular relaxation; TAPSE, tricuspid annular plane systolic excursion; TEA, thoracic epidural anesthesia; TEE, transthoracic echocardiography; TTE, transthoracic echocardiography; TV E', early diastolic velocity of the tricuspid annulus; TV S', systolic velocity of the tricuspid annulus; V₂₅ and V₁₀₀, volume intercept of end-systolic pressure–volume relation, quantified at pressure 25 and 100 mm Hg, respectively.
 ↑, increased; ↓, decreased; ■ or =, no effect; ~, unclear effect




Figure 4. Arithmetic means (\pm s.d.) of the maximal elastance (E_{max}) of the left ventricle.

*indicates $P < 0.001$ versus Group 3 (control) and versus Group 4 (LEA). TEA, thoracic epidural anaesthesia; LEA, lumbar epidural anaesthesia. Modified with permission from Goertz et al. (Goertz et al: influence of high thoracic epidural anaesthesia on left ventricular contractility assessed using the end-systolic pressure-length relationship. *Acta Anaesthesiol Scand* 1993; 37: 38-44.

Table 3. Effects of TEA on systolic and/or diastolic cardiac function in patients with coronary artery disease

Author and Year of Publication	Data Acquisition	Human or Animal	Condition	TEA Level	Level of Analgesia	MAP	HR	CPP	CO/CI	Global LV Systolic Function	Regional LV Systolic Function	LV Diastolic Function	RV Function
Kock <i>et al.</i> ¹¹⁶ 1990	Angiocardiology and ST-segment analysis	Human N = 10	General anesthesia and B-blocker at rest and during stress-induced myocardial ischemia	T3–T5 T3–T5	At least T1–T5 At least T1–T5	■ ↓	■ ■	ND ND	ND ND	■ Global EF was unchanged after TEA	■ Regional EF was unchanged after TEA	ND ND	ND ND
Saada <i>et al.</i> ¹¹⁷ 1992	Echo (TEE) and pulmonary artery catheter	Human N = 26	General anesthesia and TEA	T6–T7 or T7–T8	ND	↓	↓	↓	↓	↑ Improved global EF after TEA	■ Improved regional EF and wall motion score after TEA	ND	ND
Berendes <i>et al.</i> ¹¹⁸ 2003	Echo (TEE)	Human N = 73	General anesthesia for CABG Control (n = 37) vs. TEA (n = 36)	C7–T1	T1–T7	■	■	ND	■	FAC was unchanged after TEA	■ Wall motion score was unchanged after TEA	ND	ND
Schmidt <i>et al.</i> ¹¹⁹ 2005	Echo (TEE) and pulmonary artery catheter	Human N = 37	Awake	T1–T2 or T2–T3	C7–T7	↓	↓	↓	↓	FAC was unchanged after TEA	■ Improved wall motion score after TEA	↑ Improved flow propagation velocity and myocardial performance index	ND
Jakobsen <i>et al.</i> ¹²⁰ 2009	Echo (TTE); tissue Doppler imaging TTE	Human N = 15	Awake	T2–T3	At least T1–T5	↓	↓	ND	↑	EF and tissue tracking score increased after TEA	ND	↑ Improved E/A ^a indicative of improved relaxation	ND
Wafaa <i>et al.</i> ¹²¹ 2011	Echo (TTE)	Human N = 48	General anesthesia Control (n = 24) vs. TEA (n = 24)	T2–T5	At least T1–T5	↓	↓	ND	ND	■ EF and FAC were unchanged after TEA	ND	~ Reported improvement of relaxation pattern after TEA however based on transmural flow patterns	ND

CABG, coronary artery bypass grafting; CI, cardiac input; CO, cardiac output; CPP, coronary perfusion pressure; E/A^a, ratio of peak early and late diastolic velocity of the mitral annulus; EF, ejection fraction; FAC, fractional area change; HR, heart rate; LV, left ventricle; MAP, mean arterial pressure; ND, not determined; RV, right ventricle; ST-segment, region between the end of the S-wave and the beginning of the T-wave on the electrocardiogram; TEA, thoracic epidural anesthesia; TTE, transthoracic echocardiography.

↑, increased; ↓, decreased; ■, no effect; ~, unclear effect

Effect on RV contractile performance

The sympathetic nervous system plays an important role in the regulation of RV function. This is illustrated by a 100% increase in contractile force of the RV after right and left stellate ganglion stimulation, both containing a significant portion of the sympathetic nerves innervating the heart¹⁷. Only a few studies assessed the effects of cardiac sympathetic inhibition by TEA on RV performance. Animal studies using load-independent parameters of contractility, did not find decreases in baseline contractility of the RV after induction of TEA during general anaesthesia^{49,50} but indicated that TEA inhibited the positive inotropic effect to increased afterload (Figure 5). This mechanism referred to as homeometric autoregulation enables the RV to maintain stroke volume without compensatory dilatation of the RV^{69, 70}. In awake patients TEA reduced RV isovolumetric acceleration, suggesting decreased RV contractility. However changes in loading conditions prevented clear conclusions regarding effects of TEA on RV contractility⁵⁹.




Figure 5. Right and left ventricular contractility during baseline, epidural anaesthesia, and acute PA occlusion in control animals and animals with a thoracic or lumbar epidural anaesthesia.

The effects of epidural anaesthesia on the slope of the preload-recrutable stroke work relationship (Mw) (A and B) and the slope of the end-systolic pressure–volume relationship (Ees) (C and D) during baseline, epidural anaesthesia (EDA), and during acute PA occlusion (PA occlusion) in the left (LV) and right ventricle (RV). Values are presented as mean (SEM). C, control animals; TEA, thoracic epidural anaesthesia; LEA, lumbar epidural anaesthesia; Ees, slope of the ESPVR; Mw, slope of the PRSW relationship; PA, pulmonary artery. *P,0.05 vs baseline; †P,0.05 vs epidural; ‡P,0.05 vs C. Reprinted with permission of Missant et al. (Missant et al: Differential effects of lumbar and thoracic epidural anaesthesia on the haemodynamic response to acute right ventricular pressure overload. *BJA* 2010; 104: 143–149).

We recently investigated the effects of TEA on RV contractility in patients during lung surgery and one lung ventilation. Using fixed rate pacing and employment of pressure-volume loop analyses load-independent indices of intrinsic RV function were obtained before and after induction of TEA during general anesthesia. Our data demonstrated TEA-induced impairment of baseline RV contractility⁷¹ as reflected by changes in the slope and volume intercept of the end-systolic pressure volume relationship (Figure 6). In addition there was a 25% to 30% reduction in stroke work. These observations slightly differed from the animal studies where TEA inhibited the increase in RV function but did not reduce baseline RV performance. However, baseline measurements in our clinical study were obtained during one lung ventilation, a condition known to induce hypoxic pulmonary vasoconstriction. It was postulated therefore that sympathetic tone and RV afterload might have been elevated already prior to the initiation of TEA. This was not the case in the animal studies where baseline values were obtained during normal ventilation^{71,72}.




Figure 6. Schematic RV pressure-volume loops based on mean end-diastolic and end-systolic pressures and volumes during one lung ventilation at baseline (black loop) and after induction of TEA (red loop). The increase in ESV_{25} and the rightward shift and more shallow slope of the ESPVR after TEA, indicate a decreased contractile performance. ESPVR, end-systolic pressure-volume relationship; EDPVR, end-diastolic pressure-volume relationship; Eed, slope of the end-diastolic pressure-volume relationship; Ees, slope of the end-systolic pressure-volume relationship; ESV_{25} , volume intercept of ESPVR at 25 mmHg; RV, right ventricle; TEA, thoracic epidural anesthesia. Reprinted with permission from Wink et al. (Wink et al: Thoracic Epidural Anesthesia Reduces Right Ventricular Systolic Function With Maintained Ventricular-Pulmonary Coupling. *Circulation* 2016; 134: 1163–1175.

Regardless of this discussion, both animal studies and clinical studies were concordant in showing that cardiac sympathectomy with high TEA directly affects RV function. These effects may not have much clinical impact in subjects with normal cardiovascular function, but could be of importance in patients with preexisting or pending RV dysfunction and pulmonary hypertension. It is interesting to note that epidural analgesia was found an important contributing factor to major perioperative complications in patients undergoing pneumonectomy⁷, although that evidence appeared not robust enough to support a change in practice. Similarly, in a secondary analysis of the POISE study, Leslie et al found evidence for increased cardiovascular problems in high-risk patients receiving neuraxial block⁸. It is clear that prospective outcome studies in high-risk patients are urgently needed to address this issue. For some patients undergoing major surgery who are at risk for RV failure or for those being treated with TEA who develop sudden RV failure in the postoperative period, potent alternative analgesic techniques such as paravertebral blocks could also provide a solution.

Summary:

- Cardiac sympathetic blockade by TEA directly reduces RV contractility.
- The clinical importance of this effect is not known at present.

Effect on LV and RV diastolic function

Ventricular relaxation and compliance are important determinants of diastolic function, an essential component of cardiac pump performance. Impaired relaxation and/or reduced compliance of the ventricle have been associated with increased perioperative risk for 30-day cardiovascular events and long-term cardiovascular mortality⁷³. The effects of TEA on diastolic function have not been well established let alone the relevance of these temporary effects on clinical outcome. In normal conditions, contraction and relaxation are functionally coupled. Sympathetic stimulation of cardiac beta1 receptors causes a rise in c-AMP which enhances calcium release during systole but also facilitates removal of the excess calcium during diastole^{16,23-25,74}. It would be reasonable to expect that if TEA-induced cardiac sympathectomy causes a mild decrease in contractile performance, it would also decrease relaxation proportionally. However, most studies reported unchanged LV diastolic function after TEA in patients with normal cardiovascular status but used load dependent parameters^{67,68}. We recently studied the effect of TEA on diastolic RV and LV function in different age groups using tissue Doppler imaging^{75,76}. Peak velocity of the mitral and tricuspid annular motion during early diastole were used to assess LV and RV diastolic function⁷⁷. Baseline diastolic function was significantly lower in older patients but TEA did not reduce diastolic function in any of the age groups⁵⁹. Animal studies using invasive load independent measurement techniques also demonstrated that diastolic function of the LV and RV remained unchanged after TEA⁴⁹. Using a similar technique in patients undergoing lung surgery, we found that the decrease in RV inotropic state following TEA was accompanied by a discrete reduction in RV relaxation properties^{71,78}. TEA effects on diastolic function are mild and well tolerated in healthy patients. Whether this also applies to patients with established diastolic

dysfunction remains to be investigated. This latter group of patients is particularly sensitive to preload changes and for that reason needs specific attention when TEA is applied.

Summary:

- The direct effect of TEA on diastolic function is minimal.
- Patients with diastolic dysfunction are extremely sensitive to changes in preload which almost invariably occur with TEA. Particular attention is required to prevent the hemodynamic consequences of preload reduction following TEA in this subset of patients.

Effects in the healthy coronary system

Coronary arteries are innervated by the parasympathetic and sympathetic nervous system. Parasympathetic stimulation results in coronary vasodilatation⁷⁹. Sympathetic alpha-adrenoceptor-mediated coronary vasoconstriction has been demonstrated^{80, 81} but adrenergic activation of the heart will also induce beta- adrenoceptor-mediated coronary vasodilatation⁸². Indirectly, adrenergic activation raises myocardial oxygen demand resulting in increased myocardial blood flow via local vasodilatory mechanisms⁸³. These sympathetic-mediated mechanisms compete with local metabolic vasodilatation, making control of myocardial perfusion a complex phenomenon⁸⁰.

Hirabayashi and colleagues assessed TEA effects on coronary circulation in healthy dogs⁶⁶. Coronary perfusion pressure, coronary blood flow as well as systemic arterial pressure decreased after TEA and LEA. Interestingly, TEA only increased calculated coronary vascular resistance. The authors suggest that the increased coronary vascular resistance is an autoregulatory response to the decreased myocardial oxygen demand after TEA as a consequence of a lower arterial pressure, heart rate and myocardial contractility. Their data are consistent with an earlier animal study that reported decreases in arterial pressure and increases of coronary diastolic pressures after induction of TEA⁸⁴. Whether these TEA-induced changes in diastolic pressure and resistance in coronary arteries are autoregulatory responses to decreased myocardial oxygen demand or direct effects of blockade of sympathetic efferents to the coronary arteries remains unclear. A recent clinical study in patients with normal cardiovascular physiology failed to demonstrate any effect of TEA on myocardial blood flow in rest. On the other hand, during sympathetic stimulation by the cold pressor test there was a 70% increase in myocardial blood flow in the control group whereas myocardial blood flow in the TEA group remained the same⁸⁵. However, with a mean difference in increase of rate pressure product between the TEA and control group of 2215 (mmHg/min) the increase in cardiac work and myocardial oxygen demand, if any, in the TEA group was substantially lower. Therefore, the lack of an augmented myocardial flow response during the cold pressor test could have been an autoregulatory response as suggested in animal studies. While unique for the fact that it is one of the only studies in men on this subject, the data should be interpreted with caution as myocardial contrast echocardiography is not an accurate technique to quantify myocardial perfusion⁸⁶.

Summary:

TEA effects on normal coronary arteries are primarily governed by the reduction in myocardial oxygen demand.

Cardiovascular effects during stress

The concept of cardiac sympathetic innervation being primarily essential in the state of exercise and not in rest is illustrated in heart transplant patients. Reinnervation of the surgically denervated heart occurs only in some cardiac transplant patients. Cardiac pump performance at rest, as measured by global and regional ejection fraction, does not differ between patients with and without reinnervation. However, the group with reinnervation has a significantly better chronotropic and inotropic response to exercise, resulting in better exercise performance⁸⁷. Several animal studies suggest that cardiovascular effects of TEA are more pronounced during stress. TEA had no or minimal effect on HR in baseline conditions yet the substantial increase of HR during raised RV afterload was blunted following TEA^{49, 50, 88, 89}. In addition TEA prevented the increase in contractility of the RV and LV to acutely raised RV afterload resulting in a decrease of CO and SV^{49, 50} (**Figure 4**). In dogs the cardiovascular response to severe hypoxemia was almost completely abolished by high epidural anesthesia⁹⁰. Although the level of neuroaxial block was not determined in the latter study, HR increase to hypoxia was clearly suppressed, indicating involvement of the cardiac sympathetic efferents in epidural blockade. Several clinical studies demonstrated that TEA significantly reduced increases in blood pressure and/or HR following laryngoscopy and intubation⁹¹⁻⁹³ whereas baseline values were only minimally changed by TEA. In these studies TEA involved blockade of the sympathetic innervation to the heart as well as to the adrenal glands. In a study by Dohi and colleagues cardiac sympathetic blockade by TEA or adrenal sympathetic blockade by LEA did not attenuate the circulatory response to laryngoscopy or intubation⁹⁴.

These contrasting results suggest that sympathetic innervation both to the heart and adrenal glands contribute to the circulatory response following laryngoscopy and intubation. Kirno et al also showed more pronounced effects of TEA in humans during stress. They reported significant TEA-induced reductions in the cardiac norepinephrine spillover compared to a control after, but not prior to the surgical stress of sternotomy⁹⁵. TEA exercise studies in healthy volunteers studies showed that increases in HR during exercise were blunted following TEA and the decrease in HR appeared to be more substantial with increasing workloads^{52, 96}. Interestingly TEA did not completely abolish the HR response to stress and/or exercise suggesting either incomplete cardiac sympathetic blockade by TEA or involvement of the adrenal glands. Ottesen and colleagues studied the effects of selective blockade of the sympathetic cardiac segments by TEA in rest and during physical exercise in volunteers using a pulmonary artery catheter. During maximal exercise, SV was maintained and CO decreased only because of a reduction in HR after TEA⁹⁶. The preservation of SV does not indicate preservation of cardiac function however as higher end-diastolic volumes in the presence of reduced ejection fraction would also preserve SV.

Unfortunately, intrinsic ventricular function was not assessed in this study, hence comparisons between TEA effects on ventricular function during rest and during stress could not be made.

In conclusion, both experimental and clinical studies suggest that cardiovascular effects of TEA are more pronounced during stress/exercise than in resting conditions. Data from TEA studies performed in resting conditions do not provide information on its role in the perioperative period, which is typically characterized by surgical and hemodynamic stress. The reduction in cardiac metabolic demands and blunting of the stress response have generally been considered beneficial properties of cardiac sympathectomy with TEA. This is undoubtedly the case for patients with ischemic heart disease but may not apply for other subgroups where the endogenous sympathetic stress response is required to restore cardiovascular homeostasis. Interestingly, systematic reviews failed to show a beneficial effect on cardiovascular outcome in patients treated with TEA.

Summary:

- TEA effects are more pronounced during elevated sympathetic tone.
- Elevation of sympathetic tone is an established short term survival mechanism to preserve cardiovascular homeostasis in the face of hemodynamic disruption. As such sympatholysis by TEA could interfere with this endogenous defense mechanism when hemodynamic challenges occur in the perioperative setting.

Effects of TEA in cardiovascular disease

Ischemic heart disease

Coronary blood flow is regulated primarily by change in myocardial oxygen demand⁸³ induced by variations in wall tension, contractile state and heart rate. In addition, large coronary epicardial coronaries and coronary resistance vessels are densely innervated by the sympathetic nervous system⁸⁰. Experimental animal studies have demonstrated that TEA improves endocardial blood flow during acute myocardial infarction⁸⁴, reduces myocardial acidosis and ischemia after coronary artery occlusion^{97, 98}, reduces myocardial ischemic injury and infarct size⁹⁹⁻¹⁰¹ and improves recovery from myocardial stunning¹⁰². In patients with coronary artery disease TEA has been demonstrated to increase myocardial oxygen availability¹⁰³ and to improve myocardial oxygen balance by reducing heart rate, preload and afterload without affecting coronary perfusion pressure¹⁰⁴. Reiz and colleagues showed that in patients with coronary artery disease TEA reduces coronary vascular resistance and myocardial oxygen consumption¹⁰⁵. This cardioprotective role of TEA is further supported by observations that TEA compared to controls decreased loading conditions of the heart and myocardial oxygen demand following sternotomy⁹⁵.

Coronary atherosclerosis and endothelial dysfunction are associated with an exaggerated response to coronary alpha-adrenergic activation that result in a reduced coronary blood flow response during sympathetic stimulation¹⁰⁶⁻¹⁰⁸. Cardiac sympathetic inhibition by TEA therefore might improve coronary function in patients with coronary artery disease. Indeed, TEA resulted in an increased luminal diameter in stenotic epicardial coronary arteries but not in the non-stenotic epicardial coronaries¹⁰⁹. Whether this resulted in increased myocardial blood flow, a phenomenon referred to as reverse coronary steal, is unknown since myocardial blood flow was not measured in this study. In a more recent study by Nygard and colleagues the effects of TEA on myocardial blood flow were assessed in patients with coronary artery disease. While in patients without TEA myocardial blood flow was unchanged during sympathetic stimulation, patients with TEA demonstrated increases in myocardial blood flow at all vascular territories. After sympathetic stimulation by the cold pressor test coronary vascular resistance increased in the group without TEA and decreased in stenotic and non-stenotic vessels with TEA. TEA induced changes in myocardial blood flow were less than 10% at rest whereas 17-100% increases in myocardial blood flow were shown during the cold pressor test¹¹⁰. These data suggest that coronary sympathetic innervation is of minor importance at rest, yet plays an important role during sympathetic stimulation^{111, 112}.

TEA has been used in patients with ischemic heart disease for the treatment of refractory angina pectoris reducing the incidence of myocardial ischemia, decreasing the number and duration of ischemic episodes, producing symptomatic relief of angina and improving quality of life¹¹³⁻¹¹⁵. Besides improving myocardial oxygen balance these TEA induced results may at least partially be attributable to the pain relief obtained by blockade of spinal afferents. The improvement in myocardial oxygen balance after TEA may also affect myocardial function in patients with coronary artery disease. In awake patients with coronary artery disease global and regional wall motion during stress-induced myocardial ischemia has been shown to improve after induction of TEA¹¹⁶. These results were confirmed in patients with coronary artery disease during general anesthesia and TEA. Despite lower coronary perfusion pressures after the induction of TEA segmental wall motion was unchanged in patients with coronary artery disease whereas segmental wall motion decreased in the patient group without coronary artery disease¹¹⁷. TEA significantly improved regional left ventricular wall motion and reduced ischemia and coronary risk in patients with coronary artery disease in another study¹¹⁸. Schmidt and co-workers report improved diastolic and maintained systolic LV function after TEA in coronary artery disease patients¹¹⁹. Another study using a derivative of tissue Doppler imaging demonstrated TEA-induced improvements of diastolic and systolic LV function in patients with ischemic heart disease¹²⁰.

Clinical studies have been performed assessing the effect of TEA on myocardial damage determined by the amount of postoperative Troponin in coronary artery bypass graft patients. The potential myocardial protective effect of TEA is supported by decreases in postoperative

cardiac Troponin I and T after cardiac surgery^{55, 118, 121} whereas other studies failed to demonstrate an effect of TEA on postoperative Troponin as a marker of myocardial damage¹²²⁻¹²⁶. A potential influence of TEA on the incidence of perioperative myocardial infarction is favored by some studies¹²⁷⁻¹²⁹, but remains to be clarified. A recent meta-analysis by Svircevic and colleagues found no significant effect of TEA on the incidence of perioperative myocardial infarction¹³⁰.

Although 2731 patients from 28 studies were included the authors concluded that the meta-analysis was underpowered and estimated the need for a sample size of 10,000 patients to obtain statistical significance for the reported reduction in the incidence of myocardial infarction from 3.8% to 2.8% after TEA. In fact, the majority of meta-analyses available today failed to show a significant clinical impact of TEA on cardiovascular outcome. This is in contrast with the well documented effects of TEA on the cardiovascular system as reported in physiology studies. The apparent discrepancy may be related to the fact that clinical studies have not zoomed-in on specific risk groups but included very heterogeneous populations instead. As a result, any potential beneficial effect in a particular subgroup, as well as any potentially detrimental effect of TEA in a specific risk population, may go unnoticed and even cancel out a statistical effect on outcome. Ideally, randomized controlled trials should focus on specific patient populations, guided by the observations from pathophysiology studies, to better define the benefits and risks of TEA and optimize its application in clinical practice.

Summary:

TEA improves coronary function and myocardial oxygen balance in patients with ischemic heart disease which results in increased myocardial performance and a reduction of the number and duration of ischemic episodes.

Pulmonary Hypertension

Acute pulmonary hypertension is a frequently encountered phenomenon in cardiothoracic surgery and during hypoxic pulmonary vasoconstriction in the critically ill. This is important since pulmonary hypertension may result in RV failure and RV function has been shown to be an important determinant of outcome¹⁰⁻¹². Jahn and colleagues demonstrated that in a model of ovine pulmonary embolism induction of TEA contrary to lumbar epidural anaesthesia (LEA) improves hemodynamic variables mainly as a result of a decrease in pulmonary vascular resistance. Effects of TEA on right ventricular contractility were not measured and extension of neural blockade was not assessed. Hemodynamic deterioration caused by pulmonary embolism was reduced by TEA and aggravated by LEA^{88, 89}. The authors suggest that reduction of sympathetic outflow to the heart and lungs is the underlying mechanism for the beneficial results of TEA during pulmonary hypertension. They also suggest that increased sympathetic tone in the unblocked thoracic spinal levels associated with LEA is responsible for the hemodynamic aggravation during pulmonary hypertension. This is in contrast with the results of earlier experimental studies which demonstrated that in respiratory distress syndrome, during

pulmonary artery constriction and in pulmonary artery embolism, RV contractility increases proportionally to an increase in RV afterload (homeometric autoregulation). In anesthetized pigs, induction of TEA has been shown to abolish this inotropic response to acutely raised RV afterload (**Figure 4**). Combined with an increase in pulmonary vascular resistance induction of TEA resulted in a significant decrease in cardiac output^{49, 50}. Moreover lumbar epidural anaesthesia had no effect on the hemodynamic response to pulmonary hypertension (**Figure 4**). Results indicate that the sympathetic nervous system might have an important role in the described inotropic response of the RV to pulmonary hypertension. In humans, however, TEA did not affect the native positive inotropic response of the RV to increased afterload (**Figure 5**)⁷¹. Cardiac output was maintained. Of importance is that all patients were paced at a constant rate to accurately assess cardiac contractility. This way HR reduction by TEA was prevented and cardiovascular effects might have been more profound without pacing. Overall, cardiac sympathetic blockade by TEA reduces RV contractile performance. The clinical importance of this finding is unknown. The role of the RV has been neglected for a long time and it has never been investigated whether temporary changes in RV function whether drug induced or epidural induced change outcome. In cardiovascular healthy patients during normal circumstances this is not a safety concern. However, during conditions of acutely raised RV afterload TEA might interfere with the capacity of the RV to adapt to increases in afterload resulting in decreases in CO and cardiovascular collapse.

Summary:

High TEA limits cardiac reserve and the capacity of the RV to adapt to increases in RV afterload, which can decrease cardiac output.

Conclusion

There has been renewed interest in the cardiovascular effects of TEA since the latest review on this subject was published ten years ago⁹. The beneficial hemodynamic effects and cardioprotective properties of TEA, demonstrated in a number of experimental studies, did not translate in better cardiac outcome for patients undergoing surgery who were treated with TEA. More recent exploratory studies even suggested that TEA was associated with increased cardiovascular problems in high risk patients. Mechanisms underlying such a potential harmful effect and characteristics of high risk populations remain speculative however.

Recent experimental and clinical studies have added information on the complex interaction between TEA- induced sympatholysis and cardiovascular homeostasis. Using more advanced methodology they clearly demonstrated that cardiac sympathetic blockade by high TEA reduces LV as well as RV contractility. This direct effect of TEA is well tolerated in healthy subjects because concomitant arteriolar vasodilation and the subsequent decrease in LV afterload facilitates cardiac ejection. Hence overall pump performance and cardiac output are preserved provided that alterations in preload are accounted for. Such a compensatory decrease in afterload however

does not occur in the pulmonary circulation and a direct reduction of RV inotropic state can have more impact on pump performance.

The cardiovascular effects of TEA as studied in baseline conditions at low sympathetic tone do not fully reveal the impact of sympathectomy on cardiovascular homeostatic mechanisms when activated by surgical stress or by hemodynamic disruptions such as hypovolemia and bleeding. TEA attenuates baroreflex function and may unmask primary cardiac reflexes to altered volume status. It is not known to what extent the attenuation of endogenous cardiovascular reflexes affects outcome in the overall population treated with TEA.

Finally, the effects of cardiac sympathectomy with high TEA vary with the pathophysiologic substrate. In patients with coronary artery disease the use of TEA improves myocardial oxygen balance and produces relief of angina. TEA was consistently shown to enhance LV diastolic and systolic function in this population.

In conclusion, the conviction that TEA has beneficial hemodynamic effects may not apply to all patients. While protective in particular pathophysiological conditions such as ischemic heart disease, cardiac sympathectomy may also attenuate the hearts capacity to respond to hemodynamic challenges in particular subgroups. This should be considered whenever patients treated with TEA develop hemodynamic instability in the perioperative setting. Since alternative analgesic strategies with equivalent efficacy are now available, it appears prudent to use a more restrictive approach towards recommending high TEA for patients at risk for perioperative RV failure.

Reference list

1. Ballantyne JC, Carr DB, deFerranti S, Suarez T, Lau J, Chalmers TC, Angelillo IF, Mosteller F. The comparative effects of postoperative analgesic therapies on pulmonary outcome: cumulative meta- analyses of randomized, controlled trials. *Anesth Analg* 1998;86:598-612.
2. Block BM, Liu SS, Rowlingson AJ, Cowan AR, Cowan JA, Jr., Wu CL. Efficacy of postoperative epidural analgesia: a meta-analysis. *JAMA* 2003;290:2455-2463.
3. Arndt JO, Hock A, Stanton-Hicks M, Stuhmeier KD. Peridural anesthesia and the distribution of blood in supine humans. *Anesthesiology* 1985;63:616-623.
4. Baron JF, Payen D, Coriat P, Edouard A, Viars P. Forearm vascular tone and reactivity during lumbar epidural anesthesia. *Anesth Analg* 1988;67:1065-1070.
5. Freise H, Van Aken HK. Risks and benefits of thoracic epidural anaesthesia. *Br J Anaesth* 2011;107:859-868.
6. Wijeyesundera DN, Beattie WS, Austin PC, Hux JE, Laupacis A. Epidural anaesthesia and survival after intermediate-to-high risk non-cardiac surgery: a population-based cohort study. *Lancet* 2008;372:562-569.
7. Powell ES, Cook D, Pearce AC, Davies P, Bowler GM, Naidu B, Gao F; UKPOS Investigators. A prospective, multicentre, observational cohort study of analgesia and outcome after pneumonectomy. *Br J Anaesth* 2011;106:364-370.
8. Leslie K, Myles P, Devereaux P, Williamson E, Rao-Melancini P, Forbes A, Xu S, Foex P, Pogue J, Arrieta M, Bryson G, Paul J, Paech M, Merchant R, Choi P, Badner N, Peyton P, Sear J, Yang H. Neuraxial block, death and serious cardiovascular morbidity in the POISE trial. *Br J Anaesth* 2013;111:382-390.
9. Clemente A, Carli F. The physiological effects of thoracic epidural anesthesia and analgesia on the cardiovascular, respiratory and gastrointestinal systems. *Minerva Anesthesiol* 2008;74:549-563.
10. Haddad F, Doyle R, Murphy DJ, Hunt SA. Right ventricular function in cardiovascular disease, part II: pathophysiology, clinical importance, and management of right ventricular failure. *Circulation* 2008;117:1717-1731.
11. Reed CE, Dorman BH, Spinale FG. Mechanisms of right ventricular dysfunction after pulmonary resection. *Ann Thorac Surg* 1996;62:225-231.
12. Okada M, Ota T, Okada M, Matsuda H, Okada K, Ishii N. Right ventricular dysfunction after major pulmonary resection. *J Thorac Cardiovasc Surg* 1994;108:503-511.
13. Palma JA, Benarroch EE. Neural control of the heart: recent concepts and clinical correlations. *Neurology* 2014;83:261-271.
14. Bonica JJ. Autonomic innervation of the viscera in relation to nerve block. *Anesthesiology* 1968 July;29:793-813.

15. Winter J, Tanko AS, Brack KE, Coote JH, Ng GA. Differential cardiac responses to unilateral sympathetic nerve stimulation in the isolated innervated rabbit heart. *Auton Neurosci* 2012;166:4-14.
16. Zhou W, Yamakawa K, Benharash P, Ajjola O, Ennis D, Hadaya J, Vaseghi M, Shivkumar K, Mahajan A. Effect of stellate ganglia stimulation on global and regional left ventricular function as assessed by speckle tracking echocardiography. *Am J Physiol Heart Circ Physiol* 2013;304:H840-H847.
17. Randall WC, Szentivanyi M, Pace JB, Wechsler JS, Kaye MP. Patterns of sympathetic nerve projections onto the canine heart. *Circ Res* 1968;22:315-323.
18. Yanowitz F, Preston JB, Abildskov JA. Functional distribution of right and left stellate innervation to the ventricles. Production of neurogenic electrocardiographic changes by unilateral alteration of sympathetic tone. *Circ Res* 1966;18:416-428.
19. Zarse M, Plisiene J, Mischke K, Schimpf T, Knackstedt C, Gramley F, Mühlenbruch G, Waldmann M, Schmid M, Hatam N, Graf J, Schuster D, Hanrath P, Pauza D, Schauerte P. Selective increase of cardiac neuronal sympathetic tone: a catheter-based access to modulate left ventricular contractility. *J Am Coll Cardiol* 2005;46:1354-1359.
20. Vaseghi M, Zhou W, Shi J, Ajjola OA, Hadaya J, Shivkumar K, Mahajan A. Sympathetic innervation of the anterior left ventricular wall by the right and left stellate ganglia. *Heart Rhythm* 2012;9:1303-1309.
21. Schlack W, Thamer V. Unilateral changes of sympathetic tone to the heart impair left ventricular function. *Acta Anaesthesiol Scand* 1996;40:262-271.
22. Ajjola OA, Howard-Quijano K, Scovotti J, Vaseghi M, Lee C, Mahajan A, Shivkumar K. Augmentation of cardiac sympathetic tone by percutaneous low-level stellate ganglion stimulation in humans: a feasibility study. *Physiol Rep* 2015;3:1-10.
23. Burwash IG, Morgan DE, Koilpillai CJ, Blackmore GL, Johnstone DE, Armour JA. Sympathetic stimulation alters left ventricular relaxation and chamber size. *Am J Physiol* 1993;264:R1-R7.
24. Henning RJ, Khalil I. Autonomic nervous stimulation affects left ventricular relaxation more than left ventricular contraction. *J Auton Nerv Syst* 1989;28:15-25.
25. Henning RJ, Levy MN. Effects of autonomic nerve stimulation, asynchrony, and load on dP/dt_{max} and on dP/dt_{min} . *Am J Physiol* 1991;260:H1290-H98.
26. Heesch CM. Reflexes that control cardiovascular function. *Am J Physiol* 1999;277:S234-S43.
27. Furnival CM, Linden RJ, Snow HM. Reflex effects on the heart of stimulating left atrial receptors. *J Physiol* 1971;218:447-463.
28. Chapter: Wouters P: Cardiovascular control mechanisms, *Core Topics in Cardiac Anesthesia*, 2nd edition. Edited by Mackay JH, Arrowsmith JE. Cambridge University Press, 2012, pp 28-31.
29. Armour JA, Huang MH, Pelleg A, Sylven C. Responsiveness of in situ canine nodose ganglion afferent neurones to epicardial mechanical or chemical stimuli. *Cardiovasc Res* 1994;28:1218-1225.

30. Jacobsen J, Sofelt S, Brocks V, Fernandes A, Warberg J, Secher NH. Reduced left ventricular diameters at onset of bradycardia during epidural anaesthesia. *Acta Anaesthesiol Scand* 1992;36:831-836.
31. Sander-Jensen K, Marving J, Secher NH, Hansen IL, Giese J, Warberg J, Bie P. Does the decrease in heart rate prevent a detrimental decrease of the end-systolic volume during central hypovolemia in man? *Angiology* 1990;41:687-695.
32. Hainsworth R. Cardiovascular control from cardiac and pulmonary vascular receptors. *Exp Physiol* 2014;99:312-319.
33. McMahan NC, Drinkhill MJ, Hainsworth R. Vascular responses to stimulation of carotid, aortic and coronary artery baroreceptors with pulsatile and non-pulsatile pressures in anaesthetized dogs. *Exp Physiol* 1996;81:969-981.
34. Bonnet F, Szekely B, Abhay K, Touboul C, Boico O, Saada M. Baroreceptor control after cervical epidural anesthesia in patients undergoing carotid artery surgery. *J Cardiothorac Anesth* 1989;3:418-424.
35. Takeshima R, Dohi S. Circulatory responses to baroreflexes, Valsalva maneuver, coughing, swallowing, and nasal stimulation during acute cardiac sympathectomy by epidural blockade in awake humans. *Anesthesiology* 1985;63:500-508.
36. Dohi S, Tsuchida H, Mayumi T. Baroreflex control of heart rate during cardiac sympathectomy by epidural anesthesia in lightly anesthetized humans. *Anesth Analg* 1983;62:815-820.
37. Goertz A, Heinrich H, Seeling W. Baroreflex control of heart rate during high thoracic epidural anaesthesia. A randomised clinical trial on anaesthetised humans. *Anaesthesia* 1992 ;47:984-987.
38. Licker M, Spiliopoulos A, Tschopp JM. Influence of thoracic epidural analgesia on cardiovascular autonomic control after thoracic surgery. *Br J Anaesth* 2003;91:525-531.
39. Tanaka M, Goyagi T, Kimura T, Nishikawa T. The effects of cervical and lumbar epidural anesthesia on heart rate variability and spontaneous sequence baroreflex sensitivity. *Anesth Analg* 2004;99:924-929.
40. Lipman RD, Salisbury JK, Taylor JA. Spontaneous indices are inconsistent with arterial baroreflex gain. *Hypertension* 2003;42:481-487.
41. Barbieri R, Triedman JK, Saul JP. Heart rate control and mechanical cardiopulmonary coupling to assess central volume: a systems analysis. *Am J Physiol Regul Integr Comp Physiol* 2002;283:R1210-R20.
42. Sancetta SM, Lynn RB, Simeone FA, Scott RW, Heckman G, Janouskovec H. Studies of Hemodynamic changes in humans following induction of low and high spinal anesthesia. *Circulation* 1952;6:559-571.
43. Crystal GJ, Salem MR. The Bainbridge and the "reverse" Bainbridge reflexes: history, physiology, and clinical relevance. *Anesth Analg* 2012;114:520-532.
44. Baron JF, Decaux-Jacolot A, Edouard A, Berdeaux A, Samii K. Influence of venous return on baroreflex control of heart rate during lumbar epidural anesthesia in humans. *Anesthesiology* 1986;64:188-193.

45. Robinson BF, Epstein SE, Beiser GD, Braunwald E. Control of heart rate by the autonomic nervous system. Studies in man on the interrelation between baroreceptor mechanisms and exercise. *Circ Res* 1966;19:400-411.
46. Chapter: Veering BT, Cousins MJ: Epidural neural blockade, Neural blokade in clinical anesthesia and pain medicine, 4th edition. Edited by Cousins MJ, Carr DB, Horlocker T.T., Bridenbaugh P.O. Philadelphia, Wolters Kluwer/Lippincott Williams & Wilkins, 2009, pp 241-295.
47. Goertz AW, Seeling W, Heinrich H, Lindner KH, Schirmer U. Influence of high thoracic epidural anesthesia on left ventricular contractility assessed using the end-systolic pressure-length relationship. *Acta Anaesthesiol Scand* 1993;37:38-44.
48. Magnusdottir H, Kirno K, Ricksten SE, Elam M. High thoracic epidural anesthesia does not inhibit sympathetic nerve activity in the lower extremities. *Anesthesiology* 1999;91:1299-1304.
49. Rex S, Missant C, Segers P, Wouters PF. Thoracic epidural anesthesia impairs the hemodynamic response to acute pulmonary hypertension by deteriorating right ventricular-pulmonary arterial coupling. *Crit Care Med* 2007;35:222-229.
50. Missant C, Claus P, Rex S, Wouters PF. Differential effects of lumbar and thoracic epidural anaesthesia on the haemodynamic response to acute right ventricular pressure overload. *Br J Anaesth* 2010;104:143-149.
51. Wink J, Wolterbeek R, Aarts LP, Koster SC, Versteegh MI, Veering BT. Upper thoracic epidural anaesthesia: effects of age on neural blockade and cardiovascular parameters. *Acta Anaesthesiol Scand* 2013;57:767-775.
52. Wattwil M, Sundberg A, Arvill A, Lennquist C. Circulatory changes during high thoracic epidural anaesthesia--influence of sympathetic block and of systemic effect of the local anaesthetic. *Acta Anaesthesiol Scand* 1985;29:849-855.
53. Tanaka K, Harada T, Dan K. Low-dose thoracic epidural anesthesia induces discrete thoracic anesthesia without reduction in cardiac output. *Reg Anesth* 1991;16:318-321.
54. Owczuk R, Steffek M, Wujtewicz MA, Szymanowicz W, Twardowski P, Marjanski T, Wojciechowski J, Zienciuk A, Rzyman W, Wujtewicz M. Effects of thoracic epidural anaesthesia on cardiac repolarization. *Clin Exp Pharmacol Physiol* 2009;36:880-883.
55. Loick HM, Schmidt C, van Aken H, Junker R, Erren M, Berendes E, Rolf N, Meissner A, Schmid C, Scheld HH, Möllhoff T. High thoracic epidural anesthesia, but not clonidine, attenuates the perioperative stress response via sympatholysis and reduces the release of troponin T in patients undergoing coronary artery bypass grafting. *Anesth Analg* 1999;88:701-709.
56. Funayama T, Aida S, Matsukawa T, Okada K, Kumazawa T. Systemic, but not pulmonary, hemodynamics are depressed during combined high thoraco-cervical epidural and general anesthesia in dogs. *Can J Anaesth* 2003;50:454-459.
57. Rooke GA. Cardiovascular aging and anesthetic implications. *J Cardiothorac Vasc Anesth* 2003;17:512- 523.

58. Holman SJ, Bosco RR, Kao TC, Mazzilli MA, Dietrich KJ, Rolain RA, Stevens RA. What constitutes an effective but safe initial dose of lidocaine to test a thoracic epidural catheter? *Anesth Analg* 2001;93:749-754.
59. Wink J, Veering BT, Aarts LP, Wouters PF. Effect of increasing age on the haemodynamic response to thoracic epidural anaesthesia: an observational study. *Eur J Anaesthesiol* 2014;31:597-605.
60. Fleg JL, Schulman S, O'Connor F, Becker LC, Gerstenblith G, Clulow JF, Renlund DG, Lakatta EG. Effects of acute beta-adrenergic receptor blockade on age-associated changes in cardiovascular performance during dynamic exercise. *Circulation* 1994;90:2333-2341.
61. Robotham JL, Takata M, Berman M, Harasawa Y. Ejection fraction revisited. *Anesthesiology* 1991;74:172-183.
62. Suga H, Sagawa K, Shoukas AA. Load independence of the instantaneous pressure-volume ratio of the canine left ventricle and effects of epinephrine and heart rate on the ratio. *Circ Res* 1973;32:314-322.
63. A'roch R, Gustafsson U, Johansson G, Poelaert J, Haney M. Left ventricular strain and peak systolic velocity: responses to controlled changes in load and contractility, explored in a porcine model. *Cardiovasc Ultrasound* 2012;10:10-22.
64. Connelly KA, Royse C, Royse AG. Tissue Doppler Em and instantaneous end-diastolic stiffness: validation against pressure-volume loops in patients undergoing coronary artery bypass surgery. *Heart Lung Circ* 2011;20:223-230.
65. Hotvedt R, Refsum H, Platou ES. Cardiac electrophysiological and hemodynamic effects of beta- adrenoceptor blockade and thoracic epidural analgesia in the dog. *Anesth Analg* 1984;63:817-824.
66. Hirabayashi Y, Shimizu R, Fukuda H, Saitoh K, Igarashi T. Effects of thoracic vs. lumbar epidural anaesthesia on systemic haemodynamics and coronary circulation in sevoflurane anaesthetized dogs. *Acta Anaesthesiol Scand* 1996;40:1127-1131.
67. Niimi Y, Ichinose F, Saegusa H, Nakata Y, Morita S. Echocardiographic evaluation of global left ventricular function during high thoracic epidural anesthesia. *J Clin Anesth* 1997;9:118-124.
68. Shiga T. Thoracic epidural blockade preserves left ventricular early diastolic filling assessed by transesophageal echocardiography. *J Anesth* 1998;12:7-12.
69. de Vromen M, Cardozo RH, Steendijk P, van BF, Baan J. Improved contractile performance of right ventricle in response to increased RV afterload in newborn lamb. *Am J Physiol Heart Circ Physiol* 2000;278:H100-H105.
70. Wauthy P, Pagnamenta A, Vassalli F, Naeije R, Brimiouille S. Right ventricular adaptation to pulmonary hypertension: an interspecies comparison. *Am J Physiol Heart Circ Physiol* 2004;286:H1441-H1447.
71. Wink J, de Wilde RB, Wouters PF, van Dorp EL, Veering BT, Versteegh MI, Aarts LP, Steendijk P. Thoracic Epidural Anesthesia Reduces Right Ventricular Systolic Function With Maintained Ventricular-Pulmonary Coupling. *Circulation* 2016; 134:1163-1175.

72. Talbot NP, Balanos GM, Dorrington KL, Robbins PA. Two temporal components within the human pulmonary vascular response to approximately 2 h of isocapnic hypoxia. *J Appl Physiol* (1985) 2005;98:1125-1139.
73. Flu WJ, van Kuijk JP, Hoeks, Kuiper R, Schouten O, Goei D, Elhendy A, Verhagen HJ, Thomson IR, Bax JJ, Fleisher LA, Poldermans D. Prognostic implications of asymptomatic left ventricular dysfunction in patients undergoing vascular surgery. *Anesthesiology* 2010;112:1316-1324
74. Korner PI. Integrative neural cardiovascular control. *Physiol Rev* 1971;51:312-367.
75. Mor-Avi V, Lang RM, Badano LP, Belohlavek M, Cardim NM, Derumeaux G, Galderisi M, Marwick T, Nagueh SF, Sengupta PP, Sicari R, Smiseth OA, Smulevitz B, Takeuchi M, Thomas JD, Vannan M, Voigt JU, Zamorano JL. Current and evolving echocardiographic techniques for the quantitative evaluation of cardiac mechanics: ASE/EAE consensus statement on methodology and indications endorsed by the Japanese Society of Echocardiography. *Eur J Echocardiogr* 2011;12:167-205.
76. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF 3rd, Dokainish H, Edvardsen T, Flachskampf FA, Gillebert TC, Klein AL, Lancellotti P, Marino P, Oh JK, Popescu BA, Waggoner AD. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2016;29:277-314.
77. Firstenberg MS, Greenberg NL, Main ML, Drinko JK, Odabashian JA, Thomas JD, Garcia MJ. Determinants of diastolic myocardial tissue Doppler velocities: influences of relaxation and preload. *J Appl Physiol* (1985) 2001;90:299-307.
78. Zile MR, Baicu CF, Gaasch WH. Diastolic heart failure--abnormalities in active relaxation and passive stiffness of the left ventricle. *N Engl J Med* 2004;350:1953-1959.
79. Feigl EO. Parasympathetic control of coronary blood flow in dogs. *Circ Res* 1969;25:509-519.
80. Feigl EO. Neural control of coronary blood flow. *J Vasc Res* 1998;35:85-92.
81. Raizner AE, Chahine RA, Ishimori T, Verani MS, Zacca N, Jamal N, Miller RR, Luchi RJ. Provocation of coronary artery spasm by the cold pressor test. Hemodynamic, arteriographic and quantitative angiographic observations. *Circulation* 1980;62:925-932.
82. Miyashiro JK, Feigl EO. Feedforward control of coronary blood flow via coronary beta-receptor stimulation. *Circ Res* 1993;73:252-263.
83. Klocke FJ. Coronary blood flow in man. *Prog Cardiovasc Dis* 1976;19:117-166.
84. Klassen GA, Bramwell RS, Bromage PR, Zborowska-Sluis DT. Effect of acute sympathectomy by epidural anesthesia on the canine coronary circulation. *Anesthesiology* 1980;52:8-15.
85. Bulte CS, Boer C, Hartemink KJ, Kamp O, Heymans MW, Loer SA, de Marchi SF, Vogel R, Bouwman RA. Myocardial Microvascular Responsiveness During Acute Cardiac Sympathectomy Induced by Thoracic Epidural Anesthesia. *J Cardiothorac Vasc Anesth* 2017;31:134-141.
86. Thomas JD. Myocardial contrast echocardiography perfusion imaging: still waiting after all these years. *J Am Coll Cardiol* 2013;62:1362-1364.

87. Bengel FM, Ueberfuhr P, Schiepel N, Nekolla SG, Reichart B, Schwaiger M. Effect of sympathetic reinnervation on cardiac performance after heart transplantation. *N Engl J Med* 2001;345:731-738.
88. Jahn UR, Waurick R, Van Aken H, Hinder F, Booke M, Bone HG, Schmidt C, Meyer J. Thoracic, but not lumbar, epidural anesthesia improves cardiopulmonary function in ovine pulmonary embolism. *Anesth Analg* 2001;93:1460-1465.
89. Jahn UR, Waurick R, Van AH, Hinder F, Meyer J, Bone HG. Therapeutic administration of thoracic epidural anesthesia reduces cardiopulmonary deterioration in ovine pulmonary embolism. *Crit Care Med* 2007;35:2582-2586.
90. Peters J, Kutkuhn B, Medert HA, Schlaghecke R, Schuttler J, Arndt JO. Sympathetic blockade by epidural anesthesia attenuates the cardiovascular response to severe hypoxemia. *Anesthesiology* 1990;72:134-144.
91. Kumar A, Anand R, Wadhawan S, Rahal A. Effects of thoracic epidural anaesthesia on the haemodynamic response during induction and endotracheal intubation and on induction dose of propofol. *J Anaesth Clin Pharmacol* 2010;26:213-218.
92. Licker M, Farinelli C, Klopfenstein CE. Cardiovascular reflexes during anesthesia induction and tracheal intubation in elderly patients: the influence of thoracic epidural anesthesia. *J Clin Anesth* 1995;7:281-287.
93. Nakatani T, Saito Y, Sakura S, Kanata K. Haemodynamic effects of thoracic epidural anaesthesia during induction of anaesthesia: an investigation into the effects of tracheal intubation during target-controlled infusion of propofol. *Anaesthesia* 2005;60:530-534.
94. Dohi S, Nishikawa T, Ujike Y, Mayumi T. Circulatory responses to airway stimulation and cervical epidural blockade. *Anesthesiology* 1982;57:359-363.
95. Kirno K, Friberg P, Grzegorzczak A, Milocco I, Ricksten SE, Lundin S. Thoracic epidural anesthesia during coronary artery bypass surgery: effects on cardiac sympathetic activity, myocardial blood flow and metabolism, and central hemodynamics. *Anesth Analg* 1994;79:1075-1081.
96. Ottesen S. The influence of thoracic epidural analgesia on the circulation at rest and during physical exercise in man. *Acta Anaesthesiol Scand* 1978;22:537-547.
97. Tsuchida H, Omote T, Miyamoto M, Namiki A, Ichihara K, Abiko Y. Effects of thoracic epidural anesthesia on myocardial pH and metabolism during ischemia. *Acta Anaesthesiol Scand* 1991;35:508-512.
98. Fujita S, Tsuchida H, Kanaya N, Kokita N, Kawamata M, Namiki A, Ichihara K. Effects of thoracic epidural anesthesia on changes in ischemic myocardial metabolism induced by intracoronary injection of endothelin in dogs. *J Cardiothorac Vasc Anesth* 1996;10:903-908.
99. Vik-Mo H, Ottesen S, Renck H. Cardiac effects of thoracic epidural analgesia before and during acute coronary artery occlusion in open-chest dogs. *Scand J Clin Lab Invest* 1978;38:737-746.

100. Groban L, Zvara DA, Deal DD, Vernon JC, Carpenter RL. Thoracic epidural anesthesia reduces infarct size in a canine model of myocardial ischemia and reperfusion injury. *J Cardiothorac Vasc Anesth* 1999;13:579-585.
101. Davis RF, DeBoer LW, Maroko PR. Thoracic epidural anesthesia reduces myocardial infarct size after coronary artery occlusion in dogs. *Anesth Analg* 1986;65:711-717.
102. Rolf N, Van de Velde M, Wouters PF, Mollhoff T, Weber TP, Van Aken HK. Thoracic epidural anesthesia improves functional recovery from myocardial stunning in conscious dogs. *Anesth Analg* 1996;83:935-940.
103. Lagunilla J, Garcia-Bengochea JB, Fernandez AL et al. High thoracic epidural blockade increases myocardial oxygen availability in coronary surgery patients. *Acta Anaesthesiol Scand* 2006;50:780-786.
104. Blomberg S, Emanuelsson H, Ricksten SE. Thoracic epidural anesthesia and central hemodynamics in patients with unstable angina pectoris. *Anesth Analg* 1989;69:558-562.
105. Reiz S, Nath S, Rais O. Effects of thoracic epidural block and prenalterol on coronary vascular resistance and myocardial metabolism in patients with coronary artery disease. *Acta Anaesthesiol Scand* 1980;24:11-16.
106. Nabel EG, Ganz P, Gordon JB, Alexander RW, Selwyn AP. Dilation of normal and constriction of atherosclerotic coronary arteries caused by the cold pressor test. *Circulation* 1988;77:43-52.
107. Zeiher AM, Drexler H, Wollschlaeger H, Saurbier B, Just H. Coronary vasomotion in response to sympathetic stimulation in humans: importance of the functional integrity of the endothelium. *J Am Coll Cardiol* 1989;14:1181-1190.
108. Heusch G, Baumgart D, Camici P, Chilian W, Gregorini L, Hess O, Indolfi C, Rimoldi O. alpha-adrenergic coronary vasoconstriction and myocardial ischemia in humans. *Circulation* 2000;101:689-694.
109. Blomberg S, Emanuelsson H, Kvist H, Lamm C, Pontén J, Waagstein F, Ricksten SE. Effects of thoracic epidural anesthesia on coronary arteries and arterioles in patients with coronary artery disease. *Anesthesiology* 1990;73:840-847.
110. Nygard E, Kofoed KF, Freiberg J, Holm S, Aldershvile J, Eliassen K, Kelbaek H. Effects of high thoracic epidural analgesia on myocardial blood flow in patients with ischemic heart disease. *Circulation* 2005;111:2165-2170.
111. Di Carli MF, Tobes MC, Mangner T, Levine AB, Muzik O, Chakroborty P, Levine TB. Effects of cardiac sympathetic innervation on coronary blood flow. *N Engl J Med* 1997;336:1208-1215.
112. Lorenzoni R, Rosen SD, Camici PG. Effect of alpha 1-adrenoceptor blockade on resting and hyperemic myocardial blood flow in normal humans. *Am J Physiol* 1996;271:H1302-H1306.
113. Olausson K, Magnusdottir H, Lurje L, Wennerblom B, Emanuelsson H, Ricksten SE. Anti-ischemic and anti-anginal effects of thoracic epidural anesthesia versus those of conventional medical therapy in the treatment of severe refractory unstable angina pectoris. *Circulation* 1997;96:2178-2182.

114. Richter A, Cederholm I, Fredrikson M, Mucchiano C, Traff S, Janerot-Sjoberg B. Effect of long-term thoracic epidural analgesia on refractory angina pectoris: a 10-year experience. *J Cardiothorac Vasc Anesth* 2012;26:822-828.
115. Gramling-Babb P, Miller MJ, Reeves ST, Roy RC, Zile MR. Treatment of medically and surgically refractory angina pectoris with high thoracic epidural analgesia: initial clinical experience. *Am Heart J* 1997;133:648-655.
116. Kock M, Blomberg S, Emanuelsson H, Lomsky M, Stromblad SO, Ricksten SE. Thoracic epidural anesthesia improves global and regional left ventricular function during stress-induced myocardial ischemia in patients with coronary artery disease. *Anesth Analg* 1990;71:625-630.
117. Saada M, Catoire P, Bonnet F, Delaunay L, Gormezano G, Macquin-Mavier I, Brun P. Effect of thoracic epidural anesthesia combined with general anesthesia on segmental wall motion assessed by transesophageal echocardiography. *Anesth Analg* 1992;75:329-335.
118. Berendes E, Schmidt C, Van Aken H, Hartlage MG, Wirtz S, Reinecke H, Rothenburger M, Scheld HH, Schlüter B, Brodner G, Walter M. Reversible cardiac sympathectomy by high thoracic epidural anesthesia improves regional left ventricular function in patients undergoing coronary artery bypass grafting: a randomized trial. *Arch Surg* 2003;138:1283-1290.
119. Schmidt C, Hinder F, Van Aken H, Theilmeier G, Bruch C, Wirtz SP, Bürkle H, Gühs T, Rothenburger M, Berendes EH. The effect of high thoracic epidural anesthesia on systolic and diastolic left ventricular function in patients with coronary artery disease. *Anesth Analg* 2005;100:1561-1569.
120. Jakobsen CJ, Nygaard E, Norrild K, Kirkegaard H, Nielsen J, Torp P, Sloth E. High thoracic epidural analgesia improves left ventricular function in patients with ischemic heart. *Acta Anaesthesiol Scand* 2009;53:559-564.
121. Wafaa G, Ahmed AAA-A, Soad Sayed El-Gaby, Hanna Fouad Mohamed. Effect of high thoracic epidural analgesia on left ventricular function in patients with coronary artery disease undergoing elective non-cardiac surgery. *Research Journal of Cardiology* 2011;4:28-37.
122. Barrington MJ, Kluger R, Watson R, Scott DA, Harris KJ. Epidural anesthesia for coronary artery bypass surgery compared with general anesthesia alone does not reduce biochemical markers of myocardial damage. *Anesth Analg* 2005;100:921-928.
123. Kendall JB, Russell GN, Scawn ND, Akrofi M, Cowan CM, Fox MA. A prospective, randomised, single-blind pilot study to determine the effect of anaesthetic technique on troponin T release after off-pump coronary artery surgery. *Anaesthesia* 2004;59:545-549.
124. Jakobsen CJ, Bhavsar R, Nielsen DV, Ryhammer PK, Sloth E, Greisen J. High thoracic epidural analgesia in cardiac surgery. Part 1--high thoracic epidural analgesia improves cardiac performance in cardiac surgery patients. *J Cardiothorac Vasc Anesth* 2012;26:1039-1047.
125. Caputo M, Alwair H, Rogers CA et al. Myocardial, inflammatory, and stress responses in off-pump coronary artery bypass graft surgery with thoracic epidural anesthesia. *Ann Thorac Surg* 2009;87:1119-1126.

126. Palomero Rodriguez MA, Suarez GL, Villar AF, Varela CC, Moreno GL, Criado JA. Thoracic epidural anesthesia decreases C-reactive protein levels in patients undergoing elective coronary artery bypass graft surgery with cardiopulmonary bypass. *Minerva Anesthesiol* 2008;74:619-626.
127. Stenger M, Fabrin A, Schmidt H, Greisen J, Erik MP, Jakobsen CJ. High thoracic epidural analgesia as an adjunct to general anesthesia is associated with better outcome in low-to-moderate risk cardiac surgery patients. *J Cardiothorac Vasc Anesth* 2013;27:1301-1309.
128. Scott NB, Turfrey DJ, Ray DA, Nzewi O, Sutcliffe NP, Lal AB, Norrie J, Nagels WJ, Ramayya GP. A prospective randomized study of the potential benefits of thoracic epidural anesthesia and analgesia in patients undergoing coronary artery bypass grafting. *Anesth Analg* 2001;93:528-535.
129. Beattie WS, Badner NH, Choi P. Epidural analgesia reduces postoperative myocardial infarction: a meta-analysis. *Anesth Analg* 2001;93:853-858.
130. Svircevic V, van Dijk D, Nierich AP, Passier MP, Kalkman CJ, van der Heijden GJ, Bax L. Meta-analysis of thoracic epidural anesthesia versus general anesthesia for cardiac surgery. *Anesthesiology* 2011;114:271-282.