



Universiteit
Leiden
The Netherlands

The use of eHealth in rehabilitation after stroke

Wentink, M.M.

Citation

Wentink, M. M. (2019, October 15). *The use of eHealth in rehabilitation after stroke*. Retrieved from <https://hdl.handle.net/1887/79515>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/79515>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/79515> holds various files of this Leiden University dissertation.

Author: Wentink, M.M.

Title: The use of eHealth in rehabilitation after stroke

Issue Date: 2019-10-15

The use of eHealth in rehabilitation after stroke

Manon Wentink

ISBN 978-94-6375-507-8

Cover by Nettl, Leiderdorp

Layout by Manon Wentink and Jos Hendrix

Printed by Ridderprint BV, www.ridderprint.nl

The studies in this thesis were financially supported by Revalidatiefonds Nederland, grant nr. R2011184 and Fonds NutsOhra, grant nr. 1202-006 (Chapters 2 and 3) and Nationaal Regieorgaan Praktijkgericht Onderzoek SIA, grant nr.2014-01-46PRO (Chapters 4, 5 and 6).

Printing of this thesis was financially supported by Basalt and the Vereniging voor Oefentherapeuten Cesar en Mensendieck.

© Manon Wentink, 2019

The use of eHealth in rehabilitation after stroke

Proefschrift

ter verkrijging van

de graad van Doctor aan de Universiteit Leiden,

op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker,

volgens besluit van het College voor Promoties te verdedigen op

dinsdag 15 oktober 2019 klokke 11.15 uur

door

Manon Maria Wentink

geboren te Noorder-Koggenland
in 1987

Promotiecommissie

Promotores: Prof. dr. T.P.M. Vliet Vlieland

Co-promotores: Dr. J.J.L. Meesters
Dr. A.J. de Kloet

Overige leden: Prof. dr. A.W.M. Evers (Leids Universitair Medisch Centrum)
Dr. J. Verhoef (Hogeschool Leiden)
Prof. dr. J.M.A. Visser-Meily (Universitair Medisch Centrum, Utrecht)

Contents

Chapter 1.	General introduction	7
Chapter 2.	The effects of an 8-week computer-based brain training programme on cognitive functioning, QoL and self-efficacy after stroke <i>Neuropsychological Rehabilitation 2016;26(5-6):847-865</i>	23
Chapter 3.	Adherence of stroke patients with an online brain training program: the role of health professionals' support <i>Topics in Stroke Rehabilitation 2018;25(5):359-365</i>	47
Chapter 4.	The patient perspective on the use of information and communication technologies and e-health in rehabilitation <i>Disability and Rehabilitation: Assistive Technology 2018;13(7):620-625</i>	63
Chapter 5.	How to improve eRehabilitation programs in stroke care? A focus group study to identify requirements of end-users. <i>BMC Medical informatics and Decision Making 2019;19(1):145.</i>	79
Chapter 6.	What is important in e-health interventions for stroke rehabilitation? A survey study among patients, informal caregivers and health professionals. <i>International journal of telerehabilitation 2018;10(1):15-28</i>	101
Chapter 7.	Teachers' and students' perceptions about eHealth education in the curriculum of exercise and physical therapy: a focus groups study on barriers and facilitators. <i>Accepted: BMC Medical Education (London)</i>	121
Chapter 8.	Summary and general discussion	137
Appendix		
	Samenvatting en discussie	157
	Publications	169
	Curriculum Vitae	171
	Dankwoord	173



General introduction



Definition and epidemiology

Stroke, or a cerebrovascular accident (CVA), occurs when a clot in the blood vessel blocks the blood flow to the brain cells (ischemic stroke) or when a blood vessel in the brain breaks or ruptures (hemorrhagic stroke). Subsequently, brain cells are deprived of oxygen and glucose, causing damage to the brain tissues [1]. Abilities controlled by these brain tissues, such as memory and/or speech control, can be lost. As a consequence, persons who survive a stroke can face long-term disability with a considerable impact on their lives and those of their relatives [2].




Figure 1. Images of the occurrence of an ischemic and hemorrhagic stroke [2].

After ischemic heart disease, stroke is the second leading cause of death in the world [3]. According to the World Stroke Organization (WSO), 1 in 6 people worldwide will have a stroke in their lifetime, 15 million people worldwide suffer a stroke each year and 5.8 million people die from it [4]. In the Netherlands, the incidence of stroke was estimated to be 42.300 in 2016 and the prevalence in that year was estimated to be 320.000 in the community based population [5].

Overall, stroke incidence has mainly shown either declining time trends or stable rates in high income countries [6]. On the other hand, the absolute number of stroke is expected to increase because of the ageing population and declining mortality rates, largely as a result of better control of high blood pressure and faster and better treatment (e.g. intravenous thrombolytic therapy and stroke services) [7;8]. Consequently, the overall stroke burden across the globe, in terms of the number of people affected by or who remain disabled from stroke, has increased [9]. Estimates from the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD 2010) ranked stroke as the disease with the third-highest burden in disability, expressed in disability-adjusted life-years (DALY's), and the burden is projected to further rise from around 38 million DALYs globally in 1990 to 61 million DALYs in 2020 [10].

Consequences of stroke regarding functioning, disability and health

Stroke can have a broad and significant impact on a persons’ physical, cognitive and social functioning. Especially cognitive impairment is common after stroke. A cohort study among 395 patients in the Netherlands showed that more than half of the stroke patients (66 percent) suffer from cognitive impairment two months post-stroke [12]. Although improvement is seen at six months post-stroke (prevalence of 50 percent), cognitive impairment still remains highly prevalent up to 6 months after stroke [12]. This often causes lifelong problems in activities of daily life and participation in society [11].

The International Classification of Functioning, Disability and Health (ICF) of the World Health Organization (WHO) is a helpful framework to classify a person’s functioning and disability and plan delivery of care (Figure 1) [13]. The ICF is based on the biopsychosocial model and provides a coherent view of different perspectives of health: biological, individual and social. As the diagram indicates, in the ICF disability and functioning are viewed as outcomes of interactions between *health conditions* (diseases, disorders and injuries) *personal* [14;15] and *contextual factors* [13]. Because the ICF is very extensive, disease specific core sets (ICF Core Sets) have been developed for a number of chronic health conditions, including stroke, in order to provide an overview of categories of post-stroke disability [16]. The Brief ICF Core Set represents a selection of ICF domains or categories which can serve as a minimal standard for reporting of functioning and health in clinical studies and clinical encounters (Table 1-3) [17].




Figure 2. The model of the International Classification of Functioning, Disability and Health (ICF) shows the relationship between the different ICF components [18].

Table 1. International Classification of Functioning, Disability and Health (ICF) – categories of the components included in the Brief ICF Core Set for Stroke.

Body functions
b110 Consciousness functions
b114 Orientation functions
b140 Attention functions
b144 Memory functions
b167 Mental functions of language
b730 Muscle power functions
Body structures
s110 Structure of brain
s730 Structure of upper extremity
Activities and participation
d310 Communicating with – receiving – spoken messages
d330 Speaking
d450 Walking
d510 Washing oneself
d530 Toileting
d540 Dressing
d550 Eating
Environmental factors
e310 Immediate family
e355 Health professionals
e580 Health services, systems and policies

Medical management after stroke

Each year approximately 9.600 people in the Netherlands die because of stroke [19]. Those who survive a stroke are treated with acute care in a hospital (intensive care, medium care, stroke unit and/or neurology ward). The stay in the hospital is generally short and ends when the patient is medically stable [20]. Afterwards, about 60-65% is discharged to their homes. If needed, treatment can be provided by health professionals close to home (primary care) or in a rehabilitation center (outpatient care) [21]. For 25-30% of the patients, mostly older patients with multiple impairments, a nursing home for specialized geriatric rehabilitation or long stay is required. Ten percent, often young patients with potential for recovery and higher participation goals, are eligible for rehabilitation in a specialized rehabilitation center [22]. A flow of the patient journey after stroke is presented in Figure 3.




Figure 3. Patient journey after stroke describing the possible pathways of a stroke patient through the healthcare system in the Netherlands.

Specialized medical rehabilitation after stroke

Approximately 3.200 patients in the Netherlands use specialized medical rehabilitation stroke care yearly [22]. Medical specialist stroke rehabilitation is a comprehensive, multi-dimensional process including interventions that aim to facilitate restitution or substitution of limitations in impairment, activity or participation caused by stroke [23]. Rehabilitation typically follows a process of assessment of needs, goal setting, therapy and reassessment [24]. This process involves the patient, informal caregivers and various health care providers (e.g. physicians, physical and occupational therapists, speech-language therapists, psychologists and social worker). For each patient, a tailored treatment plan is defined, dependent on the type and severity of the impairments and the patients' goals. The length of these clinical pathways differs from 3 till 26 weeks [20]. After the rehabilitation period, about 90% of the patients are able to live independently [21]. The costs related to medical specialized rehabilitation were estimated at €147 million in 2015 [25]. The general health care costs for stroke were estimated at nearly €2.3 billion euros in 2011, 2.5% of the total healthcare costs [26]. As a result, stroke is one of ten most expensive diseases in the Netherlands.

eHealth in rehabilitation care (eRehabilitation)

Information and Communication Technology (ICT) is increasingly used in health care in general (eHealth) and in rehabilitation (eRehabilitation). eRehabilitation is applied for a number of (potential) purposes: 1) support of self-management and self-ownership [27], 2) improving access for (stroke) patients to health care services as well as bridging between services [28;29;30], 3) increasing the doses of treatment and more effective distribution

in time [27], 4) monitoring compliance and progress [27], 5) structuring treatment and collecting treatment related data, 6) offering opportunities for more cost-effective interventions in the future and coping with ageing society [31;32] and 7) fulfilling a need in the digitizing society, patients and health care. With the different goals for eHealth, the nature of the technology employed in health care may vary largely, and there is a demand for their classification. An example of such a classification is published in the 'Co-creation eHealthbook' (Table 2) [33].

Table 2. Classification of eHealth applications for each technology [Wentzel, 2014; translated from Dutch].

Technology	Explanation
<i>Web applications and portals</i>	Applications that are offered to users via a web browser such as patient portals or educational portals for health professionals.
<i>Mobile applications</i>	Applications that are offered via a mobile device such as a smartphone or tablet.
<i>Electronic patient records</i>	Systems of a mostly medical-administrative nature in which healthcare providers register medical patient data within their own care organization.
<i>Health-sensors and wearable devices</i>	A category of devices that are often used in the home situation of patients to measure (vital body) functions, collect the results and possibly pass them on to a health professional.
<i>Video communication (telecommunication)</i>	Applications in which a visual dimension is added to the usual forms of telecommunication with the aim to strengthen contact between the care recipient and the care provider and to increase the communication possibilities.
<i>Domotics</i>	A general term for the application of electronics for home automation. It is often a combination of environment-aware sensors and actuators (devices that can influence the environment) with which the living environment in a home can be regulated or things in the home can be automatically operated.
<i>Robotics</i>	Machines that can perform and take over certain tasks such as offering structure in the day.
<i>Medical integration networks</i>	Electronic networks on which medical information is exchanged, such as medication data and prescriptions or radiological images.
<i>General integration networks</i>	Electronic networks of a more general nature for the exchange of data between cooperating (business) partners, for example about orders.
<i>Business intelligence and big data</i>	Business intelligence systems are focused on analyzing structured and unstructured data to provide information that can be used for decision support. If this is done within the field of care, this is called 'medical intelligence'. An analysis of very large amounts of data from different sources is called 'big data'.
<i>Serious gaming</i>	Use of game technology by applying a game element to 'serious matters', such as exercises in the context of psychological or physical treatment.

In addition, Krijgsman et al. 2012 describes three dimension to structure eHealth initiatives: 1) *the setting of the care process in which the eHealth application is used* (e.g. in the public health domain, as part of a care process, or in the support of a care process); 2) *the persons who use the eHealth application* (e.g. for health care professionals, for communication between patient and health care professional or patients between themselves; 3) *the technological basis of the eHealth application* (e.g. which platform or application type) [34].

Application of eRehabilitation after stroke

An established application of eHealth in stroke care management is *video communication*. Video patient-practitioner conferences are recommended in guidelines for acute stroke management in order to enroll a greater number of patients in therapies (e.g. correct diagnosis, promptly decide thrombolytic therapy and guide transfers to appropriate centers) [28;35]. More recently, video communication is applied for continuity of care [28]. Interactive patient-practitioner video consultations, as well as web portals, are used for both medical and educational services and stroke secondary prevention and monitoring [28].

In addition, 'off the shelf' *mobile and web applications* (e.g. exercises to train cognitive, speech and/or physical functioning) [36;37], *video games* (e.g. Nintendo Wii and Balance Board, Microsoft Kinect, etc.) [38] and *health sensors* (e.g. Fit-Bit, smartwatch, etc.) [39], developed for commercial purposes, are applied for stroke rehabilitation. Several applications have been developed for therapy purposes in general (e.g. Physitrack, Minddistrict, etc.) or for stroke specific (i.e. Care4Stroke, etc.). These technologies, mostly used on the initiative of patients and practitioners as a supplement to traditional care [28], are being applied in order to increase doses of treatment, monitor compliance and/or improve rehabilitation outcomes. *Robotics*, like exoskeleton types, also deliver high-intensity training for stimulation of motor disorders caused by stroke [40].

Furthermore, *electronic patient records* and *general/medical integration networks* have facilitated data-collection and data-analysis of aspects of stroke care in past years (e.g. team response time and time to other specific aspects of the care protocol, treatment outcomes, etc.). Registering accurate data and continue comparing data with other medical centers will further improve quality of stroke care and assist to identify patients who are eligible for clinical trials in the stroke population in the future [28].

Evidence for effectiveness of stroke eRehabilitation

This thesis concerns research on cognitive and comprehensive eRehabilitation interventions in patients with stroke. In general, in medical specialist rehabilitation, comprehensive multidisciplinary eRehabilitation that covers multiple aspects of stroke management (e.g. physical and cognitive functioning) is highly recommended and might become available in the future [28], but is not yet studied in clinical trials.

The effects of cognitive eHealth interventions on cognitive functioning in stroke patients are summarized in one meta-analysis of Laver et al. (2017) [41]. The aim of this meta-analysis was to determine the efficacy of virtual reality compared with an alternative intervention or no intervention on multiple outcome measures (i.e. upper limb function and activity, gait and balance, global motor function, activity limitation, participation restriction, quality of life and adverse events), including cognitive functioning. They searched different databases (e.g. Cochrane Stroke Group Trials Register (April 2017), CENTRAL, MEDLINE, Embase), trials registries and reference lists. A total of 72 studies (n=2470) were included, from which only three studies evaluated the effect of virtual reality interventions on cognition. These three trials [42;43;44] did not allow analysis, so that there are no results available on the effect of virtual reality on cognitive functioning. Other meta-analysis or systematic reviews of eHealth interventions on cognitive functioning in stroke patients are absent.

However, recently two RCTs addressing this question were performed [45;46]. Faria et al. (2016) studied the potential benefits of virtual reality based cognitive rehabilitation after stroke through simulated activities of daily living as compared to conventional therapy only [45]. The intervention involved a virtual simulation of a city where memory, attention, visuo-spatial abilities and executive functions tasks are integrated in the performance of several daily routines. A between groups analysis showed significantly greater improvements in global cognitive functioning, attention and executive functions in patients with stroke when comparing virtual reality to conventional therapy.

Van de Ven et al. (2017) investigated whether computer-based cognitive flexibility training improved subjective cognitive functioning and quality of life after stroke [46]. In the trial, adults (30±80 years old) who had suffered a stroke within the last 5 years were assigned to either an intervention group (n = 38), active control group (i.e., mock training; n=35), or waiting list control group (n=24). It was found both training groups improved on training tasks and all groups improved on executive functioning tasks, attention, reasoning, and psychomotor speed. However, the amount of improvement in executive and general cognitive functioning in the intervention group was similar to that of both control groups (active control and waiting list). Therefore, this improvement was likely due to training-unspecific effects. Overall, there is a lack of studies assessing the effect of eHealth intervention for cognitive functioning and quality of life after stroke.

Uptake of stroke eRehabilitation

Uptake of eRehabilitation can be described as a planned process and systematic introduction of eRehabilitation with the aim that it is given a structural place in rehabilitation care [47]. According to the theory of Grol and Wensing (2004), implementation can be influenced by several barriers and facilitators at different levels: Innovation, Individual health care professional, Individual Patient, Social context, Organizational context and Political and economic context [48]. For instance, for eRehabilitation a potential barrier at the level of the Innovation might be the variability in resources between different virtual rehabilitation networks [28]. Moreover, eHealth interventions often do not match with the requirements of intended users (e.g. patients and health care professionals), impairing the adoption of eHealth interventions [49;50]. At the level of the Individual health care professional, there are barriers to technology adoption related to the absence of a composite set of knowledge and skills among health care professionals regarding the use of eHealth [51]. Subsequently, there is a demand for new knowledge, skills and attributes among (future) health care professionals and new initiatives in education of health professions [52].

Outline of this thesis

Given the lack of knowledge the aims of this thesis are to:

- I. Evaluate the outcome and process of an eHealth intervention for cognitive stroke rehabilitation.
- II. Explore the (readiness for) use of eHealth among patients in rehabilitation.
- III. Investigate the requirements of patients, informal caregivers, health professionals, teachers and students regarding the use of eHealth interventions in stroke rehabilitation.

Chapter 2 presents a randomized controlled trial (RCT) to determine the effect of an online serious brain training programme on multiple aspects of cognitive functioning in comparison to a control intervention in patients with self-perceived cognitive impairments 12–36 months after stroke (aim I).

Chapter 3 describes the adherence of patients with the online brain training programme by comparing two types of health professionals' supervision (aim I).

In *Chapter 4* the perspective of patients on use of eHealth in rehabilitation care is described by exploring their usage of common devices and investigating their preferences regarding usage of these devices in rehabilitation care (aim II).

Chapter 5 describes the results of a qualitative study, in which the requirements of patients with stroke, their informal caregivers and health professionals for the accessibility, usability

and content of comprehensive eRehabilitation after stroke were identified (aim III). The identified requirements were prioritized in *Chapter 6* by using a quantitative study design in order to assess which requirements found in the qualitative research are most important (aim III).

Chapter 7 provides an overview of the requirements (barriers and facilitators) according to students and teachers for the uptake of eHealth in curricula of two allied health professions, based on qualitative focus groups (aim III).

In *Chapter 8* a discussion is conducted on/elaborates the overall findings and recommendations are provided.

Reference list

1. World Health Organization. *Health topics - Stroke, Cerebrovascular accident*. https://www.who.int/topics/cerebrovascular_accident/en/ (accessed 21 January 2019).
2. Brain Foundation of the Netherlands [in Dutch: Hersenstichting]. *Beroerte*. <https://www.hersenstichting.nl/alles-over-hersenen/hersenaandoeningen/beroerte> (accessed 21 January 2019).
3. World Health Organization. *The top 10 causes of death*. <https://www.who.int/en/news-room/fact-sheets/detail/the-top-10-causes-of-death> (accessed 21 January 2019).
4. World Stroke Organization. *Facts and Figures about Stroke*. <https://www.world-stroke.org/component/content/article/16-forpatients/84-facts-and-figures-about-stroke> (accessed 21 January 2019).
5. National Institute for Public Health and the Environment [in Dutch: Rijksinstituut voor Volksgezondheid en Milieu (RIVM)]. *Beroerte, Cijfers & Context, Huidige situatie - Prevalentie en nieuwe gevallen van beroerte*. <https://www.volksgezondheidenzorg.info/onderwerp/beroerte/cijfers-context/huidige-situatie#node-prevalentie-en-nieuwe-gevallen-van-beroerte> (accessed 21 January 2019).
6. Thrift AG, Thayabaranathan T, Howard G, Howard VJ, Rothwell PM, Feigin VL, Norrving B, Donnan GA, Cadilhac DA. Global stroke statistics. *International Journal of Stroke*. 2017;12(1):13-32.
7. Schweikert A, Kahles T, Nedeltchev K. Intravenous Thrombolysis for Acute Ischemic Stroke. In *Stroke Revisited: Diagnosis and Treatment of Ischemic Stroke 2017* (pp. 49-58). Springer, Singapore.
8. Kunst AE, Amiri M, Janssen F. The decline in stroke mortality: exploration of future trends in 7 Western European countries. *Stroke*. 2011;42(8):2126-30.
9. Feigin VL, Norrving B & Mensah, GA. Global burden of stroke. *Circulation research*. 2017;120(3):439-448.
10. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, Moran AE, Sacco RL, Anderson L, Truelsen T, O'Donnell M. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *The Lancet*. 2014;383(9913):245-55.
11. Nijssse B, Visser-Meily JM, van Mierlo ML, Post MW, de Kort PL, van Heugten CM. Temporal evolution of poststroke cognitive impairment using the montreal cognitive assessment. *Stroke*. 2017;48(1):98-104.
12. Cumming TB, Brodtmann A, Darby D, Bernhardt J. The importance of cognition to quality of life after stroke. *Journal of psychosomatic research*. 2014;77(5):374-9.
13. World Health Organization. *Classifications - ICF Application and Training Tools*. <http://www.who.int/classifications/icf/icfaptraining/en/index.html> (accessed 21 January 2019).
14. Verberne DPJ1,2, Post MWM3,4, Köhler S1, Carey LM5,6, Visser-Meily JMA3,7, van Heugten CM1,2,8. Course of Social Participation in the First 2 Years After Stroke and Its Associations With Demographic and Stroke-Related Factors. *Neurorehabil Neural Repair*. 2018;32(9):821-833.
15. van Mierlo ML1, Schröder C, van Heugten CM, Post MW, de Kort PL, Visser-Meily JM. The influence of psychological factors on health-related quality of life after stroke: a systematic review. *Int J Stroke*. 2014;9(3):341-8.
16. Geyh S, Cieza A, Schouten J, Dickson H, Frommelt P, Omar Z, Kostanjsek N, Ring H, Stucki G. ICF Core Sets for stroke. *Journal of rehabilitation medicine*. 2004;36(0):135-41.
17. World Health Organization - ICF Research Branch. *Brief ICF Core Set for Stroke*. <https://www.icf-research-branch.org/download/send/12-cardiovascularandrespiratoryconditions/198-brief-icf-core-set-stroke-18categ> (accessed 21 January 2019).
18. World Health Organization. *International Classification of Functioning, Disability and Health: ICF*. Geneva: WHO; 2001.
19. National Institute for Public Health and the Environment [in Dutch: Rijksinstituut voor Volksgezondheid en Milieu (RIVM)]. *Beroerte, Cijfers & Context, Sterfte - Sterfte beroerte naartype*. <https://www.volksgezondheidenzorg.info/onderwerp/beroerte/cijfers-context/sterfte#node-sterfte-beroerte-naartype> (accessed 21 January 2019).
20. Groeneveld IF, Proper KI, Van der Beek AJ, Van Mechelen W. Sustained body weight reduction by an individual-based lifestyle intervention for workers in the construction industry at risk for cardiovascular disease: results of a randomized controlled trial. *Preventive medicine*. 2010;51(3-4):240-246.

21. Rehabilitation Netherlands [in Dutch: Revalidatie Nederland]. *Revalidatie Factsheet - Revalidatie na een beroerte*. <https://www.revalidatie.nl/userfiles/File/publicaties/factsheet-beroerte-2012.pdf> (accessed 21 January 2019).
22. Poos MJJC, Blokstra A, van der Noordt M. Hoeveel zorg gebruiken patiënten met beroerte en wat zijn de kosten? [Healthcare use and expenses by stroke patients.] [Internet] Bilthoven: RIVM; 2014 [updated 2014 Jun 3; cited 2015 Sep 15]. Available from: <http://www.nationaalkompas.nl/gezondheid-en-ziekte/ziektenen-aandoeningen/hartvaatstelsel/beroerte/welke-zorg-gebruikenpatiënten-en-kosten/>.
23. Clark B, Whittall J, Kwakkel G, Mehrholz J, Ewings S, Burrage J. Time spent in rehabilitation and effect on measures of activity after stroke. *Cochrane Database of Systematic Reviews*. 2017(3).
24. Langhorne P, Bernhardt J, Kwakkel G. Stroke rehabilitation. *The Lancet*. 2011 May 14;377(9778):1693-702.
25. National Institute for Public Health and the Environment [in Dutch: Rijksinstituut voor Volksgezondheid en Milieu (RIVM)]. *StatLine - Kosten van ziekten 2015*. <https://statline.rivm.nl/#/RIVM/nl/dataset/50040NED/table?graphType=Table&ts=1512975518824> (accessed 21 January 2019).
26. National Institute for Public Health and the Environment [in Dutch: Rijksinstituut voor Volksgezondheid en Milieu (RIVM)]. *Kosten van zorg voor beroerte*. <https://www.volksgezondheidenzorg.info/onderwerp/beroerte/kosten/kosten#node-kosten-van-zorg-voor-beroerte> (accessed 21 January 2019).
27. Nam HS, Park E, Heo JH. Facilitating stroke management using modern information technology. *Journal of stroke*. 2013 Sep;15(3):135.
28. Dumitrascu OM, Demaerschalk BM. Telestroke. *Current cardiology reports*. 2017;19(9):85.
29. Tenforde AS, Hefner JE, Kodish-Wachs JE, Iaccarino MA, Paganoni S. Telehealth in physical medicine and rehabilitation: a narrative review. *PM&R*. 2017 May 1;9(5):S51-8.
30. Dorsey ER, Topol EJ. State of telehealth. *New England Journal of Medicine*. 2016;375(2):154-61.
31. Switzer J, Demaerschalk B, Xie J, Fan L, Villa K, Wu E. Cost-Effectiveness of Hub-and-Spoke Telestroke Networks for the Management of Acute Ischemic Stroke (P05. 243). *Neurology*. 2012 78(1 Supplement):05-243.
32. Nelson RE, Okon N, Lesko AC, Majersik JJ, Bhatt A, Baraban E. The cost-effectiveness of telestroke in the Pacific northwest region of the USA. *J Telemed Telecare*. 2016;22(7):413-21.
33. Wentzel MJ, Beerlage-de Jong N, Sieverink F, et al. Slimmer eHealth ontwikkelen en implementeren met de CeHRes Roadmap. In: Politiek C, Hoogendijk R, editors. CO-CREATIE EHEALTHBOEK eHealth, technisch kunstje of pure veranderekunde? The Netherlands: Nictiz; 2014.
34. Krijgsman J, Peeters J, Waverijn G, Lettow B, van der Hoek L, de Jong J, Friele R, van Gennip L. *Rapportage eHealth-doelstellingen 2016*. Den Haag: Nictiz, Utrecht: Nivel.
35. Jauch EC, Saver JL, Adams Jr HP, Bruno A, Connors JJ, Demaerschalk BM, Khatri P, McMullan Jr PW, Qureshi AI, Rosenfield K, Scott PA. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2013;44(3):870-947.
36. Singer J, Levine SR. Stroke and technology: prescribing mHealth apps for healthcare providers, patients and caregivers—a brief, selected review. *Future Neurology*. 2016;11(2):109-112.
37. Pugliese M, Ramsay T, Johnson D, Dowlathshahi D. Mobile tablet-based therapies following stroke: A systematic scoping review of administrative methods and patient experiences. *PloS one*. 2018;13(1):e0191566.
38. Ranjan A, Joshi P, Kalore S, Mithari P. Effects Based on Serious Gaming for Rehabilitation. *International Research Journal of Engineering and Technology*. 2017;4:1737-1740.
39. Li HT, Huang JJ, Pan CW, Chi HI, Pan MC. Inertial sensing based assessment methods to quantify the effectiveness of post-stroke rehabilitation. *Sensors*. 2015;15(7):16196-209.
40. Chang, W. H., & Kim, Y. H. (2013). Robot-assisted therapy in stroke rehabilitation. *Journal of stroke*, 15(3), 174.
41. Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. *Stroke*. 2018;49(4):e160-1.
42. Piemonte MEP, Oliveira TP, Miranda C, Muzzi J, Perez DB. Motor-cognitive intervention based on Nintendo Wii Fit games to improve balance and cognitive functions in patients with stroke: a randomized controlled trial. *Personal communication (to be presented at the Neurorehabilitation Congress) 2014*.
43. NCT02857803. A randomised controlled trial comparing the impact of virtual reality, paper and pencil and conventional methods on stroke rehabilitation
44. NCT02553993. Comparing the cognition effects of two exergame systems and traditional weight shifting training in patients with chronic stroke: a pilot randomized comparison trial

45. Faria AL, Andrade A, Soares L, i Badia SB. Benefits of virtual reality based cognitive rehabilitation through simulated activities of daily living: a randomized controlled trial with stroke patients. *Journal of neuroengineering and rehabilitation*. 2016;13(1):96.
46. Van De Ven RM, Buitenweg JI, Schmand B, Veltman DJ, Aaronson JA, Nijboer TC, Kruiper-Doesborgh SJ, van Bennekom CA, Rasquin SM, Ridderinkhof KR, Murre JM. Brain training improves recovery after stroke but waiting list improves equally: A multicenter randomized controlled trial of a computer-based cognitive flexibility training. *PloS one*. 2017;12(3):e0172993.
47. The Netherlands Organisation for Health Research and Development [in Dutch: Nederlandse organisatie voor gezondheidsonderzoek en zorginnovatie (ZonMw)]. Met het oog op toepassing; beleidsnota implementatie ZON 1997-1999. Den Haag. 1997 in ZonMw. *Evaluatie Implementatiemonitor ZonMw - Onderzoek naar de relatie tussen implementatieplannen en hun uitkomsten*. <https://www.nivel.nl/sites/default/files/bestanden/Implementatiemonitor-ZonMw.pdf> (accessed 21 January 2019).
48. Grol R, Wensing M: What drives change? Barriers to and incentives for achieving evidence-based practice. *Med J Aust* 2004, 180: S57-S60.
49. Hamid A, Sarmad A. Evaluation of e-health services: user's perspective criteria. *Transform Gov* 2008;2(4):243-255.
50. Esser PE, Goossens RH. A framework for the design of user-centred teleconsulting systems. *J Telemed Telecare* 2009;15(1):32-39.
51. Barakat A, Woolrych RD, Sixsmith A, Kearns WD, Kort HS. eHealth technology competencies for health professionals working in home care to support older adults to age in place: outcomes of a two-day collaborative workshop. *Medicine 2.0*. 2013;2(2).
52. Kaljouw M, Van Vliet K. Naar nieuwe zorg en zorgberoepen: de contouren. Diemen, Zorginstituut Nederland. 2015.



The effects of an 8-week computer-based brain training programme on cognitive functioning, QoL and self-efficacy after stroke

M.M. Wentink
M.A.M. Berger
A.J. de Kloet
J.J.L. Meesters
G.P.H. Band
R. Wolterbeek
P.H. Goossens
T.P.M. Vliet Vlieland

Neuropsychological Rehabilitation 2016;26:5-6, 847-865.
Poster presentation: World Congress International Brain Injury
Association, 20 March 2014, San Francisco, USA.



Abstract

Cognitive impairment after stroke has a direct impact on daily functioning and quality of life (QoL) of patients and is associated with higher mortality and healthcare costs. The aim of this study was to determine the effect of a computer-based brain training programme on cognitive functioning, QoL and self-efficacy compared to a control condition in stroke patients. Stroke patients with self-perceived cognitive impairment were randomly allocated to the intervention or control group. The intervention consisted of an 8-week brain training programme (Lumosity Inc.[®]). The control group received general information about the brain weekly. Assessments consisted of a set of neuropsychological tests and questionnaires. In addition, adherence with trained computer tasks was recorded. No effect of the training was found on cognitive functioning, QoL or self-efficacy when compared to the control condition, except for very limited effects on working memory and speed. This study found very limited effects on neuropsychological tests that were closely related to trained computer tasks, but no transfers to other tests or self-perceived cognitive failures, QoL or self-efficacy. These findings warrant the need for further research into the value of computer-based brain training to improve cognitive functioning in the chronic phase after stroke.

Introduction

The crude incidence of stroke in Western countries in the past decades is estimated to be 1.14 cases per 1000 persons per year for first ever strokes and 3.5 cases per 1000 persons per year for all strokes (Zhang et al., 2012). About 20–50% of the patients die within the first month (Truelsen et al., 2006). For the patients who survive, the disease impact is considerable, with the estimated proportions of patients with cognitive impairments being 50% directly after stroke (Douiri, Rudd, & Wolfe, 2013) and about 30% three years after a first stroke (Patel, Coshall, Rudd, & Wolfe, 2003). Cognitive impairment has a direct impact on daily functioning and quality of life (QoL) of patients and their relatives (Paker, Buğdaycı, Tekdöş, Kaya, & Dere, 2010). It is also associated with greater rates of institutionalisation (Pasquini, Leys, Rousseaux, Pasquier, & Henon, 2007), higher mortality (Tatemichi et al., 1994) and higher healthcare costs (Claesson, Lindén, Skoog, & Blomstrand, 2005).

Cognitive rehabilitation after stroke is usually based on specific learning strategies with differences between compensatory approaches: (1) focused on learning strategies to improve performance on cognitive tasks and restitution-based approaches, or (2) focused on training and stimulation of impaired cognitive functions for recovery. Among this last approach, computer-based cognitive rehabilitation (CBCR) has been shown to significantly improve performance on cognitive tasks assessing the specific domains trained within CBCR programmes (Lynch, 2002; Owen et al., 2010; Smith et al., 2009). CBCR interventions are based on principles that the impaired system(s) can be restored or at least improved by structured drills and practice using tasks that contain similar elements to the target skill (e.g., attention span, reaction time) (Lynch, 2002). A widely used CBCR intervention in stroke patients is CogMed®, a training targeting the cognitive domain of working memory (WM) by games aimed at remembering multiple stimuli at the same time, during short delays and in a unique order. Other examples of CBCR interventions are BrainGymmer® and Lumosity®, both designed to improve people's general cognitive performance with various games and mental exercises in multiple cognitive domains.

Regarding the effectiveness of CBCR interventions after stroke, several randomised controlled trials (RCTs) have been published in the last decade. Three RCTs specifically examined the effect of a computer-based attention (Barker-Collo et al., 2009) or WM training (Cogmed) (Åkerlund, Esbjörnsson, Sunnerhagen, & Björkdahl, 2013; Westerberg et al., 2007) as compared to standard care and no intervention. In Barker-Collo et al. (2009), 78 stroke patients received 30 hours of computerised attention training (intervention group) or standard care alone (control group) 6 months after stroke. Statistically significant effects were seen regarding a test combining visual and auditory attention scores, either separately or combined, but this was not reflected in statistically significant improvement on other attention measures, such as the Trail Making Task A and B, or in wider outcomes.

Westerberg et al. (2007) concluded that the treatment group improved significantly more than the passive control group on tests that measured WM and subjective cognitive failures. Effect sizes between groups were respectively 0.83 on the Span Board ($p = .05$), 1.58 on the Digit Span ($p < .01$), and 0.80 on the Cognitive Failures Questionnaire (CFQ) ($p < .01$). Åkerlund et al. (2013) also showed that results on the Digit span after WM training differed significantly compared to the results of usual rehabilitation at the same time, with greater improvement in the training group (Digit Span forward $p = .04$, Digit Span backward $p < .01$).

A review by Laver, George, Thomas, Deutsch, and Crotty (2015) pointed out that it is still unclear how effective virtual reality may be for cognitive rehabilitation after stroke. Akinwuntan et al. (2005) studied the effect on fitness to drive of a 5-week 15-hour simulator-based training (experimental group) compared to a driving-related cognitive tasks (control group) among 83 stroke patients. At 6–9 months post-stroke, significant differences between both groups were found in “road sign recognition” and “fitness to drive” in favour of the experimental subjects.

Overall the interpretation of the results of RCTs is hampered by relatively small sample sizes with 18, 47, and 78 participants in total (Åkerlund et al., 2013; BarkerCollo et al., 2009; Westerberg et al., 2007). Moreover, none of these trials included a training focused on multiple cognitive domains, although the literature suggests that targeting a variety of cognitive functions is needed to promote generalisation of abilities beyond the trained task and to achieve an outcome that also impacts on daily life (Buitenweg, Murre, & Ridderinkhof, 2012; Green & Bavelier, 2008; Sohlberg & Mateer, 2001). In addition, the effects on QoL have not been studied (Barker-Collo et al., 2009). Thus, there is a need for large studies focusing not only on WM and attention, but also on other cognitive domains and on QoL.

The aim of this study was to determine the effect of a CBCR intervention on multiple aspects of cognitive functioning, QoL, and self-efficacy, and compare it to a control intervention in patients with self-perceived cognitive impairments 12–36 months after stroke. Additionally, within the intervention group the aim was to study adherence with the intervention. The hypothesis of the study was that a CBCR training targeting multiple cognitive domains leads to greater improvements on processing speed, flexibility and fluid intelligence in stroke patients than the provision of information through the internet (Rebok et al., 2014; Van der Oord, Ponsioen, Geurts, Ten Brink, & Prins, 2012).

Materials, methods and patient characteristics

Design

This study had a randomised, controlled design and took place in The Netherlands between January 2013 and September 2013 at the Rijnlands Rehabilitation Centre in Leiden and Sophia Rehabilitation in The Hague. The study was approved by the Medical Ethical Review Board of the Leiden University Medical Centre (P 12.190). All participants gave written informed consent prior to participation and were rewarded with a gift card of 50 euros for participating in the trial, with no difference between those who completed the games and those who did not. The CONSORT (Consolidated Standards of Reporting Trials) guidelines were used for adequate reporting of the study (Moher et al., 2010).

Recruitment and inclusion

Inclusion criteria for participation in the study were: age between 45 and 75 years, diagnosed with stroke 12–36 months ago, having self-perceived cognitive impairments (extracted from the checklist accompanying the recruitment letter), having access to the internet, being able to visit the rehabilitation centre, and having time to participate. Exclusion criteria were: antidepressant use, receiving actual treatment for cognitive impairments, severe aphasia, lack of computer skills, and not being proficient in Dutch. In addition, participants with psychological disorders in need of treatment, for example depression, and patients with physical disorders known to impact cognition were excluded.

Potentially eligible patients were identified by first searching the electronic patient registers of the rehabilitation centres for patients who were aged between 45 and 75 years and who had had a stroke 12–36 months ago. The chosen age range constitutes the largest proportion of patients admitted to rehabilitation and to prevent age-specific bias (e.g., stroke patient 18 years old versus 75 years old). Onset of stroke was “not more recently than 12 months” to minimise the influence of natural recovery and “not longer ago than 36 months” to prevent the risk of impaired learning due to age group specific comorbidities (e.g., Alzheimer’s). Potential participants received a letter with information about the study, a checklist concerning the other four inclusion criteria (having self-perceived cognitive impairments, having access to internet, being able to visit the rehabilitation centre and having time to participate) and a form to indicate their willingness to participate in the study. All responding patients received a telephone call from one of the researchers (MW, IV, AK) to make sure they met the inclusion criteria and did not meet the exclusion criteria. Also they were asked to state their age (years), sex and educational level (categorised as low: lower technical and vocational training; medium: secondary technical and vocational training; and high: higher technical and vocational training and university) (Centraal Bureau voor de Statistiek [CBS], 2006).

Randomisation

Included patients were subsequently randomised by an administrative assistant who was not involved in the study. A blocked randomisation scheme (blocks of six), with stratification for age and education level (extracted from the recruitment form and checked by examining the electronic patient registers), made up by a random digit generator (Microsoft Excel 2010), was used to allocate the patients either to the intervention group or control group. Stratification for age and education level was based on findings in a study of Patel et al. (2003) who studied the relationship between different variables and cognitive functioning among 645 stroke patients. As the majority of patients was of the same ethnic origin, stratification for that variable was not performed. The variable "age" was divided into two categories (≤ 58 and > 58), since the median age was 58 years. The variable "education level" was divided into the categories "low", "moderate" and "high", by using a standardised classification system (CBS, 2006).

Blinding

The randomisation sequence was concealed (blinded) from research personnel, so that assessors were not aware of whether a subject was randomised to the intervention or control group.

Intervention and control conditions

The conditions compared in the study consisted of a CBCR training (intervention) versus weekly information about the brain (control). All participants received a user identification and password to log on to a website specifically designed for this study (www.spelenderwijsbeter.nl). After log-on, the website provided either access to the training or the information. All participants were given the contact details of a research assistant, whom they could contact by telephone in case of difficulties using the website or, in case of participants in the intervention group, with the training software.

Intervention

The training consisted of gaming at home during a period of 8 weeks, at least 5 days per week, approximately 15–20 minutes per day, resulting in a requested play time of 600 minutes. The training software was supplied by Lumosity Inc.[®]. This software was chosen because it targets more cognitive domains than other CBCR interventions (e.g., Cogmed), since effective training should include games in several cognitive areas, including flexibility (Buitenweg et al., 2012). Moreover, frequent switches prevent boredom and thereby therapy compliance is stimulated. The recently developed training software of BrainGymmer[®] was

not available when this study took place.

Sixteen games were used and five cognitive domains were targeted: attention, speed, memory, flexibility and problem solving. Three games were randomly assigned to the participant per session. The duration of each game was approximately five minutes. After finishing all sessions of a single game, a new game started after pressing the "next" button. When all three games were completed, participants received feedback that the session for that day was finished. Participants were able to play longer by selecting games from the menu themselves after finishing the training session.

With each game, all participants began at the same level of difficulty. The difficulty level was then raised or lowered depending on the performance in the previous round of the respective game. The software provided feedback about game scores. Furthermore, participants were instructed to complete an extra game session when they missed a session and/or were not able to play five days a week. After eight weeks, participants still had access to the games, but were instructed to temporarily quit playing.

Control

The participants in the control group received weekly information about stroke after log-on at the website of the study. The information provision was not interactive, it provided unidirectional explanations about brain differences between men and woman, the influence of stress on brain function and possible difficulties with living with a damaged brain. Each week, during a period of 8 weeks, new information (text or a video clip) was added to the website. The information was accessible for the participants during the period of 8 weeks. The total duration of the control intervention was on average 70 minutes per person.

Assessments

All assessments were done before the intervention (t0), and 8 and 16 weeks thereafter (t1 and t2). At all time points, the assessments consisted of a set of electronic questionnaires and computer-based performance tests. With respect to the questionnaires, participants received an e-mail with a digital link to the questionnaires one week before each planned assessment, with the request to complete it at home. All computerised performance tests, which took approximately one hour to complete, were conducted in the rehabilitation centre. Tests were administered in a fixed order and every patient received the same instructions. Assistance during testing was provided by the principal investigator (MB) and two students and consisted of logging the participant on to the computer, setting up the tests, providing support in case of technical problems, explaining the test procedure if necessary and saving test results.

Sociodemographic and disease characteristics

Characteristics recorded at baseline by means of a questionnaire were: living situation (alone/together), daily functioning (dependent/independent), and participation in paid work (yes/no). In addition, the affected hemisphere (left/right/other), type of stroke (infarction/haemorrhage), time between stroke and enrolment, and time spent in rehabilitation centre (in months) were recorded from the medical records. The group of responders was compared to the group of non-responders regarding their age and gender using the Mann Whitney U test, the Chi-square test.

Adherence

Data about participants' total play time (in days and minutes) were provided by the software of the training programme.

Primary outcome measures

The primary outcome measures included five neuropsychological tests and the Cognitive Failures Questionnaire (CFQ). Secondary outcomes included a measure of QoL and of self-efficacy. As patients in the intervention group were discouraged from playing the games after 8 weeks, and under the assumption that a present effect would diminish afterwards, time point 1 was chosen as the primary endpoint for both primary and secondary outcome measures.

Attention and flexibility

The Trail Making Test (TMT; Reitan & Wolfson, 1985) was used to assess attention and flexibility. It consists of two parts (TMT-A and TMT-B): In TMT-A the participants were asked to draw lines sequentially connecting 25 encircled numbers distributed on the screen. In TMT-B task requirements were similar, however the participants must alternate between numbers and letters (e.g., 1, A, 2, B, 3, C, etc.). The scores represent the amount of time required to complete task A and B (attention) and the mean difference between time A and B (flexibility) (Strauss, Sherman, & Spreen, 2006). Lower scores for attention and higher scores for flexibility indicate better functioning. In addition, the number of correct connections made by the patients was scored. Both tests were proven to be valid indicators of organic brain damage (Lezak, Howieson, & Loring, 2004).

Working memory

The Block Span Task (Corsi, 1972) and Digit Span Task (Wechsler, 1945) were used to assess working memory with two subtests: blocks/digits forward (sequential order) and blocks/

digits backward (reversed order). The scores consist of the highest number of blocks/digits a participant can correctly reproduce, so that higher scores indicate better functioning. A number of studies have published data on the Block Span Task in stroke patients and concluded that it can be used effectively to assess visuospatial short-term memory in patients with brain damage (Chechacz, Rotshtein, & Humphreys, 2014; De Renzi, Faglioni, & Previdi, 1977; Kessels, Van Zandvoort, Postma, Kappelle, & De Haan, 2000).

Speed and flexibility

With the Eriksen Flanker Task (Eriksen & Schultz, 1979) participants are instructed to respond as quickly as possible to a central target stimulus, which was flanked by either four congruent (e.g., $\leftarrow\leftarrow\leftarrow\leftarrow$) or four incongruent (e.g., $\leftarrow\leftarrow\rightarrow\leftarrow$) stimuli (Eriksen, 1995). The score of the test is the reaction time in milliseconds (speed) and the mean difference between reaction time incongruent and congruent (resistance to interference, flexibility) (Levin & Cross, 2004). Better functioning is indicated by lower scores for speed and higher scores for flexibility.

Fluid intelligence

The Raven Standard Progressive Matrices (SPM; Raven, 1958) is a multiple-choice test consisting of incomplete figures, in which participants are asked to choose the correct missing part depicted in one of the six alternatives. Three versions of equal difficulty with 20 items each were derived from the original 60-item version to assess fluid intelligence at the three time points in a counterbalanced order. The score of the test is the number of correct items translated in IQ scores. Higher scores indicate better functioning. Internal consistency coefficients tend to cluster around .90 for adults (Llabre, 1984) and retest reliability correlations are in the range of .70 and .90 (Llabre, 1984).

Cognitive Failures Questionnaire (CFQ)

The CFQ (Broadbent, Cooper, FitzGerald, & Parkes, 1982) in a Dutch version (Merckelbach, Muris, Nijman, & de Jong, 1995) was used to measure self-perceived cognitive failure. The CFQ includes 25 questions, for example: "Do you fail to notice signposts on the road?". The 5-point scale ranges from 0 (never) to 4 (very often). The total score ranges from 0 to 100, with higher scores indicating less cognitive failure. Broadbent et al. (1982) reported that the questionnaire has high test-retest correlation and high internal consistency. This was confirmed in a Dutch study by Merckelbach et al. (1995), in which the test-retest reliability (.83) and internal validity (Cronbach's $\alpha = .81$) were high. The questionnaire was also completed by the spouse or a person related to the participant.

Secondary outcome measures

Stroke Specific Quality of Life Scale (SS-QoL-12)

A Dutch and short version of the SS-QoL was used to measure health-related QoL (HRQoL) (Post et al., 2011). The self-rating 5-point scale consisted of 12 items. The total score ranges from 12 to 60, with higher scores indicating better function (Williams, Weinberger, Harris, Clark, & Biller, 1999). The questionnaire has been validated in patients with haemorrhage and ischaemic stroke (Post et al., 2011).

General Self-Efficacy Scale (GSES)

A Dutch version of the GSES, a 10-item self-rating scale (Schwarzer & Jerusalem, 1995), was used to assess participants' belief in the ability to respond to and cope with novel or difficult situations and unexpected setbacks or obstacles. The scale ranged from 1 (not at all true) to 4 (exactly true), resulting in a sum score ranging from 10 to 40, with higher scores indicating greater self-efficacy.

Data analysis

The target sample size was based on the ability to detect a difference of 25% in the proportion of patients (35% in the intervention group and 10% in the control group) showing a 25% improvement of the Raven SPM. Although larger than the effects seen in previous similar studies (Jaeggi, Buschkuhl, Jonides, & Perrig, 2008; Karbach & Kray, 2009; Klingberg et al., 2005; Klingberg, Forssberg, & Westerberg, 2002; Schmiedek, Lövdén, & Lindenberger, 2010; Shipstead, Redick, & Engle, 2012; Van Muijden, Band, & Hommel, 2012), this difference was considered to be clinically relevant. Based on a power of .80 to detect a significant difference (2-sided $p = .05$), 43 patients would be required for each study group. To compensate for an expected dropout rate of 15%, we planned to enrol at least 50 patients in each study group.

Adherence of participants in the intervention group was analysed using descriptive statistics and presented as the mean with the range or numbers and percentages. Change scores of endpoint measures over time between t0 and t1 and t1 and t2 in both the intervention and control groups were computed with the 95% confidence interval and analysed by means of paired t-tests or Wilcoxon Signed Rank test, where appropriate. Differences in the change scores between the groups were analysed by independent t-tests or Mann-Whitney U tests, where appropriate.

In addition, a linear mixed model with patient number as a random factor and group and measurement moments interaction as fixed factors were used to determine effects over the total period (t0–t1–t2). Corrections were made for differences in type of stroke

between the intervention and control group. Moreover, Spearman correlation coefficients were calculated to determine correlations between subjective cognitive impairment (CFQ) at baseline and change scores on neuropsychological tests between t0 and t1 and between play time on WM games and change scores on neuropsychological tests between t0 and t1.

All effectiveness analyses were based on intention to treat, meaning all available data were used. Per protocol analysis was performed, including the participants from the intervention group who played at least 600 minutes. For all analyses, the level of statistical significance was set at .05. All analyses were done with the software SPSS version 21.

Results

According to the registers, 889 patients who were aged between 45 and 75 years and had had a stroke between 12–36 months ago, were identified and invited to participate (Figure 1). Of these, 146 patients (64 in Leiden, 82 in The Hague) indicated they fulfilled the inclusion criteria and were interested in the study, of whom 142 were screened for eligibility (4 could not be reached). From these 142 patients, 27 were excluded because they either did not fulfil one or more inclusion criteria or met one or more of the exclusion criteria, so that 115 patients were included and randomised. Directly after randomisation, five patients refused further participation. Eventually, 110 patients were present at the first assessment day and included in the analysis (53 intervention and 57 control). Of these, 107 participants (97%) completed the study, 50/53 (94%) in the treatment group and 57/57 (100%) in the control group.

Demographic characteristics

The demographic and clinical characteristics of the 110 study participants are shown in Table 1. Participants' characteristics were similar in the intervention and control groups at baseline. However, in the intervention group significantly more patients had had a haemorrhage (21/53; 40%) compared to the control group (13/57; 23%) ($p = .02$).

No significant differences were found between age and gender for the group of 146 responders (median age 59, range 46–74; male $n = 69$, 63%) compared to the group of 743 non-responders (median age 61, range 45–75; male $n = 459$, 59%), $p = .62$ for age and $p = .66$ for gender, respectively.




Figure 1. Trial profile.

Adherence to the intervention

Seven out of 53 patients in the intervention group failed to play games due to technical problems with their computer (n = 3), lack of motivation (n = 2) or not feeling well (n = 2). The number of days and total play time of the remaining 46 patients who actually played are shown in Table 2. The median number of days these patients played was 45 (range = 4–63) and the median total time played in minutes was 562 (range = 63–1264).

Two patients logged on only once and three patients between three and five times, so their play time was very limited. Reasons for not playing, according to the participants' personal coaches were: having difficulties playing the games due to health status, not sufficiently motivated, or technical problems with the computer. A group of seven patients played more than the 600 minutes requested. Two patients (4%) in the intervention group completed 2–3 training sessions during the follow-up period (8–16 weeks) despite explicit discouragement to do so.

Table 1. Baseline demographic and clinical characteristics of the 110 stroke patients.

	Intervention group (n = 53)	Control group (n = 57)
Age in years; median (range)	59 (46–74)	59 (46–73)
Sex, male n (%)	34 (64)	35 (61)
Type of stroke Infarction n (%)	29 (55)	44 (77)
Haemorrhage n (%)	21 (40)	13 (23)
Unknown ^a n (%)	3 (5)	0 (0)
Location of stroke		
Hemisphere left n (%)	23 (43)	28 (49)
Hemisphere right n (%)	26 (49)	26 (47)
Basal ganglia n (%)	3 (6)	0 (0)
Unknown ^a n (%)	1 (2)	2 (4)
Time from stroke onset to randomisation in months; mean \pm SD	26 \pm 9.1	25 \pm 7.4
Time spent in rehabilitation centre in months; mean \pm SD	6 \pm 3.6	5 \pm 3.4
Educational status^b		
Low n (%)	17 (32)	20 (35)
Intermediate n (%)	18 (34)	18 (32)
High n (%)	18 (34)	19 (33)
Living alone n (%)	13 (25)	15 (25)
Independent in daily functioning n (%)	52 (98)	54 (95)
Participation in paid work n (%)	14 (26)	20 (32)
Subjective cognitive failure (CFQ); median (IQR)	63 (19)	64 (21)

^a No data were available for medical status.

^b Low: lower technical and vocational training; medium: secondary technical and vocational training; and high: higher technical and vocational training and university.

Primary outcome measures

Intention-to-treat analysis

Baseline and follow-up scores of primary outcomes are shown in Table 3(a). At baseline, no significant differences were found between the groups. A significant difference between the groups of changes between t0 and t1 was found regarding the number of correct items in favour of the intervention group on the Block Span Forward Test ($p = .02$) and the reaction time incongruent on the Eriksen Flanker Test ($p < .01$). No differences in improvement were seen between the two groups on other outcome measures.

Within both groups significant improvements were found between t0 and t1 regarding the cognitive domain attention for the parameters Time B and Items Correct B, but not for

Time A and Items Correct A. Moreover, participants in the intervention group improved between t0 and t1 within the cognitive domain flexibility (TMT-A/TMT-B), reaction time congruent (Eriksen Flanker Task) and fluid intelligence (Raven SPM, Peck score), whereas no significant changes were seen within the control group. In addition, there were no statistically significant differences between the groups taking into account all time points (T0-T1-T2) (mixed model analysis of variance).

The following correlations were found between baseline scores of subjective cognitive impairment (CFQ) and mean differences between t0 and t1 on the neuropsychological tests: TMT A Spearman, $r = .180$ ($p = .07$), and TMT B, $r = .308$ ($p < .01$), Block Span forward, $r = -.181$ ($p = .06$), and backward, $r = -.079$ ($p = .42$), Digit Span forward, $r = -.184$ ($p = .06$), and backward, $r = -.295$ ($p < .01$), Flanker incongruent, $r = .168$ ($p = .09$), and congruent, $r = .158$ ($p = .11$), and the Raven SPM, $r = -.158$ ($p = .15$). No significant correlations were found between play time on WM games and change scores on neuropsychological tests between t0 and t1.

Table 2. Total days and minutes played during the 8-week training period in the intervention group (46 out of 53 patients).

Characteristics	Median (IQR) ^a	
Participants who played	46	
Total time played (days)	45	(4–63)
Total time played (minutes)	528	(63–1264)
Total time played per cognitive domain (minutes):		
Attention	60	(3–408)
Speed	53	(3–139)
Memory	232	(26–646)
Flexibility	111	(6–349)
Problem solving	81	(1–265)

^aIQR's are expressed as the net result of 75th percentile minus 25th percentile.

Table 3a. Baseline, follow-up and change scores for attention, flexibility, memory, speed, fluid intelligence and cognitive failure in the intervention versus the control group.

	T0			T0-T1			T1-T2		
				Within group	Between groups		Within group	Between groups	
	Median (IQR) ^a			Mean difference (95% CI) ^b	p-value ^c		Mean difference (95% CI) ^b	p-value ^c	
Working memory									
Items forward (Block Span Task) (0-9 items)	I: 3 (2)	0.7	(0.3,1.1)	0.02	-0.1	(-0.4,0.3)	0.59		
	C: 4 (2)	-0.1	(-0.6,0.4)		0.2	(-0.2,0.6)			
Items backward (Block Span Task) (0-8 items)	I: 3 (2)	0.3	(-0.2,0.8)	0.45	-0.2	(-0.7,0.2)	0.56		
	C: 3 (4)	0.1	(-0.3,0.6)		-0.1	(-0.6,0.4)			
Items forward (Digit Span Task) (0-9 items)	I: 5 (2)	0.1	(-0.6,0.5)	0.23	0.4	(-0.1,0.8)	0.43		
	C: 5 (3)	0.5	(-0.0,0.1)		0.0	(-0.3,0.3)			
Items backward (Digit Span Task) (0-8 items)	I: 4 (2)	0.1	(-0.5,0.8)	0.28	0.1	(-0.5,0.7)	0.25		
	C: 3 (3)	0.6	(0.1,1.1)		-0.2	(-0.7,0.2)			
Attention									
Time A (TMT-A) (0-2880s)	I: 78 (49)	-7	(-21.6,7.8)	0.54	-5	(-17.2,8.1)	0.87		
	C: 74 (42)	-19	(-39.8,1.9)		8	(-12.3,27.7)			
Time B (TMT-B) (0-2880s)	I: 169 (142)	-73	(-134.1,-11.5)	0.10	-11	(-27.5,4.9)	0.40		
	C: 144 (202)	-58	(-102.3,-12.8)		25	(-18.6,68.0)			
Items correct A (TMT-A) (0-24)	I: 24 (0)	0.6	(-0.3,1.5)	0.85	0.1	(-0.7,0.9)	0.11		
	C: 24 (0)	0.7	(-0.4,1.8)		-0.3	(-0.7,0.1)			
Items correct B (TMT-B) (0-24)	I: 17 (13)	2.5	(0.7,4.2)	0.38	1.1	(-0.3,2.6)	0.30		
	C: 20 (15)	1.6	(0.2,2.9)		-0.2	(-1.4,1.1)			
Speed (Flanker Task)									
Reaction time congruent (0-2000 milliseconds)	I: 770 (258)	-51	(-97.1,-5.4)	0.08	-16	(-38.8,6.6)	0.48		
	C: 816 (285)	-4	(-57.9,50.7)		-42	(-86.0,2.5)			
Reaction time incongruent (0-2000 milliseconds)	I: 797 (268)	-63	(-118.9,-7.4)	0.00	-14	(-38.2,9.3)	0.08		
	C: 833 (342)	9	(-49.5,67.1)		-46	(-91.1,-0.9)			
Flexibility									
Difference time A/time B (TMT-A/TMT-B)	I: -80 (125)	66	(6.5,124.7)	0.18	7	(-13.0,26.9)	0.56		
	C: -61 (145)	34	(-15.7,84.5)		-17	(-65.9,31.8)			
Difference reaction time congruent/incongruent (Flanker Task)	I: -31 (39)	10	(-9.4,29.0)	0.62	1	(-9.0,11.5)	0.56		
	C: -33 (62)	-12	(-39.8,15.0)		4	(-11.9,20.5)			
Fluid intelligence (Raven SPM)									
Peck-score (0-150)	I: 110 (19)	6	(1.3,9.8)	0.47	-3	(-7.4,1.0)	0.34		
	C: 111 (17)	3	(-1.4,7.3)		-1	(-5.7,3.1)			
Cognitive failure (CFQ)									
Total score (participant) (0-100)	I: 63 (19)	-0.5	(-3.9,2.9)	0.14	-0.9	(-5.5,3.8)	0.87		
	C: 64 (21)	1.4	(-1.0,3.6)		-1.6	(-4.9,1.7)			
Total score (relatives) (0-100)	I: 63 (35)	-0.2	(-4.4,3.9)	0.75	1.5	(-2.8,5.8)	0.81		
	C: 59 (19)	3.2	(-2.0,8.5)		-0.05	(-5.6,5.5)			

^a IQRs are expressed as the net result of 75th percentile minus 25th percentile. All analyses were done with adjustment for significant differences in baseline characteristics between patients.

^b Values are the difference in mean change values between groups (treatment effect with 95% confidence interval).

^c p-values were analysed using Mann-Whitney U test or mixed model analyses where appropriate.

Table 3b. Baseline and follow-up scores of variables attention, flexibility, working memory, and speed with significance between groups per protocol (n = 19).

	T0		T0-T1			T1-T2		
			Within group		Between groups	Within group		Between groups
	Median (IQR)^a		Mean difference (95% CI)^b		p-value^c	Mean difference (95% CI)^b		p-value^c
Working memory								
Items forward (Block Span Task)	I: 4 (3)	0.9	(0.4,1.8)		.09	-0.3 (-0.4,0.3)		.66
(0-9 items)	C: 4 (2)	-0.1	(-0.6,0.4)			0.2 (-0.2,0.6)		
Items backward (Block Span Task)	I: 3 (2)	0.4	(-0.2,0.7)		.42	-0.2 (-0.8,0.2)		.35
(0-8 items)	C: 3 (4)	0.1	(-0.3,0.6)			-0.1 (-0.6,0.4)		
Items forward (Digit Span Task)	I: 5 (2)	0.1	(-0.8,0.5)		.50	0.4 (-0.1,0.7)		.21
(0-9 items)	C: 5 (3)	0.5	(-0.0,0.1)			0.0 (-0.3,0.3)		
Items backward (Digit Span Task)	I: 4 (3)	0.1	(-0.5,0.7)		.40	0.2 (-0.6,0.7)		.88
(0-8 items)	C: 3 (3)	0.6	(0.1,1.1)			-0.2 (-0.7,0.2)		
Attention								
Time A (TMT-A)	I: 76 (45)	-27	(-45.2,-3.8)		.69	-9 (-14.1,10.1)		.93
(0-2880s)	C: 74 (42)	-19	(-39.8,1.9)			8 (-11.5,34.7)		
Time B (TMT-B)	I: 169 (143)	-81	(-160.0,-3.0)		.04	-12 (-14.9,3.7)		.30
(0-2880s)	C: 144 (202)	-58	(-102.3,-12.8)			25 (-18.6,68.0)		
Items correct A (TMT-A)	I: 24 (0)	0.6	(-0.3,1.5)		.35	0.1 (-0.9,1.1)		.20
(0-24)	C: 24 (0)	0.7	(-0.4,1.8)			-0.3 (-0.7,0.1)		
Items correct B (TMT-B)	I: 17 (15)	2.5	(0.7,4.2)		.45	1.4 (-0.3,2.6)		.12
(0-24)	C: 20 (15)	1.6	(0.2,2.9)			-0.2 (-1.4,1.1)		
Speed (Flanker Task)								
Reaction time congruent	I: 696 (244)	-59	(-110.1,-13.2)		.09	-22 (-40.2,7.8)		.48
(0-2000 milliseconds)	C: 816 (285)	-4	(-57.9,50.7)			-42 (-86.0,2.5)		
Reaction time incongruent	I: 719 (266)	-71	(-125.0,-5.4)		.07	-19 (-36.6,12.3)		.08
(0-2000 milliseconds)	C: 833 (342)	9	(-49.5,67.1)			-46 (-91.1,-0.9)		
Flexibility								
Difference time A/time B	I: -91 (119)	71	(12.2,136.9)		.00	-14 (-3.6,32.8)		.56
(TMT-A/TMT-B)	C: -61 (145)	34	(-15.7,84.5)			-17 (-65.9,31.8)		
Difference reaction time congruent	I: -26 (32)	15	(-5.4,32.2)		.62	4 (-11.5,9.9)		.56
/incongruent (Flanker Task)	C: -33 (62)	-12	(-39.8,15.0)			4 (-11.9,20.5)		
Fluid intelligence (Raven SPM)								
Peck-score	I: 107 (21)	7	(1.0,9.3)		.47	-3 (-7.4;1.0)		.34
(0-150)	C: 111 (17)	3	(-1.4,7.3)			-1 (-5.7;3.1)		
Cognitive failure (CFQ)								
Total score (participant)	I: 63 (19)	-0.4	(-4.0,3.19)		.14	-0.5 (-3.6,2.3)		.87
(0-100)	C: 64 (21)	1.4	(-1.2,3.4)			-1.6 (-4.9,1.7)		
Total score (relatives)	I: 63 (35)	-0.2	(-4.2,3.2)		.75	0.5 (-2.8,3.6)		.81
(0-100)	C: 59 (19)	3.2	(-2.0,8.5)			-0.05 (-5.6,5.5)		

^a IQRs are expressed as the net result of 75th percentile minus 25th percentile. All analyses were done with adjustment for significant differences in baseline characteristics between patients.

^b Values are the difference in mean change values between groups (treatment effect with 95% confidence interval).

^c *p*-values were analysed using Mann-Whitney U test or mixed model analyses where appropriate.

Per protocol analysis

Comparisons of change scores between the intervention and control group ($n = 57$), while including only the 19 patients who played at least 600 minutes in the intervention group are shown at Table 3(b). A significant difference between groups was found at 8 weeks regarding attention (TMT-B, Time B, $p = .04$) and flexibility (TMT-A/TMT-B, Difference time A and time B, $p < .01$), both in favour of the intervention group. In the intention-to-treat (ITT) analysis no significant differences were found between groups regarding the TMT. However, effects found in the ITT analysis between groups did not remain significant in the per protocol analysis when compared to the control group regarding Block Span items correct forward ($p = .07$) and reaction time incongruent on the Flanker Task ($p = .09$).

When compared to ITT analysis, the magnitude of improvements in the per-protocol analysis increased within the intervention group between t0 and t1 regarding speed (Eriksen Flanker Task, reaction time congruent; reaction time incongruent), working memory (Block Span backward), flexibility (difference time A/time B; difference reaction time congruent/incongruent), fluid intelligence (Raven SPM) and subjective cognitive failure (CFQ).

Secondary outcome measures

Baseline and follow-up scores of secondary outcomes are shown in Table 4. At baseline, no significant differences were found between both groups and no differences were seen in change scores between the two groups. In addition, there were no statistically significant difference between the groups taking into account all time points (T0-T1-T2) (mixed model analysis of variance).

Table 4. Baseline and follow-up scores of QoL and self-efficacy with significance in the intervention versus the control group.

		T0	T0-T1		T1-T2	
		Median (IQR) ^a	Mean difference (95% CI) ^b	<i>p</i> -value ^c	Mean difference (95% CI) ^b	<i>p</i> -value ^c
QoL ^d	I:	3.9 (1.5)	0.0 (−0.2,0.2)	.76	0.1 (−0.1,0.3)	.45
Total score (0–5)	C:	3.8 (1.3)	0.0 (−0.1,0.2)		0.0 (−0.1,0.1)	
Self-efficacy ^e	I:	31 (10)	1.0 (0.3,2.2)	.15	0.52 (−0.6,1.6)	.86
Total score (0–40)	C:	30 (8)	−0.6 (−1.8,0.6)		0.29 (−1.1,1.7)	

^aIQRs are expressed as the net result of 75th percentile minus 25th percentile. All analyses were done with adjustment for significant differences in baseline characteristics between patients.

^bValues are the difference in mean change values between groups (treatment effect with 95% confidence interval).

^c*p*-values were analysed using Mann-Whitney U test or mixed model analyses where appropriate.

^dMeasured with the Stroke Specific Quality of Life Questionnaire (SSQoL).

^eMeasured with the General Self-efficacy Scale (GSES).

Discussion

In this randomised study an 8-week computer-based brain training programme was compared with providing information to patients with cognitive complaints after stroke. In general only performances on cognitive function tests that were similar to the games included in the intervention improved in the CBCR training group compared to the control group. However, no near transfer effect was found to tasks such as the Digit Span Task or Trail Making Task.

The effects found in our study on the WM tests; effect size 0.23 regarding the Span board and 0.11 regarding the Digit Span between groups, were smaller than other studies using CBCR training primarily focused on one cognitive function among stroke patients. Westerberg et al. (2007), who compared a computerised 5-week WM training (Cogmed) on various WM tasks with a passive control group, reported significant improvements in favour of the intervention group on the WM tests; effect size Span Board 0.83 and effect size Digit Span 1.58. Åkerlund et al. (2013) also found a significant difference on the Digit Span Task in favour of the group who received WM training in the sub-acute phase compared to usual rehabilitation. Smith et al. (2009) evaluated the effect of a 40-hour CBCR training targeting speed and information processing among healthy adults and found an effect size of 0.26 between groups on the Digit Span Backward in favour of the intervention group. Based on these results it can be concluded that the effects of the CBCR training were limited in this group of patients.

Moreover, other CBCR studies found improvements of daily functioning after training. Westerberg et al. (2007) reported a significant change on the CFQ ($p < .01$) and Åkerlund et al. (2013) found less depressive symptoms after WM training between groups ($p = .03$), in both studies in favour of the intervention group. In the current study, no far transfer effects were shown, i.e., no significant difference was found on cognitive functioning by means of the CFQ, QoL scale or self-efficacy scale. Although results of the studies cannot be easily compared to the current study, since the total training period was lower in our study, they indicate that interventions primarily focused on one cognitive function instead of multiple cognitive domains might be more effective for stroke patients. However, in line with the results in our study, Barker-Collo et al. (2009) found very limited effects on tests that were related to the trained tasks and no result in a near or far transfer.

There are a number of potential explanations for the observed lack of effect. Apart from the intervention not being effective, it remains unclear to what extent responsiveness of the outcome measures also played a role, although the neuropsychological tests have proven ability to measure changes over time in studies among stroke patients (Åkerlund et al., 2013; Barker-Collo et al., 2009; Broadway & Engle, 2010; Chechlacz et al., 2014; Kessels et al., 2000; Nys et al., 2006; Sanchez-Cubillo et al., 2009; Spikman, 2001; Westerberg et al., 2007).

Second, it is questionable whether the dose of the training was enough to achieve responses (dose-response relation). Seven out of the 53 people in the intervention group did not follow the training and only 19 out of the 53 patients, who actually played, completed the training (playing > 600 minutes). This was why a per protocol analysis was done. Indeed, within the intervention group the magnitude of change was larger in the per protocol analysis than with the ITT analysis. However, the difference with the control condition did not reach statistical significance, most likely due to a lack of power because of the limited sample size.

Of note was the finding that some significant improvements were also seen in the control group (time B and items correct B). This might be explained by a learning effect on the neuropsychological tests, since patients became more familiar with using the computer while testing more frequently. For instance, the TMT requires skills moving the mouse on the screen to connect items. By using strategies, for example, using the non-affected hand, scores can increase. Although the primary focus of our study was to investigate an intervention based on restitution approaches, it cannot be ruled out that patients developed strategies that had positive effects on the outcomes. Unfortunately, the average time spent on the control intervention was not available at the level of the individual patient.

Overall, significant differences between both groups were found on 2 out of the 17 outcome measures used in the study. Since multiple outcomes are used, appearances of significant differences can occur from chance alone. However, no corrections for multiple testing were performed for the reason that as the number of tests increases, the p-value that has to be exceeded to achieve statistical significance decreases markedly, lowering the statistical power (Armstrong, 2014; Nakagawa, 2004).

The observation that only 19/53 completed the intervention as it was intended indicates that the feasibility of a programme in the present form is limited. Non-users were also found in other CBCR studies among stroke patients (Connor & Standen, 2012; Cruz et al., 2013), where it was suggested that CBCR training is not well used by all patients, probably due to high demands of training and the motivation needed to complete the training. On the other hand, the fact that 46 out of 53 patients (partly) followed the training (93%), indicates that there is a need among some stroke patients for online training in the chronic phase after stroke. More research is needed to determine patient characteristics that can impair or improve compliance and which stroke patients can benefit the most from CBCR or cognitive rehabilitation in general.

A limitation of the study is that participants were selected on the basis of subjective cognitive impairment and that scores on the CFQ showed that the level of impairment varied widely among patients. Selection bias could have occurred when included patients had lower ability to improve because they were objectively not impaired. Moreover, patients with aphasia or patients unable to use a computer were excluded. This implies that the

patients who were selected were in better health.

A second limitation is that all patients (age: 45–75 years, CVA: 12–36 months ago) were invited by letter to participate in the trial by their treating physician, but were only asked to return the letter if they were willing to participate. Therefore, it is not clear why non-responders did not wish to participate.

In addition, the sample size was based on the assumption that the intervention would lead to a relatively large difference in neuropsychological function as compared to the control group. This assumption was probably too optimistic, leading us to the conclusion that in future research even larger studies are needed to detect changes over time and/or differences between groups. With more knowledge about the clinical relevance of improvements becoming available in stroke research, future sample size calculations will be better underpinned, using empirical data.

A strength of the study is that it was one of the first to examine the effect of CBCR on multiple cognitive functions and also to assess the transferability to QoL and self-efficacy after stroke. Additionally, it is one of the few RCT CBCR studies with a large sample and follow-up that has investigated long-term effects. Furthermore, only three participants did not complete the follow-up assessments, so the drop-out rate was relatively low.

Overall, the results of this study and similar research imply that CBCR interventions targeting one cognitive domain are more effective for stroke patients in terms of near and far transfer effects compared to those targeting multiple cognitive domains. In order to improve daily activities of stroke patients, computer tasks need to be closely related to the impaired task itself. Thus, CBCR training needs to be tailored and adapted to each patient's individual profile. It would appear important to support stroke patients with CBCR training, since training is not well used by all patients. It is possible patients benefit more when they learn how to use strategies in their training and when motivated by clinicians (Lynch, 2002). Further research is needed to determine if CBCR training can improve cognitive functioning in chronic stroke patients. Brain mapping techniques, such as functional magnetic resonance imaging, might be helpful in identifying effects on brain plasticity (Carter et al., 2010).

Reference list

- Åkerlund, E., Esbjörnsson, E., Sunnerhagen, K. S., & Björkdahl, A. (2013). Can computerized working memory training improve impaired working memory, cognition and psychological health? *Brain Injury*, 27(13–14), 1649–1657.
- Akinwuntan, A. E., De Weerd, W., Feys, H., Pauwels, J., Baten, G., Arno, P., & Kiekens, C. (2005). Effect of simulator training on driving after stroke: A randomized controlled trial. *Neurology*, 65(6), 843–850.
- Armstrong, R. A. (2014). When to use the Bonferroni correction. *Ophthalmic and Physiological Optics*, 34(5), 502–508.
- Barker-Collo, S. L., Feigin, V. L., Lawes, C. M., Parag, V., Senior, H., & Rodgers, A. (2009). Reducing attention deficits after stroke using attention process training a randomized controlled trial. *Stroke*, 40(10), 3293–3298.
- Broadbent, D. E., Cooper, P. F., FitzGerald, P., & Parkes, K. R. (1982). The Cognitive Failures Questionnaire (CFQ) and its correlates. *British Journal of Clinical Psychology*, 21, 1–16.
- Broadway, J. M., & Engle, R. W. (2010). Validating running memory span: Measurement of working memory capacity and links with fluid intelligence. *Behavior Research Methods*, 42(2), 563–570.
- Buitenweg, J. I., Murre, J. M., & Ridderinkhof, K. R. (2012). Brain training in progress: A review of trainability in healthy seniors. *Frontiers in Human Neuroscience*, 6, 79–89.
- Carter, A. R., Astafiev, S. V., Lang, C. E., Connor, L. T., Rengachary, J., Strube, M. J., & Corbetta, M. (2010). Resting interhemispheric functional magnetic resonance imaging connectivity predicts performance after stroke. *Annals of Neurology*, 67(3), 365–375.
- Centraal Bureau voor de Statistiek (CBS). (2006). The Dutch Standard Classification of Education, SOI 2006 [PDF file]. Retrieved from: <https://www.cbs.nl/en-gb/background/2008/24/the-dutch-standardclassification-of-education-soi-2006>
- Chechacz, M., Rotshtein, P., & Humphreys, G. W. (2014). Neuronal substrates of corsi block span: Lesion symptom mapping analysis in relation to attentional competition and spatial bias. *Neuropsychologia*, 64, 240–251.
- Claesson, L., Lindén, T., Skoog, I., & Blomstrand, C. (2005). Cognitive impairment after stroke—Impact on activities of daily living and costs of care for elderly people. *Cerebrovascular Diseases*, 19(2), 102–109.
- Connor, B. B., & Standen, P. J. (2012, September). So much technology, so little time: Factors affecting use of computer-based brain training games for cognitive rehabilitation following stroke. In P. M. Sharkey & E. Klinger (Eds.), *Proceedings of the 9th International Conference on Disability, Virtual Reality and Associated Technologies* (pp. 53–59). Laval: The University of Reading.
- Corsi, P. M. (1972). Human memory and the medial temporal region of the brain (Unpublished doctoral dissertation). McGill University, Montreal, Quebec, Canada.
- Cruz, V. T., Pais, J., Bento, V., Mateus, C., Colunas, M., Alves, I., Rocha, N. P. (2013). A rehabilitation tool designed for intensive web-based cognitive training: Description and usability study. *JMIR Research Protocols*, 2(2), e59. doi:10.2196/resprot.2899
- De Renzi, E., Faglioni, P., & Previdi, P. (1977). Spatial memory and hemispheric locus of lesion. *Cortex*, 13(4), 424–433.
- Douiri, A., Rudd, A. G., & Wolfe, C. D. (2013). Prevalence of poststroke cognitive impairment South London stroke register 1995–2010. *Stroke*, 44(1), 138–145.
- Eriksen, C. W. (1995). The flankers task and response competition: A useful tool for investigating a variety of cognitive problems. *Visual Cognition*, 2, 101–118.
- Eriksen, C. W., & Schultz, D. W. (1979). Information processing in visual search: A continuous flow conception and experimental results. *Perception & Psychophysics*, 25, 249–263.
- Green, C. S., & Bavelier, D. (2008). Exercising your brain: A review of human brain plasticity and training induced learning. *Psychology and Aging*, 23(4), 692–701.

- Jaeggi, S. M., Buschkuhl, M., Jonides, J., & Perrig, W. J. (2008). Improving fluid intelligence with training on working memory. *Proceedings of the National Academy of Sciences*, 105(19), 6829–6833.
- Karbach, J., & Kray, J. (2009). How useful is executive control training? Age differences in near and far transfer of task-switching training. *Developmental Science*, 12(6), 978–990.
- Kessels, R. P., Van Zandvoort, M. J., Postma, A., Kappelle, L. J., & De Haan, E. H. (2000). The Corsi blocktapping task: Standardization and normative data. *Applied Neuropsychology*, 7(4), 252–258.
- Klingberg, T., Fernell, E., Olesen, P. J., Johnson, M., Gustafsson, P., Dahlström, K., ... Westerberg, H. (2005). Computerized training of working memory in children with ADHD—a randomized, controlled trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44(2), 177–186.
- Klingberg, T., Forssberg, H., & Westerberg, H. (2002). Training of working memory in children with ADHD. *Journal of Clinical and Experimental Neuropsychology*, 24(6), 781–791.
- Laver, K. E., George, S., Thomas, S., Deutsch, J. E., & Crotty, M. (2015). Virtual reality for stroke rehabilitation: An abridged version of a Cochrane review. *European Journal of Physical and Rehabilitation Medicine*, 51 (4), 497–506.
- Levin, D. Z., & Cross, R. (2004). The strength of weak ties you can trust: The mediating role of trust in effective knowledge transfer. *Management Science*, 50(11), 1477–1490.
- Lezak, M. D., Howieson, D. B., & Loring, D.W. (2004). *Neuropsychological assessment* (4th ed.). New York: Oxford University Press.
- Llabre, M. M. (1984). Standard progressive matrices. In D. J. Keyser & R. C. Sweetland (Eds.), *Test critiques* (Vol. 1, pp. 595–602). Kansas City, MO: Test Corporation of America.
- Lynch, B. (2002). Historical review of computer-assisted cognitive retraining. *The Journal of Head Trauma Rehabilitation*, 17(5), 446–457.
- Merckelbach, H., Muris, P., Nijman, H., & de Jong, P. J. (1995). Self-reported cognitive failures and neurotic symptomatology. *Personality and Individual Differences*, 20, 715–724.
- Moher, D., Hopewell, S., Schulz, K. F., Montori, V., Gøtzsche, P. C., Devereaux, P. J., Elbourne, D., Egger, M., & Altman, D. G. (2010). CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomised trials. *Journal of Clinical Epidemiology*, 63(8), e1–e37.
- Nakagawa, S. (2004). A farewell to Bonferroni: The problems of low statistical power and publication bias. *Behavioral Ecology*, 15(6), 1044–1045.
- Nys, G. M. S., Van Zandvoort, M. J. E., Van Der Worp, H. B., De Haan, E. H. F., De Kort, P. L. M., Jansen, B. P.W., & Kappelle, L. J. (2006). Early cognitive impairment predicts long-term depressive symptoms and QoL after stroke. *Journal of the Neurological Sciences*, 247(2), 149–156.
- Owen, A. M., Hampshire, A., Grahn, J. A., Stenton, R., Dajani, S., Burns, A. S., Ballard, C. G. (2010). Putting brain training to the test. *Nature*, 465(7299), 775–778.
- Paker, N., Buğdaycı, D., Tekdöş, D., Kaya, B., & Dere, Ç. (2010). Impact of cognitive impairment on functional outcome in stroke. *Stroke Research and Treatment*, 2010, 1–5.
- Pasquini, M., Leys, D., Rousseaux, M., Pasquier, F., & Henon, H. (2007). Influence of cognitive impairment on the institutionalisation rate 3 years after a stroke. *Journal of Neurology, Neurosurgery & Psychiatry*, 78 (1), 56–59.
- Patel, M., Coshall, C., Rudd, A. G., & Wolfe, C. D. (2003). Natural history of cognitive impairment after stroke and factors associated with its recovery. *Clinical Rehabilitation*, 17(2), 158–166.
- Post, M. W. M., Boosman, H., van Zandvoort, M. M., Passier, P. E. C. A., Rinkel, G. J. E., & Visser-Meily, J. M. A. (2011). Development and validation of a short version of the Stroke-Specific QoL Scale. *Journal of Neurology, Neurosurgery and Psychiatry*, 82, 3(2010), 283–286.
- Raven, J. C. (1958). *Mill Hill vocabulary scale: MHV; Raven's standard progressive matrices: SPM*. London: HK Lewis.
- Rebok, G. W., Ball, K., Guey, L. T., Jones, R. N., Kim, H. Y., King, J. W., ... & Willis, S. L. (2014). Ten-year effects of the advanced cognitive training for independent and vital elderly cognitive training trial on cognition and everyday functioning in older adults. *Journal of the American Geriatrics Society*, 62(1), 16–24.

- Reitan, R., & Wolfson, D. (1985). The Halstead-Reitan neuropsychological test battery, Tucson, AZ: Neuropsychology Press.
- Sanchez-Cubillo, I., Perianez, J. A., Adrover-Roig, D., Rodriguez-Sanchez, J. M., Rios-Lago, M., Tirapu, J. E. E. A., & Barcelo, F. (2009). Construct validity of the Trail Making Test: Role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. *Journal of the International Neuropsychological Society*, 15(3), 438–450.
- Schmiedek, F., Lövdén, M., & Lindenberger, U. (2010). Hundred days of cognitive training enhance broad cognitive abilities in adulthood: Findings from the COGITO study. *Frontiers in Aging Neuroscience*, 2, 2–27.
- Schwarzer, R., & Jerusalem, M. (1995). Generalized self-efficacy scale. Measures in health psychology: A user's portfolio. Causal and Control Beliefs, 1, 35–37.
- Shipstead, Z., Redick, T. S., & Engle, R. W. (2012). Is working memory training effective? *Psychological Bulletin*, 138(4), 628–654.
- Smith, G. E., Housen, P., Yaffe, K., Ruff, R., Kennison, R. F., Mahncke, H. W., & Zelinski, E. M. (2009). A cognitive training program based on principles of brain plasticity: Results from the Improvement in Memory with Plasticity-based Adaptive Cognitive Training (IMPACT) Study. *Journal of the American Geriatrics Society*, 57(4), 594–603.
- Sohlberg, M. M., & Mateer, C. A. (Eds.). (2001). *Cognitive rehabilitation: An integrative neuropsychological approach*. New York: Guilford Press.
- Spikman, J. M. (2001). *Attention, mental speed and executive control after closed head injury: Deficits, recovery and outcome* (Doctoral dissertation). University of Groningen.
- Strauss, E., Sherman, E. M. S., & Spreen, O. (2006). *A compendium of neuropsychological tests: Administration, norms, and commentary*. New York: Oxford University Press.
- Tatemichi, T. K., Desmond, D. W., Stern, Y., Paik, M., Sano, M., & Bagiella, E. (1994). Cognitive impairment after stroke: Frequency, patterns, and relationship to functional abilities. *Journal of Neurology, Neurosurgery & Psychiatry*, 57(2), 202–207.
- Truelsen, T., Piechowski-Józwiak, B., Bonita, R., Mathers, C., Bogousslavsky, J., & Boysen, G. (2006). Stroke incidence and prevalence in Europe: A review of available data. *European Journal of Neurology*, 13, 581–598.
- Van der Oord, S., Ponsioen, A. J. G. B., Geurts, H. M., Ten Brink, E. L., & Prins, P. J. M. (2012). A pilot study of the efficacy of a computerized executive functioning remediation training with game elements for children with ADHD in an outpatient setting: Outcome on parent- and teacher-rated executive functioning and ADHD behavior. *Journal of Attention Disorders*. doi:1087054712453167
- Van Muijden, J., Band, G. P., & Hommel, B. (2012). Online games training aging brains: Limited transfer to cognitive control functions. *Frontiers in Human Neuroscience*, 6, 141–154.
- Wechsler, D. (1945). A standardized memory scale for clinical use. *The Journal of Psychology*, 19(1), 87–95.
- Westerberg, H., Jacobaeus, H., Hirvikoski, T., Clevberger, P., Östensson, M. L., Bartfai, A., & Klingberg, T. (2007). Computerized working memory training after stroke - A pilot study. *Brain Injury*, 21(1), 21–29.
- Williams, L. S., Weinberger, M., Harris, L. E., Clark, D. O., & Biller, J. (1999). Development of a stroke-specific QoL scale. *Stroke*, 30(7), 1362–1369.
- Zhang, Y., Chapman, A. M., Plested, M., Jackson, D., & Purroy, F. (2012). The incidence, prevalence, and mortality of stroke in France, Germany, Italy, Spain, the UK, and the US: A literature review. *Stroke Research and Treatment*, 2012, 1–11.



Adherence of stroke patients with an online brain training program: the role of health professionals' support

M.M. Wentink
J.J.L. Meesters
M.A.M. Berger
A.J. de Kloet
E. Stevens
G.P.H. Band
C.H. Kromme
R. Wolterbeek
P.H. Goossens
T.P.M. Vliet Vlieland

Topics in Stroke Rehabilitation, 2018;25(5):359-365.

Poster presentation: Dutch Congress of Rehabilitation Medicine,
11 November 2016, Maastricht, Netherlands.



Abstract

Background: Computer-based cognitive rehabilitation is used to improve cognitive functioning after stroke. However, knowledge on adherence rates of stroke patients is limited.

Objective: To describe stroke patients' adherence with a brain training program using two frequencies of health professionals' supervision.

Methods: This study is part of a randomized controlled trial comparing the effect of the brain training program (600 min playtime with weekly supervision) with a passive intervention in patients with self-perceived cognitive impairments after stroke. Patients randomized to the control condition were offered the brain training after the trial and received supervision twice (vs weekly in intervention group). Adherence was determined using data from the study website. Logistic regression analyses were used to examine the impact of supervision on adherence.

Results: 53 patients allocated to the intervention group (group S8; 64% male, mean age 59) and 52 patients who were offered the intervention after the trial (group S2; 59% male, mean age 59) started the brain training. The median playtime was 562 min (range 63–1264) in group S8 vs. 193 min (range 27–2162) in group S2 ($p < 0.001$, Mann Whitney U).

Conclusions: The overall adherence of stroke patients with a brain training was low and there are some implications that systematic, regular interaction with a supervisor can increase training adherence of stroke patients with a restitution-focused intervention performed at home.

Introduction

Although stroke mortality rates in the past two decades have decreased, according to the World Health Federation stroke is still the second leading cause of death in the world. In 2010, the absolute number of people with first stroke was 16.9 million and the number of stroke survivors was 33 million.¹

Among the survivors of stroke, 22–50%^{2,3} experience cognitive impairment, such as aphasia, neglect, reduced processing speed and impaired attention,⁴ with direct consequences for dependency in activities of daily living and functional outcomes.⁵

Neurorehabilitation after stroke is focused on compensational strategy training and restitution-focused training.⁶ Compensational training aims to compensate for the lost function using remaining intact functions. Restitution-focused treatments consist of frequent repetition or stimulation of the affected function by high-intensity training.^{7,8} Therefore, therapists often prescribe intensive exercise regimes for patients.⁹ However, a study found that only 31% of patients actually performed exercises as recommended.¹⁰

Recently, computer-based cognitive rehabilitation (CBCR) programs, especially serious brain games, have emerged as a tool for restitution-focused treatment in stroke patients. It is expected that serious brain training helps patients in recovering from a stroke by making training more fun, as monotony of repeated motions is decreased and direct feedback about performance is provided.^{9,11,12} However, results about the effect of restitution-focused computer training are still conflicting.^{13–15} Studies are often hampered by low adherence rates,^{16–19} although this is one of the main requirements for success of an intervention as well as improved patient outcomes.²⁰ Laver et al. concluded in their review that studies should provide more detail in their reporting of adherence of stroke patients with CBCR interventions.¹⁵ Moreover, the impact of the extent of supervision on stroke patients' adherence with restitution-focused interventions is unknown. It was found in a review of Kelders et al. that frequency of interaction with a counselor was a significant predictor for adherence with web-based health interventions in different patient groups ($p < 0.001$).²¹

The goal of this study was to contribute to a better understanding of the impact of supervision on stroke patients' training adherence with restitution-focused interventions, as a potential factor to increase adherence, ultimately leading to better treatment outcomes for those recovering from stroke. The aim of the study was to describe stroke patients' adherence with a home-based 8-week brain training program (Lumosity Inc.[®]) by comparing two frequencies of health professional's supervision. The hypothesis of the study was that a CBCR training with more supervision would lead to higher training adherence in stroke patients.²¹


Materials & methods

Study design

The present study on adherence was part of a randomized controlled trial (RCT) evaluating the effectiveness of an 8-week CBCR program on cognitive functioning, quality of life (QoL) and self-efficacy as compared to a passive intervention.²² In this study no effect of the CBCR program were found on cognitive functioning, quality of life or self-efficacy when compared to the control group, except for very limiting effects on working memory and speed. A profile of the study is shown in Figure 1.

The current study compares patients in the original intervention group who received supervision eight times during the CBCR intervention period (S8) vs. the original control group who underwent the CBCR intervention after the original RCT and received supervision twice during the intervention period (S2). For the present analysis the data from all patients who agreed to take part in the program were used.

Patients in the S2 group received weekly information about stroke during the period that the S8 group received their intervention. The information provision was not interactive, it provided unidirectional information about brain differences between men and woman, the influence of stress on brain function and possible difficulties with living with a damaged brain. No new information was provided in these brain facts that were not already extensively addressed during previous rehabilitation treatment. Each week, during a period of 8 weeks, new information (text or a video clip) was added to the website. The study was approved by the Medical Ethical Review Board of The Leiden Medical Center (P 12.190). The CONSORT (Consolidated Standards of Reporting Trials) guidelines were used for adequate reporting of the study.²²



*Weekly contacts with a supervisor during the 8-week training period (S8)

^Contacts with a supervisor twice (S2)

Figure 1. Study profile.

Recruitment and inclusion

Inclusion criteria for participation in the study were: age between 45 and 75 years, diagnosed with stroke 12–36 months ago, having self-perceived cognitive impairments (extracted from a checklist accompanying the recruitment letter), having access to the Internet, being able to visit the rehabilitation center and having time to participate. Exclusion criteria were: antidepressant use; receiving actual treatment for cognitive impairments; severe aphasia; lack of computer skills; not being proficient in Dutch; participants with psychological disorders in need of treatment; patients with physical disorders known to impact cognition. Patients were recruited from the participating rehabilitation centers.

The recruitment procedure is described in more detail in a previous report.²³ In total, 142 patients meeting the inclusion criteria were screened for eligibility, of whom 53 were eventually randomised to the S8 group and 57 patients to the S2 group. 50 patients (94%) in the treatment group and 57 patients (100%) in the control group completed the study. Of the S2 group, 52 accepted the offer to participate in the program after the trial was completed. A flow chart of the inclusion is shown in Figure 2.

Intervention

The CBCR intervention was a home-based brain training program with duration of 8 weeks. The duration of 8 weeks for both the intervention and the follow-up were based on clinical expertise by the research team and the health professionals involved in the project team. All participants received a user identification and password to log on to a website providing access to the brain training (www.spelenderwijsbeter.nl). The training software was supplied by Lumosity Inc.[®]. This program was selected because it targets multiple cognitive domains and adapts the level of difficulty of games to patients' own abilities. In total, 16 games were used targeting five cognitive domains: attention, speed, memory, flexibility, and problem solving.

The minimum requested total playtime was 600 min. Patients were encouraged to complete at least one session a day (approximately 15–20 min) on at least 5 days per week. Each session, random selections of three games were assigned to the participant, each game lasting about five minutes. Patients were able to play longer after finishing the training session. Furthermore, participants were instructed to complete an extra game session when they missed one game session and/or were not able to play 5 days a week. With each game, all patients started at the same level of difficulty. The difficulty level was then raised or lowered depending on the performance in the previous round of the respective game. The software provided feedback about game scores and how much games were completed. Patients could receive reminders for training by email or a text message for mobile phone.




Figure 2. Flow of inclusion.

Supervision

During the training period, patients from the S8 group received digital support by a supervisor. Supervisors were three health care providers (a psychologist, physical therapist, and occupational therapist) from the two rehabilitation centers that participated in the study. The supervision consisted of a short meeting with their own supervisor at the first assessment day in the rehabilitation center. Moreover, digital support was provided to patients weekly during the 8-week training period by their supervisor by telephone if training adherence was lower than five times a week.

A structured plan with instructions and a timeline was provided to the supervisors. Moreover, during a meeting at the start of the intervention and evaluations during the intervention these instructions were discussed with the supervisors. Supervisors were instructed that contact was aimed at: (1) providing assistance needed to solve problems impairing a patient to play (e.g. help with software problems, explaining game instructions, install a reminder for training appointments, etc.), (2) providing strategies to achieve or improve adherence by using the PlanDo-Check-Act method and (3) encourage patients to increase training frequency using motivational interviewing. Moreover, patients were able to contact their supervisor themselves by email or telephone anytime they needed assistance, for instance, in case of problems with training software or questions about a certain game.

Patients from the S2 group received supervision twice during the training period: a short meeting with their supervisor at the start of the brain training and contact by telephone once, after 4 weeks of training. They were encouraged to contact the supervisor by email or telephone in case they experienced difficulties using the training.

Assessments

The main outcome of interest in the current study was training adherence during the intervention period. Moreover, patient characteristics were used to determine which variables predict adherence. These data were retrieved from medical records and online questionnaires.

Adherence

Training adherence was measured by registering the patients' frequency of logging on to the website of the study during the total training period of 8 weeks and for each week. This was done in order to determine whether patients played 40 times in total (five times a week during 8 weeks) as was required. Data were automatically recorded by the software of the website and therefore gathered independently of the provider of the software. In addition, the patients' playtime, expressed as the minutes played during the 8-week training period for all cognitive domains together and for each cognitive domain (attention, memory, speed, flexibility, problem solving) were registered. These data were provided by Lumosity Inc.[®]. Logging into the brain training and not playing any game was not registered as playtime.

Patient characteristics

Demographic characteristics included gender, age (years) and level of education (low: primary and lower vocational education; middle: secondary and middle vocational education; high: higher vocational and university education).²⁴ In addition, living situation (alone/together with spouse or other(s)), daily functioning (dependent/independent) and participation in paid work (yes/no) were recorded. Stroke characteristics included the affected hemisphere (left/right/other), type of stroke (infarction/hemorrhage), time between stroke and enrollment and length of stay in the rehabilitation center (in months).

Physical and psychological characteristics included Health related quality of life (HRQoL), measured with a Dutch version of the short Stroke Specific Quality of Life Questionnaire (SSQoL) with higher scores indicating better quality of life (range 12–60),²⁵ self-perceived cognitive failures, measured with a Dutch version of the 25-item Cognitive Failures Questionnaire (CFQ) with higher scores indicating less cognitive failure (range 0–100)²⁶ and self-efficacy, measured with a Dutch version of the general self-efficacy scale (SES) with higher scores indicating greater self-efficacy (range 10–40).^{27,28}

Analysis

Patients' baseline characteristics and adherence with the CBCR program were analysed using descriptive analyses. Data were presented as the number with percentage, median with the range or mean with SD. It was tested if variables were normally distributed by means of the Kolmogorov-Smirnov normality-test. Differences between the S8 group and S2 group were analysed with independent t-tests, Mann-Whitney U tests or Chi-square test, where appropriate.

Logistic regression analyses were used to examine the impact of the extent of supervision on training adherence. Adherence (high and moderate versus low) was dichotomized to be used as dependent variable. Moderate/high adherence was defined as ≥ 300 min, as this was half of the required amount of total playtime (600 min). Low adherence was defined as < 300 min of total playtime. All cognitive domains were included. The impact of supervision (group S8/ group S2) was included as independent variable, while adjusting for the following potential confounders: age, sex, educational level, type of stroke, affected hemisphere, cognitive failure (CFQ), quality of life (QoL), and self-efficacy (SES). Moreover, multiple logistic regression analyses with a stepwise backward selection procedure were executed in order to determine which variables predict adherence. Variables with the highest *P* value were removed one by one from the prediction model, until all remaining variables were $p < 0.05$. All analyses were performed with the SPSS statistical software package (version 21).

Results

Patient characteristics

The analysis on adherence concerned 53 patients who were allocated to group S8 and 52 patients in group S2, who agreed to participate in the CBCR program after the trial ended (Figure 2). Characteristics of the 105 patients included in the study are presented in Table 1. The mean age of all patients was 59 (range 46–74) and 66 (63%) of the patients were male. Patient characteristics were similar for group S8 (baseline) and group S2 ($t = 16$ weeks), except for more patients who had a hemorrhage in group S8 (21/53; 40%) compared to group S2 (11/52; 21%) ($p = 0.04$).

Adherence

A number of 21 out of the 105 (20%) included patients failed to complete any game. Reasons for not playing at all, as recorded by the supervisor, were: technical problems with the computer ($n = 6$), lack of motivation ($n = 6$), health problems ($n = 4$), vacation ($n = 3$), hospital stay ($n = 1$), finding training too difficult ($n = 1$). From the remaining 84 patients who completed at least one training session, 46 patients (87%) were in the S8 group and 38 (73%) in the S2 group ($p < 0.01$, Chi-Square). Data about adherence for both groups are presented in Table 2.

The median total play time was 424 min (range 27–216; 71%) in the total population. The median playtime in group S8 was 528 min (range 63–1264; 88%) vs. 193 min (range 27–2162; 32%) in group S2 ($p < 0.001$ Mann Whitney U). In the total group, 24 out of 84 patients (29%) played ≥ 600 min. In groups S8 and S2, 19/53 (36%) and 5/52 (10%), of the patients played ≥ 600 min ($p < 0.001$, Chi-Square). The median frequency of logging into the website during the training period was 66 (1–164) in group S8 and 47 (1–318) in group S2 ($p < 0.001$ Mann Whitney U). A number of seven patients from group S2 did not receive interaction with their supervisor after 4 weeks of brain training, since they did not respond to any of the calls.

The Odds Ratio of being in the moderate and high adherence group was 3.4 [95% CI 1.98–5.96] ($p = 0.00$) for patients who received weekly supervision (group S8), -0.9 [95% CI 0.86 –1.02] ($p = 0.06$) for older age (each year of life), 1.65 [95% CI 0.78–3.46] ($p = 0.19$) for higher self-perceived quality of life (SSQoL), 1.0 [95% CI 0.99–1.05] ($p = 0.26$) for higher subjective cognitive failure (CFQ), 1.1 [95% CI 0.92–1.08] ($p = 0.50$) for higher self-efficacy (SES) and 1.8 [0.67–4.60] ($p = 0.25$) for patients with a high educational level.

Table 1. Baseline characteristics of patients with stroke who participated in an 8-week cbcr program, presented for all patients and per group.

	All patients (n=105)	Group 1 (n=53)	Group 2 (n=52)	Between groups (p-value)^a
Age in years; median (range)	59 (46-74)	59 (46-74)	59 (46-73)	0.85
Sex, male;	66 (63)	34 (64)	32 (62)	
Type of stroke:				
Infarction, n (%)	68 (67)	29 (55)	41 (79)	
Haemorrhage, n (%)	34 (32)	21 (40)	11 (21)	0.04
Unknown ‡, n (%)	3 (6)	3 (5)	0 (0)	
Location of stroke:				
Hemisphere left, n (%)	50 (48)	23 (43)	27 (52)	
Hemisphere right, n (%)	50 (48)	26 (49)	24 (46)	0.21
Basal ganglia, n (%)	3 (3)	3 (6)	0 (0)	
Unknown ^b , n (%)	2 (2)	1 (2)	1 (2)	
Time from stroke onset to randomization in months; mean (SD)	25 (8.2)	26 (9.1)	25 (7.5)	0.32
Time spent in rehabilitation centre in months; mean (SD)	5.8 (3.6)	6 (3.6)	5 (3.4)	0.23
Educational status:^c				
Low, n (%)	36 (34)	17 (32)	19 (36)	
Intermediate, n (%)	33 (32)	18 (34)	15 (29)	0.94
High, n (%)	36 (34)	18 (34)	18 (35)	
Living alone, n (%)	26 (25)	13 (25)	13 (25)	0.84
Independent in daily functioning, n (%)	101 (96)	52 (98)	54 (95)	0.35
Participation in paid work, n (%)	32 (31)	14 (26)	18 (35)	0.33
Subjective cognitive failure ^d ; median (IQR)	63 (20)	63 (19)	63 (20)	0.62
Quality of life ^e median (IQR)	3.7 (2.5)	3.7 (2.4)	3.8 (2.4)	0.57
Self-efficacy ^f ; median (IQR)	30 (15)	28 (14)	30 (15)	0.83

^a Differences between the groups were analysed with independent t-tests, Mann-Whitney U tests or Chi-square test, where appropriate.

^b No data were available for medical status.

^c Low: lower technical and vocational training; median: secondary technical and vocational training; and high: higher technical and vocational training and university.

^d Measured with the Cognitive Failures Questionnaire (CFQ): range 0-100; higher scores indicating less cognitive failure.

^e Measured with the Stroke Specific Quality of Life Questionnaire (SSQoL): range 12-60; higher scores indicating better quality of life.

^f Measured with the General Self Efficacy Scale (GSES): range 10-40; higher scores indicating greater self-efficacy.

Table 2. Training adherence with an 8-week cbc program, presented for all stroke patients and per group^a.

	All patients (n = 84)	Group 1 (n = 46)	Group 2 (n = 38)	Between groups (P-value)^b
Total time played (minutes):	424 (27–2162)	528 (63–1264)	193 (27–2162)	<0.001
Total time played per cognitive domain (minutes), %:				
Attention	58 (2–466) (14)	60 (3–408) (11)	37 (2–466) (19)	0.002
Speed	49 (1–139) (12)	53 (3–139) (10)	6 (1–48) (3)	<0.001
Memory	168 (4–646) (40)	232 (26–646) (44)	95 (4–645) (49)	<0.001
Flexibility	87 (4–1009) (21)	109 (6–349) (20)	50 (4–1009) (26)	0.001
Problem solving	53 (1–265) (13)	81 (1–265) (15)	7 (2–168) (3)	<0.001
Frequency of logging on to the website (number):	50 (1–318)	66 (1–164)	47 (1–318)	<0.001

^a Presented as the median total play time (in minutes) with the range and percentage of the total play time, unless indicated otherwise.

^b Differences between the groups were analysed with independent t-tests, Mann–Whitney U tests or Chi-square test, where appropriate.

Discussion

The aim of this study was to compare two different types of support on training adherence in chronic stroke patients. This study found that only 24 out of 105 patients (23%) were able to complete the required amount of playtime (600 min) of brain training. Training adherence was significantly higher for the patients in group S8 (median 528 min, range 63–1264) compared to patients in group S2 (median 193 min, range 27–2162). A small proportion of patients (5/52, 10%), who had interaction with a supervisor twice (group S2), were able to complete the training. Therefore, there are some implications that systematic, regular interaction with a supervisor can increase stroke patients' training adherence with a restitution-focused intervention performed at home.

The observation of non-adherence to a CBCR training program in patients with stroke is in line with findings of other studies.^{16–19} The adherence rate for all patients found in the current study was 29% (36% in group S8 vs. 10% in group S2), but cannot be compared to those studies because of lack of adequate reporting of training adherence.¹⁵ Compared to an average adherence rate of 50% found in a review of 101 publications about Web-based interventions in different areas (chronic conditions, lifestyle, and mental health) adherence was low,²¹ which confirms non-adherence with CBCR programs is a problem

among patients with stroke.

Although a number of patient characteristics were examined with respect to their association with training adherence, apart from the extent of supervision, no other predictors were found in the current study. Therefore, it remains unclear for which patients with stroke a CBCR program in its current form is most suitable. The intensity of training (600 min within 8 weeks) might be too demanding for stroke patients. It should be further investigated if lower intensity of training can improve adherence among patients with stroke.

The current study has a number of limitations. First, patients in the S8 group were allocated to the intervention group during the RCT and probably received in general more attention (outside the weekly contact) compared to patients from group S2. Second, interactions between the supervisors and patients were not logged (e.g. topics of conversation and duration of contacts) and could therefore not be verified. It cannot be truly concluded that patients who received more support from supervisors showed greater levels of adherence than those who received very little support.

Third, there should be the same time difference between being informed about the study and the start of the intervention in both groups. Patients in the S8 group started the training straight away when it was still exciting and novel, while patients in the S2 group had to wait 16 weeks when excitement could have waned. On the other hand, those patients were probably still motivated to participate in the brain training, given the option to start the training themselves (self-selection). Moreover, the control group has been given some information about stroke at the time when the intervention group was undergoing the brain training procedure. This passive intervention could have had limited impacts on the patients. But the information was very general and previously received by patients during their rehabilitation process.

In conclusion, the overall adherence with an online brain training was low and it seems that serious brain training is not suitable for all initially motivated chronic stroke patients. Moreover, despite a number of methodological limitations in the study design, this study provides ground to further investigate the effect of the extent of supervision on training adherence of stroke patients with restitution-focused training. Although only little or no effects were found on cognitive functioning, self-efficacy and quality of life of the restitution-focused brain training in the overall study,²³ this seems important since low adherence rates undermine the effect of interventions. A future study is recommended comparing three groups operating simultaneously with one intervention group receiving weekly support, a second intervention group receiving only two episodes of support and a control group.

Disclosure statement

No potential conflict of interest was reported by the authors.

Reference list

1. Feigin VL, Roth GA, Naghavi M, et al. Global burden of stroke and risk factors in 188 countries, during 1990–2013: A systematic analysis for the Global Burden of Disease Study. *Lancet Neurol.* 2013;2014:245–255.
2. Douiri A, Rudd AG, Wolfe CD. Prevalence of poststroke cognitive impairment: South London stroke register 1995–2010. *Stroke.* 2013;44(1):138–145.
3. Gargano JW, Reeves MJ. Sex differences in stroke recovery and strokespecific quality of life: Results from a statewide stroke registry. *Stroke.* 2007;38(9):2541–2548.
4. Cumming TB, Marshall RS, Lazar RM. Stroke, cognitive deficits, and rehabilitation: Still an incomplete picture. *Int J Stroke.* 2013;8(1):38–45.
5. Appelros P, Nydevik I, Terént A. Living setting and utilisation of ADL assistance one year after a stroke with special reference to gender differences. *Disabil Rehabil.* 2006;28(1):43–49.
6. van de Ven RM, Murte JM, Veltman DJ, Schmand BA. Computer-based cognitive training for executive functions after stroke: A systematic review. *Front Human Neurosci.* 2016; 10:150.
7. Hamzei F, Liepert J, Dettmers C, Weiller C, Rijntjes M. Two different reorganization patterns after rehabilitative therapy: An exploratory study with fMRI and TMS. *NeuroImage.* 2006;31:710–720. doi:10.1016/j.neuroimage.2005.12.035.
8. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: A systematic review. *Lancet Neurol.* 2009;8:741–754.
9. Alankus G, Lazar A, May M, Kelleher C. Towards customizable games for stroke rehabilitation. Proc SIGCHI Conf Human Factors Comput Syst. 2010: 2113–2122.
10. Shaughnessy M, Resnick BM, Macko RF. Testing a model of poststroke exercise behavior. *Rehabil Nursing Official J Assoc Rehabil Nurses.* 2006;1:15–21.
11. Lumsden J, Edwards EA, Lawrence NS, Coyle D, Munafò MR. Gamification of cognitive assessment and cognitive training: A systematic review of applications and efficacy. *JMIR Serious Games.* 2016;4(2).
12. Cicerone KD, Langenbahn DM, Braden C, et al. Evidence-based cognitive rehabilitation: Updated review of the literature from 2003 through 2008. *Arch Phys Med Rehabil.* 2011;92(4):519–530.
13. Poulin V, Korner-Bitensky N, Dawson DR, Bherer L. Efficacy of executive function interventions after stroke: A systematic review. *Top Stroke Rehabil.* 2012;19(2):158–171.
14. das Nair R, Cogger H, Worthington E, Lincoln NB. Cognitive rehabilitation for memory deficits after stroke – An updated review. *Stroke.* 2017;48(2):e28–e29.
15. Laver KE, George S, Thomas S, Deutsch JE, Crotty M. Virtual reality for stroke rehabilitation. *Cochrane Library.* 2015.
16. Cruz VT, Pais J, Bento V, et al. A rehabilitation tool designed for intensive web-based cognitive training: Description and usability study. *JMIR Res Protoc.* 2013;2(2):59.
17. Bergquist TF, Yutis M, Sullan MJ. Satisfaction with cognitive rehabilitation delivered via the internet in persons with acquired brain injury. *Int J Telerehabil.* 2014;6(2):39.
18. Connor BB, Standen PJ. So much technology, so little time: Factors affecting use of computer-based brain training games for cognitive rehabilitation following stroke. Proc 9th Int Conf Disabil Virtual Reality Assoc Tech. 2012: 53–59.
19. Åkerlund E, Esbjörnsson E, Sunnerhagen KS, Björkdahl A. Can computerized working memory training improve impaired working memory, cognition and psychological health? *Brain Inj.*
20. Duncan PW, Horner RD, Reker DM, et al. Adherence to postacute rehabilitation guidelines is associated with functional recovery in stroke. *Stroke.* 2002;33(1):167–178.
21. Kelders SM, Kok RN, Ossebaard HC, Van Gemert-Pijnen JE. Persuasive system design does matter: A systematic review of adherence to webbased interventions. *J Med Internet Res.* 2012;14(6):152.
22. Moher D, Hopewell S, Schulz KF, et al. CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomised trials. *J Clin Epidemiol.* 2010;63(8):e1–e37.
23. Wentink MM, Berger MAM, de Kloet AJ, et al. The effects of an 8-week computer-based brain training programme on cognitive functioning QoL and self-efficacy after stroke. *Neuropsychol Rehabil.* 2016;26(5):847–865.

24. Centraal Bureau voor de Statistiek (CBS). The Dutch Standard Classification of Education, SOI 2006 [PDF file]; 2006. <https://www.cbs.nl/en-gb/background/2008/24/the-dutch-standardclassificationof-education-soi-2006>.
25. Post MWM, Boosman H, van Zandvoort MM, Passier PECA, Rinkel GJE, Visser-Meily JMA. Development and validation of a short version of the stroke-specific QoL scale. *J Neurol Neurosurg Psychiatry*. 2011;82:283–286.
26. Broadbent DE, Cooper PF, FitzGerald P, Parkes KR. The cognitive failures questionnaire (CFQ) and its correlates. *Br J Clin Psychol*. 1982;21:1–16.
27. Merckelbach H, Muris P, Nijman H, de Jong PJ. Self-reported cognitive failures and neurotic symptomatology. *Personality Individ Differ*. 1995;20:715–724.
28. Schwarzer R, Jerusalem M. Generalized self-efficacy scale. *Meas Health Psychol User's Portfolio. Causal Control Beliefs*. 1995;1:35–37.



The background of the page is a light gray with a pattern of overlapping hexagons. Some hexagons are solid, while others are outlines. Plus signs (+) are scattered throughout the pattern, some inside hexagons and some outside. The overall aesthetic is clean and modern, suggesting a medical or technological theme.

The patient perspective on the use of information and communication technologies and e-health in rehabilitation

M.M. Wentink
E. Prieto
A.J. de Kloet
T.P.M. Vliet Vlieland
J.J.L. Meesters

*Disability and Rehabilitation:
Assistive Technology* 2018;13(7):620-625.



Abstract

Introduction: Success of e-health relies on the extent to which the related technology, such as the electronic device, is accepted by its users. However, there has been limited research on the patients' perspective on use of e-health-related technology in rehabilitation care.

Objective: To explore the usage of common electronic devices among rehabilitation patients with access to email and investigate their preferences regarding their usage in rehabilitation.

Methods: Adult patients who were admitted for inpatient and/or outpatient rehabilitation and were registered with an email address were invited to complete an electronic questionnaire regarding current and preferred use of information and communication technologies in rehabilitation care.

Results: 190 out of 714 invited patients completed the questionnaire, 94 (49%) female, mean age 49 years (SD 16). 149 patients (78%) used one or more devices every day, with the most frequently used devices were: PC/laptop (93%), smartphone (57%) and tablet (47%). Patients mostly preferred to use technology for contact with health professionals (mean 3.15, SD 0.79), followed by access to their personal record (mean 3.09, SD 0.78) and scheduling appointments with health professionals (mean 3.07, SD 0.85).

Conclusion: Most patients in rehabilitation used one or more devices almost every day and wish to use these devices in rehabilitation.

Introduction

In The Netherlands, approximately 90,000 persons are admitted to specialized rehabilitation each year due to illness, an accident or a congenital disease [1]. Multidisciplinary inpatient and outpatient rehabilitation is one of the most expensive health care sectors in the Netherlands [2]. Rising healthcare costs and decreasing number of health professionals [3] as well as the increasing number of patients having access to and using the Internet warrant the need for innovative and efficient rehabilitation strategies.

E-health allows cost-effective disease management as well as patients' empowerment and health promotion [4]. The definition of e-health is "the use of new Information and communication technologies (ICT), mostly internet technology, to improve or support health and health care" [4]. Examples of e-Health in rehabilitation include virtual reality, computer games, assistive technology and online communication tools. Especially, long-term medical care needs could be addressed at significant lower expenditures [5], by means of improved accessibility to rehabilitation programs for clients with mobility impairments [6–8], expanded continuity of care and increased self-management by promoting personalized care, choice and personal autonomy [9]. Indeed, a number of studies showed that e-health is acceptable for subgroup of patients in rehabilitation care [9–11] and has the potential to support management of chronic conditions such as Alzheimer's/ dementia [12], diabetes [13,14] and COPD [15,16].

Despite the many advantages, limited uptake and non-use of ehealth interventions is still a common problem in health care [7,8]. The Normalization Process Theory explain problems with embedding an innovation in practice by the complex interplay between the new technology, individual actions and context [17]. In line with the Diffusion of Innovations theory of Rogers, the fit between the

needs of individuals and groups and the e-health services is of utmost importance [18]. Thus, identifying patients' preferences is required before development and implementation of e-health [18–20]. Indeed, in previous studies it was found that adoption of ehealth was associated with a persons' positive attitude toward technology, self-efficacy and perceived usefulness [19–21]. Moreover, Vankatesh's Unified Theory of Acceptance and Use of Technology (UTAUT) explains current performance and use behaviour explain a large proportion of the variance in the intention to use a new technology [22,23]. In addition, a few studies identified patients' preferences for e-health in rehabilitation by using a qualitative research design [7,24,25]. These studies found that patients with cancer and diabetes were willing to use ICT in (rehabilitation) care for self-monitoring of symptoms, web-based physical exercise programs [7], communication with peers [24,25] and access to their health record [7]. However, a lack of studies with a quantitative design impaired generalization of results and so far it is unclear what usage preferences are for other patient

groups in rehabilitation care.

In summary, for the rehabilitation setting specifically it is unclear which ICT devices are most commonly used by patients in rehabilitation and what their needs and preferences are regarding e-health services needs to be delivered.

This paper aims to contribute to future research and use of ICT tools in rehabilitation care. Therefore, we aim (1) to explore usage of ICT devices among rehabilitation patients and (2) to investigate patients' perspective to incorporate this technology in the rehabilitation process.

Methods

Study design

This cross-sectional study, involving a one-time online survey, was conducted between March 2014 and May 2014 among (former) patients who had been admitted to a Rehabilitation Centre in The Netherlands. The study protocol was presented to the Medical Ethical Committee of the Leiden University Medical Center. They judged the study as non-medical research according to the Medical Research Involving Human Subjects Act.

Patients

Patients were invited to participate in the study if they met the following criteria: (1) 18 years and older, (2) admitted for inpatient and/or outpatient rehabilitation between 2008 and 2013 and (3) being registered with an email address to select patients with access to and using ICT. Potentially eligible patients were first identified by searching the electronic patient registers of the rehabilitation centre. All eligible patients received an email with information about the study and an invitation to fill in the survey by using the digital link.

Survey

An online questionnaire was developed in collaboration with patient representatives in order to measure usage preferences of electronic devices in rehabilitation health care. The self-developed questionnaire comprised a maximum of 61 questions that aimed to identify current possession and use of ICT devices (maximum of 27 questions) and desired usage of ICT devices in the rehabilitation process in the future (maximum of 15 questions).

In addition, 19 questions were about following socio-demographics and disease characteristics: gender, age, living status (living alone or living with partner/family), educational level (low: up to and including lower technical and vocational training; medium: up to and including secondary technical and vocational training; and high: up to and including higher technical and vocational training and university) [26] and occupational

status (student, employed, unemployed and disabled or retired).

For 27 out of 61 questions, patients were able to select one or more answers from a given set of options questions (a minimal of two and a maximum of eight options), 14 questions required an open answer and 10 questions required the answer “Yes” or “No”. Moreover, 10 questions that were used to examine user preferences were initially measured on a 4-point Likert scale (1 ¼ totally disagree, 2 ¼ disagree, 3 ¼ agree and 4 ¼ totally agree). These items were used to calculate the mean per item in order to make a ranking from highest till lowest preference.

The questionnaire was pilot tested among 59 patients from the rehabilitation centre for completeness, feasibility, readability and presentation (e.g., perceived question difficulties, response errors, screen layout, etc.). The pilot testing led to minor changes in the wording and format of the final questionnaire.

A total of three reminders were sent for participation in the study. Patients received the first reminder 2 weeks after invitation. The second reminder was sent 1 week after the first reminder. All data from the online survey were collected anonymously.

Statistical analysis

Patient characteristics, possession and use of ICT devices and user preferences of ICT in rehabilitation were analyzed using descriptive statistics and presented as numbers with percentages, means with standard deviations (SD) or medians with ranges (Inter Quartile Range; IQR), i.e., 25th percentile–75th percentile) where appropriate. Results about possession and use of ICT devices were presented for the total group and for different age categories based on the 25% percentile distribution of age. Age was divided in four categories: (1) 18–36 years, (2) 37–51 years, (3) 52–61 years and (4) 62 > years. The group of eligible patients was compared to the group of responded patients regarding their age and gender using the independent t-tests and the Mann Whitney U test, the Chi-square test. All statistical analyses were performed using Statistical Packages for the Social Sciences (IBM SPSS 22.0 for Windows).

Results

According to the registers, 714 patients who were aged 18 years old, received outpatient or inpatient rehabilitation between 2008 and 2013 and were registered with an e-mail address, were identified and invited to participate in the study by email (Figure 1). A number of 233 patients filled in the questionnaire, from which 43 questionnaires were incomplete, resulting in a total of 190 completed questionnaires out of 714 invited patients (27%).




Figure 1. Flow of patients.

Patient characteristics

Table 1 shows that the 190 patients who completed the questionnaire had a mean age of 49 years ($SD=16$), 94 (49%) were female and 52 (27%) were living alone. The majority of patients followed either a “higher education” (41%) or “middle education” (41%) and 19% received “lower education”. Most patients were diagnosed with “acquired brain injury” (42%), followed by “neuromuscular disease” (13%) and “orthopaedics” (6%). The majority of the patients (73%) had outpatient rehabilitation treatment, 20 patients (10%) had inpatient rehabilitation, 26 patients (14%) had had both and two patients (1%) received a short rehabilitation program for heart rehabilitation.

The eligible patients had a mean age of 44 years ($SD=19$) and 341 (49%) were female. No significant differences were found for age and gender between the group of eligible patients and group of responded patients.

Table 1. Patient characteristics of 190 patients in rehabilitation care who participated in this cross-sectional study.

Characteristics	
Age in years (mean, SD)	49 (16)
Female gender (number, %)	94 (49)
Living status (number, %)	
Living alone	52 (27)
Living with partner/family	138 (73)
Educational level (number, %) ^a	
Lower	36 (19)
Middle	77 (40)
Higher	77 (40)
Type of rehabilitation (number, %)	
Inpatient	20 (10)
Outpatient	142 (73)
Both	26 (14)
None ^b	2 (1)
Diagnostic group (number, %)	
Acquired Brain Injury (stroke, tumour, trauma)	79 (42)
Cardiological	5 (3)
Chronic pain	9 (5)
Hand injury	4 (2)
Neuromuscular disease	24 (13)
Orthopaedics	13 (6)
Spinal cord injury	12 (6)
Trauma surgery (amputation)	4 (2)
Other	40 (21)

^a Low: up to and including lower technical and vocational training; medium: up to and including secondary technical and vocational training; and high: up to and including higher technical and vocational training and university.

^b A short heart rehabilitation program.

Possession and usage of ICT devices

Possession and use of ICT devices among the 190 patients who responded to the questionnaire are shown in Table 2 for the total group and four categories of age. The most frequently possessed ICT device was a computer/laptop (93%), followed by a smartphone (57%), tablet (47%), game console (16%), smart TV (15%) and e-reader (14%). Five patients (3%) possessed no ICT device at all. The mean number (SD) of possessed ICT devices per patient was 2.8 (SD ¼ 1.5).

In the first age category (1st quartile of patients between 18–37 years old), the mean number (SD) of possessed ICT devices was 3.1 (SD ¼ 1.6). The most possessed ICT devices

were the laptop (75%) and smartphone (75%). In the second age category (38–51 years old), the mean number of ICT devices was 2.9 (SD 1.5), with most possessed ICT devices being the smartphone (62%), the tablet (62%) and the pc (62%). In the third age category (52–61 years old), the mean number of ICT devices was 3.2 (SD 1.5). The two most possessed devices were the laptop (75%) and the pc (71%). In the fourth age category (62 > years old), the mean number of possessed ICT devices was 1.9 (SD 1.1). The two most possessed ICT devices were the pc (57%) and the laptop (47%).

A number of 149/190 (78%) responded to use their ICT devices all 7 days of the week, whereas two patients (1%) reported to use their devices less than once a week. From the patients who used their ICT device all 7 days a week, 36 patients (77%) were from the first age category (18–37 years old), 38 patients (91%) from the second age category, 37 (77%) from the third age category (52–61 years old) and 38 (75%) from the fourth age category (62 > years old).

The computer/laptop was the most frequently used ICT device by patients for e-mail (166/190 patients, 94%), to search for information (162/190 patients, 92%), for support (e.g., scheduling, banking and route planning) (137/190 patients, 78%) and social media (104/190, 59%). For physical and mental exercise the PC/ laptop was used by 39 patients (22%), followed by a tablet by 21 patients (24%), a smartphone by 12 patients (11%) and a game console by 8 patients (26%).

Table 2. Usage of ICT devices among 190 patients in rehabilitation care^a.

	All age categories	18–36 years	37–51 years	52–61 years	>62 years
Possession of ICT devices:^b					
PC/Laptop	176 (93)	46 (98)	37 (88)	47 (98)	46 (87)
Tablet	89 (47)	19 (40)	26 (62)	27 (56)	17 (32)
Smartphone	109 (57)	35 (76)	29 (69)	20 (63)	15 (28)
E-reader	27 (14)	5 (11)	9 (21)	8 (17)	5 (9)
Game console	31 (16)	18 (38)	5 (12)	7 (15)	1 (2)
Smart TV	29 (15)	9 (19)	5 (12)	11 (23)	4 (8)
No one of above	5 (3)	1 (2)	0 (0)	1 (2)	3 (6)
The frequency of use of any ICT device:					
Less than once a week	2 (1)	0 (0)	1 (2)	0 (0)	1 (2)
1–3 days per week	11 (6)	4 (9)	1 (2)	3 (6)	3 (6)
4–6 days a week	26 (14)	7 (15)	2 (5)	8 (17)	9 (18)
7 days per week	149 (78)	36 (77)	38 (91)	37 (77)	38 (75)

Table 2. Continued.

	All age categories	18–36 years	37–51 years	52–61 years	>62 years
Use of ICT device to search for information:^b					
PC/Laptop	162 (92)	40 (87)	36 (87)	44 (94)	42 (91)
Tablet	73 (82)	11 (58)	23 (86)	25 (93)	14 (82)
Smartphone	76 (40)	27 (77)	18 (62)	20 (67)	11 (73)
Use of ICT device for email:^b					
PC/Laptop	166 (94)	42 (91)	36 (97)	45 (96)	43 (93)
Tablet	62 (70)	11 (58)	18 (69)	19 (70)	14 (82)
Smartphone	84 (77)	27 (77)	22 (76)	24 (80)	11 (73)
Use of ICT device for social media:^b					
PC/Laptop	104 (59)	37 (80)	26 (70)	24 (51)	17 (37)
Tablet	47 (53)	14 (74)	10 (39)	13 (48)	10 (59)
Smartphone	61 (56)	29 (83)	14 (48)	14 (47)	4 (27)
Use of ICT device for physical and mental exercise:^b					
PC/Laptop	39 (22)	12 (26)	8 (22)	13 (28)	6 (13)
Tablet	21 (24)	6 (32)	5 (19)	6 (22)	4 (24)
Smartphone	12 (11)	5 (14)	5 (17)	2 (7)	0 (0)
Game console	8 (26)	5 (14)	2 (7)	0 (0)	1 (2)
Use of ICT device for support (e.g., scheduling, banking, route planning):^b					
PC/Laptop	137 (78)	34 (74)	31 (84)	37 (79)	35 (76)
Tablet	56 (63)	9 (47)	17 (65)	19 (70)	11 (65)
Smartphone	91 (83)	28 (80)	25 (86)	26 (87)	12 (80)

^a Indicated as the number of patients possessing the device (%).^b Patients could give more than one answer to each question.

Usage preferences of ICT devices in future rehabilitation

A top 10 of usage preferences of ICT devices for rehabilitation in the future are shown in Table 3. It was found that patients (highly) prefer to have digital contact with a health professional (mean 3.15, SD 0.79). Second, patients want to have digital access to their personal record in which both the patient and the health professional can make notes (mean, 3.09, SD 0.78). Third, digital scheduling of appointments with a health professional was highly preferred by most patients (mean 3.07, SD 0.85).

About 108 out of 190 patients (58%) agreed they want to fill in digital questionnaires about quality of care (mean 3.03, SD 0.84). Moreover, 93 patients (49%) agreed and 49 patients (26%) highly agreed they want to do exercises at home using a computer (mean 2.97, SD 0.80). 79/190 patients agreed (42%) and 34/190 patients (18%) highly agreed they

want to use ICT for self-measurement of health status (e.g., blood pressure) and forwarding the results to a health professional, while 62 patients (33%) disagreed and 13 patients (7%) highly disagreed. Contact with other patients (peers) was a preference of 90/190 patients (47%), but not for 55/ 190 patients (29%).

Less important user preferences for future rehabilitation were digital participation in group therapy under supervision of a health professional (mean 2.36, SD 0.86), receiving information and latest news from the rehabilitation centre (mean 2.56, SD 0.91) and getting support from health professionals at home (mean 2.61, SD ¼ 0.80).

Table 3. Patients' ranking of usage preferences of ICT devices in future rehabilitation^a.

Factor	Totally disagree	Disagree	Agree	Totally agree	Mean (SD)
1. Having contact with a health professional	9 (4.7)	19 (10)	95 (50)	65 (34)	3.15 (0.79)
2. Access to health record to make notes	9 (4.7)	22 (11.6)	101 (53.2)	56 (29.5)	3.09 (0.78)
3. Schedule appointments with health professional	12 (6.3)	26 (13.7)	87 (45.8)	63 (33.2)	3.07 (0.85)
4. Fill in questionnaires about quality of care	11 (5.8)	21 (11.1)	108 (56.8)	48 (25.2)	3.03 (0.84)
5. Exercises to do at home	8 (4.2)	38 (20)	93 (48.9)	49 (25.8)	2.97 (0.80)
6. Self-measurement and forwarding results to a health professional	13 (6.8)	62 (32.6)	79 (41.6)	34 (17.9)	2.71 (0.84)
7. Contact with other patients	18 (9.5)	55 (28.9)	90 (47.4)	25 (13.2)	2.65 (0.83)
8. Support from health professionals at home	15 (7.9)	65 (34.2)	86 (45.3)	22 (11.6)	2.61 (0.80)
9. Information and latest news from the rehabilitation centre	18 (9.5)	30 (15.8)	52 (27.4)	15 (7.9)	2.56 (0.91)
10. Participating in group therapy under supervision of a health professional	28 (14.7)	85 (44.7)	55 (28.9)	20 (10.5)	2.36 (0.86)

^a Data are presented as the number with percentages (%) unless indicated otherwise.

Discussion

This cross-sectional survey explored the usage of ICT devices and usage preferences to incorporate ICT devices in rehabilitation treatment among 190 adult rehabilitation patients from a rehabilitation centre in the Netherlands with access to email. We found that more than 90% of the responded patients used at least one ICT device, from which a computer/laptop most frequently, followed by a smartphone and then a tablet. Younger patients were found to use more devices than older patients, but older patients use their ICT devices with the same frequency as younger patients. Patients used their devices for e-mail, finding information, support (e.g., scheduling, banking, route planning), social media and physical

and mental exercises. Patients' usage preferences of ICT devices in rehabilitation were: having contact with health professionals (telecommunication), have access to their health record and scheduling appointments with health professionals.

Population based studies by the Dutch Central Agency for Statistics (2015) showed that in 2014 80% of the households in The Netherlands were possessing a laptop, 65% a PC, 63% a tablet and 74% a smartphone [27]. Comparing these data to the current study, possession of a computer/PC (93%) is slightly higher, although possession of a tablet (47%) and smartphone (57%) is lower among the responded patients in rehabilitation. Purposes of using ICT devices found in the current study were almost similar compared to the general population [28]. A more recent study amongst patients with multiple sclerosis showed that 86% (44/51) used a mobile phone [29]. From patients with cardiovascular disease, Buys et al. found most patients (97%) had a mobile phone, from which 64% owned a smartphone [30]. In the current study, possession of a smartphone was slightly lower (57%).

A few studies investigated patient preferences for use of ICT in (rehabilitation) health care using a qualitative research design [7,24,25]. Therefore, frequency data found in the current study could not be compared to other literature. First, this study showed that 101 out of 190 patients (53%) highly prefer to have insight in their personal record and also want to use it to make notes for health professionals. Although growing patient demand to online personal health record (PHR) access was already recognized in other studies [25,31–33], it is still not widely adopted. A study among 283 individuals over the age of 18 found that individual factors (satisfaction with provider, belief of the tool to be empowering) and environmental factors (communication tactics, technology characteristics and management support) influence intentions to use a PHR [34]. These aspects should be taken into account with the implementation of PHRs. Moreover, development of such systems should focus on patients and their families as well as on physicians and other healthcare professionals [33].

Second, communication with peers was identified as patient preference among 90/190 patients (47%) and also found in other studies [24–25]. Available evidence suggests that online peer-to-peer support interventions might be beneficial for users [35]. More research is needed to further investigate how ICT tools can be used to fulfil to patients' preferences for peer support. Third, we found a group of patients in rehabilitation preferred to use ICT to schedule appointments with their clinicians. This was not identified as a preference in previous studies. Moreover, a sub group of patients wanted to use ICT for self-monitoring and physical exercises, which is in line with preferences of patients with cancer [7] and cardiac patients [30].

Transferability of the findings for usage of devices to other contexts might be impaired, because participants were a subgroup from the general population of patients in rehabilitation and their treatment process might have influenced their preferences for e-health in future rehabilitation. More importantly, participants were from two rehabilitation

centres in one region in one country. There may be large differences in access to and usage of ICT among countries, hampering the generalizability. As an example, the highest proportion of households with internet access in Europe in 2016 was recorded in the Netherlands and Luxembourg (97%). By contrast, the lowest rate was found in Bulgaria (64%) [36].

A limitation of the current study is that only patients with an email address were invited to fill in the electronic questionnaire. However, we aimed to identify usage and preferences in patients who have access to ICT and having an email address suggests the latter. Future studies should also include patients in rehabilitation with low level of access to ICT [10], lack of ICT experience [10,22,37] and personal traits for ICT utilization (e.g., age and health condition) [38] in order to better understand how to enable all patients to benefit from of e-health.

In addition, only 190 out of 714 patients responded (27%) and data about possession and usage of ICT devices is approximately 2.5 years old. However, we compared it with data from 2015 and the availability of data like ours is scarce in health care and rehabilitation in particular. Moreover, to the best of our knowledge, this is one of the few studies investigating the use of ICT devices and usage preferences amongst patients in rehabilitation. Since a quantitative study design was used a high number of patients participated in the study (n ¼ 190).

This study found that the majority of patients in rehabilitation with a registered email address used one or more ICT devices every day of the week (younger patients more often than older patients). The most frequently used devices were a computer/laptop, smartphone and tablet. According to patients, e-health in rehabilitation needs to include online access to their health record, communication with peers and scheduling appointments with health professionals. To better assist patients with e-health in rehabilitation care in the future, further research is needed about how the preferences identified in the current study could be implemented in rehabilitation care by using the most commonly used ICT devices (computer/laptop, smartphone and tablet).

Acknowledgements

We gratefully want to thank all the patients that participated in this study.

Disclosure statement


No potential conflict of interest was reported by the author(s).

Reference list

1. Revalidatie Nederland. Revalidatie Nederland. [cited 2010 Oct 11]. Available from: <http://www.revalidatie.nl/revalideren/home-r>.
2. Slobbe LCJ, Smit JM, Groen J, et al. Kosten van ziekten in Nederland 2007: Trends in de Nederlandse zorguitgaven 1999-2010. RIVM rapport 2011;270751023.
3. Arbeidsmarkt Zorg en Welzijn (AZW). Arbeidsmarktprognoses van VOV-personeel in Zorg en Welzijn 2013-2017. Zoetermeer, 2015.
4. Wentzel MJ, Beerlage-de Jong N, Sieverink F, et al. Slimmer eHealth ontwikkelen en implementeren met de CeHRes Roadmap. In: Politiek C, Hoogendijk R, editors. CO-CREATIE EHEALTHBOEK eHealth, technisch kunstje of pure veranderkunde? The Netherlands: Nictiz; 2014.
5. Ueckert F, Goerz M, Ataian M, et al. Empowerment of patients and communication with health care professionals through an electronic health record. *Int J Med Informat.* 2003; 70:99–108.
6. Lindberg B, Nilsson C, Zotterman D, et al. Using information and communication technology in home care for communication between patients, family members, and healthcare professionals: a systematic review. *Int J Telemed Appl.* 2013;2013:461829.
7. Timmerman JG, Tonis TM, Dekker-van Weering MG, et al. Co-creation of an ICT-supported cancer rehabilitation application for resected lung cancer survivors: design and evaluation. *BMC Health Serv Res.* 2016; 16:1.
8. Broens TH, Vollenbroek-Hutten MM, Hermens HJ, et al. Determinants of successful telemedicine implementations: a literature study. *J Telemed Telecare.* 2007; 13:303–309.
9. Brennan DM, Mawson S, Brownsell S. Telerehabilitation: enabling the remote delivery of healthcare, rehabilitation, and self management. *Stud Health Technol Inform.* 2009; 145:231–248.
10. Stendal K. How do people with disability use and experience virtual worlds and ICT: a literature review. *J Virtual Worlds Res* 2012; 5:1.
11. Matthew-Maich N, Harris L, Ploeg J, et al. Designing, implementing, and evaluating mobile health technologies for managing chronic conditions in older adults: a scoping review. *JMIR mHealth Uhealth.* 2016; 4:e29.
12. De LG, Brivio E, Sautter SW. Supporting autobiographical memory in patients with Alzheimer's disease using smart phones. *Appl Neuropsychol.* 2011;18:69–76.
13. Durso SC, Wendel I, Letzt AM, et al. Older adults using cellular telephones for diabetes management: a pilot study. *Med Surg Nurs.* 2003;12:313–317.
14. Rollo ME, Ash S, Lyons-Wall P, et al. Trial of a mobile phone method for recording dietary intake in adults with type 2 diabetes: evaluation and implications for future applications. *J Telemed Telecare.* 2011;17:318–323.
15. Nguyen HQ, Gill DP, Wolpin S, et al. Pilot study of a cell phone-based exercise persistence intervention postrehabilitation for COPD. *Int J Chron Obstruct Pulmon Dis.* 2009;4:301–313.
16. Liu W, Wang C, Lin H, et al. Efficacy of a cell phone-based exercise programme for COPD. *Eur Respir J.* 2008;32: 651–659.
17. May C, Finch T. Implementing, integrating and embedding practices: an outline of normalization process theory. *Sociology* 2009;43:535–554.
18. Rogers EM. The diffusion of innovation. 4th ed. New York: Free Press; 1995.
19. Young R, Willis E, Cameron G, et al. "Willing but unwilling": attitudinal barriers to adoption of home-based health information technology among older adults. *Health Informatics J.* 2013; 20:127–135.
20. Porter CE, Donthu N. Using the technology acceptance model to explain how attitudes determine Internet usage: the role of perceived access barriers and demographics. *J Bus Res.* 2006; 59:999–1007.
21. Or CKL, Karsh BT. A systematic review of patient acceptance of consumer health information technology. *J Am Med Inform Assoc.* 2009;16:550–560.
22. Venkatesh V, Morris MG, Davis GB, et al. User acceptance of information technology: toward a unified view. *MIS Q.* 2003;27:425–478.
23. Venkatesh V. Determinants of perceived ease of use: integrating control, intrinsic motivation, and emotion into the technology acceptance model. *Informat Syst Res.* 2000;11: 342–365.

24. Armstrong N, Hearnshaw H, Powell J, et al. Stakeholder perspectives on the development of a virtual clinic for diabetes care: qualitative study. *J Med Internet Res*. 2007; 9:e23.
25. Meijer W, Ragetlie P. Empowering the patient with ICTtools: the unfulfilled promise. *Stud Health Technol Inform*. 2007; 127:199–218.
26. Centraal Bureau voor de Statistiek (CBS). The Dutch Standard Classification of Education, SOI 2006 [cited 2016 Oct 9]. Available from: <https://www.cbs.nl/en-gb/background/2008/24/the-dutch-standardclassification-of-education-soi-2006>.
27. Centraal Bureau voor de Statistiek (CBS). Trendbreukanalyse ICT 2014 [cited 2016 Oct 11]. Available from: <https://www.cbs.nl/NR/rdonlyres/97A03A1B-0503-49A0-9181-AAF5B04183D2/0/trendbreukanalyseict2014.pdf>.
28. Centraal Bureau voor de Statistiek (CBS). ICT-vaardigheden van Nederlanders [cited 2016 Nov 11]. Available from: <https://www.cbs.nl/-/.../2016st06-ict-vaardigheden-vannederlanders.pdf>.
29. Van Kessel K, Babbage DR, Reay N, et al. Mobile technology use by people experiencing multiple sclerosis fatigue: survey methodology. *JMIR mHealth Uhealth*. 2017; 5:e6.
30. Buys R, Claes J, Walsh D, et al. Cardiac patients show high interest in technology enabled cardiovascular rehabilitation. *BMC Med Inform Decis Mak*. 2016; 16:95.
31. de Lusignan S, Mold F, Sheikh A, et al. Patients' online access to their electronic health records and linked online services: a systematic interpretative review. *BMJ Open*. 2014;4:e006021.
32. Liu CF, Tsai YC, Jang FL. Patients' acceptance towards a web-based personal health record system: an empirical study in Taiwan. *Int J Environ Res Public Health*. 2013;10:5191–5208.
33. Ose D, Baudendistel I, Pohlmann S, et al. Personal health records on the Internet. A narrative review of attitudes, expectations, utilization and effects on health outcomes. *Z Evid Fortbild Qual Gesundheitswes*. 2017;122:9–21.
34. Agarwal R, Anderson C, Zarate J, et al. If we offer it, will they accept? Factors affecting patient use intentions of personal health records and secure messaging. *J Med Internet Res*. 2013; 15:e43.
35. Eysenbach G, Powell J, Englesakis M, et al. Health related virtual communities and electronic support groups: systematic review of the effects of online peer to peer interactions. *BMJ* 2004;328:1166.
36. Eurostat Statistics explained. Digital economy and society access. Digital economy and society statistics – households and individuals [cited 2017 July 13]. Available from: http://ec.europa.eu/eurostat/statisticsexplained/index.php/Digital_economy_and_society_statistics_-_households_and_individuals.
37. Dunnebeil S, Sunyaev A, Blohm I, et al. Determinants of physicians' technology acceptance for e-health in ambulatory care. *Int J Med Inform*. 2012; 81:746–760.
38. Heart T, Kalderon E. Older adults: are they ready to adopt health-related ICT? *Int J Med Informatics*. 2013; 82: e209–e231.





How to improve eRehabilitation programs in stroke care? A focus group study to identify requirements of end-users.

M. Wentink
L. van Bodegom-Vos
B. Brouns
H. Arwert
S. Houdijk
P. Kewalbansing
L. Boyce
T. Vliet Vlieland
A. de Kloet
J. Meesters

BMC Medical informatics and Decision Making 2019;19(1):145.



Abstract

Background: A user-centered design approach for eHealth interventions improves their effectiveness in stroke rehabilitation. Nevertheless, insight into requirements of end-users (patients/informal caregivers and/or health professionals) for eRehabilitation is lacking. The aim of this study was to identify end-user requirements for a comprehensive eHealth program in stroke rehabilitation.

Methods: Eight focus groups were conducted to identify user requirements; six with patients/informal caregivers and two with health professionals involved in stroke rehabilitation (rehabilitation physicians, physiotherapists, occupational therapists, psychologists, team coordinators, speech therapist). The focus groups were audiotaped and transcribed in full. Direct content analysis was used to identify the end-user requirements for stroke eHealth interventions concerning three categories: accessibility, usability and content.

Results: In total, 45 requirements for the accessibility, usability and content of a stroke eRehabilitation program emerged from the focus groups. Most requirements concerned content (27 requirements), followed by usability (12 requirements) and accessibility (6 requirements). Patients/informal caregivers and health professionals each identified 37 requirements, respectively, with 29 of them overlapping.

Conclusions: Requirements between stroke patients/informal caregivers and health professionals differed on several aspects. Therefore, involving the perspectives of all end users in the design process of stroke eRehabilitation programs is needed to achieve a user-centered design.

Trial registration: The study was approved by the Medical Ethical Review Board of the Leiden University Medical Center [P15.281].

Introduction

Stroke, or a cerebrovascular accident (CVA), often occurs when a clot in the blood vessel blocks the blood flow to the brain cells (ischemic stroke) or when a blood vessel in the brain breaks or ruptures (hemorrhagic stroke). Subsequently, brain cells are deprived of oxygen and glucose, causing damage to the brain tissues. Although stroke mortality rates have decreased in Western countries, the prevalence of stroke is increasing [1].

Stroke survivors can experience lasting impairments with disruption of psychological and social well-being, including activities of daily life, cognitive and emotional functioning and social relationships [2,3]. Therefore, stroke rehabilitation is a comprehensive, multi-dimensional process including multiple interventions aimed at individual treatment goals in impairment, activity or participation [4], which involves both the patient, their informal caregivers and various health professionals (e.g. physicians, physical and occupational therapists, speech-language pathologists, psychologists) [5].

EHealth is proposed as a useful tool to improve efficiency and quality of rehabilitation care [6,7]. Ehealth is defined as ‘the use of Information and Communication Technology (ICT) to improve or support interventions in health care’. Consequently, (the effectiveness of) use of eHealth in rehabilitation, also known as eRehabilitation, after stroke has become a research area of interest [8-12]. Examples of stroke eRehabilitation programs used in studies are serious brain games, virtual reality or telerehabilitation [8,13,14].

Despite widespread agreement about the potentials of eRehabilitation, eHealth interventions often do not match with the requirements of intended users (e.g. patients and health care professionals), impairing their adoption in health care [15]. Therefore, a ‘user-centered design approach’, in which the requirements/needs of end users are taken into account at each stage of the design of a new product, intervention or service is highly recommended [16-22]. Requirements of end users involved in stroke care (patients, their informal caregivers and different health professionals) can be identified by means of qualitative research (e.g. interviews, focus groups, brainstorm sessions) [23,24].

Studies assessing requirements/needs of intended users of an eRehabilitation program [17, 25-32] found concerning the content, that interventions should be adapted to the patients’ own circumstances [25,30,31] (including personal goals [17,27]), should deliver rewarding feedback [32] and need to demonstrate outcomes on training performances [25,27,28,31]. Moreover, eRehabilitation programs must be user-friendly [17,25,28,31] (e.g. size of buttons, colors, information delivery, instructions etc.).

However, requirements should not only be identified for the content and usability of eRehabilitation programs, but also for their accessibility in order to enable successful adoption of eRehabilitation in health care [33,34]. This is important since easy access of eHealth technology (accessibility) allows users to start using it and the extent to which

the technology can be used allows users to achieve specified goals (usability) so that users benefit from the services provided (content). Moreover, it is argued that identifying user requirements for eHealth should go beyond functional and technical requirements and also needs to consider requirements for accessibility and acceptability [34,35,46].

In addition, the requirements in these studies mainly addressed only one aspect of stroke recovery (e.g. hand function, upper limb rehabilitation, weight shifting) or one technology tool (e.g. a game, robotica) [25,27-31] and not to a comprehensive eRehabilitation program in which multiple modalities are delivered. Although there have been some studies focusing on this area [37-39], requirements of intended users for comprehensive eRehabilitation that covers multiple aspects of stroke management are rather unknown.

Therefore, the aim of this study was to identify the requirements of end users (patients, their informal caregivers and health professionals) for the content, usability and accessibility of a comprehensive eRehabilitation program in stroke care.

Methods

Design

To identify the requirements for eRehabilitation in stroke care, a qualitative focus group study was employed with end users. In this study end users were patients with stroke, their informal caregivers and health professionals involved in stroke care. Focus groups are a useful method to gather information about perceptions of participants and to identify perceived requirements of subgroups [40]. The study took place between January 2016 and March 2016 in two rehabilitation centers in the Netherlands, both providing inpatient and outpatient multidisciplinary stroke care: Rijnlands Rehabilitation Center (RRC) in the city of Leiden and Sophia Rehabilitation (SR) in the area of The Hague.

Recruitment and inclusion

Patients and informal caregivers

Patients were recruited based on the following criteria: >18 years, diagnosed with stroke, completed rehabilitation which started after June 2011. From a group of circa 2.700 potential participants which are treated in one of the two rehabilitation centers, 200 patients of each rehabilitation center were randomly selected (Figure 1). Those 400 patients received a letter with information about the study and an invitation to participate from their former rehabilitation physician. Invitations to patients were directed to the informal caregiver as well, which could be a partner, child, parent or friend who is involved in the daily life of the patient. In addition, a group of five former stroke patients from SR (innovation partners),

who come together on a regular basis to discuss the newest innovations in rehabilitation, were invited.

The invitation included a self-developed questionnaire concerning marital status (single, married, divorced, widow/widower), daily activities ((un)paid job, household tasks, student), education level (low: up to and including lower technical and vocational training, medium: up to and including secondary technical and vocational training, and high: up to and including higher technical and vocational training and university), impairments as a consequence of stroke (physical, communication, cognition), use of ICT-devices (smartphone, tablet, laptop, pc) and the purpose of this use (applications, email, information, games, exercises). From the patients/informal caregivers that indicated their willingness to participate, we purposively selected both man and women, young (<20 age) and older (>70 age) patients, patients with less (e.g. using a computer only for mail) and more experience with digital devices (e.g. using a smartphone/tablet for applications) and patients with communication, cognitive and physical impairments. Patients with aphasia or severe cognitive problems were asked to bring their informal caregiver in order to help them represent their perspective.

Health professionals

Health professionals from the two rehabilitation centers (n=56, 29 at Sophia Rehabilitation, 27 at Rijnlands Rehabilitation Centre) were invited to participate by e-mail and selected based on the following criteria: a practiced and certified health professional (rehabilitation physician, physical therapist, occupational therapist, speech therapist and/or a psychologist) with \geq two years of working experience in multidisciplinary stroke care or working as a coordinator of a multidisciplinary team. A selection was made based on availability and profession, so that each profession was represented.

Data collection

Focus groups

It was planned to execute four focus groups at the two Rehabilitation centers, two with patients/informal caregivers and two with health professionals. Separate groups were organized with patients/informal caregivers vs health professionals in order to allow patients/informal caregivers to speak freely about experiences in the rehabilitation center and professionals to share their opinions about delivery of care. The aimed group size was 6 to 8 participants, although higher invitation rates were used to account for participants who would decline last minute [41].

A moderator (MW; Msc, female), assistant (BB; Msc, female/HB; Msc, female) and observer (SH; physiotherapist, male/PK; MD, female) conducted the focus groups. The assistant contributed with questions, made sure all participants were involved in the discussion

and managed the tape-recorders and time. The observer observed and took notes. The moderator and assistant had no involvement in patient care and a master's degree in Health Sciences/Human Movement Sciences including education about conduct of interviews. The moderator was trained in communication skills (listening, summarizing and disquisition). Patients/informal caregivers received travel costs reimbursement and were rewarded for participating with a gift card of 10 euro.

Interview guide

A semi-structured interview guide was developed with open-ended questions concerning three categories of eRehabilitation: accessibility, usability and content. These categories were based on research findings. Existing eHealth frameworks as a theory and guidance for the focus groups were considered by the research team, for instance the 'Technology Acceptance Model' [42], the 'Comprehensive Health Technology Assessment Framework' [43] and the 'Evaluation of e-health services: user's perspective criteria' [44]. However, most frameworks focus on the evaluation process [43-48] instead of the development process of eHealth technologies [49-52], and although the frameworks for development of eHealth admitted user requirements should be identified in an early stage, none of these frameworks described which aspects to explore.

In relation to the focus in this study, accessibility was defined by the research team as "easy access to eRehabilitation for all end users, including patients with disabilities as a consequence of stroke". Usability was defined as "the extent to which the eRehabilitation service can be used by the specified users (patients, informal caregivers and health professionals) to achieve specified goals with effectiveness, efficiency and satisfaction (e.g. recovery after stroke) in a specific context of use (e.g. during their stay in the rehabilitation center and/or at home)" [35]. Content was defined as "everything end users want to include in eRehabilitation (e.g. services, information, applications, etc.) to achieve specified goals for eRehabilitation in their rehabilitation process."

Examples of questions included were: "what ICT devices would you like to use for eRehabilitation?" (accessibility), "what aspects would make eRehabilitation easy to use?" (usability) and "what elements of care should be included in an eRehabilitation program?" (content). Prompts were included in the interview guide (e.g. example of eRehabilitation, pictures, etc.) to facilitate participants in verbalizing thoughts.

The interview guide was tested in a pilot focus group with a group of former stroke patients. No adjustments were made to the interview guide and therefore data from this focus group were included in the analysis. The focus groups lasted two hours, including breaks, and were audiotaped and transcribed in full.

Participants

Out of the 400 patients and their informal caregivers (200 at RRC/200 at SR) invited to participate in this study, 53 patients (27 at SR/26 at RRC) and 22 informal caregivers (11 at SR/11 at RC) responded. Reasons for non-response were not recorded. Of the 53 responded patients, 32 were invited to participate of whom 27 were present at the focus groups (Figure 1). Five innovation partners were also present, so that a total of 32 patients participated. Of these patients, 15 had an informal caregiver that participated with them in the focus groups. Seven out of these 15 caregivers were required to support a patient with aphasia or severe cognitive problems. In total 56 health professionals (29 at SR/27 at RCC) were invited, from which 22 responded (11 at SR/11 at RRC). Nine professionals were not able to attend the focus groups, so that eventually 13 professionals participated in the study (7 at SR/6 at RRC).

In total, eight focus groups were conducted; six with patients/informal caregivers and two with health professionals. The characteristics of all patients, informal caregivers and health professionals are presented in Table 1.




Figure 1. Flow of inclusion

Table 1. Participants of the focus groups, including the pilot focus group, exploring end-user requirements for eRehabilitation in stroke care.

	Patients	Informal caregivers	Health professionals
Number of participants	321	15	13
Gender, male; number (%)	19 (59)	4 (27)	3 (23)
Age in years; mean (SD)	57 (15)	61 (10)	-
Time since stroke in months; mean (SD)	28 (14)	-	-
Physical impairment; number (%)	20 (63)	-	-
Problems with communication; number (%)	16 (50)	-	-
Cognitive impairment; number (%)	24 (75)	-	-
Using digital devices (laptop, tablet, smartphone) in daily life; number (%)	32 (100)	-	-
Purpose of using digital devices; number (%)^a:			
Access to email	18 (56)		
Access to applications	15 (47)		
Searching information	10 (31)		
Playing games	14 (44)		
Doing exercises	8 (25)		
Profession; number (%):			
Physiotherapist	-	-	3 (23)
Psychologist	-	-	1 (8)
Occupational therapist	-	-	3 (23)
Speech therapist	-	-	1 (8)
Rehabilitation physician	-	-	4 (31)
Team coordinator	-	-	1 (8)

^a Patients could give more than one answer to each question.

Ethical issues and approval

Informed consent was obtained from all participants. Participants were informed that all their comments were confidential and would be used to improve rehabilitation treatment. It was explicitly mentioned that participation would not affect future treatment in the rehabilitation center. Collected data were reported in such way that persons could not be identified. Only researchers involved in the data analysis had access to the data. The study was approved by the Medical Ethical Review Board of the Leiden University Medical Center [P15.281]. COREQ guidelines were used for adequate reporting of the study [53].

Data analysis

The audio-tapes of the focus groups were transcribed in full. Directed content analysis was

used [54], in which the three predetermined categories of the interview guide (accessibility, usability and content) were used as guidance for analyzing the data. First, two (MW, BB) out of four researchers (BB, MW, SH, PK) independently highlighted text that appeared to reflect a requirement for eRehabilitation (codes). These requirements were directly classified in one of the prescribed categories (accessibility, usability or content). Second, researcher MW examined the data to determine whether subcategories were needed and requirements with comparable content were merged (subcategories). A new category was added if one of the three prescribed categories were not sufficient for identified requirements. It was aimed to stop conducting focus groups when no additional categories of user requirements were found, indicating saturation [40]. In each step of the analysis, discrepancies were compared and discussed in order to reach consensus. When the two researchers (BB, MW) still disagreed, a third researcher (JM), made a final decision. The framework and illustrations were discussed with two other researchers (LB, JM). Transcripts and findings were not returned to participants for comments. The 2 software package Excel 2010 was used to organize codes, subcategories and categories.

Results

User requirements

In total, 45 user requirements (codes) for a comprehensive eRehabilitation program were identified for the three prescribed categories (accessibility, usability and content). No categories were added, since the three prescribed categories were sufficient for all identified requirements. The requirements were classified into a total number of eleven self-determined subcategories. Most subcategories and requirements were identified for Content (6 subcategories/27 requirements), followed by Usability (4 subcategories/12 requirements) and Accessibility (1 subcategory/6 requirements). The eleven subcategories are presented in Figure 2. An additional table presents the user requirements for eRehabilitation in stroke care for each (sub)category (Table 2).

A total number of 45 requirements were retrieved from the focus groups. Thirty-seven requirements were mentioned by patients/informal caregivers (6 for accessibility, 12 for usability and 19 for content) and 37 by health professionals (6 for accessibility, 9 for usability and 22 for content). Thirty-two requirements were overlapping between patients/informal caregivers and health professionals, 8 requirements were unique for patients/informal caregivers and 8 requirements were only mentioned by health professionals. The results will be further explained in the following sections by a description of the identified user requirements for each category within each category.




Figure 2. Subcategories of user requirements for the accessibility, usability and content of a stroke eRehabilitation program.

Table 2. Requirements for an eRehabilitation program in stroke care according to patients, informal caregivers and health professionals.

Categories	Subcategories	Requirements	Patients / caregivers	Professionals
Accessibility:	Access:	No internet connection is required to use eHealth interventions (offline use). (1)	X	X
		eHealth interventions are accessible without logging on each time. (2)	X	X
		Applicable to most commonly possessed ICT-devices (laptop, tablet and smartphone). (3)	X	
		Access for health professionals to the electronic patient record to stay informed about training results. (4)	X	
		Applicable on computers at the rehabilitation center and synchronization with programmes used for the electronic patient record. (5)		X
		Different eHealth interventions should be brought together in one central dashboard. (6)		X
Usability:	Product attributes (visual):	Use of pictograms, symbols and graphics. (1)	X	X
		Non-flashing and tranquil interface. (2)	X	X
		Adjustable lay-out settings (font style, font size, background and colors). (3)	X	

Table 2. Continued.

Categories	Subcategories	Requirements	Patients / caregivers	Professionals
Usability:	<i>Product attributes (auditory):</i>	Ability to listen to written text. (4)	X	X
		Sounds for alert or as feedback. (5)	X	
	<i>Product attributes (simplicity):</i>	Limited amount of open webpages as a consequence of using a service. (6)	X	X
		Limited amount of information on a single screen. (7)	X	
		Limited options on a single screen to click further to another screen. (8)	X	
	<i>Service (support):</i>	Menu with frequently asked questions (FAC). (9)	X	X
		Videos with instructions on how to use eRehabilitation. (10)	X	X
		Helpdesk. (11)	X	X
		Direct assistance at home/ workplace. (12)	X	
Content	<i>Personalized training facilities:</i>	Physical exercises. (1)	X	X
		Exercises for cognitive functioning. (2)	X	X
		Speech exercises. (3)	X	X
	<i>Tracking:</i>	Monitor activities in daily living (i.e. what activities and for how long). (4)	X	X
		A video system to record exercises at home. (5)	X	
		Monitor a patients' health status (e.g. body weight, heart rate function, etc.). (6)		X
	<i>Agenda and reminders:</i>	Insight in the rehabilitation schedule of a patient. (7)	X	X
		A reminder function for scheduled appointments. (8)	X	X
		Scheduled time to use eRehabilitation (digital training). (9)	X	X
		Scheduling appointments with health professionals on the initiative of patients and their informal caregivers. (10)	X	X
	<i>Communication:</i>	Contact with peers (patients) to share experiences on how to cope with having a stroke. (11)	X	X
		Contact with peers (care givers) to share experiences on how to cope with having a relative with stroke. (12)	X	X
		Communication between patients and their informal caregivers and health professionals from a distance (telecommunication). (13)	X	X

Table 2. Continued.

Categories	Subcategories	Requirements	Patients / caregivers	Professionals
Content	<i>Information:</i>	General information about stroke. (14)	X	X
		Hyperlinks to reliable and relevant web pages for patients with stroke and their informal caregivers. (15)	X	X
		Information about patient organizations. (16)	X	X
		Information on how to cope with consequences of stroke (psycho-education). (17)	X	X
		Descriptions on how to perform daily activities (strategy training). (18)	X	
		Insight in agreements and information discussed during a consult. (19)		X
		Insight in final reports of a patients' rehabilitation process. (20)		X
	<i>Goal setting and evaluation:</i>	Setting goals for eRehabilitation. (21)	X	X
		Evaluation of goals for eRehabilitation. (22)	X	X
		Feedback about training results (i.e. insight in what is trained, the number of completed training sessions and training outcomes). (23)	X	X
		Feedback on goals (i.e. when a goal is accomplished). (24)	X	X
		Use of clinical assessments for goal setting and goal evaluation. (25)		X
		Use of valid questionnaires for goal setting and goal evaluation. (26)		X
		Compare training outcomes of a single patient with those of other patients. (27)		X

Accessibility

Access

Most patients are interested in eRehabilitation, but not all patients want to use it for their recovery, because it is not suitable for them (e.g. lack of computer skills or disabilities impairing use of technology). This was also acknowledged by health professionals. [Professional_5: *"It is not realistic to strive for every patient to use eHealth. You can offer it to the people who are willing to use it and have the required skills."*].

Easy access is important according to all end users to establish effective eRehabilitation interventions. [Patient_11: *"If getting into the program fails the first time you try, then you will be done with it soon."*]. Requirements for easy access were: *no internet connection is needed* (requirement 1) and *logging on is only required once* (requirement 2).

Furthermore, patients/informal caregivers want eRehabilitation to be *applicable to most possessed ICT devices* (requirement 3) and that *their health professionals have access to their electronic patient record to stay informed about training results* (requirement 4).

For professionals it was important that eRehabilitation is *applicable on computers at the rehabilitation center and synchronizes with programs used for the electronic patient record* (requirement 5). Moreover, they stated that *different eHealth interventions should be brought together in one central dashboard* (requirement 6). [Professional_7: "I would like to argue that people only have one account and that all facilities are directly available via one digital environment."].

Usability

Product attributes (visual)

Visual disabilities related to stroke were mentioned (e.g. neglect) and resulted in a list of requirements regarding visual attributes, i.e. *use of pictograms, symbols and graphics* (requirement 1) and *a non-flashing and tranquil interface* (requirement 2). [Patient_18: "I prefer a light and calm background. Lots of colors and flashing lights onscreen often cause me a headache."].

In addition, patients/informal caregivers mentioned *lay-out settings should be adjustable* (requirement 3), so that this can be adapted to the patients' own preferences. [Caregiver_4: "I have seen so many differences between patients with stroke. No one is the same. So, I can imagine type of letters, colors and so on need to be adjustable."].

Product attributes (auditory)

As a consequence of cognitive and speech disabilities (e.g. aphasia), end users mentioned eRehabilitation interventions would be more suitable for stroke patients when *being able to listen to written text* (e.g. instruction of exercises) (requirement 4). [Caregiver_12: "It would be very helpful for my father if he can listen to instructions instead of reading it himself."].

In addition, *sounds for alerts or as feedback* (requirement 5) was a requirement of patients/informal caregivers. They prefer alerts as a reminder (e.g. alarm for training) or for fun (direct feedback whilst training). However, some patients also mentioned irritation and fatigue as negative side effects of sounds.

Product attributes (simplicity)

End users required simple eRehabilitation interventions to increase usability: *limited options on a single screen to click further to another screen* (requirement 6) and *a limited amount of information on a single screen* (requirement 7).

Patients/informal caregivers also required *a limited amount of open webpages as a*

consequence of using a service (requirement 8) to prevent patients from getting lost. [Caregiver_9: *"If I look at my husband when he uses the computer, I think it should be very simple. So not too much text, pictures, things you can click on... otherwise he has no idea what he is doing"*].

Service (support)

Support on how to use eRehabilitation was considered crucial for usability according to all end users. Several requirements were mentioned: *menu with frequently asked questions* (requirement 9), *videos with instructions* (requirement 10), *helpdesk* (requirement 11) and *direct support on location* (requirement 12). [Caregiver_4: *"If you're at home and you'll get stuck, you want to be able to ask someone directly for help"*].

Content

Personalized training facilities

End users want eRehabilitation to include tailored training facilities for recovery after stroke, i.e. *physical* (requirement 1), *cognitive* (requirement 2) and *speech exercises* (requirement 3). [Patient_19: *"If I came to know one thing, it is that no person who have had a stroke is the same, so you should be able to compose it in a way that it applies to you."*]. Moreover, training facilities need to deliver constant personalized feedback (e.g. symbols, sounds, etc.) to prevent boredom with training and increase fun and accordingly stimulate training adherence and rehabilitation outcomes.

Tracking

All end users mentioned activity trackers as an eRehabilitation tool to *monitor daily activities* (requirement 4). [Caregiver_11: *"He forgot to count taking a shower, having breakfast, going up and down the stairs... Then this device measured all activities and he became aware that he actually did a lot. That explained why he was so tired."*].

Patient/informal caregivers also mentioned *a video system to record exercises* (requirement 5) and the ability to send these recordings to their health professional for feedback. A requirement of health professionals was *monitoring of a patients' health status* (e.g. *body weight, heart rate function, etc.*) (requirement 6).

Agenda and reminders

All end users preferred a digital agenda which includes: *a patients' rehabilitation schedule* (requirement 7), *a reminder function for scheduled appointments* (requirement 8) and *scheduled time to use eRehabilitation (digital training)* (requirement 9). This was found important in case of cognitive impairments and difficulties with time management after

stroke.

Furthermore, patients/informal caregivers want to be *able to make appointments with health professionals on their own initiative* (requirement 10). Professionals required a limit in the number of appointments. [Professional_2: *"I would like if patients can schedule an appointment with me, but only within restrictions. I certainly do not want them to schedule appointments with me every week."*].

Communication

End users required digital communication tools (e.g. chat room, video chat, etc.) in order to facilitate *contact with peers to share experiences on how to cope with having a (relative with) stroke* (requirement 11). [Professional_5: *"There must be a digital function that allows patients and caregivers who have come to know each other in the center, to stay in contact if they want to"*]. Moreover, communication tools can provide *communication between health professionals and patients and their informal caregivers from a distance (telecommunication)* (requirement 12).

Information

According to all end users provision of information should include: *general information about stroke* (requirement 13), *hyperlinks to reliable and relevant web pages* (requirement 14), *information about patient organizations* (requirement 15) and *information on how to cope with consequences of stroke (psycho-education)* (requirement 16).

In addition, patients/informal caregivers required to get *information on how to complete daily activities (strategy training)* (requirement 18). Professionals want patients/informal caregivers to have insight in *agreements and information discussed during a consult* (requirement 19) and *final reports of their rehabilitation* (requirement 20). [Professional_7: *"Patients easily forget what I have discussed with them during a consult. It would be great if they can have access to this information later. To be honest: To me it is quit strange that some information is still not digitally available."*].

Individual goal setting and evaluation

Requirements mentioned by all end users were: *setting individual goals for eRehabilitation* (requirement 21), *evaluation of these goals* (requirement 22) by getting *feedback about training results* (i.e. insight in what is trained, number of completed training sessions and training outcomes) (requirement 23) and receiving *feedback on goals* (e.g. receiving a digital medal when a goal is accomplished) (requirement 24).

In addition, requirements of health professionals were: *use of clinical assessments to set and evaluate goals* (requirement 25), *use of valid questionnaires to set and evaluate goals* (requirement 26) and *comparing training outcomes of a single patient with those of other*

patients (requirement 27). However, they find this irrelevant for patients [Professional_1: *"Comparison of scores is especially useful to me, but not for patients. I am thinking of the prognosis and comparison with the average."*].

Discussion

The aim of the study was to identify end-user requirements for the accessibility, usability and content of a comprehensive eRehabilitation program for stroke care. In total 45 user requirements were identified, which were grouped into eleven subcategories. Most requirements of end-users concerned the content of eRehabilitation (27 requirements), followed by usability (12 requirements) and then accessibility (6 requirements).

User requirements were quite similar between patients/informal caregivers and health professionals, but also showed differences in perspectives. For instance, professionals required that eHealth programs are able to run on the computer at their workplace, whereas patients and caregivers mainly want to their smartphone or tablet. This implies that eHealth interventions should be designed in such a way that both requirements are met. Other studies incorporating multiple perspectives [32,56] did not specifically mention differences between end users, impairing comparability of results.

Compared to previous literature, this study identified new requirements for stroke eRehabilitation interventions. For **Accessibility**, it was found that offline eRehabilitation interventions, brought together in one digital dashboard, need to be directly availability after logging on once. To our knowledge, this was the first qualitative study that found requirements concerning accessibility of stroke eRehabilitation programs.

Identified requirements for **Usability** found in the current study that can be added to the literature were: use of icons/symbols, non-flashing and tranquil interface, ability to listen to written text, methods for simplicity (e.g. limited amount of information on a single screen) and support (a helpdesk, video instructions, etc.). Similar to other studies, all patients had different requirements for lay-out [28,31]. Thus, design solutions should be tailored to a range of users or need to include lay-out options that users can choose from according to their preferences.

Identified requirements for **Content** of eRehabilitation that can be added to the literature were: digital agenda, tracking systems, communication tools, provision of information and goal setting and evaluation. This rather broad range of requirements can be explained by the fact that this study aimed to identify user requirements for a comprehensive eRehabilitation program, instead of a single intervention. Similarities with previous studies were found regarding training facilities, i.e. adaptation to patients' own preferences and capabilities [17,25,27,28,30-32] and provision of (rewarding) feedback [25,27,28].

A limitation of the study is participants with a certain interest in technology and eRehabilitation were probably more likely to respond, causing response bias and reduced generalizability. Therefore, we used purposive sampling based on the purpose of the use of ICT-devices to capture a broad range of perspectives. The group of patients that participated in the study (responders) did not significantly differ in age and gender from the group non-responders. Another limitation is that the chosen study methodology does not allow for comparisons between subgroups of focus group participants (e.g. different technological abilities), since requirements were studied on the level of the group (patients/informal caregivers and health professionals) instead of the individual participant. However, it would be interesting to know if there are differences in requirements of subgroups and this should be studied in the future. In addition, we could not aim for data saturation amongst health professionals. Data saturation was reached after six focus groups with patients/informal caregivers, but due to practical considerations this was not possible for health professionals. Differences in results between patients/informal caregivers and health professionals may have resulted from this imbalance.

In conclusion, user requirements for an eRehabilitation program for stroke care were identified addressing three categories: content, usability and accessibility. Requirements were to some extent different between stroke patients/ informal caregivers and health professionals. Therefore, involving perspectives of all end users in the design process of eHealth is needed to increase their effectiveness in rehabilitation care. The results in the current study can be used in future studies that apply a user-centered design approach to identify requirements for new eHealth interventions for stroke rehabilitation.

Declarations

Ethics approval and consent to participate: All participants gave written informed consent prior to participation. The study was approved by the Medical Ethical Review Board of the Leiden University Medical Center [P15.281].

Consent for publication: Not applicable.

Availability of data and material: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: Stichting Innovatie Alliantie supported this project financially (Grant 2014-046PRO).

Authors' contributions: BB and MW conducted the focus groups and analysed the data. JM, LBV and TV were a major contributor in writing the manuscript. AK, HA, LB contributed to the organisation of the focus groups and all authors read and approved the final manuscript.

Acknowledgements: We thank all the patients, caregivers and healthcare professionals

participating in this project. We are grateful to the consortium partners for their effort in sharing expertise. We also thank Preschana Kewalbansing, Sander Houdijk, Liesbeth Boyce and Hanneke Braakhuis for their contribution to this study.


Reference list

1. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, et al. Global and regional burden of stroke during 1990–2010: findings from the Global Burden of Disease Study 2010. *The Lancet*. 2014;383(9913):245–255.
2. Martin C, Dellatolas G, Viguier D, Willadino-Braga L, Deloche G. Subjective experience after stroke. *Applied Neuropsychology* 2002;9:148–158.
3. Wilkinson PR, Wolfe CDA, Warburton FG, Rudd AG, Howard RS, Roos-Russell RW, Beech RR. A long-term follow-up of stroke patients. *Stroke* 1997;28:507–512.
4. Clark B, Whittall J, Kwakkel G, Mehrholz J, Ewings S, & Burrridge J. Time spent in rehabilitation and effect on measures of activity after stroke. *Cochrane Database of Systematic Reviews* 2017;(3).
5. Winstein CJ, Stein J, Arena R, Bates B, Cherney LR, Cramer SC, et al. Guidelines for adult stroke rehabilitation and recovery. *Stroke* 2016;47(6):e98–e169.
6. Eysenbach, G. What is eHealth?. *Journal of medical Internet research* 2001;3(2):e20.
7. Bashshur RL, Shannon G, Krupinski EA, & Grigsby J. Sustaining and realizing the promise of telemedicine. *Telemedicine and eHealth* 2013;19(5):339–345.
8. Teasell RW, Foley NC, Bhogal SK, & Speechley, MR. An evidence-based review of stroke rehabilitation. *Topics in stroke Rehabilitation* 2015.
9. Laver KE, George S, Thomas S, Deutsch JE, & Crotty M. Virtual reality for stroke rehabilitation: an abridged version of a Cochrane review. *European journal of physical and rehabilitation medicine* 2015; 51(4):497–506.
10. Rubin MN, Wellik KE, Channer DD, Demaerschalk BM. Systematic review of telestroke for post-stroke care and rehabilitation. *Current atherosclerosis reports* 2013;15(8):1–7.
11. Saposnik G, Levin M & Stroke Outcome Research Canada (SORCan) Working Group. Virtual reality in stroke rehabilitation a meta-analysis and implications for health professionals. *Stroke* 2011;42(5):1380–1386.
12. De Mauro A. Virtual reality based rehabilitation and game technology. *EICS4 Med* 2011;1:48–52.
13. Webster D, Celik O. Systematic review of Kinect applications in elderly care and stroke rehabilitation. *Journal of neuroengineering and rehabilitation* 2014;11(1):108.
14. Schwamm LH, Chumbler N, Brown E, Fonarow GC, Berube, D, Nystrom K, et al. Recommendations for the Implementation of Telehealth in Cardiovascular and Stroke Care: A Policy Statement From the American Heart Association. *Circulation* 2016.
15. Anrich B, Mayora O, Bardram J, Tröster G. Pervasive healthcare: paving the way for a pervasive, user-centered and preventive healthcare model. *Methods Inf Med* 2010;49(1):67–73.
16. Holtzblatt K, Wendell JB, Wood S. Introduction in *Rapid Contextual Design*, San Francisco, CA: Morgan Kaufmann Publishers, 2005.
17. Sivan M, Gallagher J, Holt R, Weightman A, Levesley M, Bhakta B. Investigating the international classification of functioning, disability, and health (ICF) framework to capture user needs in the concept stage of rehabilitation technology development. *Assistive Technology* 2014;26(3):164–173.
18. van Gemert-Pijnen JE, Nijland N, van Limburg M, Ossebaard HC, Kelders SM, Eysenbach, G. et al. A holistic framework to improve the uptake and impact of eHealth technologies. *Journal of medical Internet research* 2011;13(4).
19. Eslami M, Firoozabadi M, Homayounvala E. User preferences for adaptive user interfaces in health information systems. *Universal Access Inf Soc*. 2018;17(4):875–83.
20. Schnall R, Rojas M, Bakken S, Brown W, Carballo-Dieguez A, Carry M, Travers J. A user-centered model for designing consumer mobile health (mHealth) applications (apps). *J Biomed Inf*. 2016;60:243–51.
21. Kushniruk A. Evaluation in the design of health information systems: application of approaches emerging from usability engineering. *Computers in biology and medicine* 2002;32(3):141–149.
22. Fouladinejad F, Roberts JR. A computer aided learning package for training users and maintainers of medical equipment. Paper presented at the Proceedings of the 18th Annual International Conference of the IEEE, “Bridging Disciplines for Biomedicine,” Amsterdam, the Netherlands, 1996.

23. Shah SG, Robinson I. User involvement in healthcare technology development and assessment: Structured literature review. *International Journal of Health Care Quality Assurance Incorporating Leadership in Health Services* 2006;19:500–515.
24. Shah SG, Robinson I, AlShawi S. Developing medical device technologies from users' perspectives: A theoretical framework for involving users in the development process. *International Journal of Technology Assessment in Health Care* 2009;25:514–521.
25. Nasr N, Leon B, Mountain G, Nijenhuis SM, Prange G, Sale P, et al. The experience of living with stroke and using technology: opportunities to engage and co-design with end users. *Disability and Rehabilitation: Assistive Technology* 2016;11(8):653–660.
26. Ehn M, Hansson P, Sjölander M, Boman IL, Folke M, Sommerfeld D, et al. Users perspectives on interactive distance technology enabling home-based motor training for stroke patients. In *pHealth* 2015;5:145–152.
27. Mawson S, Nasr N, Parker J, Zheng H, Davies R & Mountain G. Developing a personalised self-management system for post stroke rehabilitation; utilising a user-centred design methodology. *Disability and Rehabilitation: Assistive Technology* 2014;9(6):521–528.
28. Parker J, Mawson S, Mountain G, Nasr N, Davies R & Zheng H. The provision of feedback through computer-based technology to promote self-managed post-stroke rehabilitation in the home, *Disability and Rehabilitation: Assistive Technology* 2014.
29. Lu EC, Wang R, Huq R, Gardner D, Karam P, Zabjek K, et al. Development of a robotic device for upper limb stroke rehabilitation: A user-centered design approach. *Paladyn, Journal of Behavioral Robotics* 2011;2(4):176–184.
30. Lange B, Flynn S, Proffitt R, Chang CY, Rizzo A. Development of an interactive game-based rehabilitation tool for dynamic balance training. *Topics in stroke rehabilitation* 2010;17(5):345–352.
31. Zheng H, Davies R, Zhou H, Hammerton J, Mawson SJ, Ware PM, et al. SMART project: application of emerging information and communication technology to home-based rehabilitation for stroke patients. *International Journal on Disability and Human Development* 2006;5(3):271.
32. Mountain G, Ware P, Hammerton J, Mawson S, Zheng H, Davies R, et al. The SMART project: a user led approach to developing applications for domiciliary stroke rehabilitation. *Designing accessible technology* 2006;135–144.
33. Miñón R, Moreno L, Martínez P, & Abascal J. An approach to the integration of accessibility requirements into a user interface development method. *Science of Computer Programming* 2014;86:58–73.
34. Kim Y & Crowston K. Technology adoption and use theory review for studying scientists' continued use of cyber-infrastructure. *Proceedings of the American Society for Information Science and Technology* 2011;48(1),1–10.
35. International Organization for Standardization. ISO 9241-210 Ergonomics of human-system interaction - Part 210. Human-centred design for interactive systems. 2010.
36. Pagliari C. Design and evaluation in eHealth: challenges and implications for an interdisciplinary field. *J Med Internet Res* 2007;9(2):e15.
37. Hebert M. Telehealth success: evaluation framework development. *Stud Health Technol Inform.* 2001;84(Pt 2):1145–9.
38. Koch S. Improving quality of life through eHealth—the patient perspective. *Stud Health Technol Inform;* 2012 ;25–9.
39. Rinkus S, Walji M, Johnson-Throop KA, Malin JT, Turley, JP, Smith JW & Zhang J. Human-centered design of a distributed knowledge management system. *Journal of Biomedical Informatics*, 2005;38(1),4–17.
40. Davoody N, Koch S, Krakau I & Hägglund M. Post-discharge stroke patients' information needs as input to proposing patient-centred eHealth services. *BMC medical informatics and decision making*, 2016;16(1), 66.
41. Kitzinger J. Qualitative research. Introducing focus groups. *BMJ: British medical journal* 1995;311(7000):299.
42. Kitzinger J. Focus group research: Using group dynamics to explore perceptions, experiences and understandings. In: Holloway I, ed. *Qualitative research in health care*. 1st ed. Berkshire, England: Open University Press; 2005:56–70.
43. Francis JJ, Johnston M, Robertson C, Glidewell L, Entwistle V, Eccles MP. What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychol Health*. 2010;25:1229–45.

44. Venkatesh V, Morris MG, Davis GB, et al. User acceptance of information technology: toward a unified view. *MIS Q* 2003;27:425–478.
45. Kazanjian A, Green CJ. Beyond effectiveness: the evaluation of information systems using A Comprehensive Health Technology Assessment Framework. *Comput Biol Med* 2002;32(3):165-177.
46. Hamid A, Sarmad A. Evaluation of e-health services: user's perspective criteria. *Transforming government: people, process and policy* 2008;2(4):243-255.
47. Kaufman D, Roberts WD, Merrill J, Lai TY, Bakken S. Applying an evaluation framework for health information system design, development, and implementation. *Nurs Res* 2006;55(2 Suppl):S37-42.
48. Eysenbach G. A framework for evaluating e-health: systematic review of studies assessing the quality of health information and services for patients on the Internet. *J Med Internet Res* 2000;2(suppl2).
49. Eng TR, Gustafson DH, Henderson J, Jimison H, Patrick K. Introduction to evaluation of interactive health communication applications. Science Panel on Interactive Communication and Health. *Am J Prev Med* Jan 1999;16(1):10-15.
50. Esser P, Goossens R. A framework for the design of user-centered teleconsulting systems. *J Telemed Telecare* 2009;15(1):32-39.
51. Gulliksen J, Göransson B, Boivie I, Blomkvist S, Persson J, Cajander Å. Key principles for user-centred systems design. *Behaviour and Information Technology*. 2003;22(6):397-409.
52. Oinas-Kukkonen H, Harjuma M. Persuasive systems design: key issues, process model, and system features. *Commun Assoc Inf Syst* 2009;24(1):28.
53. Resnicow K, Strecher V, Couper M, Chua H, Little R, Nair V, et al. Methodologic and design issues in patient-centered e-health research. *Am J Prev Med* 2010 Jan;38(1):98-102.
54. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International journal for quality in health care* 2007;19(6):349-357.
55. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res* 2005 Nov;15(9):1277-1288.
56. Francis JJ, Johnston M, Robertson C, Glidewell L, Entwistle V, Eccles MP. What is an adequate sample size? 39 Operationalising data saturation for theory-based interview studies. *Psychol Health* 2010;25:1229-1245.



The background of the page is a light gray with a pattern of hexagons and medical symbols. Some hexagons are solid gray, while others are white with a gray outline. Medical symbols, including plus signs and a cross, are scattered throughout. A large, dark gray, stylized number '6' is positioned in the bottom right corner, partially overlapping the hexagonal pattern.

What is important in e-health interventions for stroke rehabilitation? A survey study among patients, informal caregivers and health professionals

M. Wentink
L. van Bodegom-Vos
B. Brouns
H. Arwert
T. Vliet Vlieland
A. de Kloet
J. Meesters

International Journal of Telerehabilitation 2018;10(1):15-28.

Abstract

Incorporating user requirements in the design of e-rehabilitation interventions facilitates their implementation. However, insight into requirements for e-rehabilitation after stroke is lacking. This study investigated which user requirements for stroke e-rehabilitation are important to stroke patients, informal caregivers, and health professionals. The methodology consisted of a survey study amongst stroke patients, informal caregivers, and health professionals (physicians, physical therapists and occupational therapists). The survey consisted of statements about requirements regarding accessibility, usability and content of a comprehensive stroke e-health intervention (4-point Likert scale, 1=unimportant/4=important). The mean with standard deviation was the metric used to determine the importance of requirements. Patients (N=125), informal caregivers (N=43), and health professionals (N=105) completed the survey. The mean score of user requirements regarding accessibility, usability and content for stroke e-rehabilitation was 3.1 for patients, 3.4 for informal caregivers and 3.4 for health professionals. Data showed that a large number of user requirements are important and should be incorporated into the design of stroke e-rehabilitation to facilitate their implementation.

Introduction

Stroke is a major health problem in Europe for which the incidence is expected to increase from 1.1 million per year in 2000 to 1.5 million per year in 2025 (Truelsen, et al., 2006). Patients suffering from stroke may experience multiple disabilities and require comprehensive rehabilitation. Overall, an increase is expected in the need for rehabilitation post stroke, not only because of the rising incidence, but also since, due to the improvement of the initial medical treatment, more patients now survive a stroke (Feigin et al., 2016). Comprehensive rehabilitation is delivered by various health professionals from different disciplines (e.g., physical therapists, occupational therapists, speech-language pathologists, psychologists, and social workers), with therapy aimed at individual treatment goals involving the patient and his or her informal caregiver (Winstein et al. 2016).

Due to developments in society and health care, including limited resources for the delivery of comprehensive rehabilitation, Information and Communication Technologies (ICT) play an important role in the delivery of rehabilitation care. 'The use of ICT, mostly internet technology, to improve or support health and health care, is known as e-health (Wentzel, Beerlage-de Jong, & Sieverink, 2014). E-rehabilitation refers to the application of e-health in rehabilitation care (e.g., serious brain games, virtual reality and telerehabilitation). Although many e-rehabilitation interventions have been tested regarding their effectiveness, the use of e-rehabilitation by end users remains low (Brewer, McDowell, & Worthen-Chaudharim, 2007; Lum, Reinkensmeyer, Mahoney, Rymer, & Burgar, 2002).

Implementation of e-health is influenced by its complexity, the adaptability of the technology to fit the local context, and its compatibility with existing systems, work practices, and costs (Ross, Stevenson, Lau, & Murray, 2016). End user input in the design and development of e-health technologies (i.e., user-centered design approach) is a way to overcome such barriers (Goldstein et al., 2014; Pagliari 2007; Ross et al., 2016; van Gemert-Pijnen et al., 2011).

Prior qualitative research (via interviews and focus groups) on end users' requirements for stroke e-rehabilitation (Ehn et al., 2015; Lange, Flynn, Proffitt, Chang, & Rizzo, 2010; Mawson et al., 2014; Mountain et al., 2006; Nasr et al., 2016; Parker et al., 2014; Zheng et al., 2006) found that interventions should be tailored (Lange et al., 2010; Nasr et al., 2016; Zheng et al., 2006); need to involve goal setting (Mawson et al., 2014; Sivan et al., 2014); must be easy to use; and should provide feedback about training performances (Mawson et al., 2014; Mountain et al., 2006; Nasr et al., 2016; Parker et al. 2014; Zheng et al., 2006).

However, quantitative studies regarding user requirements for e-rehabilitation after stroke are scarce. Thus far, one study used a quantitative survey among 233 health professionals in stroke care to rank the importance of the requirements that were identified in a previous qualitative study of Lu et al., (2011). However, this study was concerned with

only one aspect of stroke recovery (upper limb rehabilitation), and one technology tool (robot). Moreover, only health professionals, mainly occupational therapists and physical therapists, completed the survey whereas patients and their informal caregivers were not involved.

Thus, it remains unclear what requirements are most important for the comprehensive delivery of e-rehabilitation interventions (e.g., an app with upper limb exercises, brain games and/or telecommunication) including all potential end users, (i.e., patients, informal caregivers and health professionals). Therefore, this study aims to prioritize the requirements for stroke e-rehabilitation according to patients, informal caregivers, and health professionals. This is relevant for the application of user-centered design and accordingly the development and implementation of effective e-health interventions in stroke rehabilitation.

Patients and Materials

Design and setting

This cross-sectional study, involving a one-time, online survey, was conducted in June 2016 among (former) patients who had been admitted to Sophia Rehabilitation Centre (the Hague) and Rijnlands Rehabilitation Centre (Leiden) in The Netherlands, their informal caregivers, and healthcare professionals (rehabilitation physicians, psychologists, physical therapists and managers). The study was approved by the Medical Ethical Review Board of the Leiden University Medical Center [P15.281].

Study population

Patients and informal caregivers

Patients and informal caregivers were recruited by identifying potentially eligible patients in the electronic patient registries of the two rehabilitation centres, based on the following criteria: older than 18 years, diagnosed with stroke, rehabilitation started after June 2011 and rehabilitation was completed. Four hundred patients (200 in Leiden and 200 in The Hague) were randomly selected by assigning a number to every patient using a random number generator and subsequently selecting the first four-hundred patients and their informal caregivers.

Health professionals

Health professionals were selected if they were a practicing health professional (i.e., rehabilitation physicians, physical therapists, or psychologists) with at least two years of working experience in a multidisciplinary team for stroke patients. Health professionals

were randomly selected from the Dutch medical address book (which includes most professionals in The Netherlands), the Dutch Association of Rehabilitation Physicians (VRA: Nederlandse Vereniging van Revalidatieartsen) and the Royal Dutch Society of Physical therapy (KNGF: Koninklijk Nederlands Genootschap voor Fysiotherapie). If an email address was missing, other methods (e.g., internet, telephone calls) were used. We aimed to invite at least 300 health professionals.

Patients and health professionals received an email about the study including a digital link to the survey. Informal caregivers (e.g., partner, family member, etc.) were invited to fill in the questionnaire in the email directed to the patients. Thus, it remains unclear whether the patients had an informal caregiver and if so, whether they passed on the invitation. If the invited health professional stated that he or she was not involved in stroke care, they were asked to invite colleagues to fill out the survey. Non-responders received two reminders, each with an in-between period of 1.5 weeks.

Survey development

The content of the survey was based on a previous qualitative study, in which a framework for end user requirements for e-rehabilitation in stroke care was established (Figure 1). The framework comprises 45 identified requirements, classified into eleven self-determined categories and organized by three self-determined key themes: 'accessibility', 'usability', and 'content'. Accessibility refers to "easy access to e-rehabilitation for all end users, including patients with disabilities as a consequence of stroke." Usability is "the ease with which end users can use e-rehabilitation interventions for recovery after stroke during their stay in the rehabilitation center and/or at home."




Figure 1. Key themes end-user requirements for e-health interventions in stroke rehabilitation.

Content was defined as “everything end users want to include in e-rehabilitation (e.g., services, interventions, information, applications, etc.) to achieve specified goals for e-rehabilitation in their rehabilitation process.”

The user requirements identified for patients/ informal caregivers and health professionals were translated into neutral statements for the survey. Each survey consisted of two parts: (1) socio-demographic and disease characteristics, and (2) a list of user requirements for accessibility, usability, and content for patients/ informal caregivers and health professionals. The survey was pilot tested amongst two health professionals and three patients who were undergoing treatment in the rehabilitation center for recovery after stroke. The survey was tested for feasibility, readability and presentation (e.g. perceived question difficulties, response errors, screen layout, etc.). The pilot testing led to minor changes in the wording and format of the final survey.

Survey content

Socio-demographic (and disease) characteristics

The age and gender of patients, informal caregivers, and health professionals were recorded. In addition, patients were asked to provide the following information: education level (low [no or only primary education], intermediate [prevocational secondary education, senior secondary vocational training, senior secondary general education, preuniversity education], high [higher professional education or university (bachelor, master, or PhD degree)]); living status (living alone/ living together); employment (paid job/ no paid job); time after stroke (in months); and self-perceived impairments as a consequence of stroke (cognitive, physical, communication). Health professionals were asked about their discipline; region (north, middle, and/or south of the Netherlands); work setting (primary care, rehabilitation centre, general hospital); years of work experience; and estimated average number of new stroke patients per month. Moreover, they were asked whether they used e-health in routine stroke rehabilitation (yes, no).

User requirements

Forty-five requirements for the three themes ‘accessibility’ (8 requirements), ‘usability’ (12 requirements) and ‘content’ (25 requirements) of a comprehensive e-health intervention after stroke were identified in the qualitative study and were transformed into neutral statements for the survey. A total of 39/45 requirements were directly transformed and 6/45 requirements were divided into 2 or more statements, resulting in 15 additional statements for the survey (52 statements). The 52 statements were included in the survey for patients. There were 2/52 statements that were accidentally missing in the survey for caregivers, resulting in 50 statements in the survey for caregivers. In the survey of health professionals,

a number of 7/52 statements were asked from the perspective of a patient next to their own perspective, resulting in 7 additional statements. There were 11/52 statements derived from the qualitative study were only applicable for patients and caregivers, so eventually 48 statements (52+7-11) were included in the survey of health professionals.

All participants were asked to rate the importance of the given statements on a 4-point Likert scale (1=unimportant, 2=rather unimportant, 3=rather important, 4=important). These scores were used to calculate the mean in order to make a ranking from highest to least important requirements.

Analysis

Respondents were included in the analyses if they completed ≥ 90 percent of survey. Socio-demographic and disease characteristics were analyzed using descriptive statistics and presented as numbers with percentages, means with standard deviations (SD), or medians with ranges (Inter Quartile Range; IQR), i.e., 25th percentile–75th percentile), where appropriate.

To quantify the importance of requirements for accessibility, usability and content of e-rehabilitation interventions as perceived by respondents, descriptive analysis was used. The mean with the standard deviation (SD) for each statement were reported to discriminate between and prioritize the statements used in the survey items. Means provide the most accurate insight in the importance of the requirements. Scores on statements per subgroup (patients, informal caregivers and health professionals) are presented in separate tables for each theme: Accessibility, Usability and Content. In addition, the mean score of all statements were provided per subgroup. All statistical analyses were performed using Statistical Packages for the Social Sciences (IBM SPSS 22.0 for Windows).

Ethical issues and approval

Participants filled in the survey anonymously implying that patient's, informal caregiver's and health professional's characteristics were not traceable, (e.g., age instead of date of birth). Immediately after filling in the survey, participants were thanked for their willingness to participate. Participants did not receive results of the study, since they filled in the survey anonymously.

Results

Response

Of the 400 invited patients, 32 had no valid email address; the survey was completed by 125 out of 368 invited patients (34%). Additionally, 43 informal caregivers, and 105 health professionals completed the survey (Figure 2). Reasons for nonresponse were not verified.




Figure 2. Flow of inclusion.

Socio-demographic (and disease) characteristics

The characteristics of the 273 responders are shown in Table 1. Respondents included 72/125 (58%) patients, 16/43 (37%) informal caregivers and 25/105 (24%) health professionals. The mean age of the patients was 58 years (SD 11.4), of the informal caregivers 58 years (SD 12.0) and of the health professionals 42 years (SD 10.5). In total, 41/105 (39%) of the health professionals were physical therapists, 15/105 (14%) were psychologists, 47/105 (45%) were physicians and 2/105 (2%) did not mention their discipline. Seventy-five out of 105 (71%) responding professionals worked in a rehabilitation center.

Table 1. Characteristics of participating patients, informal caregivers and health professionals.

Characteristics	Patients (n=125)	Caregivers (n=43)	Professionals (n=105)
Age, years (mean, SD)	58 (11.4)	58 (12.0)	42 (10.5)
Sex, no. male (%)	72 (58)	16 (37)	25 (24)
Education, no. (%)			
Low	21 (17)	-	-
Intermediate	46 (37)	-	-
High	57 (46)	-	-
Living status, no. living alone (%)	22 (18)	5 (12)	-
Employment, no. with a paid job (%)	42 (34)	21 (49)	-
Work region, no. (%)			
North	-	-	20 (20)
Middle	-	-	63 (60)
South	-	-	21 (20)
Health professional discipline, no. (%)			
Physical therapist	-	-	41 (39)
Psychologist	-	-	15 (14)
Physician	-	-	47 (45)
Unknown ^a	-	-	2 (2)
Work setting, no. (%)			
Health centre in primary care	-	-	10 (10)
Rehabilitation centre	-	-	75 (71)
General hospital	-	-	34 (32)
Work experience, no. years (%)			
>0-5	-	-	25 (23.8)
>6-10	-	-	28 (26.7)
>11-15	-	-	14 (13.3)
>15	-	-	37 (35.2)
Estimated average number of new stroke patients per month; no. (%)			
>0-5	-	-	47 (46)
>6-10	-	-	33 (32)
>11-15	-	-	11 (11)
>15	-	-	11 (11)
Time after stroke, months (mean, SD)	30.6 (29.2)		

Table 1. Continued.

Characteristics	Patients (n=125)	Caregivers (n=43)	Professionals (n=105)
Self-perceived impairments, no. yes (%)			
Cognitive impairments	81 (65)		-
Physical impairments	84 (67)		-
Aphasia	48 (38)		-
Use of any device in daily life, no. yes (%)	113 (90)	41 (95)	
Use of device, no. yes (%)			
Smartphone	85 (68)	33 (77)	-
Tablet	62 (50)	30 (70)	-
Laptop	71 (57)	30 (70)	-
Computer (PC)	54 (43)	20 (47)	
Use of digital rehabilitation tools, no. yes (%)	-	-	40 (38)

^a Health professionals who did not mention their discipline.

Requirements

The mean score of all user requirements regarding accessibility, usability and content for stroke e-rehabilitation was 3.1 for patients, 3.4 for informal caregivers and 3.4 for health professionals. For patients, the mean score (SD) for the least important requirement was 2.4 (1.1) and for the most important 3.6 (0.8). For caregivers, the mean score (SD) for the least important requirement was 2.8 (1.1) and for the most important 3.8 (0.4). For health professionals, the mean score (SD) for the least important requirement was 2.4 (1.0) and for the most important 3.9 (0.4).

Accessibility

Two requirements for accessibility to e-rehabilitation after stroke were found to be the most important according to all end users: *e-rehabilitation is applicable to most commonly possessed ICT-devices, e.g., laptop, tablet and smartphone* (patients: mean 3.5, SD 0.9; informal caregivers: mean 3.5, SD 0.7; professionals: mean 3.6, SD 0.6) and *access for health professionals to the electronic patient record to stay informed about training results* (patients: mean 3.3, SD 1.0; informal caregivers: mean 3.5, SD 0.9; professionals: mean 3.5, SD 0.7) (see table 2a).

Usability

Categories for usability were: visual appeal, auditory appeal, simplicity and support. Two requirements regarding the category 'support' were found to be most important according to all end users: *videos with instructions on how to use e-rehabilitation* (patients: mean 3.3, SD 1.0; informal caregivers: mean 3.7, SD 0.9; professionals: mean 3.7, SD 0.6) and *a menu with frequently asked questions for patients* (patients: mean 3.1, SD 1.0; informal caregivers: mean 3.7, SD 0.9; professionals: mean 3.7, SD 0.6) (see table 2b).

Moreover, three requirements showed a mean score higher than the mean score on all statements for both patients and informal caregivers: *limited options on a single screen to click further to another screen* within the category simplicity (patients: mean 3.1, SD 1.1; informal caregivers: mean 3.4, SD 1.0), *non-flashing and tranquil interface* (patients: mean 3.3, SD 0.8; informal caregivers: mean 3.8, SD 0.4) and *adjustable font style and font size settings* (patients: mean 3.0, SD 1.1; informal caregivers: mean 3.6, SD 0.7) within the category visual appeal.

Content

Categories for content were: training facilities, tracking, agenda/ reminders, communication, information and goal setting/ evaluation. A relatively large number of requirements for content showed higher mean scores than the mean score on all statements by all end users, e.g., *insight in agreements made during a consult* in the category information (patients: mean 3.5, SD 0.9; informal caregivers: mean 3.6, SD 0.8; professionals: mean 3.7, SD 0.6), *insight in final reports of a patients' rehabilitation process* in the category information (patients: mean 3.6, SD 0.7; informal caregivers: mean 3.7, SD 0.8; professionals: mean 3.4, SD 0.8) and *physical exercises* in the category training facilities (patients: mean 3.4, SD 1.0; informal caregivers: mean 3.7, SD 0.8, professionals: mean 3.6, SD 0.6).

Table 2a. Importance of requirements for **accessibility** of stroke e-rehabilitation according to end users (n=273).

Category	Requirement	End user		
		Patients (n=125)	Caregivers (n=43)	Professionals (n=105)
		Mean (SD)	Mean (SD)	Mean (SD)
The mean score of all statements per subgroup		3.3	3.5	3.4
Access	Applicable to most commonly possessed ICT-devices.	3.5 (0.9)	3.5 (0.7)	3.6 (0.6)
Access	No internet connection is required to use e-health interventions (offline use).	3.2 (1.0)	3.5 (1.0)	3.1 (0.9)
Access	Different e-health interventions are accessible without logging into the system each time.	3.1 (1.0)	3.3 (0.9)	3.5 (0.8)
Access	Access for health professionals to the electronic patient record to stay informed about training results.	3.3 (1.0)	3.5 (0.9)	3.5 (0.7)

Table 2b. Importance of requirements for **usability** of stroke e-rehabilitation according to end users (n=273).

Category	Requirement	End user		
		Patients (n=125) Mean (SD)	Caregivers (n=43) Mean (SD)	Professionals (n=105) Mean (SD)
The mean score of all statements per subgroup		2.9	3.4	3.5
Visual appeal	Adjustable settings: background.	2.6 (1.1)	3.0 (0.1)	-
Visual appeal	Adjustable settings: colors.	2.5 (1.1)	3.3 (0.1)	-
Visual appeal	Adjustable settings: page lay-out.	2.7 (1.1)	3.3 (0.9)	-
Visual appeal	Adjustable settings: font style and font size.	3.0 (1.1)	3.6 (0.7)	-
Visual appeal	Use of pictograms, symbols and graphics.	2.7 (1.1)	3.3 (1.0)	-
Visual appeal	Non-flashing and tranquil interface.	3.3 (0.8)	3.8 (0.4)	-
Auditory appeal	Ability to listen to written text.	2.7 (1.1)	3.4 (1.0)	-
Auditory appeal	Sounds for alert or as feedback.	2.7 (1.1)	3.3 (0.9)	-
Simplicity	Limited amount of open webpages as a consequence of using a service.	2.8 (1.0)	3.6 (0.9)	3.5 (0.7)
Simplicity	Limited amount of information on a single screen.	3.3 (1.0)	Missing ^a	-
Simplicity	Limited options on a single screen to click further to another screen.	3.1 (1.1)	3.4 (1.0)	-
Support	Direct assistance at home.	3.3 (1.08)	3.1 (1.1)	-
Support	Video for patients with instructions on how to use e-rehabilitation.	3.3 (1.0)	3.7 (0.9)	3.7 (0.6)
Support	Video for professionals with instructions on how to use e-rehabilitation.	-	-	3.1 (0.8)
Support	Menu with frequently asked questions for patients (FAQ).	3.1 (1.0)	3.7 (0.9)	3.7 (0.6)
Support	Menu with frequently asked questions for professionals (FAQ).	-	-	3.3 (0.7)
Support	A helpdesk for patients.	2.9 (1.1)	3.5 (0.9)	3.9 (0.4)
Support	A helpdesk for professionals.	-	-	3.5 (0.6)

^a This requirement was accidentally missing in the survey for caregivers.

Table 2c. Importance of requirements for **content** of stroke e-rehabilitation according to end users (n=273).

Category	Requirement	End user		
		Patients (n=125) Mean (SD)	Caregivers (n=43) Mean (SD)	Professionals (n=105) Mean (SD)
The mean score of all statements per subgroup		3.0	3.2	3.2
Training facilities	Exercises for cognitive functioning.	3.6 (0.9)	3.8 (0.4)	3.2 (0.9)
Training facilities	Physical exercises.	3.4 (1.0)	3.7 (0.8)	3.6 (0.6)
Training facilities	Speech exercises.	2.9 (1.3)	3.5 (1.0)	3.3 (1.0)
Tracking	Monitor activities in daily living:			
	Insight in completed activities	2.5 (1.1)	3.1 (1.1)	3.2 (0.8)
	Duration of completed activities	3.1 (1.0)	3.6 (0.7)	3.3 (0.7)
Tracking	A video system to record exercises at home.	2.4 (1.1)	2.8 (1.1)	3.1 (0.9)
Tracking	Monitor a patients' health status:			
	Body weight	2.9 (1.1)	3.2 (1.0)	2.5 (0.9)
	Heart rate	2.9 (1.1)	Missing ^a	2.5 (0.8)
Agenda/ reminders	Insight in the rehabilitation schedule of a patient.	3.2 (1.1)	3.4 (1.0)	3.6 (0.7)
Agenda/ reminders	A reminder function for scheduled appointments.	2.9 (1.0)	3.4 (1.0)	3.7 (0.6)
Agenda/ reminders	Scheduled time to use e-rehabilitation (digital training).	3.2 (1.1)	3.3 (1.0)	3.3 (0.8)
Agenda/ reminders	Appointments with healthcare professionals:			
	Make a request for an appointment	3.1 (1.0)	3.3 (1.0)	2.9 (0.9)
	Schedule an appointment themselves	3.0 (1.0)	3.2 (1.0)	2.5 (1.0)
Communication	Contact for care givers to share experiences on how to cope with having a relative with stroke.	2.6 (1.1)	2.7 (1.0)	3.6 (0.6)
Communication	Contact for patients to share experiences on how to cope with having a stroke.	2.8 (1.0)	3.0 (1.0)	3.5 (0.6)
Communication	Communication between patients/ caregivers and professionals from a distance (telecommunication).	2.5 (1.1)	3.0 (1.0)	2.9 (0.9)
Information	General information about stroke.	3.4 (0.8)	3.4 (0.9)	3.7 (0.5)
Information	Hyperlinks to reliable and relevant webpages for patients and caregivers.	3.2 (0.9)	3.4 (0.5)	3.6 (0.5)
Information	Information about patient organizations.	3.3 (1.0)	3.0 (1.0)	3.7 (0.6)
Information	Information on how to cope with consequences of stroke (psycho-education).	2.8 (1.0)	3.7 (0.8)	3.6 (0.6)
Information	Descriptions on how to perform daily activities (strategy training).	2.4 (1.2)	3.1 (0.9)	3.3 (0.7)
Information	Insight after a consult with a health professional in:			
	Agreements that were made	3.5 (0.9)	3.6 (0.8)	3.7 (0.6)
	Information that was discussed	3.5 (0.8)	3.7 (0.7)	3.4 (0.8)
Information	Insight in final reports of a patients' rehabilitation process.	3.6 (0.7)	3.7 (0.8)	3.4 (0.8)

Table 2c. Continued.

Category	Requirement	End user		
		Patients (n=125) Mean (SD)	Caregivers (n=43) Mean (SD)	Professionals (n=105) Mean (SD)
Goal setting/ evaluation	Setting goals for e-rehabilitation (shared decision making).	3.4 (0.9)	3.7 (0.8)	3.4 (0.7)
Goal setting/ evaluation	Evaluation of goals for e-rehabilitation.	3.4 (0.9)	3.7 (0.7)	3.4 (0.7)
Goal setting/ evaluation	Feedback about training results for patients:			
	Insight in what is trained	3.2 (1.0)	3.6 (0.6)	2.3 (1.1)
	The number of completed sessions	3.1 (0.9)	3.6 (0.6)	3.5 (0.7)
	Training outcomes	3.2 (1.0)	3.7 (0.5)	3.5 (0.7)
Goal setting/ evaluation	Feedback about training results for professionals:	-	-	3.2 (0.8)
	Insight in what is trained	-	-	3.2 (0.8)
	The number of completed sessions	-	-	3.3 (0.8)
	Training outcomes			
Goal setting/ evaluation	Feedback on when a goal for e-rehabilitation is accomplished.	3.3 (0.9)	3.6 (0.9)	3.7 (0.6)
Goal setting/ evaluation	Use of clinical assessments for goal setting and evaluation.	3.5 (0.9)	3.6 (0.7)	3.3 (0.8)
Goal setting/ evaluation	Use of valid questionnaires for goal setting and evaluation.	3.3 (0.9)	3.5 (0.8)	3.4 (0.8)
Goal setting/ evaluation	Compare training outcomes of a single patient with those of other patients.	2.4 (1.2)	2.9 (1.0)	2.4 (1.0)

^a This requirement was accidentally missing in the survey for caregivers.

Discussion

The aim of this study was to make an inventory and prioritize the requirements for stroke e-rehabilitation according to patients, informal caregivers, and health professionals. Relatively large mean scores for user requirements regarding accessibility, usability and content for a comprehensive e-health intervention after stroke were found for each subgroup (patients 3.1, informal caregivers 3.4 and health professionals 3.4). Moreover, similarities and differences were found between the perspectives of patients, informal caregivers, and health professionals about the importance of the requirements.

To our knowledge one previous study used a quantitative survey in stroke care to discover the importance of the requirements that were identified in a previous qualitative study of Lu et al. (2011). Similar to the findings from the perspective of health professionals in this study, provision of feedback for patient and therapist and the tool being useful for stroke patients were found to be important requirements. However, comparison of the findings between both studies is hampered. Lu et al. (2011) focused on the user' requirements regarding

a robot for upper limb rehabilitation, while our study was concerning a comprehensive e-health intervention using multiple tools. Moreover, their survey was conducted among 233 health professionals while our study also included other user groups (i.e., patients and their informal caregivers).

Overall, requirements prioritized in this study were both similar and different compared to previous qualitative studies that identified user requirements for an e-health intervention. An important requirement regarding accessibility found in this study was the ability to use e-rehabilitation on multiple digital devices (smartphone, tablet, laptop). This corresponds to requests identified in previous literature that e-health be integrated in familiar and existing tools/applications, (not replacing them) (Matthew-Maich et al., 2016); is available alongside the work of health professionals (Mountain et al., 2006); is easy to set-up (Sivan et al., 2014; Zheng et al., 2006); and is suitable to the constant modification of the environment (Ross et al., 2016).

An important requirement found in this study regarding usability was a non-flashing and tranquil interface. This is in line with a previous study of Parker et al., (2014) that found participants preferred a simpler looking screen without additional background pictures. In contradiction to previous studies in which design settings needed to be changeable for adjustment to a patient's needs (Parker et al., 2014; Zheng et al., 2006), this study found changeable lay-out, background and color settings were less important. It can be added to the literature that incorporation of support for use of e-rehabilitation programmes (e.g., helpdesk, FAQ menu, videos with instructions) are highly important. These requirements should be integrated in e-rehabilitation designs to increase acceptance of e-rehabilitation for stroke patients, who often suffer from disabilities impairing usage.

Regarding content, the following important requirements were similar to previous literature: general information (McKevitt, Redfern, Mold, & Wolfe, 2004; Peoples, Satink, & Steultjens, 2011; Reed, Wood, Harrington, & Paterson, 2012; Salter, Hellings, Foley, & Teasell, 2008); goal setting and evaluation (Lu et al., 2011; McKevitt et al., 2004; Parke et al., 2015); and providing feedback (Hochstenbach-Waelen & Seelen, 2012; Lu et al., 2011; Mawson et al., 2015; Mountain et al., 2006; Nasr et al., 2016; Parker et al., 2014; Zheng et al., 2006). In contradiction, telecommunications in stroke care are rapidly developing worldwide because of their importance (Blacquièr et al., 2017), although this was found a less important requirement in the current study according to all end users. A broad range of requirements regarding content of comprehensive e-health interventions can be added to the literature (e.g., exercises for cognitive and physical functioning, hyperlinks to webpages, a reminder function, etc.), since this study prioritized user requirements for a comprehensive e-health intervention instead of a single e-health intervention addressing one aspect of stroke recovery with one technology tool.

Furthermore, similarities were found in perspectives of patients, informal caregivers,

and health professionals about the importance of requirements. The requirements of physical exercises, insight in information discussed during a consult, insight in final reports of the rehabilitation process and setting and evaluation of goals for e-rehabilitation were considered relatively important by all end users. However, notable differences were also found between the subgroups. The required exercises for cognitive functioning were important for patients and informal caregivers, whilst this was a less important requirement for health professionals. In addition, health professionals found contact with peers for caregivers and patients important, although patients and informal caregivers found this less important. Moreover, psycho-education was found to be a relatively important requirement by health professionals and informal caregivers, whereas patients seemed to find this less important. Therefore, differences in the importance of user requirements should be identified so that e-health interventions can be designed in such a way that requirements of different users are met.

As to our knowledge, this is the first study that prioritized a set of requirements for e-rehabilitation amongst multiple subgroups (patients, informal caregivers, and health professionals) and in which informal caregivers were treated as a separate group of end users. Differences in the importance of requirements for comprehensive e-health interventions for recovery after stroke between patients, informal caregivers, and professionals were not previously identified in the literature.

A limitation of the study is that selection bias might have occurred since the survey was distributed by mail, and only patients and their informal caregivers with an email address were able to fill in the survey. As a consequence, the perspective of patients and their informal caregivers with least experience with digital devices might be missing. However, we aimed to identify user requirements for e-rehabilitation, so knowledge and understanding of ICT and e-health were desirable. Moreover, informal caregivers of patients were invited to fill in the questionnaire in the invitation mail directed to the patients. If the invited health professional stated that he or she was not involved in stroke care, they were asked to invite colleagues to fill out the survey. Therefore, the actual amount of invited informal caregivers and health professionals and the response rates are unknown.

In summary, we prioritized requirements for accessibility, usability, and content of comprehensive e-health interventions from the perspective of patients, informal caregivers, and health professionals. It was found that a relatively large amount of user requirements were found important by each separate group and by all subgroups. These results can be used by developers, researchers and health professionals to apply user-centered design to develop effective e-health interventions and accordingly to enable their acceptance and adoption in stroke rehabilitation. However, more research is needed to identify which requirements are most important to optimize implementation, usage and adaptation of e-health in stroke rehabilitation.

Acknowledgements


We gratefully thank all the patients, their informal caregivers and healthcare professionals who participated in this study.

Reference list

1. Blacquiére, D., Lindsay, M. P., Foley, N., Taralson, C., Alcock, S., Balg, C., Ghanem, A. (2017). Canadian stroke best practice recommendations: Telestroke best practice guidelines update 2017. *International Journal of Stroke*, 12, 886-895.
2. Brewer, B. R., McDowell, S. K., & Worthen-Chaudhari, L. C. (2007). Poststroke upper extremity rehabilitation: A review of robotic systems and clinical results. *Topics in stroke rehabilitation*, 14, 22-44. <http://www.tandfonline.com/doi/abs/10.1310/tsr1406-22>
3. Casaubon, L. K., Boulanger, J. M., Blacquiére, D., Boucher, S., Brown, K., Goddard, T., ... Lavoie, P. (2015). Canadian stroke best practice recommendations: Hyperacute stroke care guidelines, update 2015. *International Journal of Stroke*, 10, 924-940.
4. Ehn, M., Hansson, P., Sjölander, M., Boman, I. L., Folke, M., Sommerfeld, D., ... Palmcrantz, S. (2015). Users perspectives on interactive distance technology enabling home-based motor training for stroke patients. *pHealth*, 2015(5), 145-152.
5. Feigin, V. L., Roth, G. A., Naghavi, M., Parmar, P., Krishnamurthi, R., Chugh, S., ... Estep, K. (2016). Global burden of stroke and risk factors in 188 countries, during 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *The Lancet Neurology*, 15, 913-924. [http://www.thelancet.com/journals/lanneur/article/PIIS1474-4422\(16\)30073-4/fulltext](http://www.thelancet.com/journals/lanneur/article/PIIS1474-4422(16)30073-4/fulltext)
6. Goldstein, D. H., Phelan, R., Wilson, R., Ross-White, A., VanDenKerkhof, E. G., Penning, J. P., & Jaeger, M. (2014). Brief review: Adoption of electronic medical records to enhance acute pain management. *Canadian Journal of Anesthesia/Journal Canadien d'Anesthésie*, 61, 164-179. <https://www.ncbi.nlm.nih.gov/pubmed/24233770>
7. Hochstenbach-Waelen, A., & Seelen, H. A. (2012). Embracing change: Practical and theoretical considerations for successful implementation of technology assisting upper limb training in stroke. *Journal of Neuroengineering and Rehabilitation*, 9(1), 52. doi: <https://doi.org/10.1186/1743-0003-9-52>
8. Lange, B., Flynn, S., Proffitt, R., Chang, C. Y., & Rizzo, A. (2010). Development of an interactive game-based rehabilitation tool for dynamic balance training. *Topics in Stroke Rehabilitation*, 17, 345-352. <https://www.ncbi.nlm.nih.gov/pubmed/21131259>
9. Lu, E.C., Wang, R.H., Hebert, D., Boger, J., Galea, M.P., & Mihailidis, A. (2011). The development of an upper limb stroke rehabilitation robot: identification of clinical practices and design requirements through a survey of therapists. *Disability and Rehabilitation: Assistive Technology*, 6, 420-431. <https://www.ncbi.nlm.nih.gov/pubmed/21184626>
10. Lum, P., Reinkensmeyer, D., Mahoney, R., Rymer, W.Z., & Burgar, C. (2002). Robotic devices for movement therapy after stroke: Current status and challenges to clinical acceptance. *Topics in stroke rehabilitation*, 8, 40-53. <https://www.ncbi.nlm.nih.gov/pubmed/14523729>
11. Matthew-Maich, N., Harris, L., Ploeg, J., Markle-Reid, M., Valaitis, R., Ibrahim, S., ... Isaacs, S. (2016). Designing, implementing, and evaluating mobile health technologies for managing chronic conditions in older adults: A scoping review. *Journal of Medical Internet Research, mHealth and uHealth*, 4(2). <https://www.ncbi.nlm.nih.gov/pubmed/27282195>
12. Mawson, S., Nasr, N., Parker, J., Zheng, H., Davies, R., & Mountain, G. (2014). Developing a personalised self-management system for post stroke rehabilitation; utilising a user-centred design methodology. *Disability and Rehabilitation: Assistive Technology*, 9, 521-528. <https://www.ncbi.nlm.nih.gov/pubmed/24131371>
13. McKeivitt, C., Redfern, J., Mold, F., & Wolfe, C. (2004). Qualitative studies of stroke: A systematic review. *Stroke*, 35, 1499-1505. <https://www.ncbi.nlm.nih.gov/pubmed/15105517>
14. Mountain, G. W. P. M., Ware, P., Hammerton, J., Mawson, S., Zheng, H., Davies, R., ... Eccleston, C. (2006). The SMART project: A user led approach to developing applications for domiciliary stroke rehabilitation. *Designing Accessible Technology*, 135-144. <http://opus.bath.ac.uk/6402/>
15. Murray, J., Ashworth, R., Forster, A., & Young, J. (2003). Developing a primary care-based stroke service: A review of the qualitative literature. *British Journal of General Practice*, 53, 137-142.
16. Nasr, N., Leon, B., Mountain, G., Nijenhuis, S. M., Prange, G., Sale, P., & Amirabdollahian, F. (2016). The experience of living with stroke and using technology: Opportunities to engage and co-design with end users. *Disability and Rehabilitation: Assistive Technology*, 11, 653-660. <https://www.ncbi.nlm.nih.gov/pubmed/25879304>
17. Pagliari, C. (2007). Design and evaluation in eHealth: Challenges and implications for an interdisciplinary field. *Journal of Medical Internet Research*, 9(2). <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1913937/>

18. Parke, H. L., Epiphaniou, E., Pearce, G., Taylor, S. J., Sheikh, A., Griffiths, C. J., ... Pinnock, H. (2015). Self-management support interventions for stroke survivors: a systematic meta-review. *PLoS One*, *10*(7), e0131448. <https://www.ncbi.nlm.nih.gov/pubmed/26204266>
19. Parker, J., Mawson, S., Mountain, G., Nasr, N., Davies, R., & Zheng, H. (2014). The provision of feedback through computer-based technology to promote self-managed post-stroke rehabilitation in the home. *Disability and Rehabilitation: Assistive Technology*, *9*, 529-538. <https://www.ncbi.nlm.nih.gov/pubmed/24131369>
20. Peoples, H., Satink, T., & Steultjens, E. (2011). Stroke survivors' experiences of rehabilitation: A systematic review of qualitative studies. *Scandinavian Journal of Occupational Therapy*, *18*, 163-171. URL: <https://www.ncbi.nlm.nih.gov/pubmed/20701431>
21. Reed, M. C., Wood, V., Harrington, R., & Paterson, J. (2012). Developing stroke rehabilitation and community services: A meta-synthesis of qualitative literature. *Disability and Rehabilitation*, *34*(7), 553-563. <https://www.ncbi.nlm.nih.gov/pubmed/21985186>
22. Ross, J., Stevenson, F., Lau, R., & Murray, E. (2016). Factors that influence the implementation of e-health: A systematic review of systematic reviews (an update). *Implementation Science*, *11*(1), 146. <https://www.ncbi.nlm.nih.gov/pubmed/27782832>
23. Salter, K., Hellings, C., Foley, N., & Teasell, R. (2008). The experience of living with stroke: A qualitative meta-synthesis. *Journal of Rehabilitation Medicine*, *40*, 595-602. <https://www.ncbi.nlm.nih.gov/pubmed/19020691>
24. Sivan, M., Gallagher, J., Holt, R., Weightman, A., Levesley, M., & Bhakta, B. (2014). Investigating the international classification of functioning, disability, and health (ICF) framework to capture user needs in the concept stage of rehabilitation technology development. *Assistive Technology*, *26*, 164-173. <https://www.ncbi.nlm.nih.gov/pubmed/26131797>
25. Truelsen, T., Piechowski-Jóźwiak, B., Bonita, R., Mathers, C., Bogousslavsky, J., & Boysen, G. (2006). Stroke incidence and prevalence in Europe: A review of available data. *European journal of Neurology*, *13*, 581-598. <https://www.ncbi.nlm.nih.gov/pubmed/16796582>
26. van Gemert-Pijnen, J. E., Nijland, N., van Limburg, M., Ossebaard, H. C., Kelders, S.M., Eysenbach, G., & Seydel, E. R. (2011). A holistic framework to improve the uptake and impact of eHealth technologies. *Journal of Medical Internet Research*, *13*(4). <https://www.ncbi.nlm.nih.gov/pubmed/22155738>
27. Wentzel, M.J., Beerlage-de Jong, N., Sieverin, K.F., & van Gemert-Pijnen, J.E.W.C. (2014). *Slimmer e-health ontwikkelen en implementeren met de CeHRes Roadmap*. In: C Politiek & R Hoogendijk (Eds.), CO-CREATIE E-HEALTHBOEK e-health, technisch kunstje of pure veranderekunde? https://www.utwente.nl/en/bms/ehealth/documents/workshop-2014/eHealth-Research_in_Beeld.pdf
28. Winstein, C.J., Stein, J., Arena, R., Bates, B., Cherney, L. R., Cramer, S.C., ... Lang, C.E. (2016). Guidelines for adult stroke rehabilitation and recovery. *Stroke*, *47*(6), e98-e169.
29. Zheng, H., Davies, R., Zhou, H., Hammerton, J., Mawson, S. J., Ware, P., ... Mountain, G. A. (2006). SMART project: Application of emerging information and communication technology to home-based rehabilitation for stroke patients. *International Journal on Disability and Human Development*, *5*, 271-276. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.537.4063&rep=rep1&type=pdf>





Teachers' and students' perceptions about eHealth education in the curriculum of exercise and physical therapy: a focus groups study on barriers and facilitators.

M. Wentink
P. Siemonsma
L. van Bodegom-Vos
A. de Kloet
J. Verhoef
T. Vliet Vlieland
J. Meesters

Accepted: *BMC Medical Education.*



Abstract

Background: Despite the growing importance of eHealth it is not consistently embedded in the curricula of functional exercise and physical therapy education. Insight in barriers and facilitators for embedding eHealth in education is required for the development of tailored strategies to implement eHealth in curricula. This study aims to identify barriers/facilitators perceived by teachers and students of functional exercise/physical therapy for uptake of eHealth in education.

Method: A qualitative study including six focus groups (two with teachers/four with students) was conducted to identify barriers/facilitators. Focus groups were audiotaped and transcribed in full. Reported barriers and facilitators were identified, grouped and classified using a generally accepted framework for implementation comprising levels of: innovation, individual teacher/student, social context, organizational context and political and economic factors.

Results: Teachers (n=11) and students (n=24) of functional exercise/physical therapy faculties of two universities of applied sciences in the Netherlands participated in the focus groups. A total of 109 barriers/facilitators were identified during the focus groups. Most found on the level of the innovation (n=26), followed by the individual teacher (n=22) and the organization (n=20). Teachers and students identified similar barriers/facilitators for uptake of eHealth in curricula: e.g. unclear concept of eHealth, lack of quality and evidence for eHealth, (lack of) capabilities of students/teachers on how to use eHealth, negative/positive attitude of students/teachers towards eHealth.

Conclusion: The successful uptake of eHealth in the curriculum of functional exercise/physical therapists needs a systematic multi-facetted approach considering the barriers and facilitators for uptake identified from the perspective of teachers and students. A relatively large amount of the identified barriers and facilitators were overlapping between teachers and students. Starting points for developing effective implementation strategies can potentially be found in reducing those overlapping barriers and facilitators.

Registration: The study protocol was a non-medical research and no registration was required. Participants gave written informed consent.

Introduction

Information and Communication Technologies (ICT; e.g. home automation, online training and games) are increasingly used to improve or support health and health care (*eHealth*) [1], such as. Physical therapists use eHealth interventions to support patients in maintaining independency in daily functioning but also for health care processes and services (e.g. telemedicine and electronic patient files) [2]. The role of technology in health care is growing and there is an urgent need for future health professionals who are able to work competently and confidently with eHealth.

The availability of technology in health care is growing and there is an urgent need for health professionals who can use eHealth competently and confidently in clinical practice. This means that there is large responsibility for the institutions for the education of future health professionals to ensure that students acquire knowledge, skills and attributes to work with eHealth, and this requires revision of curricula of education [3,4]. Students should be actively taught how to find, understand, apply and appraise eHealth innovations [5] to constantly update their skills and knowledge [2,6,7]. Ideally, students should become early adaptors and lead eHealth initiatives in settings where eHealth adoption is still low [13].

Despite the growing importance of eHealth in the work field, curricula are currently underdeveloped in teaching eHealth in the field of e.g. dietetics, nursing, occupational therapy, physiotherapy, psychology, or social work [5-9]. Therefore, a systematic approach to design, teach, assess or accredit eHealth education in the curriculum is needed. In the literature, a number of barriers for the uptake of eHealth education in curricula were identified: outdated and rigid curricula with narrow focus on technology [7], teachers' limited experience with and knowledge of the emerging field of eHealth [2,9,10] and health care teachers not feeling confident with technology [2,11,12].

There are several gaps in the knowledge of uptake of eHealth education in the curriculum. First, research about eHealth education predominantly comes from the medical and nursing literature [13]. However, the paramedical education for physical and functional exercise therapy, should also equip students to confidently use of eHealth, especially since eHealth innovations are increasingly used in daily practices of the physical and exercise therapist (e.g. Fysiogaming, eExercise, activity tracking, etc.). Since the work and patient groups of physical therapists differ from those of nurses, barriers and facilitators for eHealth education might also differ. Second, there is a need for more in-depth knowledge of barriers and facilitators for the uptake of eHealth in education, in terms of the factors that may critically influence uptake' [18]. Third, studies mainly focused on single groups of teachers, students or professionals, but not on the perspective of multiple groups.

This study aims to provide insight in the barriers and facilitators for uptake of eHealth in the education for physical therapy and functional exercise therapy, more specific to answer

the research question: what are the barriers and facilitators perceived by teachers and students for implementing eHealth in education?

Methods and materials

Design

A qualitative study was conducted among teachers and students to explore the perceived barriers and facilitators for uptake of eHealth education in the curricula of the education for physical therapy and functional exercise therapy. In this study eHealth was defined as 'the use of new Information and communication technologies (ICT) to improve or support health and health care' [1]. Given this definition, the technology employed in health care varies largely, e.g. web and mobile applications, electronic patient records, health-sensors and wearable devices, telecommunication, home automation and robotics and serious gaming [1]. 'EHealth education' was defined as teaching how to provide treatments using technology.

Focus groups were conducted to collect data that contributes to a better understanding of teachers' and students' attitudes, experiences with and expectations of eHealth in education [Kitzinger, 2006]. Participants were informed that data would remain confidential and would be anonymously used for scientific research and for improvement of eHealth education. The CONSolidated criteria for REporting Qualitative research (COREQ) guidelines were used for adequate reporting of the study [14].

Recruitment and inclusion

Teachers and students were recruited from two departments teaching functional exercise therapy and physical therapy in a four years full-time program, in the Netherlands: (1) Functional Exercise Therapy, Faculty of Health, University of Applied Sciences in Amsterdam (HvA) and (2) Physical Therapy, Division of Health Care, University of Applied Science in Leiden (HSL). Functional Exercise Therapy and Physical Therapy have similarities (both focus on restoring activities of daily life by means of exercises), but also have differences and are seen as two different paramedical health professions in the Netherlands. Two different health care educations from two different universities of applied sciences were included to ensure a diversity in the population of this study to improve the transferability of the findings.

Teachers were able to participate if they have taught for at least 2 years and met one of the following criteria: 1) Teachers from the departments of functional exercise therapy or physical therapy were included if they and were working, 2) OR working as a researcher in the field of eHealth. Students were able to participate if they were in year 3 or 4 of the study

and completed successfully their placement. Thus both teachers and students were able to reflect on the eHealth education in the curriculum as well as on requirements for successful use of eHealth in clinical practice. Teachers and students were invited to participate via the internal web page and a short oral presentation. Those who were willing to participate, received an email with study information, an informed consent form, and were invited for the focus group.

Focus groups

The focus groups took place in October and November 2016 at the Universities of Applied Sciences in Amsterdam and in Leiden. Separate groups were organized for teachers and students to ensure that both groups could talk freely about their experiences with eHealth in education. Group size was 5-8 participants to include a diversity of opinions and perspectives, and to allow optimal interaction between participants [15]. The focus groups were conducted by 1) a moderator (MW, female), 2) an assistant (student 1, female) who supported the moderator and managed the tape-recorders and time, and 3) an observer (student 2, female) who took notes and made sure every participant was given the opportunity to speak freely. The moderator has a master's degree in Health Sciences and functional exercise therapy and had formal training in conducting focus groups. The moderator was a colleague of some of the participating teachers and a former teacher of some of the participating students. Participants did not receive reimbursement for their participation.

An interview guide was developed based on the implementation model of Grol and Wensing. This model was chosen, since it offers a framework to identify and categorize barriers and facilitators for the uptake of innovations within a specific context, in this case the uptake of eHealth education in the curriculum. The framework includes six levels: Innovation (e.g. advantages, feasibility, accessibility, attractiveness of eHealth), Individual (e.g. motivation, awareness, knowledge, skills and attitude of students and teachers), Social context (e.g. opinion of colleagues, work culture), Organizational context (e.g. organization of the curriculum, capacities, resources, structures) and Political and Economic factors (e.g. financial arrangements, regulations, policies). By including questions according to each level of the framework the research team aimed to contribute to the need for more in-depth knowledge of factors (barriers and facilitators) that may critically influence the uptake of eHealth in education [13].

At the beginning of each focus group a brief description of eHealth was given. Open-ended questions within each level were asked to facilitate interactions and in depth discussion between the participants about eHealth education [16]. Examples of questions are: "What do you need in order to be able to use eHealth in education?" or "Why would you use eHealth in your lessons?". Prompts were used (e.g. pictures expressing emotions) to

facilitate participants in verbalizing thoughts. The interview guide was discussed and pilot tested in a group of students. The focus groups were planned to last approximately one hour and were aimed to continue until data saturation was reached (not more than two new subthemes retrieved from the focus groups).

Ethical issues and approval

All participants gave written informed consent prior to participation. Participants were informed that their statements were confidential, would be used to improve eHealth education and would not affect their position as a teacher or student. Although the study protocol was a non-medical research, we aimed to conduct the study conform the guidelines for 'Good Clinical Practice' (GCP).

Data analysis

Focus groups were audiotaped, transcribed in full and analyzed using direct content analysis. The implementation framework of Grol and Wensing (2004) is often used for implementing interventions and innovations in health care. Because the framework is highly structured and generally accepted in the field of implementation it was used in this study to structure and describe barriers and facilitators for implementing eHealth in education from the perspective of both students and teachers [17]. First step in the analyses was to identify barriers and facilitators for each level of the framework (innovation, individual teacher, individual student, social context, organizational context and political and economic factors) by initial coding of quotes. Second, quotes with comparable content for barriers and facilitators were categorized into (sub)themes. These (sub)themes were further analyzed and categorized into main themes. Data analysis was performed by two students who independently coded and categorized the data. Each step of the data analyses was discussed among the students until consensus was reached. The completed analyses was verified by a third researcher [MW]. Again, discrepancies were discussed among students and researcher until consensus was reached. Microsoft Office Excel was used for data analysis.

Results

Participants

Eleven teachers and 26 students indicated their willingness to participate and were invited for the focus groups. Two students were not present, since they had forgotten the appointment. A total of six focus groups was conducted, two with teachers (n=11) and four with students (n=24). Table 1 presents participants' characteristics.

Table 1. Characteristics of the participants in the focus groups.

Characteristics	Teachers (n=11)	Students (n=24)
Age, median (range)	38 (29-52)	23 (20-25)
Gender; male, yes (%)	5 (45)	15 (63)
Year in study, number (%)		
Third year	-	13 (54)
Fourth year	-	11 (46)
Profession, number (%) ^a		
Exercise therapy	3 (27)	-
Physical therapy	7 (64)	-
Health Sciences / Human Movement Sciences	4 (36)	-
Working experience in years, number (%)		
2-3	2 (18)	-
3-4	4 (36)	-
>4	5 (46)	-
Working as a health professional, yes (%)	7 (64)	-
Working as a researcher, yes (%)	4 (36)	-

^a Multiple answers possible.

Framework

In the first step of the analyses, a total number of 109 barriers and facilitators (codes) were retrieved from the six focus groups with teachers and students, from which 44 were overlapping between teachers and students, 27 were identified by teachers and 38 by students. Next, the barriers/facilitators were organized into 51 subthemes (see Additional file 1) and 14 main themes within the six levels of the framework of Grol and Wensing (see Table 2).

In the following sections, the main themes within each level of the framework will be discussed, first those themes identified by both teachers and students, then those for teachers only and then those for students only.

The innovation

Unclear concept of eHealth was mentioned by both teachers and students. Both groups report it is important for students' to learn to motivate what, when and how to use eHealth in the treatment process of a patient, and to learn to match available eHealth with the preferences of a patient and to provide support to patients with using eHealth. Both students and teachers referred to eHealth by mentioning applications, electronic patient records and digital exercises. Other available eHealth applications (e.g. virtual reality, robotics/ house automation) were barely mentioned. When discussing eHealth tools, both teachers and students wondered whether they were aware of all the possibilities.

Table 2. Results on codes, themes and levels of perceived barriers (B) and facilitators (F) for eHealth education according to teachers and students.

Codes (n= 109)	Themes	Level	Teachers		Students	
			B	F	B	F
26	Unclear concept of eHealth	Innovation	X		X	
	Lack of a quality mark and evidence for eHealth services.		X		X	
17	Capabilities of students on how to use eHealth	Individual student	X	X	X	X
	Attitude/behavior of students towards eHealth			X	X	X
22	Capabilities of teachers on how to use eHealth	Individual teacher	X	X	X	
	Attitude/behavior of teachers towards eHealth (education)		X	X	X	
14	Inefficient use of expertise	Social context	X		X	
	Communities of practice			X		
	Interprofessional collaboration/education		X	X	X	X
20	(Lack of) a shared vision within the organization	Organizational context	X	X		
	Situational factors (e.g. lack of time, slow curricula changes)		X	X		
10	Financial aspects (e.g. no reimbursement, time and money investment)	Economic and political context	X		X	
	Role of the government (e.g. quality mark for eHealth, reimbursement)			X		X
	Role of profession bodies (e.g. provision of education for therapists)					X

Lack of a quality mark and evidence for eHealth services was reported by both teachers and students. Both groups related the lack of a quality mark (i.e. applicability, usability, content, privacy, safety) and the relative absence of evidence for eHealth interventions to a lack of eHealth in education. *Teacher K: It is tricky. You can never implement something new in education if you waiting for the scientific evidence. The paradox with eHealth is that you are not going to know until you give it a try.*

The individual student

Capabilities to use eHealth was rated highly by teachers, since they agreed that current generations of students are in general competent with using technology. However, this does not imply that students are also able to innovate health care through eHealth and apply this in their professional work. *Teacher A1: "Although students are quite skilled in technical issues, I am often disappointed in their innovativeness."* Students do think they are capable to work with technology and eHealth, but do not use it since they are unfamiliar with the available eHealth services and have a lack of experience of applying it. *Student M1: "I think I could apply eHealth, but in fact I know quite little about the possibilities. For this reason*

I will not use it just now."

Attitude/ behavior towards eHealth towards eHealth differs between individual students, according both teachers and students. Some students are highly interested in technology and choose to get more involved, whereas others do not. When discussing use of eHealth as a (future) health professional, students' expressed it is optional, in terms of having a choice. Student L1: *"I think I will not really focus on it. I do want to know what the options are, but personally I will not devote to it for my future"*. Student J1: *"Once you have graduated you can make your own choice and decide whether you use it."*

The individual teacher

Capabilities to use eHealth, i.e. the knowledge about eHealth and skills on how to use eHealth, varies widely between individual teachers according to both the teachers and students. Some teachers admitted to having no overview of existing tools and eHealth interventions in their field of work or expressed to barely know how to use a projector, while others found themselves very competent as researchers in the domain of eHealth.

Attitude and behavior towards eHealth were rated positively towards eHealth education in general by teachers. Teacher V1: *"As a teacher I want to use eHealth in my lessons to provide 'future proof' education that is innovative and interactive. This makes learning more fun and challenging for students and they would be more enthusiastic about my lessons."* However, a barrier is that teachers feel insecure about eHealth education. Teacher K1: *"I do not feel that I know enough about eHealth, but that is what I want as a teacher before I use it in my lessons."* Students expressed that teachers have a negative attitude/behavior towards eHealth. Student S1: *"Teachers often say: find something you like, because you know better than me."* Student M2: *"If a passionate teacher puts something forward, then I am more open for it. However if a teacher says 'you are the young generation. You surely know of some app, and give it a try', then I am not. And the latter is what I have been told so far."*

The social context

Inefficient use of expertise of eHealth within the organization was mentioned as a barrier by both teachers and students. Student L1: *"Teachers do not collaborate. For example get a lecturer who knows a lot about eHealth to take over the lesson. All should benefit from it, since the less experienced teacher will catch up."* Teacher M2: *"More experienced teachers who are doing research in technology or eHealth should share their knowledge with less experienced teachers. A kind of cross-fertilization"*.

Communities of practice were recognized as a facilitator for eHealth education according to teachers. Such communities were seen as mixed groups (teachers, students, researchers and/or the workfield) of people who share their passion and learn how to do it better by interacting regularly. Teacher A1: *"we do quite a bit of research using eHealth. It would be great*

if we can get an exchange between research, education and practice and in this way create an inspiring environment around eHealth."

Interprofessional collaboration education is, by both teachers and students, regarded as a facilitator for eHealth in education if students of different professions work together (e.g. technology, ICT, media, etc.). *Student N1: "I think it would have an enormous added value if you would work on a project on eHealth together with students with an IT background; developing an app for example."*

The organizational context

(Lack of) shared vision/rationale in the university about what students should learn about eHealth is absent according to teachers. *Teacher D1: "As a school you have to decide to what degree you want to integrate this in your basic curricula. A choice can be to provide eHealth as a dedicated subject of choice. This can be a choice, but you will have to have an idea."* Teachers felt that what students learn about eHealth is too much of a coincidence, depending on a students' own interest in technology, the extent of eHealth experience during their internships/study route and the teachers they had. *"There is no clear approach for eHealth in the curricula in which connects to the professional roles (CanMeds) of the care professionals (OT/PT)."* Although there is not (yet) a clear vision about what students should learn, teachers did agree that there is a consensus within the university about the importance of eHealth education.

Situational factors, as mentioned by teachers, include the following barriers for eHealth education: lack of time for preparing lessons, failing technology, absence of didactic materials and relative slow curricula changes. Facilitators according to teachers would be: presence of ICT professionals within the organization, direct accessibility to materials (e.g. LivingLabs), (scheduled) time to prepare lessons, special interest group of teachers taking the lead and training for teachers to improve their competences.

Economic and political context

Financial aspects are expressed by both teachers and students in 10 quotes. The lack of reimbursement by health insurance companies in the Netherlands for eHealth interventions results in the absence of incentives to use eHealth in the field of work. Teachers specifically mentioned financial aspects such as the investments in technology and eHealth, needed for eHealth education, are a financial barrier for uptake.

The role of the government is to provide a definition of the future health professionals in relation to eHealth to give direction for eHealth education, according to the teachers. Students expressed that it is the role of the government to manage reimbursements for eHealth interventions and to lower the workload for health professionals. According to the students, this would facilitate health professionals to apply eHealth and consequently

students can experience eHealth during internships. Moreover, students want the government to improve the quality of eHealth by a national quality mark or at least a check list to determine quality of eHealth services.

Role of profession bodies. Students noticed during their internships eHealth is not yet imbedded in the work field. According to them, their professional organizations should facilitate uptake of eHealth in daily practice, for instance by providing education to health professionals and incorporation of eHealth in practical guidelines.

Discussion

The aim of this study was to identify barriers and facilitators for eHealth education as perceived by teachers and students of physical therapy and functional exercise therapy. Teachers and students equally contributed to the number of facilitators and barriers for the use of eHealth in professional education identified in this focus group study. Main barriers for the *innovation* were a lack of understanding the full concept of eHealth and a lack of knowledge and skills in critically appraising eHealth. For the *individual users*, the variety in knowledge and skills of individuals was a factor influencing uptake. On the level of the *organization*, identified factors for uptake were the shared sense of importance of implementing eHealth in education, a shared vision about what students should learn about eHealth and didactic materials. . *Economical* barriers were seen in the investments in technology and eHealth. Finally, *political* factors were identified in the national government to manage future reimbursements for eHealth interventions and to improve the quality of eHealth by a quality mark.

As expected, and based on the literature, we found unanimous support for implementing eHealth in education in the curricula of two departments in the Netherlands. In line with literature, we found barriers and facilitators on all levels of implementation as described by Groel and Wensing (2014) [17]. The barriers to the uptake of the eHealth *innovation* found in our study are in line with previous research: limited skills and knowledge about the eHealth intervention in both teachers and students [5,10,11,13,18], limited confidence in working with technology in health practice [6,11,13] and critically appraising and applying technique [5]. Lam (2016) mentioned: 'while students demonstrated the technical skills that would potentially enable them to engage in eHealth, they displayed a lack of understanding how these skills could be applied to professional health contexts' [13]. On the level of the *individual user*, there is a marked diversity in knowledge and skills in both teachers and students, which is in line with previous studies [5,13], and might hinder uptake of eHealth in the curricula.

On the level of *organization*, barriers and facilitators identified in our study add to the

growing consensus that the uptake of eHealth needs a multi-faceted approach and not just 'writing a new module'. In literature, static curricula with narrow focus on technology were reported as important barriers [7], and a clear need for a shared vision/rationale about what students should learn about eHealth and need didactic materials which is also reflected in literature [9]. *Political and economic* factors, i.e. the influential role of government, policies and professional bodies on uptake of eHealth in education, found in this study, were also reported elsewhere: this was phrased by Hilberts and Gray (2014) as the need for 'an education infrastructure in large-scale eHealth strategies' [18].

Strength of this study is that research in this field amongst physical therapists is relatively scarce. Another strength is the structured use of the model of Grol and Wensing to gain a more in-depth knowledge of barriers and facilitators for the uptake of eHealth in education and to enable comparison between the perspectives of teachers and students. EHealth is a relatively new and emerging field and, as a consequence, so is the implementation in education. There is surprisingly little evidence for the effectiveness of education focusing on the use eHealth [18]. Last but not least, a strength is that both teachers' and students' perspectives were included, whilst most studies focused on single groups of teachers, students or professionals. Limitations of this study are the professional relationship of the first author with some of the teachers and students. For that reason it was made very clear to the participants that their statements were confidential and not affected their position as a teacher/student. However, some bias in their responses cannot be ruled out entirely. Moreover, further focus groups might add to the data-saturation to some degree.

For uptake of eHealth in the curricula of physical and functional exercise therapy it is eminent to recognize the multi-level character of it. This study highlighted the need for a vision on eHealth at a faculty level. Besides a generally limited understanding of the width of eHealth and the expected impact of eHealth in clinical practice, this study showed that both the lack of skill in critically appraising the quality and usefulness of eHealth and the diversity in background knowledge and skills in technology need to be a point of engagement in future uptake plans. Moreover, the 'higher order' influences on both education and professional practice need to be addressed in eHealth education, i.e. the role of government, policies and professional bodies. In addition, to further enhance uptake it is strongly advised to take a structured approach by addressing the levels of uptake [17]. Using 'a clear rationale for teaching clinical informatics and a detailed list of desired competencies are an important start' [9], and keeping in mind that 'there is surprisingly little evidence about what works and doesn't work with regard to the eHealth education' [18]. Finally, more tools should be provided by the organizations itself, such as didactic materials and eHealth facilities.

This study provides insights into the many factors which influence the successful uptake of eHealth in the curricula of functional exercise and physical therapy education. This is

highly important given the fact that the application of eHealth is irreversible and health professionals do not seem to be fully equipped to work with eHealth. Uptake of eHealth needs a systematic multi-facetted approach considering factors on the level of the innovation, individual users, organization and political and economic levels. Important starting points for developing uptake strategies, for both teachers and students, are a limited knowledge of eHealth, a large diversity in eHealth skills, a lack of skills in critically appraising eHealth and to development of a clear rationale for teaching eHealth. A recommendation for further research is to re-examine the study in other health professions for a good comparison of perceived barriers and facilitators for eHealth education. Moreover, future research should provide evidence for what works and doesn't work with regard to the eHealth education.

Declarations

Ethics approval and consent to participate: All participants gave written informed consent prior to participation. The study protocol was a non-medical research and no registration was required.

Consent for publication: Not applicable.

Availability of data and material: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: Stichting Innovatie Alliantie supported this project financially (Grant 2014-046PRO).

Authors' contributions: MW conducted the focus groups and analysed the data. PS was a major contributor in writing the manuscript. JM, LBV, AK, JV and TV contributed to the study methodology. All authors read and approved the manuscript.

Acknowledgements: We thank all the students and teachers participating in this project. We are grateful for the students Simone Timmers, Linda Oud, Beau Smit, Fenna Jagtman, Naomi Lagerwerf and Bart Noordermeer who have contributed to this study.

Reference list

1. Wentzel MJ, Beerlage-de Jong N, Sieverink F, et al. Slimmer eHealth ontwikkelen en implementeren met de CeHRes Roadmap. In: Politiek C, Hoogendijk R, editors. CO-CREATIE EHEALTHBOEK eHealth, technisch kunstje of pure verandering? The Netherlands: Nictiz; 2014.
2. Van Vliet K, Grotendorst A, & Roodbol P. Anders kijken, anders leren, anders doen. Grensoverstijgend leren en opleiden in zorg en welzijn in het digitale tijdperk. Diemen, Zorginstituut Nederland; 2016.
3. Kaljouw M & van Vliet K. Naar nieuwe zorg en zorgberoepen: de contouren. Diemen, Zorginstituut Nederland; 2015.
4. Pate CL & Turner-Ferrier JE. Exploring linkages between quality, eHealth and healthcare education. *EHealth Systems Quality and Reliability: Models and Standards*, 2010; 146.
5. Stellefson M, Hanik B, Chaney B, Chaney D, Tennant B, & Chavarria EA. eHealth Literacy Among College Students: A Systematic Review With Implications for eHealth Education. *Journal of Medical Internet Research*, 2011; 13(4), e102.
6. Dattakumar A, Gray K, Henderson KB, Maeder A & Chenery H. We are not educating the future clinical health professional workforce adequately for eHealth competence: findings of an Australian study. *Studies in health technology and informatics*, 2012; 178:33-38.
7. Frenk J, Chen L, Bhutta, ZA, Cohen J, Crisp N, Evans T, ... & Kistnasamy B. Health professionals for a new century: transforming education to strengthen health systems in an interdependent world. *The lancet*, 2010; 376(9756), 1923-1958.
8. Dattakumar A, Gray K, Maeder, AJ & Butler-Henderson, K. Ehealth education for future clinical health professionals: an integrated analysis of Australian expectations and challenges. In *Medinfo*, 2012; p. 954.
9. Gray K & Sim J. Factors in the development of clinical informatics competence in early career health sciences professionals in Australia: a qualitative study. *Advances in health sciences education*. 2011; 16(1), 31-46.
10. Jacobs RJ, Lou JQ, Ownby RL, & Caballero J. A systematic review of eHealth interventions to improve health literacy. *Health informatics journal*. 2014; 22(2), 81-98.
11. Buckeridge DL & Goel V. Medical informatics in an undergraduate curricula: a qualitative study. *BMC medical informatics and decision making*. 2002; 2(1), 6.
12. Nagle L. Everything I know about informatics, I didn't learn in nursing school. *Nursing Leadership (CJNL)*. 2007; 20(3), 22-25.
13. Lam MK, Hines M, Lowe R, Nagarajan S, Keep M, Penman M & Power E. Preparedness for eHealth: Health sciences students' knowledge, skills, and confidence. *Journal of Information Technology Education: Research*, 2016; 15, 305-334.
14. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International journal for quality in health care* 2007; 19(6):349-357.
15. Kitzinger J. Focus group research: Using group dynamics to explore perceptions, experiences and understandings. In: Holloway I (ed) *Qualitative research in health care*, 1st edn. Open University Press, Berkshire, England. 2005: pp 56-70.
16. Kitzinger J. Focus groups. In: Pope C, Mays N (eds) *Qualitative research in health care*, 3rd edn. Blackwell publishing, Malden, 2006, pp 21-31.
17. Grol R, Wensing M. What drives change? Barriers to and incentives for achieving evidence-based practice. *Med J Aust*, 2004; 180:557-60
18. Hilberts S, Gray K. Education as ehealth infrastructure: considerations in advancing a national agenda for ehealth. *Adv in Health Sci Educ*. 2014; 19:115-127.



Summary & general discussion



SUMMARY

Aims of this thesis

eHealth is increasingly used in rehabilitation (eRehabilitation), yet evaluations of its effectiveness, the process of delivery and the perceptions of patients, their caregivers and health care providers are relatively scarce. This thesis aimed to:

- I. Evaluate the outcome and process of an eHealth intervention for cognitive stroke rehabilitation.
- II. Explore the (readiness for) use of eHealth among patients in rehabilitation.
- III. Investigate the requirements of patients, informal caregivers, health professionals, teachers and students regarding the use of eHealth interventions in stroke rehabilitation.

Main findings

The outcome and process of an eHealth intervention for cognitive stroke rehabilitation were addressed in Chapters 2 and 3.

Chapter 2 concerned a Randomized Controlled Trial (RCT) to evaluate the effect of an online serious brain training programme on multiple aspects of cognitive functioning, quality of life (QoL) and self-efficacy in comparison to a control intervention in stroke patients with self-perceived cognitive impairments. 110 stroke patients with self-perceived cognitive impairment 12–36 months after stroke were randomly allocated to the intervention (n=53) or control group (n=57). The intervention consisted of an 8-week brain training programme (Lumosity Inc.[®]). The control group received general information about the brain weekly. Assessments consisted of a set of neuropsychological tests and questionnaires on cognitive functioning, QoL and self-efficacy. After 8 and 16 weeks, no significant effect of the intervention was found on cognitive functioning, QoL or self-efficacy as compared to the control condition, except for small, yet statistically significant, effects on working memory and speed (mixed model analysis).

In **Chapter 3** the adherence of patients with the above mentioned 8-week brain training programme (intended use a minimum of 600 minutes of playtime) was described, with a focus on the role of health professionals' supervision. This study was part of the RCT described in Chapter 2 and used data from patients in the original intervention group (n=53) and the patients in the control group who accepted the offer to use the brain training programme after the initial trial period (n=52). Patients in the original intervention group

received supervision eight times during the training period (group 1), whereas patients in the original control group were only supervised twice (group 2). It was found that patients experienced several difficulties with tasks and conditions required to participate in the brain training (e.g. not able to cope with flashing screens, etc.). Only 24 out of the 105 patients (23%) were able to complete the intended dose of 600 minutes brain training over a period of 8 weeks, with the median playtime being 424 minutes (27-2162). The median playtime was significantly higher in group 1 (562 minutes, range 63-1264) than in group 2 (193 minutes, range 27-2162) ($p < 0.001$, Mann Whitney U). A possible explanation could be that excitement for the study waned for patients in group 2 who had to wait 16 weeks after the randomisation, whilst patients in group 1 started the training straight away after inclusion. Nevertheless, the findings may indicate that the frequency of the interaction with a supervisor can increase stroke patients' adherence with a brain training programme at home.

Given the suboptimal use of the brain training programme in Chapters 2 and 3, a descriptive study in **Chapter 4** explored the (readiness for) use of information and communication technology (ICT) devices (e.g. personal computer (PC), laptop, tablet, smartphone) among patients who were or had been admitted for rehabilitation. It also aimed to investigate patients' preferences to incorporate this technology in their rehabilitation process. For this study a cross-sectional design with a self-developed online questionnaire was used. The questionnaire comprised 61 questions about current possession and use of ICT devices, desired usage of ICT devices in the rehabilitation process in the future and socio-demographics and disease characteristics. Open answers, multiple choice and a 4-point Likert scale (1-4; totally disagree-totally agree) were used to examine current possession and user preferences. 190 out of 714 invited patients admitted for inpatient and/or outpatient rehabilitation and registered with an e-mail address, completed the questionnaire. 94 (49%) were women and 96 (51%) were men. The mean age of the participants was 49 (SD16) years. 149 out of 190 patients (78%) used one or more devices every day of the week. The most frequently used devices were: PC/laptop (93%), smartphone (57%) and tablet (47%). Most of the patients were willing to incorporate ICT devices in their rehabilitation process. The most frequently mentioned potential purposes for use in rehabilitation included: having insight in the medical health record, communication with peers and scheduled appointments with health professionals.

Chapter 4 provided insight in current and desired use of ICT devices in rehabilitation in general, but the perspectives of intended users regarding eRehabilitation after stroke are not yet explored. Therefore, requirements for use of stroke eRehabilitation were investigated and prioritized in chapters 5 and 6. **Chapter 5** describes the requirements for the accessibility, usability and content of comprehensive eRehabilitation after stroke as perceived by

patients with stroke, their informal caregivers and health professionals, as identified by a qualitative (focus group) study. In total, eight focus groups were conducted to identify user requirements; six with patients/caregivers and two with health care professionals involved in stroke rehabilitation (rehabilitation physicians, physiotherapists, occupational therapists, psychologists, team coordinators). Direct content analysis was used to identify the user requirements concerning three predefined categories: accessibility, usability and content. In total, 45 requirements emerged from the focus groups. It was concluded that the majority of requirements of patients, informal caregivers and health professionals concerned content (25 requirements), followed by usability (12 requirements) and accessibility (8 requirements). Moreover, requirements between stroke patients/informal caregivers and health care professionals differed on several aspects. For instance, a requirement of health professionals was that eRehabilitation programmes can be accessed by the computer in the rehabilitation center, whereas patients and caregivers preferred to use a smartphone or tablet.

Chapter 6 provides an overview of the most important requirements for comprehensive eRehabilitation after stroke according to larger groups of patients with stroke, their informal caregivers and health professionals (physicians, psychologists and physical therapists), as based on a quantitative (survey) study. In order to determine the importance of requirements, a questionnaire with a 4-point Likert scale (1-4; unimportant-important) was developed with statements regarding accessibility, usability and content of comprehensive eRehabilitation after stroke. 125 patients, 43 informal caregivers and 105 health professionals completed the survey. The most important requirements as perceived by the majority of all stakeholder groups were: applicability of eRehabilitation to possessed ICT-devices (e.g. tablet, smartphone, computer in rehabilitation center), support with usage (i.e. instruction videos, menu with frequently asked questions), physical exercises, general information about stroke, insight in the rehabilitation process (i.e. feedback about training results, final reports) and setting and evaluation of goals. Notable differences were also found between the stakeholder groups, for instance exercises for cognitive functioning were important for patients and informal caregivers, whilst this was a less important requirement for health professionals.

Except for matching eHealth interventions with the requirements of its users, successful adoption in stroke rehabilitation also depends on how well health professionals are prepared to use eHealth in daily practice. This readiness starts with educating health professionals to work competently and confidently with eHealth.

Chapter 7 describes the barriers and facilitators for eHealth education as perceived by teachers and students involved in the education of exercise/physical therapists. A qualitative study including six focus groups was conducted: two with teachers (n=11) and four with students (n=24), all selected from two universities of applied sciences in the Netherlands. Reported barriers and facilitators were identified, grouped and classified according to the levels of a generally accepted framework for implementation (innovation, individual teacher, individual student, social context, organizational context and political and economic factors). More barriers than facilitators were perceived for the uptake of eHealth education in the curriculum, by both educators and students. Most barriers and facilitators were identified on the level of the Innovation (eHealth, e.g. unclear concept) (n=26), followed by the Individual teacher (e.g. capabilities on how to use eHealth) (n=22) and the Organizational context (e.g. didactic materials) (n=20). Starting points for developing implementation strategies of eHealth (education) in the curriculum, for both teachers and students, can be found in reducing the barriers (e.g. limited knowledge of eHealth, lack of skills in critically appraising eHealth) and by using the facilitators (e.g. shared sense of importance of implementing eHealth in education, passionate teachers, didactic materials).

DISCUSSION

Focus of this thesis

This thesis focused on the evaluation of an eHealth intervention for cognitive stroke rehabilitation and the perspectives of different stakeholders on the uptake of eRehabilitation in general.

The studies included in this thesis showed that there was no overall effect of an online brain training programme on cognitive functioning of patients with stroke. Only performances on cognitive function tests that were similar to the games included in the intervention improved, no near transfer effect was found. Moreover, usage of the training was suboptimal and not all of the patients were able to complete it. In order to improve daily activities of stroke patients, computer tasks need to be closely related to the impaired task itself. Thus, computer-based cognitive rehabilitation (CBCR) needs to be tailored and adapted to each patient's individual profile. It would appear important to support stroke patients with CBCR training, since training is not well used by all patients. It is possible patients benefit more when they learn how to use strategies in their training and when motivated by clinicians.

However, regarding patients' ICT readiness, wishes and requirements it was also found that a relatively large amount of patients in rehabilitation wish to incorporate ICT in their rehabilitation treatment and that patients with a stroke have specific requirements regarding the accessibility, usability and content of eRehabilitation. The requirements of patients were not entirely similar to those of informal caregivers and health professionals, indicating that all perspectives of all stakeholders should be taken into account. In addition, developing tailored implementation strategies to implement eHealth in the bachelor curriculum of health professionals, based on the identified barriers and facilitators in this thesis is highly relevant to make sure that future health professionals are able to work with eHealth.

Given the abovementioned findings, this Discussion focuses on conditions that can facilitate the effect and uptake of eRehabilitation, with emphasis on rehabilitation of patients after stroke.

Part I. Evaluation of an eRehabilitation programme after stroke: outcome, process and study design.

Evidence for the effectiveness of cognitive eRehabilitation after stroke

Our finding (chapter 2 and 3) is in line with the suggesting that the evidence for the effectiveness of cognitive eRehabilitation in stroke is scanty [1-6].

A recent systematic review by Laver et al. concluded that the effect of cognitive

eRehabilitation through virtual reality on cognitive functioning in stroke patients is still unclear because of a lack of trials [2]. A recently performed RCT also found that a brain training programme, which was comparable to the training used in our study, did not improve cognitive functions, subjective cognitive functioning or quality of life in patients in the chronic phase after stroke compared to a control group (waiting list) [7]. The literature is however conflicting, as some studies have also found positive effects in patients with stroke or other acquired brain injury after brain training [8-17]. It should be noted though that effects were mostly seen for tasks (outcomes) similar to tasks in the training process (i.e. near transfer effect), rather than tasks that are dissimilar to the training (i.e. far transfer effects) [15-21]. In other words, outcomes in terms of tasks that are similar to the training are less likely to contribute to improvements in daily living than outcomes that are directly linked to activities done in daily life [6]. It is possible that cognitive eRehabilitation interventions in which the tasks are closely related to the cognitive tasks in daily life, are more effective to stimulate cognitive functioning after stroke than playing games. This is supported by a RCT of Faria et al. (2016). In this study the potential benefits of virtual reality based cognitive rehabilitation after stroke through simulated activities of daily living were compared to conventional therapy only [22]. The intervention involved a virtual simulation of a city where memory, attention, visuo-spatial abilities and executive functions tasks were integrated in the performance of several daily routines. A between groups analysis showed significantly greater improvements in global cognitive functioning, attention and executive functions in patients with stroke when comparing virtual reality to conventional therapy.

Except for the scarcity of trials on the effectiveness of stroke eRehabilitation, according to the previously mentioned systematic reviews the methodology and reporting is poor in many cases, hampering the interpretation of their findings [2;23-28].

The process of cognitive eRehabilitation programme after stroke

The RCT described in this thesis showed that only 24 out of 105 patients (23%) were able to complete the desired total number of 600 minutes of brain training over a period of 8 weeks. This is in line with other studies that concluded that intensive (eHealth)-exercise regimes after stroke are difficult to perform for stroke patients [29-32].

During the execution of the trial, patients experienced several difficulties with tasks and conditions required to participate in the brain training (e.g. no possession of the required operating system to run games on their computer, incompetency with technology, not able to cope with flashing screens, etc.). These barriers might have contributed to the relatively low usage. Other barriers were identified by Pugliese et al. for mobile tablet-based care in patients with stroke (e.g. complexity of therapy instructions, fine-motor requirements, and unreliability of internet or cellular connections) [33].

To allow successful uptake of eRehabilitation it is recommended to identify methods that

can minimize the barriers for the use of eRehabilitation and support maximize adherence of patients with stroke [33]. One of these methods, explored in this thesis, was the delivery of supervision during the eRehabilitation intervention. Two intensities of supervision offered by a health professional were compared, showing that more frequent supervision (weekly) resulted in significantly higher adherence levels than low frequent (4-weekly) supervision. This is in line with a review by Kelders et al. (2012), concluding that the frequency of interaction with a counselor was a significant predictor for adherence with web-based health interventions in different patient groups [34]. Therefore, regular interaction with a supervisor is important to increase stroke patients' adherence with eRehabilitation interventions and supports the importance of offering eHealth by means of a blended care strategy.

Study strategies to evaluate a cognitive eRehabilitation programme after stroke

Although effects of eRehabilitation interventions are often evaluated in conventionally designed clinical trials, as was done in the RCT presented in this thesis, these designs are not always appropriate to evaluate effects of eHealth [35-38]. For example, an RCT does not always represent daily health care practice because of its sometimes very strict inclusion and exclusion criteria [38]. Consequently, when the effectiveness of eHealth has been established in a "laboratory setting", the results may not be replicated in a different context, where e.g. patients with less digital skills or cognitive impairments are offered the intervention.

Another drawback of traditional research designs is that their time cycle is much longer than the speed at which eHealth develops and evolves. This can make (parts of) the innovation under study already outdated by the time the results of the RCT are published [39]. Moreover, 'early adopters' among patients and health professionals may use an eHealth innovation before the evidence is available [38], decreasing the contrast between intervention and control conditions.

All of the abovementioned aspects have implications for the design of future eRehabilitation research. First, alternative research designs than the traditional RCT are recommended in order to provide information about how an eHealth intervention works in health care practice and to make experiences of users more visible. The following (research) strategies can be considered for research in eRehabilitation:

- 1) Multiple evaluations and use of clinically meaningful outcome measures that are highly sensitive and quickly responsive to the effects being evaluated [39];
- 2) Using alternative resources to generate information, e.g. digital self-measurements, social media, online databases or personal health files, big data analysis [36;40];
- 3) Qualitative research into the experiences of patients; and
- 4) Action research. Action research, often led by a group of professionals as part of

a community of practice, is initiated to solve an immediate problem identified by professionals [41]. The goal is to investigate and solve the problem, for instance by developing guidelines, strategies and knowledge in order to improve the communities' work practices. This is done by active participation of the community of professionals itself (e.g. teachers, students, researchers, patients, health care professionals). After investigation of the problem, the group makes decisions, observes and keeps note of the consequences of changing the particular situation. So, the group participates in a change situation, whilst research is conducted simultaneously, alongside the iterative process of adapting, testing and evaluating the innovation.

To conclude with, it should be noted that the development, evaluation and implementation of stroke eRehabilitation interventions and associated research must be executed in collaboration with all relevant stakeholders to make eRehabilitation relevant and feasible for the intended users in health care [42;43]. An example of such a collaboration is a so called "Living Lab", where various stakeholders (e.g. knowledge institutions, health care organizations, practitioners, patients, financiers, innovators, established companies, startups, researchers, etc.) work together to develop, evaluate and implement eHealth. In a medical specialist rehabilitation center, Basalt rehabilitation, such a Living Lab, the SmartLab, is instituted. Here, a stepwise procedure is used to test innovations in rehabilitation for their potential added value and usability (www.medicaldeltalivinglab.nl; (<http://www.basaltrevalidatie.nl/onderzoek-innovatie/smartlab/>)). Subsequent steps involve research to systematically evaluate the effectiveness, experiences and costs, and implementation when appropriate.

Part II. Readiness and requirements for eRehabilitation after stroke: implications for the uptake, health care practice and education.

Incorporating perspectives of stakeholders in the development of stroke eRehabilitation

This thesis includes a number of studies aiming to identify the readiness and requirements for stroke eRehabilitation among stakeholders involved in stroke care. A survey study found that the possession and usage of ICT devices was relatively high among patients in rehabilitation and that most patients wished to use those devices during the rehabilitation process. Moreover, a focus group study showed that stroke patients, informal caregivers and health professionals had very specific requirements for stroke eRehabilitation, e.g. doing physical exercises, information about stroke and outcomes of rehabilitation and scheduling appointments with health professionals. These findings suggest that preferably, eHealth services should be developed where diverse purposes (e.g. telecommunication, training facilities, information, agenda, etc.) are combined in one digital platform to allow easy

access and use. Partly based on the outcomes of this qualitative study a comprehensive eRehabilitation platform for stroke was built and is currently evaluated in the Fit After Stroke 'Fast@Home' project (www.fastathome.org).

Moreover, our focus group study showed that eRehabilitation training facilities and feedback should both be adapted to patients' preferences and capabilities (tailored care) [44-50]. Offering support with daily use of an eRehabilitation programme (i.e. direct assistance, a helpdesk, videos with instructions and/or a menu with frequently asked questions) was considered a crucial element as well. These findings may imply that to develop stroke eRehabilitation programmes that match with the needs of the intended users, it is essential that eRehabilitation programmes are designed in co-creation with all relevant stakeholders ('co-design'). Requirements of patients, informal caregivers and health professionals should be incorporated in each step of the design process ('user-centered design'). This is especially important since this thesis showed that different types of stakeholders have different requirements to optimize usage of eRehabilitation. The aforementioned community of practice is a good method to establish active participation of all relevant stakeholders and thereby enhances co-creation and user-centered design in eRehabilitation.

The uptake of eRehabilitation in stroke care

The impact of a stroke varies widely among individuals with the optimal treatment depending on abilities, preferences and goals of both the patient and his or her caregiver(s) [51]. Health professionals can play a central role in this patient-centered delivery of eRehabilitation (e.g. adjustment of training facilities according to progression, feedback and motivation). They are able to link the patient to effective eRehabilitation interventions and can provide guidance with using a new eHealth service. It is recommended to further implement living labs, Communities of Practice, involvement of end-users (patient, caregiver, professional, students, designers, researchers, etc.) and co-create in ideation, testing, implementation, evaluation and upscaling.

In a qualitative study aiming to identify user requirements for stroke eRehabilitation by means of focus groups, the participating healthcare providers (rehabilitation physicians, physiotherapists, occupational therapists, psychologists, team coordinators and speech therapist) also acknowledged the importance of their role in the uptake of eRehabilitation in stroke care. Health professionals are often not aware of the opportunities of eHealth and poorly informed about available eHealth interventions [37]. It is often unclear to health practitioners how eRehabilitation can be effectively used for patients [52]. If evidence from research is available, it may not be suitable to directly support health practitioners in making clinical decisions [37;38]. Virtual reality, for instance, has emerged as a therapeutic tool facilitating motor learning for balance and gait rehabilitation in stroke patients [2]. The evidence, however, has not yet resulted in protocols or standardized guidelines and/

or a consensus regarding optimal intervention programmes (e.g. dosage and tasks) [52]. Furthermore, it is complex to integrate effective eHealth interventions in existing work processes.

Therefore, health professionals need to be supported with protocols and guidelines that provide insight which type of stroke eRehabilitation works for whom [37;53]. Moreover, an overview of applicable eHealth interventions in stroke rehabilitation should be developed based on different resources, such as evidence, experiences of users and patient consumer organizations and professional organizations [37]. Therefore, the above mentioned SmartLab is linked with the National eHealth Living Lab (NeLL), an eHealth community (patients, consumers, professionals, scientists, students, organisations) aiming to create the best eHealth solutions by sharing knowledge, contacts and experiences with each other (<https://nell.eu>).

Furthermore, this thesis showed that applicable eHealth needs to be integrated into working routines. A major challenge is realizing an integrated digital infrastructure, with patients, caregivers and healthcare providers connected in a safe, dynamic and efficient system.

eHealth in health professionals' education

Future health professionals should be able to competently and confidently work with eHealth [54;55]. However, education in eHealth is currently underrepresented in the curricula of health professions' education (e.g. dietetics, nursing, occupational therapy, physiotherapy, psychology or social work) [56-59]. In this thesis a focus group study among teachers and bachelor physical and exercise therapy students was conducted to identify barriers and facilitators for the uptake of eHealth education in the curriculum. Better integration of eHealth in the curriculum was considered important by both students and teachers who participated in the focus groups and they demonstrated their willingness to use eHealth. Nevertheless, in line with other studies in the field, their understanding of eHealth was limited [60]. Moreover, they expressed a lack of skills for critically appraising eHealth and the absence of a clear rationale for teaching eHealth. To enhance implementation in the curriculum of health care education clinical reasoning' will be the driver of the toolkit eHealth that is currently developed, based on our findings and in collaboration between three Universities of Applied Sciences.

A facilitator for the uptake of eHealth education identified in the focus groups was the optimal use of communities of practices. As described previously, in a community of practice, a mixed group (teachers, students, researchers and/or the work field) shares a passion and constantly innovates, in this case eHealth, by interacting regularly in a process of action research with all stakeholders [61]. Participation in such a community of practice benefits both teachers and students in their teaching and learning about eHealth [62]. In addition,

literature suggests that real-life experimentation of eHealth innovations in a realistic context with active user involvement (LivingLabs) are stimulating learning environments [63]. Other identified facilitators, which can be used in strategies for implementation, are the shared sense of importance of eHealth education, passionate teachers, didactic materials, collaboration with ICT professionals, direct accessibility to materials (lab with technology and eHealth), (scheduled) time for teachers to prepare lessons, special interest groups of teachers taking the lead and a national benchmark for the quality of eHealth.

In general, it is advised that eHealth education should be integrated in basic courses so that future healthcare professionals are already familiar with it, but also in postgraduate education to facilitate current use in daily practice [37].

Conclusion

In this thesis conditions that can facilitate the effect and uptake of eRehabilitation were described, with emphasis on stroke patients. The use of eRehabilitation after stroke is promising, but adoption of eRehabilitation in stroke care falls behind and their evidence regarding its effectiveness is scanty. More research is needed using appropriate study designs for evaluation of the outcomes and processes of eHealth. Moreover, development should be done in co-creation based on user requirements in order to increase the uptake in stroke care. To minimize barriers with usage, both patients and health professionals, should be supported in using eHealth and eHealth (education) needs to be integrated in the curriculum of (allied) health professions of universities for applied sciences.

Reference list

1. Ogourtsova T, Souza Silva W, Archambault PS, Lamontagne A. Virtual reality treatment and assessments for post-stroke unilateral spatial neglect: a systematic literature review. *Neuropsychological rehabilitation*. 2017;27(3):409-54.
2. Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. *Stroke*. 2018;49(4):e160-1.
3. Shin H & Kim K. Virtual reality for cognitive rehabilitation after brain injury: a systematic review. *Journal of physical therapy science*. 2015;27(9):2999-3002.
4. Spreij LA, Visser-Meily J, van Heugten CM, Nijboer TC. Novel insights into the rehabilitation of memory post acquired brain injury: a systematic review. *Frontiers in human neuroscience*. 2014;8:993.
5. Larson EB, Feigon M, Gagliardo P, Dvorkin AY. Virtual reality and cognitive rehabilitation: a review of current outcome research. *NeuroRehabilitation*. 2014;34(4):759-72.
6. van de Ven RM, Murre JM, Veltman DJ, Schmand BA. Computer-based cognitive training for executive functions after stroke: a systematic review. *Frontiers in human neuroscience*. 2016;10:150.
7. Van De Ven RM, Buitenweg JJ, Schmand B, Veltman DJ, Aaronson JA, Nijboer TC, Kruiper-Doesborgh SJ, van Bennekom CA, Rasquin SM, Ridderinkhof KR, Murre JM. Brain training improves recovery after stroke but waiting list improves equally: A multicenter randomized controlled trial of a computer-based cognitive flexibility training. *PLoS one*. 2017;12(3):e0172993.
8. De Luca R, Calabrò RS, Gervasi G, De Salvo S, Bonanno L, Corallo F, De Cola MC, Bramanti P. Is computer-assisted training effective in improving rehabilitative outcomes after brain injury? A case-control hospital-based study. *Disability and Health Journal*. 2014;7(3):356-60.
9. Gray JM, Robertson I, Pentland B, Anderson S. Microcomputer-based attentional retraining after brain damage: A randomised group controlled trial. *Neuropsychological Rehabilitation*. 1992;2(2):97-115.
10. Spikman JM, Boelen DH, Lamberts KF, Brouwer WH, Fasotti L. Effects of a multifaceted treatment program for executive dysfunction after acquired brain injury on indications of executive functioning in daily life. *Journal of the International Neuropsychological Society*. 2010(1):118-29.
11. Lundqvist A, Grundström K, Samuelsson K, Rönnerberg J. Computerized training of working memory in a group of patients suffering from acquired brain injury. *Brain Injury*. 2012;26(4-5):423-4.
12. van Vleet TM, Chen A, Vernon A, Novakovic-Agopian T, D'Esposito MT. Tonic and phasic alertness training: A novel treatment for executive control dysfunction following mild traumatic brain injury. *Neurocase*. 2015;21(4): 489-498.
13. Ruff RM, Mahaffey R, Engel J, Farrow C, Cox D, et al. Efficacy study of THINKable in the attention and memory retraining of traumatically head-injured patients. *Brain Injury*. 1994;8(1): 3±14.
14. Chen SH, Thomas JD, Glueckauf RL, Bracy OL. The effectiveness of computer-assisted cognitive rehabilitation for persons with traumatic brain injury. *Brain Injury*. 1997;11(3): 197-209.
15. Åkerlund E, Esbjörnsson E, Sunnerhagen KS, Björkdahl A. Can computerized working memory training improve impaired working memory, cognition and psychological health? *Brain Inj*. 2013;27:1649– 1657.
16. Westerberg H, Jacobaeus H, Hirvikoski T, Clevberger P, Östensson ML, Bartfai A, Klingberg T. Computerized working memory training after stroke—a pilot study. *Brain Injury*. 2007;21(1):21-9.
17. Barker-Collo SL, Feigin VL, Lawes CM, Parag V, Senior H, Rodgers A. Reducing attention deficits after stroke using attention process training: a randomized controlled trial. *Stroke*. 2009;40(10):3293-8.
18. Björkdahl A, Åkerlund E, Svensson S, Esbjörnsson E. A randomized study of computerized working memory training and effects on functioning in everyday life for patients with brain injury. *Brain injury*. 2013;27(13-14):1658-65.
19. Lundqvist A, Gerdle B, Rönnerberg J. Neuropsychological aspects of driving after a stroke—in the simulator and on the road. *Applied Cognitive Psychology: The Official Journal of the Society for Applied Research in Memory and Cognition*. 2000(2):135-50.
20. Zickefoose S, Hux K, Brown J, Wulf K. Let the games begin: A preliminary study using Attention Process Training-3 and Lumosity™ brain games to remediate attention deficits following traumatic brain injury. *Brain injury*. 2013;27(6):707-16.

21. Van Vleet TM, Chen A, Vernon A, Novakovic-Agopian T, D'Esposito MT. Tonic and phasic alertness training: a novel treatment for executive control dysfunction following mild traumatic brain injury. *Neurocase*. 2015;21(4):489-98.
22. Faria AL, Andrade A, Soares L, i Badia SB. Benefits of virtual reality based cognitive rehabilitation through simulated activities of daily living: a randomized controlled trial with stroke patients. *Journal of neuroengineering and rehabilitation*. 2016;13(1):96.
23. Iruthayarajah J, McIntyre A, Cotoi A, Macaluso S, Teasell R. The use of virtual reality for balance among individuals with chronic stroke: a systematic review and meta-analysis. *Topics in stroke rehabilitation*. 2017;24(1):68-79.
24. Palma GC, Freitas TB, Bonuzzi GM, Soares MA, Leite PH, Mazzini NA, Almeida MR, Pompeu JE, Torriani-Pasin C. Effects of virtual reality for stroke individuals based on the International Classification of Functioning and Health: a systematic review. *Topics in stroke rehabilitation*. 2017 24(4):269-78.
25. de Rooij IJ, van de Port IG, Meijer JW. Effect of virtual reality training on balance and gait ability in patients with stroke: systematic review and meta-analysis. *Physical therapy*. 2016;96(12):1905-18.
26. Gibbons EM, Thomson AN, de Noronha M, Joseph S. Are virtual reality technologies effective in improving lower limb outcomes for patients following stroke—a systematic review with meta-analysis. *Topics in stroke rehabilitation*. 2016;23(6):440-57.
27. Veerbeek JM, Langbroek-Amersfoort AC, Van Wegen EE, Meskers CG, Kwakkel G. Effects of robot-assisted therapy for the upper limb after stroke: a systematic review and meta-analysis. *Neurorehabilitation and neural repair*. 2017;31(2):107-21.
28. Rintala A, Päivärinne V, Hakala S, Paltamaa J, Heinonen A, Karvanen J, Sjögren T. Effectiveness of technology-based distance physical rehabilitation interventions for improving physical functioning in stroke: a systematic review and meta-analysis of randomized controlled trials. *Archives of physical medicine and rehabilitation*. 2018;40(4):373- 387.
29. Shaughnessy M, Resnick BM, Macko RF. Testing a model of post-stroke exercise behavior. *Rehabilitation nursing*. 2006;31(1):15-21.
30. Bergquist TF, Yutsis M, Sullan MJ. Satisfaction with cognitive rehabilitation delivered via the internet in persons with acquired brain injury. *Int J Telerehabil*. 2014;6(2):39.
31. Connor BB, Standen PJ. So much technology, so little time: Factors affecting use of computer-based brain training games for cognitive rehabilitation following stroke. *Proc 9th Int Conf Disabil Virtual Reality Assoc Tech*. 2012;53–59.
32. Cruz VT, Pais J, Bento V, et al. A rehabilitation tool designed for intensive web-based cognitive training: Description and usability study. *JMIR Res Protoc*. 2013;2(2):59.
33. Pugliese M, Ramsay T, Johnson D, Dowlatshahi D. Mobile tablet-based therapies following stroke: A systematic scoping review of administrative methods and patient experiences. *PloS one*. 2018;13(1):e0191566.
34. Kelders SM, Kok RN, Ossebaard HC, Van Gemert-Pijnen JE. Persuasive system design does matter: A systematic review of adherence to webbased interventions. *J Med Internet Res*. 2012;14(6):152.
35. Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, Moore L, O'Cathain A, Tinati T, Wight D, Baird J. Process evaluation of complex interventions: Medical Research Council guidance. *bmj*. 2015 Mar 19;350:h1258.
36. May A & Matthijsen J. *"Alternatieven voor RCT bij de evaluatie van effectiviteit van interventies!?" - eindrapportage*. ZonMw, UMC, Tranzo. Oktober 2015. file:///homedir.ad.hva.nl/wenmm/Downloads/Eindrapportage_27okt2015_FINAL_Anne_Jolanda.pdf (accessed 21 January 2019).
37. Ministry of Health, Welfare and Sport [in Dutch: Ministerie van Volksgezondheid, welzijn en sport (VWS)]. *Voortgangsrapportage e-health en zorgvernieuwing*. 2016. Den Haag: Ministerie van Volksgezondheid, Welzijn en Sport.
38. Krijgsman J, Peeters J, Waverijn G, Lettow B, van der Hoek L, de Jong J, Friele R, van Gennip L. *Rapportage eHealth-doelstellingen 2016*. Den Haag: Nictiz, Utrecht: Nivel.
39. Baker TB, Gustafson DH, Shah D. How can research keep up with eHealth? Ten strategies for increasing the timeliness and usefulness of eHealth research. *Journal of medical Internet research*. 2014;16(2).
40. Galson S & Simon G. Real-World Evidence to Guide the Approval and Use of New Treatments. *NAM Perspectives*. 2016. Discussion Paper, National Academy of Medicine, Washington, DC.
41. Adelman C. Kurt Lewin and the origins of action research. *Educational action research*. 1993;1(1):7-24.

42. Kazadi K, Lievens A, Mahr D. Stakeholder co-creation during the innovation process: Identifying capabilities for knowledge creation among multiple stakeholders. *Journal of Business Research*. 2016;69(2):525-40.
43. The Netherlands Organisation for Health Research and Development [in Dutch: de Nederlandse organisatie voor gezondheidsonderzoek en zorginnovatie (ZonMw)]. *Actieplan eHealth*. 2013. <https://www.zonmw.nl/nl/over-zonmw/ehealth-en-ict-in-de-zorg/programmas/programma-detail/actieplan-ehealth/publicaties/> (accessed 21 January 2019).
44. Sivan M, Gallagher J, Holt R, Weightman A, Levesley M, Bhakta B. Investigating the international classification of functioning, disability, and health (ICF) framework to capture user needs in the concept stage of rehabilitation technology development. *Assistive Technology* 2014;26(3):164-173.
45. Nasr N, Leon B, Mountain G, Nijenhuis SM, Prange G, Sale P, et al. The experience of living with stroke and using technology: opportunities to engage and co-design with end users. *Disability and Rehabilitation: Assistive Technology* 2016;11(8):653-660.
46. Mawson S, Nasr N, Parker J, Zheng H, Davies R. & Mountain G. Developing a personalised self-management system for post stroke rehabilitation; utilising a user-centred design methodology. *Disability and Rehabilitation: Assistive Technology* 2014;9(6):521-528.
47. Parker J, Mawson S, Mountain G, Nasr N, Davies R, Zheng H. The provision of feedback through computer-based technology to promote self-managed post-stroke rehabilitation in the home. *Disability and Rehabilitation: Assistive Technology*. 2014;9(6):529-38.
48. Lange B, Flynn S, Proffitt R, Chang CY, Rizzo A. Development of an interactive game-based rehabilitation tool for dynamic balance training. *Topics in stroke rehabilitation* 2010;17(5):345-352.
49. Zheng H, Davies R, Zhou H, Hammerton J, Mawson SJ, Ware PM, et al. SMART project: application of emerging information and communication technology to home-based rehabilitation for stroke patients. *International Journal on Disability and Human Development* 2006;5(3):271.
50. Mountain G, Ware P, Hammerton J, Mawson S, Zheng H, Davies R, et al. The SMART project: a user led approach to developing applications for domiciliary stroke rehabilitation. *Designing accessible technology* 2006;135-144.
51. Wentzel MJ, Beerlage-de Jong N, Sieverink F, et al. Slimmer eHealth ontwikkelen en implementeren met de CeHRes Roadmap. In: Politiek C, Hoogendijk R, editors. CO-CREATIE EHEALTHBOEK eHealth, technisch kunstje of pure verandering? The Netherlands: Nictiz; 2014
52. Porras DC, Siemonsma P, Inzelberg R, Zeilig G, Plotnik M. Advantages of virtual reality in the rehabilitation of balance and gait: Systematic review. *Neurology*. 2018;90(22):1017-25.
53. Porras DC, Siemonsma P, Inzelberg R, Zeilig G, Plotnik M. Advantages of virtual reality in the rehabilitation of balance and gait: Systematic review. *Neurology*. 2018;90(22):1017-25.
54. Kaljouw M & van Vliet K. Naar nieuwe zorg en zorgberoepen: de contouren. Diemen, Zorginstituut Nederland; 2015.
55. Pate CL, Turner-Ferrier JE. Exploring linkages between quality, e-health and healthcare education. *International Journal of Healthcare Delivery Reform Initiatives (IJHDRI)*. 2010;2(4):66-81.
56. Dattakumar A, Gray K, Henderson KB, Maeder A & Chenery H. We are not educating the future clinical health professional workforce adequately for eHealth competence: findings of an Australian study. *Studies in health technology and informatics*. 2012;178:33-38.
57. Dattakumar A, Gray K, Maeder, AJ & Butler-Henderson, K. Ehealth education for future clinical health professionals: an integrated analysis of Australian expectations and challenges. In *Medinfo*, 2012; p. 954.
58. Frenk J, Chen L, Bhutta, ZA, Cohen J, Crisp N, Evans T, ... & Kistnasamy B. Health professionals for a new century: transforming education to strengthen health systems in an interdependent world. *The lancet*, 2010; 376(9756), 1923-1958.
59. Stellefson M, Hanik B, Chaney B, Chaney D, Tennant B, & Chavarria EA. eHealth Literacy Among College Students: A Systematic Review With Implications for eHealth Education. *Journal of Medical Internet Research*. 2011;13(4):e102.
60. Lam MK, Hines M, Lowe R, Nagarajan S, Keep M, Penman M & Power E. Preparedness for eHealth: Health sciences students' knowledge, skills, and confidence. *Journal of Information Technology Education: Research*. 2016;15:305-334.
61. Wenger E, McDermott RA, Snyder W. Cultivating communities of practice: A guide to managing knowledge. Harvard Business Press; 2002.

62. Hien TT. Why is action research suitable for education?. VNU Journal of Science: Foreign Studies. 2016 25(2).
63. Schuurman D, De Marez L, Ballon P. The impact of living lab methodology on open innovation contributions and outcomes. Technology Innovation Management Review. 2016;1(6).



The background of the page is a light gray with a complex, abstract pattern. It features a network of thin gray lines connecting small circles, some of which are white and others are dark gray. Overlaid on this network are several hexagonal shapes, some of which are solid light gray and others are outlined. Some of the hexagons contain a plus sign (+) symbol. The overall aesthetic is clean, modern, and technical, suggesting a scientific or medical theme.

Appendix

Samenvatting en discussie

Publications

Curriculum Vitae

Dankwoord

SAMENVATTING

eHealth wordt steeds meer gebruikt in de revalidatie (eRevalidatie). Wetenschappelijk onderzoek naar de effectiviteit en de implementatie van eRevalidatie is echter relatief schaars. Dit proefschrift richt zich op:

- I. Een evaluatie van het effect en het proces van een eHealth interventie in de cognitieve revalidatie na een CVA.
- II. Het gebruik van eHealth door patiënten in de revalidatie.
- III. Het inventariseren van randvoorwaarden van patiënten, mantelzorgers, zorgprofessionals, docenten en studenten met betrekking tot het gebruik van eHealth interventies in de revalidatie na een CVA.

Algemene bevindingen

Een aanzienlijk deel van de mensen met een CVA ondervindt cognitieve klachten, zoals problemen met het geheugen, initiatief nemen of het oplossen van problemen.

Hoofdstuk 2 beschrijft een Randomized Controlled Trial (RCT) om de effectiviteit van een online cognitieve training te onderzoeken. Honderdtien patiënten die 12-36 maanden na een CVA beperkingen hadden in het cognitief functioneren werden toegewezen aan de interventie- (n=53) of controlegroep (n=57). De interventiegroep doorliep een 8 weken durende, online cognitieve training (Lumosity Inc.[®]). De controlegroep ontving wekelijks algemene informatie over cognitieve functies. De metingen werden uitgevoerd na afloop van de training (8 weken) en 8 weken daarna (16 weken). De metingen bestonden uit een set van neuropsychologische tests (werkgeheugen, aandacht, flexibiliteit in denken, reactiesnelheid en vloeibare intelligentie) en vragenlijsten die betrekking hadden op het cognitief functioneren, de kwaliteit van leven en de 'self-efficacy' (het vertrouwen dat mensen hebben in hun eigen vermogen om specifiek gedrag in verschillende omstandigheden uit te voeren). Na 8 en 16 weken werden kleine, statistisch significante effecten op werkgeheugen en reactiesnelheid gevonden. Er waren geen effecten van de cognitieve training op andere belangrijke aspecten van cognitief functioneren (aandacht, flexibiliteit), kwaliteit van leven of de self-efficacy in vergelijking met de controle interventie.

Hoofdstuk 3 beschrijft de procesevaluatie van de RCT die beschreven is in hoofdstuk 2. Er werden gegevens gebruikt van patiënten uit de oorspronkelijke interventiegroep (n=53, groep 1), aangevuld met gegevens van patiënten uit de controlegroep, die na afloop van de RCT de 8-weekse training alsnog volgden (n=52, groep 2). Patiënten in de oorspronkelijke interventiegroep kregen gedurende de trainingsperiode 8 keer supervisie

door een zorgverlener, terwijl patiënten in de oorspronkelijke controlegroep slechts twee keer supervisie hadden.

Uit het contact met de zorgprofessionals bleek dat patiënten diverse problemen ondervonden bij deelname aan de cognitieve training, zoals bijvoorbeeld overprikkeling door flikkeringen op het beeldscherm. Slechts 24 van de 105 patiënten (23%) waren in staat om de beoogde 600 speelminuten over een periode van 8 weken te voltooien. De mediane speeltijd was 424 minuten (range van 27 tot 2162 minuten) en was statistisch significant hoger in groep 1 (562 minuten, range 63 tot 1264 minuten) dan in groep 2 (193 minuten, range 27 tot 2162 minuten). Dit verschil kan mogelijk verklaard worden door een intensievere begeleiding in groep 1. De verschillen kunnen ook mede veroorzaakt zijn door de lange wachttijd voor patiënten in groep 2, die tot 16 weken na de randomisatie moesten wachten om de training te kunnen volgen. Desalniettemin lijken de bevindingen uit deze studie erop te wijzen dat interactie met een zorgverlener het gebruik van een cognitieve training door CVA-patiënten kan verhogen.

Vanwege het suboptimale gebruik van de cognitieve training werd in een beschrijvende studie in **hoofdstuk 4** het gebruik van informatie- en communicatietechnologie (ICT) onderzocht bij patiënten die waren opgenomen in een revalidatiecentrum. Ook werden hun voorkeuren voor het gebruik van ICT in het eigen revalidatieproces geïnventariseerd. Deze inventarisatie bestond uit eenmalige afname van een zelfontwikkelde online vragenlijst. De vragenlijst bestond uit 61 vragen over het bezit en gebruik van ICT middelen, het gewenste gebruik van ICT in het revalidatieproces en enkele sociaal-demografische kenmerken.

714 patiënten die klinisch of poliklinisch hadden gerevalideerd en van wie het e-mailadres bekend was werden uitgenodigd voor het onderzoek. Honderdnegentig van de 714 patiënten vulden de vragenlijst in (27%); 94 (49%) vrouwen en 96 (51%) mannen, met een gemiddelde leeftijd van 49 (Standaard Deviatie (SD) 16) jaar. Uit de resultaten bleek dat 149 van de 190 patiënten (78%) dagelijks één of meer ICT-toepassingen gebruikten. De meest gebruikte apparaten hierbij waren een: personal computer (PC)/laptop (93%), smartphone (57%) en tablet (47%). Het overgrote deel van de patiënten was bereid om ICT in het eigen revalidatieproces te gebruiken. De meest voorkomende redenen hiervoor waren: inzicht in het eigen medische dossier, contact met lotgenoten en afspraken plannen met behandelaars.

Terwijl Hoofdstuk 4 inzicht geeft in gebruik van ICT in de revalidatie in het algemeen, werden de wensen en eisen voor de inhoud en het gebruik van eRevalidatie specifiek na een CVA systematisch onderzocht in hoofdstukken 5 en 6.

Hoofdstuk 5 beschrijft de eisen die patiënten met een CVA, hun mantelzorgers en behandelaars stellen aan de toegankelijkheid, bruikbaarheid en inhoud van eRevalidatie. Deze werden onderzocht in een kwalitatieve focusgroep studie (n=60). Er werden 8 focusgroepen uitgevoerd; 6 met patiënten/mantelzorgers en 2 met zorgprofessionals die betrokken zijn bij de revalidatie na een CVA (revalidatieartsen, fysiotherapeuten, ergotherapeuten, psychologen, teamcoördinatoren). De gebruikersvereisten werden ingedeeld in 3 categorieën: 1) *toegankelijkheid*, 2) *bruikbaarheid* en 3) *inhoud*.

In totaal werden er 45 vereisten benoemd in de focusgroepen. De meeste eisen van patiënten, mantelzorgers en zorgprofessionals hadden betrekking op *inhoud* (25 vereisten), gevolgd door *bruikbaarheid* (12 vereisten) en *toegankelijkheid* (8 vereisten). De eisen van patiënten/mantelzorgers en zorgprofessionals verschilden op een aantal punten van elkaar. Zo was een vereiste van zorgprofessionals dat eRevalidatie toegankelijk moet zijn via een computer in het revalidatiecentrum, terwijl patiënten en mantelzorgers de voorkeur gaven aan het gebruik via smartphone of tablet.

Aan een kwalitatief onderzoek kunnen slechts een beperkt aantal vertegenwoordigers van patiënten, mantelzorgers en behandelaars meewerken. Om een grotere groep te kunnen bereiken werd vervolgens nog een kwantitatieve studie uitgevoerd. **Hoofdstuk 6** geeft een overzicht van de belangrijkste vereisten voor eRevalidatie na een CVA zoals vastgesteld in een grotere groep patiënten met een CVA, hun mantelzorgers en zorgverleners (revalidatieartsen, psychologen en fysiotherapeuten). Hiertoe werden de verschillende eisen voortkomend uit de kwalitatieve studie (hoofdstuk 5) verwerkt in een vragenlijst, waarbij respondenten het belang van uitspraken over de toegankelijkheid, bruikbaarheid en inhoud van eRevalidatie na een CVA scoorden op een 4-puntsschaal (1-4; onbelangrijk-belangrijk).

125 patiënten, 43 mantelzorgers en 105 behandelaars vulden de vragenlijst in. De vereisten voor eRevalidatie die zowel door patiënten, mantelzorgers als behandelaars belangrijk gevonden werden waren: het kunnen gebruiken van interventies op de meest gebruikte ICT-apparaten (bijv. tablet, smartphone, computer in revalidatiecentrum), ondersteuning bij het gebruik (bijv. instructievideo's, menu met veel gestelde vragen), begeleiding bieden bij het fysiek oefenen, algemene informatie over een CVA, inzicht in het revalidatieproces (bijv. feedback op trainingsresultaten, eindrapportages over het revalidatieproces) en het vaststellen en evalueren van revalidatiedoelen. Er waren ook opvallende verschillen tussen de groepen; zo werden oefeningen voor cognitief functioneren als zeer belangrijk aangemerkt door patiënten en mantelzorgers, terwijl behandelaars dit juist minder belangrijk vonden.

Naast de afstemming van eRevalidatie na een CVA op de wensen en behoeften van gebruikers, hangt succesvol gebruik ook af van de bereidheid, het vertrouwen en de

vaardigheden van behandelaars om eHealth in hun dagelijkse praktijk te gebruiken. De basis hiervoor wordt gelegd bij het opleiden van zorgprofessionals. Om die reden beschrijft **Hoofdstuk 7** de belemmerende en bevorderende factoren voor het gebruik van eHealth in het onderwijs vanuit het perspectief van docenten en studenten van opleidingen Oefentherapie Mensendieck en Fysiotherapie. Hiervoor werd een kwalitatieve focusgroep studie met zes focusgroepsessies uitgevoerd: twee met docenten (n=11) en vier met studenten (n=24), van twee hogescholen in Nederland (Hogeschool van Amsterdam en Hogeschool Leiden). De in de focusgroepen genoemde bevorderende en belemmerende factoren werden aan de hand van het model voor implementatie van Grol en Wensing als volgt geclassificeerd: *innovatie, individuele docent, individuele student, sociale context, organisatorische context en politieke & economische factoren*. Uit de resultaten bleek dat, door zowel docenten als studenten, meer belemmerende dan bevorderende factoren werden gevonden voor toepassing van eHealth in het curriculum. De meeste factoren hadden betrekking op de *innovatie* bijv. onduidelijkheid rondom het concept eHealth (n=26), gevolgd door de *individuele docent* (bijv. vaardigheid in het gebruik van eHealth) (n=22) en de *organisatorische context* (bijv. beschikbaarheid van lesmateriaal en technologie) (n=20). Deze bevindingen vormen de basis voor implementatiestrategieën die gericht zijn op zowel het wegnemen van belemmeringen (bijv. gebrek aan kennis en vaardigheden op het gebied van eHealth) als het benutten van bevorderende factoren (bijv. beschikbare eHealth materialen).

ALGEMENE DISCUSSIE

Focus van dit proefschrift

Dit proefschrift richtte zich op de evaluatie van een eHealth interventie voor cognitieve revalidatie na een CVA (eRevalidatie) en de perspectieven van verschillende betrokkenen over de inzet van eRevalidatie in het algemeen, maar in het bijzonder na een CVA.

De studies in dit proefschrift laten zien dat er vrijwel geen effect was van een online cognitieve training op het cognitief functioneren van patiënten met een CVA. Hoewel een verbetering gevonden werd op cognitieve taken die vergelijkbaar waren met de spellen in het trainingsprogramma werd geen effect op taken anders dan de getrainde taak en de kwaliteit van leven gevonden. Dit suggereert dat computertaken in online cognitieve training nauw verband moeten houden met de daadwerkelijke problemen die patiënten na een CVA ondervinden tijdens dagelijkse activiteiten. Omdat patiënten na een CVA heel diverse cognitieve beperkingen kunnen hebben vraagt dit om het 'op maat maken' van een cognitieve training voor elke patiënt.

Een andere bevinding in dit proefschrift was dat het gebruik van de cognitieve training tegenviel en flink varieerde tussen patiënten. Een deel gebruikte de training niet, anderen waren niet in staat om de vereiste trainingsduur te bereiken. Dit kan samenhangen met specifieke eisen van CVA-patiënten met betrekking tot de toegankelijkheid, bruikbaarheid en inhoud van eRevalidatie. Het bleek echter ook belangrijk dat patiënten goed ondersteund worden door zorgprofessionals bij het gebruik van cognitieve eHealth interventies. Om dat te kunnen doen is het essentieel dat ook het perspectief van behandelaars wordt meegenomen in de ontwikkeling van eHealth. Naast het betrekken van professionals in het werkveld is het evenzo belangrijk om ervoor te zorgen dat toekomstige zorgprofessionals adequaat kunnen werken met eHealth en in staat zijn de beroepspraktijk te innoveren. Hiervoor zijn aanpassingen van het curriculum nodig, gebaseerd op door docenten en studenten ervaren belemmerende en bevorderende factoren, zoals beschreven in dit proefschrift.

Deel I. Evaluatie van een eRevalidatie programma na een CVA: resultaat, proces en onderzoekstrategie.

Bewijs voor de effectiviteit van cognitieve eRevalidatie na een CVA

Onze bevindingen (hoofdstuk 2 en 3) komen overeen met de constatering dat het bewijs voor de effectiviteit van cognitieve eRevalidatie na een beroerte schaars is [1-6]. Zo concludeerde een recente systematische review dat het effect van eRevalidatie op cognitief functioneren middels virtual reality bij patiënten na een CVA nog steeds onduidelijk is, niet alleen door een gebrek aan studies, maar ook door de matige kwaliteit ervan [2]. Onderzoek dat wel voorhanden is laat tegenstrijdige resultaten zien. Een recent uitgevoerde RCT concludeerde dat na een cognitieve training, vergelijkbaar met de training die in onze studie werd gebruikt, de cognitieve functies, het subjectieve cognitieve functioneren en de kwaliteit van leven bij patiënten in de chronische fase na een beroerte niet verbeterde in vergelijking met een controlegroep [7]. Sommige studies vonden wel positieve effecten van cognitieve training bij patiënten met een CVA of ander hersenletsel [8-17]. Echter, deze effecten werden meestal gezien voor taken (uitkomsten) die vergelijkbaar waren met taken in de training (d.w.z. 'near transfer of learning'), en niet voor taken die daarmee niet vergelijkbaar waren (d.w.z. 'far transfer of learning') [15-21].

Het is mogelijk dat cognitieve eRevalidatie interventies waarbij de taken nauw verband houden met de cognitieve taken in het dagelijks leven effectiever zijn om het cognitief functioneren na een CVA te verbeteren dan het spelen van braingames, zoals gebruikt werden in ons onderzoek. Dit wordt ondersteund door een RCT van Faria et al. (2016). In deze studie werd het effect van virtual reality met gesimuleerde dagelijkse activiteiten op de uitvoer van cognitieve taken vergeleken met traditionele therapie [22]. De interventie betrof een virtuele simulatie van een stad waar geheugen, aandacht, visuo-ruimtelijke capaciteiten en uitvoerende functionele taken werden geïntegreerd in de uitvoering van verschillende dagelijkse routines. Analyse tussen de groepen toonde aanzienlijk grotere verbeteringen van de virtual reality training op globaal cognitief functioneren, aandacht en executieve functies bij CVA-patiënten in vergelijking met traditionele therapie.

Het proces van een cognitief eRevalidatie programma na een CVA

De RCT in dit proefschrift liet zien dat slechts 24 van de 105 patiënten (23%) het gewenste aantal van 600 minuten cognitieve training in een periode van 8 weken volbrachten. Dit enigszins teleurstellende resultaat komt overeen met andere studies die concludeerden dat intensieve (eHealth) trainingsregimes na een CVA moeilijk zijn vol te houden voor patiënten [29-32]. Tijdens de uitvoering van de training in onze studie ondervonden patiënten verschillende problemen, zoals te weinig ICT-vaardigheden, een ongeschikt besturingssysteem op de computer of overprikkeling door flikkeringen op het beeldscherm.

Dergelijke belemmeringen werden ook gevonden in een andere studie, waarin CVA-patiënten diverse problemen ervoeren bij gebruik van een applicatie op een tablet, o.a. door de complexiteit van instructies, het gebrek aan fijn motorische vaardigheden voor het gebruik en de onbetrouwbaarheid van internetverbindingen [33]. Om dergelijke belemmeringen zoveel mogelijk weg te nemen is adequate ondersteuning van belang. Uit een studie in dit proefschrift bleek dat hoogfrequente supervisie (wekelijks) resulteerde in significant hogere therapietrouw dan laagfrequente supervisie (4-wekelijks). Dit komt overeen met de bevindingen in een review van Kelders et al. (2012), waarin de frequentie van interactie met een behandelaar een significante voorspeller was voor therapietrouw met online gezondheidsinterventies in verschillende patiëntengroepen [34]. Samenvattend lijkt regelmatige interactie met een behandelaar belangrijk om het gebruik van eRevalidatie bij CVA-patiënten te vergroten. In ons focusgroep onderzoek naar vereisten voor eRevalidatie erkenden de deelnemende zorgprofessionals (revalidatieartsen, fysiotherapeuten, ergotherapeuten, psychologen, teamcoördinatoren en logopedisten) het belang van hun rol bij het introduceren en superviseren van eRevalidatie in de behandeling van CVA.

Onderzoeksstrategieën voor het evalueren van eRevalidatie na een CVA

De effecten van eRevalidatie interventies worden vaak geëvalueerd door middel van traditionele klinische studies. Een voorbeeld hiervan is de RCT die in dit proefschrift is opgenomen. Dergelijke onderzoekdesigns zijn echter niet altijd geschikt om de effecten van eHealth te evalueren [35-38]. Deelnemers aan een RCT vormen bijvoorbeeld niet altijd een goede afspiegeling van de patiëntenpopulatie in de dagelijkse praktijk, o.a. door het hanteren van specifieke selectiecriteria [38]. Wanneer de effectiviteit van eHealth is vastgesteld in een 'laboratoriumomgeving', worden dezelfde positieve resultaten dan ook niet altijd gevonden in een andere context, bijvoorbeeld wanneer de interventie wordt aangeboden aan patiënten met minder ICT-vaardigheden of minder goed getrainde professionals.

Een ander nadeel van de conventionele RCT is, dat de tijd die het duurt om deze voor te bereiden, uit te voeren en te rapporteren meestal veel langer is dan de snelheid waarmee eHealth zich ontwikkelt. Dit kan ervoor zorgen dat (delen van) de onderzochte interventie al verouderd zijn tegen de tijd dat de resultaten van de RCT worden gepubliceerd [39]. Bovendien kunnen 'early adopters' onder patiënten en zorgprofessionals een eHealth innovatie al gebruiken voordat er bewijs beschikbaar is [38], waardoor in sommige gevallen het contrast tussen interventie- en controlecondities kan vervagen. Om deze redenen kunnen andere onderzoekstrategieën worden overwogen, zoals actie-onderzoek. Deze vorm van onderzoek, vaak gecoördineerd door een groep professionals als onderdeel van een zogenaamde 'community of practice', is ontstaan om een actueel probleem in de zorgpraktijk op te lossen [41]. Hierin participeren patiënten, zorgprofessionals, ontwerpers,

onderzoekers, docenten en studenten op een actieve manier. Na onderzoek van het probleem neemt de groep beslissingen, observeert en evalueert de gevolgen van de aanpassing voor die specifieke situatie. De groep neemt dus deel aan een veranderproces, waarbij naast het continu aanpassen, testen en evalueren tegelijkertijd onderzoek naar de innovatie wordt uitgevoerd.

Tot slot moet worden opgemerkt dat de ontwikkeling, evaluatie en implementatie van eRevalidatie na een CVA en bijbehorend onderzoek moeten worden uitgevoerd in samenwerking met alle relevante betrokkenen om eRevalidatie relevant en haalbaar te maken voor de beoogde gebruikers in de gezondheidszorg [42;43]. Dit zijn, naast patiënten, mantelzorgers, zorgprofessionals, ontwerpers, onderzoekers, financiers, docenten en studenten.

Deel II. Gereedheid en vereisten voor eRevalidatie na een CVA: implicaties voor de opname in de gezondheidszorgpraktijk en het onderwijs.

Integreren van gebruikersperspectieven bij de ontwikkeling van eRevalidatie

Dit proefschrift bevat een aantal onderzoeken die gericht zijn op het bepalen van het gebruik van en vereisten voor eRevalidatie onder betrokkenen in de zorg na een CVA. Uit een enquêteonderzoek bleek dat het bezit en gebruik van ICT-apparaten relatief hoog was bij revaliderende patiënten en dat de meeste patiënten hun eigen apparatuur ook ten behoeve van hun revalidatieproces wilden gebruiken. Bovendien toonde een focusgroepstudie aan dat CVA-patiënten, mantelzorgers en behandelaars specifieke wensen hadden ten aanzien van de gewenste functionaliteiten, zoals ondersteuning van de uitvoering van fysieke oefeningen, informatie over CVA, uitkomsten van de revalidatie en het plannen van afspraken met behandelaars. Deze bevindingen suggereren dat bij voorkeur interventies moeten worden ontwikkeld waarin uiteenlopende functionaliteiten (bijv. communicatiefunctie, fysieke oefenen, informatie over zorg en aandoening, agendafunctie, enz.) worden gecombineerd in één digitaal platform. Mede op basis van de uitkomsten van dit kwalitatief onderzoek werd een uitgebreid eRevalidatie platform gebouwd gericht op het herstel na een beroerte. Dit platform wordt momenteel geëvalueerd in het Fit After Stroke @Home project (FAST@Home).

Daarnaast kwam uit onze focusgroep studie onder gebruikers in de revalidatie en uit literatuur naar voren dat digitale trainingsfaciliteiten, zoals oefeningen gericht op fysiek en cognitief functioneren en spraak, moeten worden aangepast aan de voorkeuren en mogelijkheden van individuele patiënten [44-50]. Ook het bieden van ondersteuning bij dagelijks gebruik van eRevalidatie (d.w.z. directe hulp, een helpdesk, video's met instructies en/of een menu met veel gestelde vragen) werd als een cruciaal element beschouwd.

Deze bevindingen impliceren dat het ontwerpen van eRevalidatie programma's in co-creatie met alle relevante betrokkenen moet worden gedaan ('co-design'). Door vereisten van patiënten, mantelzorgers en behandelaars mee te nemen in elke stap van het ontwerpproces ('user-centered design') sluit eRevalidatie beter aan bij de behoeften van de beoogde gebruikers. Dit is vooral belangrijk omdat dit proefschrift heeft aangetoond dat verschillende gebruikersgroepen verschillende eisen hebben. Een manier om alle partijen bij elkaar te brengen is samenwerking in een zogenaamd "Living Lab". Hierin werken verschillende belanghebbenden (bijv. kennisinstellingen, gezondheidszorgorganisaties, zorgverleners, patiënten, financiers, innovators, gevestigde bedrijven, startups, onderzoekers, enz.) samen. In een medisch specialistisch revalidatiecentrum, Basalt revalidatie, is een dergelijk Living Lab (SmartLab) opgericht. Hier wordt een stapsgewijze procedure gebruikt om innovaties in revalidatie te testen op hun potentiële toegevoegde waarde en bruikbaarheid (<http://www.basaltrevalidatie.nl/onderzoek-innovatie/smartlab>, www.medicaldeltalivinglab.nl). De vervolgstappen omvatten systematisch evaluaties van de effectiviteit, ervaringen, kosten en implementatie van eHealth in de revalidatiebehandeling.

Gebruik van eHealth in de revalidatiezorg na een CVA

De impact van een CVA varieert sterk per persoon, en welke behandeling optimaal is hangt af van de mogelijkheden, voorkeuren en doelen van zowel de patiënt als zijn/haar naaste(n) [51]. Zorgprofessionals kunnen een centrale rol spelen in het op de patiënt afgestemde aanbod van eRevalidatie, bijvoorbeeld door aanpassing van het oefenprogramma op basis van vooruitgang, feedback en motivatie. Daarnaast kunnen ze bepalen welke bewezen effectieve eRevalidatie interventies geschikt zijn voor een patiënt en begeleiding bieden bij het gebruik.

Behandelaars zijn zich echter niet altijd bewust van alle mogelijkheden die eHealth interventies bieden en hoe en wanneer zij het moeten inzetten [37;52]. Resultaten van onderzoek naar eHealth zijn vaak niet direct bruikbaar voor zorgverleners in de dagelijkse praktijk [37;38]. Zo is uit onderzoek gebleken dat virtual reality voor het verbeteren van het evenwicht en looppatroon als therapeutisch hulpmiddel kan dienen om herstel van balans en lopen bij patiënten met een CVA te bevorderen [2;52]. Deze bevindingen hebben echter hun weg nog niet gevonden naar de praktijk. Daarvoor is ten eerste een overzicht van bruikbare eHealth interventies voor herstel na een CVA nodig, gebaseerd op resultaten van wetenschappelijk onderzoek, ervaringen van gebruikers, patiëntenverenigingen en beroepsorganisaties [37]. Daarnaast moet de beschikbare kennis worden verwerkt in protocollen, praktijk richtlijnen of andere handvatten die inzicht geven in welk type eRevalidatie werkt voor wie [37;52;53]. Om deze doelen te bereiken is het eerdergenoemde SmartLab verbonden met het National eHealth Living Lab (NeLL), een eHealth-community (patiënten, consumenten, professionals, wetenschappers, studenten, organisaties) die

streeft naar het creëren van de beste eHealth oplossingen door kennis, contacten en ervaringen met elkaar te delen (<https://nell.eu>). Bovendien toonde dit proefschrift aan dat voor een optimale toepasbaarheid van eHealth interventies deze in werkroutines van zorgverleners moet kunnen worden geïntegreerd.

eHealth in het onderwijs van zorgopleidingen

Toekomstige zorgverleners moeten vakkundig en vol vertrouwen kunnen werken met eHealth [54;55]. eHealth komt echter maar mondjesmaat aan bod in de curricula van het hoger beroepsonderwijs die opleiden tot zorgberoepen (bijv. voedingsleer, verpleegkunde, fysiotherapie, psychologie of maatschappelijk werk) [56-59]. In dit proefschrift is een focusgroep studie beschreven waarin docenten en studenten aangaven dat zij een betere integratie van eHealth in het curriculum belangrijk vonden, en zeker bereid waren om eHealth te gebruiken. Desalniettemin was, in overeenstemming met andere studies in het veld, hun begrip van het concept eHealth beperkt [60]. Bovendien hadden docenten en studenten behoefte aan vaardigheden om eHealth interventies kritisch te kunnen beoordelen. Om de implementatie van eHealth in het curriculum te verbeteren wordt in het FAST@Home project momenteel de ‘toolkit eHealth’ ontwikkeld, in een samenwerkingsverband van drie hogescholen (Haagse Hogeschool, Hogeschool Leiden, Hogeschool van Amsterdam).

Een bevorderende factor voor het gebruik van eHealth in het onderwijs, die ook naar voren kwam in de focusgroepen, was het optimaal benutten van de al eerder beschreven zogenaamde ‘communities of practices’ [61]. Deelname aan een dergelijke praktijk(leer) gemeenschap stimuleert en ondersteunt zowel docenten als studenten in het onderwijzen van en leren over eHealth [62]. Daarnaast suggereert de literatuur dat het uitproberen van eHealth innovaties in een realistische context met actieve betrokkenheid van gebruikers (LivingLabs) stimulerende leeromgevingen zijn [63]. Het is dus raadzaam studenten en docenten hierin te betrekken. Daarnaast is het belangrijk dat ook artsen beter met eHealth om kunnen gaan. Opgedane kennis in het hoger onderwijs is mogelijk ook toepasbaar in het geneeskunde curriculum.

Tot slot moet eHealth niet alleen in het curriculum van de bacheloropleidingen een belangrijke plaats krijgen, maar ook in het postacademisch onderwijs, aansluitend op de visie ‘leven lang leren’ en met het doel het huidige gebruik in de dagelijkse zorgpraktijk te vergemakkelijken [37].

Conclusie

eRevalidatie biedt veel mogelijkheden om het herstel na een beroerte te bevorderen. Desondanks valt het gebruik van eHealth in de revalidatiepraktijk nog altijd tegen. Dit komt onder andere door gebrek aan bewijs voor de effectiviteit ervan. Het aantal methodologisch goed uitgevoerde studies is schaars, en de resultaten wisselend. Mogelijk zijn traditionele onderzoekdesigns niet altijd even geschikt om snel ontwikkelende interventies te evalueren. Alternatieve onderzoeksstrategieën, bijvoorbeeld actieonderzoek, zijn mogelijk passender om de effectiviteit van eHealth innovaties inzichtelijk te maken. Om eHealth interventies haalbaar te maken voor de dagelijkse zorgpraktijk is het van belang dat de ontwikkeling ervan plaatsvindt in samenwerking met alle betrokkenen ('co-creatie') en op basis van de wensen van de gebruikers. Voor de implementatie van eHealth in het zorgproces is het daarnaast essentieel om patiënten en behandelaars te ondersteunen bij gebruik en ICT een prominenter plek te geven in het curriculum van zorgopleidingen.

Publications

Voorn V, Marang-van de Mheen P, **Wentink M**, So-Osman C, Vlieland T, Koopman-van Gemert A, Nelissen R, Van Bodegom-Vos L. Frequent use of blood-saving measures in elective orthopaedic surgery: a 2012 Dutch blood management survey. *BMC musculoskeletal disorders*. 2013 Dec;14(1):230.

Hofstede S, Marang-van de Mheen P, **Wentink M**, Stiggelbout A, Vleggeert-Lankamp C, Vlieland T, van Bodegom-Vos L. Barriers and facilitators to implement shared decision making in multidisciplinary sciatica care: a qualitative study. *Implementation Science*. 2013 Dec;8(1):95.

Hofstede S, van Bodegom-Vos L, **Wentink M**, Vleggeert-Lankamp C, Vlieland T, Marang-van de Mheen P, DISC study group. Most important factors for the implementation of shared decision making in sciatica care: ranking among professionals and patients. *PloS one*. 2014 Apr 7;9(4):e94176.

Voorn V, Marang-van de Mheen P, **Wentink M**, Kaptein A, Koopman-van Gemert A, So-Osman C, Vliet Vlieland T, Nelissen R, van Bodegom-Vos L, LISBOA Study Group. Perceived barriers among physicians for stopping non-cost-effective blood-saving measures in total hip and total knee arthroplasties. *Transfusion*. 2014 Oct;54(10pt2):2598-607.

Wentink M, Berger M, de Kloet A, Meesters J, Band G, Wolterbeek R, Goossens P, Vliet Vlieland T. The effects of an 8-week computer-based brain training programme on cognitive functioning, QoL and self-efficacy after stroke. *Neuropsychological rehabilitation*. 2016 Sep 2;26(5-6):847-65.

Wentink M, Prieto E, de Kloet A, Vliet Vlieland T, Meesters J. The patient perspective on the use of information and communication technologies and e-health in rehabilitation. *Disability and Rehabilitation: Assistive Technology*. 2018 Oct 3;13(7):620-5.

Wentink M, Meesters J, Berger M, de Kloet A, Stevens E, Band G, Kromme C, Wolterbeek R, Goossens P, Vliet Vlieland T. Adherence of stroke patients with an online brain training program: the role of health professionals' support. *Topics in stroke rehabilitation*. 2018 Jul 4;25(5):359-65.

Wentink M, van Bodegom-Vos L, Brouns B, Arwert H, Vliet Vlieland T, de Kloet A, Meesters J. What is important in e-health interventions for stroke rehabilitation? A survey study among patients, informal caregivers and health professionals. *International journal of telerehabilitation*. 2018;10(1):15.

Brouns B, Meesters J, **Wentink M**, de Kloet A, Arwert H, Vliet Vlieland T, Boyce L, van Bodegom-Vos L. Why the uptake of eRehabilitation programs in stroke care is so difficult - a focus group study in the Netherlands. *Implementation Science*. 2018;13(1):133.

Wentink M, van Bodegom-Vos L, Brouns B, Arwert H, Houdijk S, Kewalbansing P, Boyce L, Vliet Vlieland T, de Kloet A, Meesters, J. How to improve eRehabilitation programs in stroke care? A focus group study to identify requirements of end-users. *BMC medical informatics and decision making*. 2019;19(1),145.

Curriculum Vitae

Manon Maria Wentink werd geboren op 10 april 1987 in de gemeente Noorder-Koggenland en is opgegroeid in Midwoud. In 2005 voltooide zij het Hoger Algemeen Voortgezet Onderwijs aan de Openbare Scholengemeenschap in Hoorn. In datzelfde jaar startte zij met de opleiding Oefentherapie Mensendieck aan de Hogeschool van Amsterdam. In 2009 studeerde zij af als oefentherapeut. In 2011 behaalde Manon haar Master of Health Sciences aan de Vrije Universiteit van Amsterdam.

Zorgpraktijk

Na haar studie oefentherapie begon Manon als waarnemend oefentherapeut in diverse settingen. Van 2010 tot 2017 was zij, samen met een oud-studiegenoot, eigenaar van de praktijk voor oefentherapie Mensendieck in Haarlem. Als praktiserend oefentherapeut ontwikkelde zij een wetenschappelijk onderbouwde interventie voor mensen met slapeloosheid. In 2012 richtte zij de onderneming SlaapSlim op en vanaf 2013 tot heden geeft zij geaccrediteerde scholingen aan zorgprofessionals over (de behandeling van) slaapproblemen.

Onderzoek

In 2012 begon zij, naast haar werkzaamheden als oefentherapeut, als onderzoeksassistent bij de afdeling Kwaliteit van Zorg in het Leids Universitair Medisch Centrum (LUMC). Daar werkte zij mee aan de Lisboa studie en de DISC studie. In 2013 startte Manon onder begeleiding van prof. Dr. Thea Vliet Vlieland als promovenda in onderzoeksproject Spelenderwijs, een samenwerking tussen Basalt (destijds Sophia Revalidatie en het Rijnlands Revalidatie Centrum), het LUMC en de Haagse Hogeschool. Later werkte zij mee aan meerdere onderzoeken waaronder het project Fast@Home, waarvan een deel van de resultaten beschreven staan in dit proefschrift.

Onderwijs

Vanaf 2014 tot heden is Manon werkzaam als docentonderzoeker voor de opleiding Oefentherapie Mensendieck aan de faculteit Gezondheid van de Hogeschool van Amsterdam. Sinds 2017 vervult zij hier ook de functie als Hoofddocent Onderzoek met als doel de verbinding tussen onderwijs en onderzoek te versterken.

Dankwoord

Het realiseren van een proefschrift heeft veel overeenkomsten met de ontwikkeling van eHealth: het beste resultaat behaal je door co-creatie! Verschillende personen hebben een bijdrage geleverd aan dit proefschrift. Deze personen ben ik dan ook bijzonder dankbaar.

Speciale dank gaat uit naar alle patiënten en diens mantelzorgers, zorgverleners, studenten en docenten, die dankzij de inzet van Basalt, de Hogeschool Leiden, de Hogeschool van Amsterdam en het LUMC aan de onderzoeken hebben meegedaan.

Mijn promotor, Thea Vliet Vlieland, en co-promotoren Arend de Kloet en Jorit Meesters, zonder wie ik dit proefschrift niet had kunnen realiseren, ben ik veel dank verschuldigd. Thea, zonder jouw deskundigheid was ik kansloos. Jorit, bedankt voor jouw opbouwende feedback, humor en geduld. Arend, door jou ben ik 'spelenderwijs' het onderzoek ingerold. Bedankt voor het vertrouwen dat je destijds in mij had en de kans die je me hebt gegeven om te promoveren.

Monique Berger, voor al je inzet voor 'Spelenderwijs', opgewerkte humeur en onuitputtelijke energie. Leti Bodegom Vos, voor alle expertise op het gebied van kwalitatief onderzoek en de snelle antwoorden bij brandende vragen. Ook bedankt dat in onder jouw leiding de eerste stappen op het gebied van wetenschappelijk onderzoek heb kunnen zetten.

Alle collega's van het kenniscentrum bij Basalt revalidatie en het LUMC. Datamanager Cedric Kromme voor het mogelijk maken van wat het soms leek onmogelijke: het ordenen van de data. Statisticus Ron Wolterbeek voor de mini-colleges statistiek en kunstwerken van syntaxen. Ilonca Kraak, management assistent en moeder overste van het kenniscentrum bij Basalt, en Anika Rabelink-Hoogstraaten van het LUMC, voor jullie eerste hulp bij administratieve zaken. Inge Verhoeven, Klaasjan van Haastrecht en Monique Jakobs voor jullie ondersteuning bij de uitvoer van 'Spelenderwijs'. Sander Houdijk, Liesbeth Boyce, Hanneke Braakhuis en Preschana Kewalbansing bij de uitvoer van Fast@Home. Het was een groot plezier met een ieder van jullie te mogen werken!

Alle co-auteurs en consortiumpartners. Dank aan John Verhoef en Petra Siemonsma in het bijzonder voor de bijdrage die zij hebben geleverd aan de focusgroep studie in het onderwijs.

Mijn lieve collega's van de Hogeschool van Amsterdam voor de oprechte belangstelling in de voortgang. Berber en Hanneke, voor jullie collegiale support. Berber, ik ben je uiteraard ook bijzonder dankbaar dat je de rol als paranimf op je wilde nemen!

Ciska, Sarah en Willemijn voor de verhelderende coaching, gezelligheid en essentiële ontspanning tussendoor. Onmisbaar! Niet alleen tijdens een promotietraject, maar ook daarna. Evelien, maatje, voor al je begrip en steun! Dat ik onderzoek wilde doen naast onze ondernemingen had ook directe gevolgen voor jou. Tanne, voor jouw luisterend oor, meeleven en humor. Ook bedankt dat je zo nadrukkelijk de rol als paranimf op je wilde nemen. Zitten we na al die jaren toch nog een keer samen in de 'schoolbanken'!

Opa, voor al uw wijsheden, in het bijzonder de leus "dom door doen!". Het heeft mij geholpen om de losse eindjes aan elkaar te knopen op die momenten dat de motivatie ontbrak. Papa en mama, voor het van jongs af aan stimuleren van mijn onderzoekend vermogen. Zonder jullie ongedwongen stimulans mijzelf te ontwikkelen en ambities waar te maken was dit niet gelukt. Lisette en Anouk, mijn geweldige zussen, voor jullie onvoorwaardelijke steun op de cruciale momenten. De mogelijkheid om dit proces met jullie te kunnen delen was essentieel om de eindstreep te halen.

Last but not least Pieter, voor alle fijne afleiding en benodigde stabiliteit. Ik kan je niet genoeg bedanken voor je eindeloze geduld, steun en begrip.

