

# **The Formation Conditions of the Wide Binary Class 0 Protostars within BHR 71**

Tobin, J.J.; Bourke, T.L.; Mader, S.; Kristensen, L.; Arce, H.; Gueth, F.; ... ; Chen, X.

## **Citation**

Tobin, J. J., Bourke, T. L., Mader, S., Kristensen, L., Arce, H., Gueth, F., … Chen, X. (2019). The Formation Conditions of the Wide Binary Class 0 Protostars within BHR 71. *Astrophysical Journal (0004-637X)*, *870*(2), 81. doi:10.3847/1538-4357/aaef87

Version: Accepted Manuscript License: [Leiden University Non-exclusive license](https://hdl.handle.net/1887/license:3) Downloaded from: <https://hdl.handle.net/1887/85104>

**Note:** To cite this publication please use the final published version (if applicable).

The Formation Conditions of the Wide Binary Class 0 Protostars within BHR 71

JOHN J. TOBIN.<sup>1, 2, 3</sup> TYLER L. BOURKE.<sup>4</sup> STACY MADER.<sup>5</sup> LARS KRISTENSEN.<sup>6</sup> HECTOR ARCE.<sup>7</sup> FRÉDÉRIC GUETH, <sup>8</sup> ANTOINE GUSDORF, <sup>9</sup> CLAUDIO CODELLA, <sup>10, 11</sup> SILVIA LEURINI, <sup>12</sup> AND  $\rm XUEPENG$   $\rm CHEN^{13}$ 

<sup>1</sup>Current Address: National Radio Astronomy Observatory, 520 Edgemont Rd., Charlottesville, VA 22903, USA; jtobin@nrao.edu

 $2$ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 W. Brooks Street, Norman, OK 73019, USA

<sup>3</sup>Leiden Observatory, Leiden University, P.O. Box 9513, 2300-RA Leiden, The Netherlands

<sup>4</sup>SKA Organization, Jodrell Bank Observatory, Lower Withington, Macclesfield, Cheshire SK11 9DL, UK

<sup>5</sup>CSIRO Astronomy and Space Sciences, Parkes Observatory, PO BOX 276, Parkes NSW 2870, Australia

<sup>6</sup>Centre for Star and Planet Formation, Niels Bohr Institute and Natural History Museum of Denmark, Copenhagen University, Øster Voldgade 5–7, DK-1350 Copenhagen K, Denmark

<sup>7</sup>Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520-8101, USA

 $8$ Institut de Radioastronomie Millimétrique (IRAM), 38406 Saint-Martin dHères, France

 $^{9}$ LERMA, Observatoire de Paris, École normale supérieure, PSL Research University, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, F-75231, Paris, France

<sup>10</sup>INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

 $11$ Univ. Grenoble Alpes, Institut de Planétologie et dAstrophysique de Grenoble (IPAG), 38401 Grenoble, France

<sup>12</sup>INAF - Osservatorio Astronomico di Cagliari, Via della Scienza 5, I-09047 Selargius (CA), Italy

<sup>13</sup> Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210034, China

#### ABSTRACT

We present a characterization of the binary protostar system that is forming within a dense core in the isolated dark cloud BHR71. The pair of protostars, IRS1 and IRS2, are both in the Class 0 phase, determined from observations that resolve the sources from 1  $\mu$ m out to 250  $\mu$ m and from 1.3 mm to 1.3 cm. The resolved observations enable the luminosities of IRS1 and IRS2 to be independently measured (14.7 and 1.7  $L_{\odot}$ , respectively), in addition to the bolometric temperatures 68 K, and 38 K, respectively. The surrounding core was mapped in  $NH<sub>3</sub>(1,1)$  with the Parkes radio telescope, and followed with higher-resolution observations from ATCA in  $NH<sub>3</sub>$  (1,1) and 1.3 cm continuum. The protostars were then further characterized with ALMA observations in the 1.3 mm continuum along with  $N_2D^+$  ( $J = 3 \rightarrow 2$ ), <sup>12</sup>CO, <sup>13</sup>CO, and  $C^{18}O$  ( $J = 2 \rightarrow 1$ ) molecular lines. The Parkes observations find evidence for a velocity gradient across the core surrounding the two protostars, while ATCA reveals more complex velocity structure toward the protostars within the large-scale gradient. The ALMA observations then reveal that the two protostars are at the same velocity in  $C^{18}O$ , and  $N_2D^+$  exhibits a similar velocity structure as NH<sub>3</sub>. However, the  $C^{18}O$ kinematics reveal that the rotation on scales <1000 AU around IRS1 and IRS2 are in opposite directions. Taken with the lack of a systematic velocity difference between the pair, it is unlikely that their formation resulted from rotational fragmentation. We

instead conclude that the binary system most likely formed via turbulent fragmentation of the core.

## 1. INTRODUCTION

Binary and multiple star systems are a frequent outcome of the star formation process at both low and high masses (e.g., [Raghavan et al.](#page-27-0) [2010;](#page-27-0) Duchêne & Kraus [2013\)](#page-27-1). While the statistics of multiple star formation in the main-sequence, pre-main sequence phase, and even the protostellar phase are becoming much better characterized [\(Kraus et al.](#page-27-2) [2011;](#page-27-2) [Raghavan et al.](#page-27-0) [2010;](#page-27-0) [Ward-Duong](#page-28-0) [et al.](#page-28-0) [2015;](#page-28-0) [Chen et al.](#page-26-0) [2013;](#page-26-0) [Tobin et al.](#page-28-1) [2016\)](#page-28-1), the physical processes that lead to multiple star formation are not well-characterized observationally. The main epoch of multiple star formation is the protostellar phase [\(Tohline](#page-28-2) [2002;](#page-28-2) [Chen et al.](#page-26-0) [2013\)](#page-26-0), when the protostar(s) are surrounded by a dense infalling cloud(s) of gas and dust. Thus, it is imperative to examine the process of multiple star formation in the protostellar phase with a combination of continuum and molecular line tracers to reveal the physical processes at work.

While it is essential to examine multiple star formation in the protostellar phase, star formation typically occurs in clusters (e.g., [Lada & Lada](#page-27-3) [2003;](#page-27-3) [Megeath et al.](#page-27-4) [2012\)](#page-27-4) increasing the complexity of the data, especially for spectral line tracers that reveal the motion of gas along the line of sight. Toward small proto-clusters in molecular clouds (e.g., NGC 1333), complex motions of the surrounding gas can complicate interpretation of kinematic data, making multiple star formation difficulty to characterize. Thus, stars forming in nearby isolated clouds, the so-called Bok Globules, can be valuable laboratories for the study of multiple star formation. This is especially true for wide multiple star systems, where separations are greater than 1000 AU, because the kinematics should be less confused along the line of sight. The BHR71 system [\(Bourke et al.](#page-26-1) [1995a;](#page-26-1) [Bourke](#page-26-2) [2001;](#page-26-2) [Chen](#page-26-3) [et al.](#page-26-3) [2008;](#page-26-3) [Yang et al.](#page-28-3) [2017\)](#page-28-3) is composed of two Class 0 protostars (André et al. [1993\)](#page-26-4) separated by  $\sim$ 16" (3200 AU; 200 pc distance), and it is an ideal system to study the physics of wide multiple star formation. The fact that both components of the binary system are in the Class 0 phase means that the protostars are very young, likely less than ∼150 kyr in age [\(Dunham et al.](#page-27-5) [2014\)](#page-27-5). Therefore, the initial conditions of their formation are not likely to have been erased by the dynamical interactions of the protostars and their effects on the surrounding cloud from their outflows [\(Arce & Sargent](#page-26-5) [2006\)](#page-26-5).

The formation of wide multiple star systems has been thought to be related to the rotation of the infalling envelope of the protostar, where greater rotation rates would lead to an increased likelihood of fragmentation. Many theoretical and numerical studies have highlighted the importance of rotation in the formation of wide and close multiple star systems [\(Larson](#page-27-6) [1972;](#page-27-6) [Burkert & Bodenheimer](#page-26-6) [1993;](#page-26-6) [Boss](#page-26-7) [1995,](#page-26-7) [2002;](#page-26-8) [Boss & Keiser](#page-26-9) [2013\)](#page-26-9). The importance of rotation is typically parametrized as  $\beta_{\rm rot}$  which is the ratio of rotational energy to gravitational potential energy. [Chen et al.](#page-26-10) [\(2012\)](#page-26-10) compiled a list of all known single and multiple protostars and calculated their  $\beta_{\rm rot}$  values, finding that most binary/multiple protostars have  $\beta_{\rm rot} > 0.01$ . However,  $\beta_{\rm rot}$  is observationally derived from measurement of velocity gradients in the cores/envelopes surrounding the protostar(s), and [Tobin et al.](#page-28-4) [\(2011,](#page-28-4) [2012a\)](#page-28-5) highlighted the possibility that these velocity gradients may not truly reflect rotation due to the possibility of large scale asymmetries in the envelope structures [\(Tobin](#page-28-6) [et al.](#page-28-6) [2010b\)](#page-28-6) and filamentary structure (e.g., André et al. [2010;](#page-26-11) [Hacar & Tafalla](#page-27-7) [2011\)](#page-27-7). Furthermore, numerical simulations of star formation in turbulent molecular clouds have shown that the turbulence can lead to the formation of multiple star systems on ∼1000 AU scales [\(Padoan & Nordlund](#page-27-8) [2002;](#page-27-8)

[Offner et al.](#page-27-9) [2010\)](#page-27-9), possibly removing the need for bulk envelope rotation to initiate multiple star formation. Therefore sensitive molecular line data tracing the gas kinematics are required, spanning orders of magnitude in spatial scales to differentiate between different scenarios of multiple star formation.

We have obtained a comprehensive dataset on BHR71 spanning the scales of the star forming core at ~1' (12000 AU) resolution down to 1".5 (300 AU) to examine the full range of scales within this isolated binary protostar system. With this dataset, we aim to determine what physical process(es) lead to the formation of at least two protostars in this system. The results will be useful for interpreting observations of more evolved multiple star systems and other forming multiple systems. The dataset includes Herschel photometry, Australia Telescope Compact Array (ATCA) continuum and molecular line observations, Parkes Radio Telescope NH<sup>3</sup> line mapping, and Atacama Large Millimeter/submillimeter Array (ALMA) observations of continuum and molecular lines. The paper is organized as follows: the observations are presented in Section 2, the observational results regarding classification, photometry, and continuum are presented in Section 3, the results from the and molecular line data are presented in Section 4, the results are discussed in Section 5, and our conclusions are presented in Section 6.

#### 2. OBSERVATIONS

#### 2.1. Parkes Radio Telescope

We observed BHR71 with the Parkes 64m radio telescope on 2012 September 06 (project P825). The 13mm receiver was used and we observed an on-the-fly (OTF) map of BHR71 covering an  $8' \times 8'$ area surrounding BHR71 in the  $NH<sub>3</sub> (1,1)$  transition using dual polarization. The usable surface of the Parkes dish is 55m at 1.3 cm, resulting in a beam size of  $~\sim 60''$ , see Table 1 for a summary. Weather was good and  $T_{sys}$  was  $\sim 100$  K. The observations were conducted shortly after the 13mm receiver was mounted and an all-sky pointing model could not be completed prior to our observations and we instead used a local pointing solution on the nearby quasar 1057-797.

The flux calibration of the data was done using the quasar 1057-797, which was the gain calibrator for the ATCA observations that were conducted a few days prior (see below). The data were reduced using the *livedata* and *gridzilla* tasks within  $AIPS++$ . The *livedata* task was used to perform baseline subtraction and scaling to the proper flux density of each spectrum in the map, while *gridzilla* was used to construct a datacube from the individual spectra in FITS format. After mapping the data, we noticed that there was a systematic offset in the central coordinates of the map; BHR71 was located  $\sim$ 1' north of the pointing center. The systematic offset likely resulted from the lack of an all-sky pointing solution prior to the observations. To recenter the map, we smoothed the ATCA data to the same resolution as the Parkes data and matched peak emission of the core. Given that some emission is resolved-out in the ATCA observations, there is likely some residual pointing uncertainty of the map on the order of  $15''$  to  $20''$ .

The FITS cube was then ingested into CLASS, which is part of the GILDAS<sup>[1](#page-3-0)</sup> software package using the CLASS function  $lmv$ . We then used the built-in  $NH<sub>3</sub>$  line fitting functions to construct the centroid velocity maps and the linewidth maps.

#### 2.2. ATCA Observations

<span id="page-3-0"></span><sup>1</sup> http://www.iram.fr/IRAMFR/GILDAS

BHR71 was observed with ATCA during 2012 in the EW367 configuration and H214 configuration, project C2665. ATCA consists of 6 antennas that are 22 meters in diameter. One antenna is fixed on a pad 6 km from the array center, the remaining 5 antennas are reconfigurable on an east-west track and a shorter north-south track. In both observations, we targeted the  $NH_3$  (1,1), (2,2), and (3,3) inversion transitions at 23.694, 23.722, and 23.870 GHz. The primary beam of the 22 meter antennas at these frequencies is  $\sim$ 144". The observations are summarized in Table 1.

The Compact Array Broadband Backend (CABB) correlator was used for our observations, providing 2 GHz of continuum bandwidth (with 2048 channels each 1 MHz wide) and we placed 1 MHz zoom bands, each with 2048 channels  $(0.5 \text{ kHz channels})$  at the frequencies of the NH<sub>3</sub>  $(1,1)$ ,  $(2,2)$ , and  $(3,3)$  inversion transitions. We used 6 zoom bands for each of NH<sub>3</sub>  $(1,1)$  and  $(2,2)$ , and 4 zoom bands for  $NH_3$  (3,3). We used 1057-797 as the complex gain calibrator, 0537-441 as the bandpass calibrator, and 1934-638 was the flux calibrator for both the H214 and EW367 observations.

We observed BHR71 on 2012 April 22 in the EW367 configuration, where the maximum baseline is 367 meters with all antennas positioned along the east-west track; the shortest baseline was ∼38 meters. The weather conditions were variable during the observation with  $T_{sys}$  typically ∼100 K. The observations were conducted for a full 12 hour synthesis track. However, there was rain during the last two hours of the observations and data from that time period were unusable. The observations were conducted as a 7-point mosaic where the goal was to obtain a constant surface brightness sensitivity over the central  $\sim$ 144'' primary beam. However, due to an error in the mosaic setup, the mosaic was not as wide in the right ascension direction as intended. This did not adversely affect the sensitivity of our map because the  $NH<sub>3</sub>$  emission from BHR71 was adequately mapped with this mosaic pattern.

BHR71 was also observed on 2012 September 24 with ATCA in the H214 configuration where 5 antennas were positioned along the north-south track with a shortest baseline length of  $\sim$ 47 meters. The observing track length was 8 hours and  $T_{sys}$  was ∼80 K throughout the observations. The weather conditions were good and the phase between antennas was stable during the observations. These observations were also conducted in a 7-point mosaic, with uniform coverage of the central  $\sim$ 144".

The data were calibrated and edited using the Australian Telescope National Facility (ATNF) version of the MIRIAD software package [\(Sault et al.](#page-28-7) [1995\)](#page-28-7). The phases and amplitudes for each baseline were inspected and edited to remove periods of high phase noise and/or amplitude deviations. We also flagged the single antenna on the 6 km baseline because of high phase noise and lack of uvcoverage at intermediate baselines.

The combined EW367 and H214 datasets were imaged using the *clean* algorithm implemented in MIRIAD. We generated both a 1.3 cm continuum image of BHR71 and channel maps centered on the NH<sub>3</sub>  $(1,1)$ ,  $(2,2)$ , and  $(3,3)$  transitions; however, only the NH<sub>3</sub>  $(1,1)$  line was well-detected. We did not combine the ATCA and Parkes data due to the positional uncertainty in the Parkes data, we instead use them independently for the spatial scales they are sensitive to.

### 2.3. ALMA Observations

BHR71 was observed by the ALMA 12m array, the ALMA Compact Array (ACA), and the Total Power Array (TPA) under the project 2013.1.00518.S. The observations were conducted in band 6, with a central tuning frequency of 225 GHz. In all observations, the correlator was configured to observe <sup>12</sup>CO  $(J = 2 \rightarrow 1)$ , <sup>13</sup>CO  $(J = 2 \rightarrow 1)$ , C<sup>18</sup>O  $(J = 2 \rightarrow 1)$ , N<sub>2</sub>D<sup>+</sup>  $(J = 3 \rightarrow 2)$ , and continuum. The continuum basebands were located at 232.5 GHz, a region without many significant

spectral lines, and 218 GHz. The 218 GHz band had significant contamination from spectral lines and was not used for continuum measurements. The continuum bands were observed in TDM mode, with 128 channels over the 2 GHz of bandwidth each. <sup>12</sup>CO and  $N_2D^+$  were observed in the same baseband, with each having 58 MHz of bandwidth and 1920 channels, providing a velocity resolution of ~0.08 km s<sup>-1</sup>. <sup>13</sup>CO and C<sup>18</sup>O were also observed in the same baseband with identical bandwidth and velocity resolution. See Table 1 for a summary of the observational setup.

The 12m array observations were conducted on 2015 January 17 in a 14-point rectangular mosaic with 34 antennas operating, covering a  $45'' \times 45''$  region centered between the two protostars. The precipitable water vapor (PWV) was 3.64 mm at the time of execution. Each mosaic point was observed for ∼62 seconds and the complete scheduling block was executed in ∼44 minutes. The observations were manually calibrated by T. Hunter at the North American ALMA Science Center (NAASC) using CASA version 4.3.1 and we used the provided reduction script to generate the calibrated data products. We then self-calibrated the data using the observed dust continuum; not all mosaic points contained a detection of the continuum source, therefore self-calibration was only applied to the mosaic points where the continuum was detected. We performed both phase and amplitude self-calibration on the data. The refined phase and amplitude solutions were then applied to the spectral line base bands. The data were then imaged using the *clean* task in CASA 4.4. We used interactive cleaning for the continuum and spectral line cubes. For the spectral line data, we manually drew masks around the emission in each velocity channel and refined the mask when needed after each iteration of clean.

The ACA observations were conducted on 2014 May 19 in a 5-point mosaic centered between the two protostars. The observation took ∼50 minutes and 9 antennas were operating. Each mosaic point was observed for 74 seconds and the science observations had a total of 12.3 minutes on-source. The PWV during these observations was 0.7 mm. These data were calibrated by B. Mason at the NAASC using the ALMA pipeline and CASA version 4.2.2, and we used the reduction script to regenerate the calibrated data product. We also self-calibrated these data, all mosaic points contained the continuum source due to the larger primary beam of the 7m antennas. The refined phase and amplitude solutions from the continuum were then applied to the spectral line data. The data were imaged using the *clean* task in interactive mode following the same methodology as outlined for the 12m data.

The TPA observations were conducted on 2015 May 2, in three executions of  $\sim$ 57 minutes each observing a  $93.6'' \times 93.6''$  on-the-fly (OTF) map using 2 TPA antennas. The PWV at the time of execution was 0.9 to 1.3 mm. We found that the pipeline-reduced data did not adequately fit the spectral baseline of the  ${}^{12}CO$  data due to the broad linewidth. We re-ran the reduction script in CASA version 4.3.1 and manually adjusted the region to fit the spectral baseline in order to avoid the high-velocity  ${}^{12}CO$  emission.

We then combined the <sup>12</sup>CO, <sup>13</sup>CO, C<sup>18</sup>O, and  $N_2D^+$  data using the methods outlined in the CASA guide for M100 using CASA version  $4.6.0^2$  $4.6.0^2$ . The  $12m+ACA$  visibilities were combined using the concat task in CASA, with the weights being properly adjusted by CASA to compensate for the lower sensitivity of the ACA data. We used the clean task to image the combined dataset in the same manner as the individual datasets. We then followed the CASA guide for M100 to combine

<span id="page-5-0"></span><sup>2</sup> https://casaguides.nrao.edu/index.php/M100 Band3 Combine 5.1

the 12m+ACA and TPA data using the feather task. The fully combined spectral line dataset had significantly reduced artifacts for all molecules where there was significant extended emission. We make use of the fully combined data  $(12m+ACA+TPA)$  in this paper, except for two  $C^{18}O$  positionvelocity (PV) diagrams and this is noted in the text and caption. Furthermore, the continuum maps are only 12m+ACA because total power data are not available.

#### 2.4. Magellan PANIC Observations

We observed BHR71 with the Magellan Baade 6.5m telescope located at Las Campanas using the PANIC (Persson's Auxilary Nasmyth Infrared Camera; [Martini et al.](#page-27-10) [2004\)](#page-27-10) near-infrared imager on 2009 January 17 and 18. The PANIC instrument uses a  $1024^2$  detector with 0.  $12$  pixels, providing a  $2' \times 2'$  field of view. The observations for BHR71 were previously presented in [Tobin et al.](#page-28-8) [\(2010a\)](#page-28-8) and we only observed H and Ks bands with this instrument. The seeing was exceptional during these observations at  $\sim$ 0".4.

#### 2.5. Cerro-Tololo ISPI Observations and Photometry

We observed BHR71 using the Infrared Side-Port Imager (ISPI; [van der Bliek et al.](#page-28-9) [2004\)](#page-28-9) on the Blanco 4m telescope at Cerro-Tololo on 2009 June 11. The ISPI observations and data reduction were described in detail in [Tobin et al.](#page-28-8) [\(2010a\)](#page-28-8), and the imager has a 10' field of view with a  $2048^2$ detector. We use the J, H, and Ks-band images from ISPI for near-infrared photometry in this paper due to the larger field of view as compared to PANIC. Also the absolute calibration of the ISPI data is expected to be more robust due to the larger number of stars available for calibration against 2MASS [\(Skrutskie et al.](#page-28-10) [2006\)](#page-28-10). We measured the photometry in the J, H, and Ks bands using a circular aperture radius of  $30''$  (6000 AU) for IRS1 and a  $5''$  (1000 AU) radius for IRS2. IRS2 was undetected in this aperture. The background was measured using the median of an off-source patch of sky given the highly extended nature of the scattered light emission. The data were calibrated using photometry from the 2MASS catalog [\(Skrutskie et al.](#page-28-10) [2006\)](#page-28-10) and the photometry are listed in Table 2. The seeing during these observations was  $\sim 0$ .

### 2.6. Spitzer Observations and Photometry

BHR71 was observed with the *Spitzer Space Telescope* using the IRAC [\(Fazio et al.](#page-27-11) [2004\)](#page-27-11), MIPS [\(Rieke et al.](#page-27-12) [2004\)](#page-27-12), and IRS instruments [\(Houck et al.](#page-27-13) [2004\)](#page-27-13). The IRAC and MIPS observations were part of the *cores 2 disks* Legacy program and the reduced data were obtained from the *cores 2 disks* data archive. The IRS observations were obtained as part of the IRS guaranteed time program and previously published in [\(Yang et al.](#page-28-3) [2017\)](#page-28-3). The MIPS 24  $\mu$ m photometry for IRS1 and IRS2 were presented in [Chen et al.](#page-26-3) [\(2008\)](#page-26-3), and we re-extracted the IRAC photometry of the extended scattered light nebulae toward BHR71 IRS1 and IRS2 at 3.6, 4.5, 5.8, and 8.0  $\mu$ m using same the methods outlined for the protostar BHR7-MMS in [Tobin et al.](#page-28-11) [\(2018\)](#page-28-11). We extracted the photometry from a 6000 AU  $(30'')$  radius aperture toward IRS1 and a 1000 AU  $(5'')$  aperture radius toward IRS2. The 6000 AU aperture for IRS1 includes flux from IRS2, but the contribution is at most ∼10% (see Table 2). We also extracted the photometry from a  $10000$  AU radius (50") centered on IRS1. We chose these apertures as they are convenient fiducial values and radiative transfer models offer the option to extract flux within different apertures around the source(s) (e.g., [Whitney et al.](#page-28-12) [2003\)](#page-28-12). The 1000 AU aperture restricts the scattered light emission to a region close to the source and is less affected by the source structure; the 6000 AU and 10000 AU apertures are useful characterizing the

total flux density of scattered light from the source(s). Due to the small contribution from IRS2 at wavelengths shorter than 24  $\mu$ m, we decided to extract the large apertures for IRS1, in addition to the small uncontaminated apertures.

#### 2.7. Herschel Observations and Photometry

We obtained the *Herschel* observations toward BHR71 as part of the program OT1<sub>-</sub>jtobin<sub>-1</sub>. BHR71 was observed with the PACS photometer [\(Poglitsch et al.](#page-27-14) [2010\)](#page-27-14) on 2011 July 28. The 100 and 160  $\mu$ m maps were obtained simultaneously with a map size of  $10' \times 10'$ , observing two orthogonal scans for 18.5 minutes each. The 70  $\mu$ m data were obtained separately, mapping a smaller,  $2' \times 2'$  region, also observing 2 orthogonal scans for 14.4 minutes each. Both sets of PACS observations were made using the medium scan speed  $(20<sup>''</sup>$  per minute) for best image quality. The SPIRE [\(Griffin et al.](#page-27-15) [2010\)](#page-27-15) observations were obtained as part of the same program as the PACS observations, on 2011 August 16. We observed a  $10' \times 10'$  region around the protostars with orthogonal scan legs for a total time of 17.25 minutes. In our subsequent analyses, we utilized the Jscanam[3](#page-7-0) products downloaded from the Herschel Science Archive.

We performed photometry on the PACS and SPIRE data using both aperture photometry and point spread function (PSF) photometry. At 70, 100, 160, and 250  $\mu$ m we used PSF photometry to measure the flux densities toward both IRS1 and IRS2. The sources are significantly blended at wavelengths longer than 100  $\mu$ m and are within the PSF wings of each other at 70 and 100  $\mu$ m. We used the IDL program starfinder [\(Diolaiti et al.](#page-26-12) [2000\)](#page-26-12) to perform PSF photometry. Due to the small mapped region, we could not measure the PSF from the data. Instead, we used the Vesta PSF measurements from the pointing refined data that we then rotated and smoothed to match the data. This enabled us to measure the photometry toward the two sources, with the positions being automatically fit at 70, 100, and 160  $\mu$ m. Due to the increased blending at 250  $\mu$ m, we specified the positions for the two sources that were identified at shorter wavelengths and extracted the 250  $\mu$ m photometry. Due to the lack of other sources in the field, we could not cross-calibrate our PSF photometry with aperture photometry to verify its accuracy. However, the *Herschel* Orion Protostar Survey (HOPS) used both aperture and PSF photometry and found close agreement between the two [\(Furlan et al.](#page-27-16) [2016\)](#page-27-16). The flux density measured using PSF photometry is expected to be systematically lower than the aperture photometry because aperture photometry includes additional extended emission surrounding the sources. PSF photometry from PACS has been characterized by Ali et al (2018, in prep.) for the HOPS protostars (e.g., [Furlan et al.](#page-27-16) [2016\)](#page-27-16), finding that the PSF photometry either agrees with the aperture photometry or is systematically low due to additional extended flux in the aperture photometry. [Magnelli et al.](#page-27-17) [\(2013\)](#page-27-17) also used PSF photometry, but created empirical PSFs from their data which were truncated to not include the full PSF wings. This resulted in their photometry being systematically low,  $\sim$ 70% of the aperture measurement at 160 µm. However, we used the full Vesta PSF to fit our data and should not have this systematic effect in our measurements.

To measure the aperture photometry, we used 40" radius apertures at 70, 100, and 160  $\mu$ m, centered between the two protostars. We chose different apertures for the Herschel PACS photometry in comparison to the shorter wavelength photometry because the aperture and color corrections are tabulated for this aperture. We also corrected for the encircled energy fraction as documented in the

<span id="page-7-0"></span><sup>&</sup>lt;sup>3</sup> This is a port of the Scanamorphos map making software [\(Roussel](#page-28-13) [2013\)](#page-28-13) into the *Herschel* Interactive Processing Environment (HIPE).

PACS Data Handbook. The PSF photometry of IRS1 and IRS2 added together is about 20% less than the aperture photometry at each wavelength, which is reasonable because the PSF photometry excludes the extended emission surrounding the sources.

For the SPIRE photometry, we used standard apertures of  $22''$ ,  $30''$ , and  $40''$  for the 250, 350, and  $500 \mu m$  bands, respectively. These apertures are again different from other wavelengths because they have their aperture and color corrections tabulated. We started with the extended source map and utilized the tabulated beam correction, color correction, and aperture correction for a source with a spectral index of 3.5 and a temperature of 30 K. These are reasonable assumptions for a protostar as the SED peaks at  $\sim$ 100 µm and the dust opacity spectral index will result in a spectral index between 3 and 4 in the sub-millimeter range. Our photometry are taken toward the region of compact emission toward the protostar locations and not from the entire extended core; thus, we do not need to worry about the proper zero-point flux calibration as in [\(Sadavoy et al.](#page-28-14) [2018\)](#page-28-14)

#### 3. OBSERVATIONAL RESULTS

The multitude of infrared and submillimeter data taken toward BHR71 are shown in Figures 1 & 2. Figure 1 shows the high-resolution ground-based near-infrared view, the *Spitzer* IRAC view, and the mid- to far-infrared view with MIPS 24  $\mu$ m, *Herschel* 70  $\mu$ m, and 100  $\mu$ m. The PANIC images are shown for H and Ks-bands because of their better image quality, while the J-band image is from ISPI. Very little emission is detected in the vicinity of IRS2 at wavelengths less than 2.2  $\mu$ m, and the image is dominated by the scattered light nebula on the blue-shifted side of the IRS1 outflow. The near-infrared images on larger scales detect the shocked- $H_2$  emission from both outflows [\(Bourke](#page-26-2) [2001\)](#page-26-2). The *Spitzer* IRAC data more obviously show the scattered light emission from both IRS1 and IRS2 in addition to outflow knots, as shown previously by [Chen et al.](#page-26-3) [\(2008\)](#page-26-3). Then the MIPS 24  $\mu$ m and Herschel PACS 70 and 100  $\mu$ m data detect the thermal mid- to far-infrared emission from the inner envelopes of both protostars. IRS2 is clearly resolved from IRS1 and also appears much redder than IRS1 with substantially less 24  $\mu$ m emission. The PACS images were previously published by [Yang et al.](#page-28-3) [\(2017\)](#page-28-3).

We show a larger field of view around BHR71 from *Spitzer* IRAC and *Herschel* SPIRE in Figure 2. The Spitzer IRAC data show the envelope surrounding BHR71 in extinction against the 8  $\mu$ m Galactic background, in addition to the extended outflow knots from both sources. The SPIRE maps show the submillimeter emission from the extended BHR71 core surrounding IRS1 and IRS2, emitting where the 8  $\mu$ m absorption is apparent. At wavelengths longer than 250  $\mu$ m IRS1 and IRS2 are no longer resolved. There is faint  $500 \mu m$  emission (appearing red) along the direction of the IRS1 outflow. This may be the result of CO  $(J = 5 \rightarrow 4)$  emission contaminating the longest wavelength SPIRE band, similar the known CO contamination to SCUBA-2 maps from the  $(J = 3 \rightarrow 2)$  and  $(J = 6 \rightarrow 5)$  lines (e.g., [Drabek et al.](#page-27-18) [2012\)](#page-27-18).

#### 3.1. SED Class of IRS1 and IRS2

The broad wavelength coverage from the near-infrared to the submillimeter also enables us to more quantitatively characterize these two protostars. To accurately distinguish the observational Class for both IRS1 and IRS2, we utilized PSF photometry from *Herschel* at 70, 100, 160, and 250  $\mu$ m, aperture photometry between 3.6  $\mu$ m and 24  $\mu$ m and millimeter flux measurements from ALMA, ATCA, SEST, and SIMBA, see Table 2. All these data are used to construct the SEDs shown in Figure [3](#page-30-0) for IRS1 and IRS2 individually, as well as the combined emission of the sources. The bolometric

luminosity for BHR71 as a whole is 17.7 L<sub>o</sub>. The more luminous source, IRS1, has L<sub>bol</sub>=14.7 L<sub>o</sub> and a bolometric temperature (T<sub>bol</sub>) = 68 K, while the fainter source IRS2 has  $L_{bol} = 1.7 L_{\odot}$  and  $T_{bol}=38$  K. Thus, while IRS1 is about  $10\times$  more luminous, both are Class 0 protostars, with IRS2 possibly being slightly less evolved. The relative inclinations of IRS1 and IRS2 could contribute to their observed  $T_{bol}$  difference, but they are both moderately inclined with respect to the line of sight (see Section 3.3) and it is difficult to determine if this would fully account for the observed difference. We calculate  $L_{bol}$  by integrating the SED using the trapezoidal integration method implemented in the IDL function tsum. Then,  $T_{bol}$  is calculated following [Myers & Ladd](#page-27-19) [\(1993\)](#page-27-19) (also using tsum) in the calculation of the average frequency of the SED.  $T_{bol}$  is the temperature of a blackbody with the same average frequency of the observed SED [\(Ladd et al.](#page-27-20) [1991;](#page-27-20) [Chen et al.](#page-26-13) [1995;](#page-26-13) [Myers et al.](#page-27-21) [1998\)](#page-27-21).

#### 3.2. ATCA and ALMA Continuum data

Continuum emission is detected toward both BHR71 IRS1 and IRS2 at 1.3 mm and 1.3 cm with ALMA and ATCA, respectively, see Figure 4. The ALMA 1.3 mm emission is clearly detected toward both protostars, with IRS1 being ∼12 times brighter at 1.3 mm than IRS2 (Table 2). There is also some extended emission away from the protostar positions, tracing the envelope, and some tenuous emission detected in between IRS1 and IRS2. The emission between IRS1 and IRS2 is possibly analogous to the continuum emission that apparently connects IRAS 16293-2422 A and B [\(Jacobsen](#page-27-22) [et al.](#page-27-22) [2018\)](#page-27-22). The well-resolved structure of IRS1 makes Gaussian fitting unreliable for this source. The major and minor axis of IRS1 out to the  $3\sigma$  level are  $\sim 9''$  and  $\sim 7''$ , respectively. IRS2, on the other hand, is more symmetric. The major and minor axis to the  $3\sigma$  level are both  $\sim 3.9$ <sup>"</sup> but Gaussian fitting yields a deconvolved major axis of 1"69 and a minor axis of 1"23 with a position angle of 87.5◦ . However, the source does not appear well-resolved and the source size from Gaussian fitting may not be overly accurate.

To measure the ALMA 1.3 mm continuum flux densities as reliably as possible, we generated a sigma-clipped map, where pixels below the  $3\sigma$  level ( $\sigma$ =0.68 mJy) are set to zero to avoid including excess noise in the measurement of the flux density. This method will, however, exclude a small amount of real emission, but will result in a more repeatable measurement of flux density. We measured the combined emission from IRS1 and IRS2 in a 35" diameter circle centered between the sources finding a flux density of  $1.41\pm0.02$  Jy. The flux density of IRS1 alone was measured within a 25" diameter circle centered on IRS1, measuring a flux density of  $1.28\pm0.01$  Jy. Finally, the flux density for IRS2 was measured in a  $10<sup>''</sup>$  diameter aperture, centered on IRS2, measuring a flux density of  $0.12\pm0.005$  Jy. Note that the uncertainties on flux densities are statistical only.

The ATCA 1.3 cm continuum data also resolve IRS1 and IRS2, though IRS2 is much fainter; IRS2 is only detected at the  $4\sigma$  level, 12.5 times fainter than IRS1. Since the sources appear point-like and not highly-resolved, we measured the flux densities by simultaneously fitting Gaussians to each source using CASA. The measured flux densities for IRS1 and IRS2 are 2.7 and 0.22 mJy, respectively. [Chen](#page-26-3) [et al.](#page-26-3) [\(2008\)](#page-26-3) also detected both IRS1 and IRS2 at 3 mm; thus, with the addition of our data we can examine the radio spectrum toward IRS1 and IRS2, shown in Figure [5.](#page-31-0) The emission from both IRS1 and IRS2 are consistent with dust-only emission out to 1.3 cm, but with a shallow spectral slope, 2.7 $\pm$ 0.2 for IRS1 and 2.9 $\pm$ 0.5 for IRS2 ( $F_\nu \propto \lambda^{-\alpha}$ ). In this formalism, the expected spectral index for dust emission is  $\alpha = (2+\beta)$  where  $\beta$  is the dust opacity spectral index and  $\beta \sim 1.8$  is expected for dust grains in the interstellar medium. Our data indicate  $\beta$  of 0.7-0.9, which is consistent with other

studies of dust opacity in protostars [\(Kwon et al.](#page-27-23) [2009;](#page-27-23) [Chiang et al.](#page-26-14) [2012\)](#page-26-14). However, we cannot rule-out some contribution from free-free emission from either protostar at 1.3 cm.

We can use the dust continuum data to estimate the inner envelope (and presumed disk) masses surrounding IRS1 and IRS2. We do this by assuming isothermal and optically thin dust emission, adopting a dust-only opacity at 1.3 mm from [Ossenkopf & Henning](#page-27-24) [\(1994\)](#page-27-24) having  $\kappa_{1.3mm}$  0.899  $\text{cm}^2 \text{ g}^{-1}$ , and assuming a dust to gas mass ratio of 1:100 [\(Bohlin et al.](#page-26-15) [1978\)](#page-26-15). We assume a dust temperature of 20 K for IRS1 and IRS2, comparable to the expected envelope temperatures at ∼1000 AU scales for the luminosities of IRS1 and IRS2 [\(Whitney et al.](#page-28-12) [2003\)](#page-28-12). Finally, employing the equation

$$
M_{dust} = \frac{D^2 F_{\lambda}}{\kappa_{\lambda} B_{\lambda} (T_{dust})} \tag{1}
$$

we estimate a total mass (dust+gas) for the inner envelopes/disks of both IRS1 and IRS2 to be 1.25  $M_{\odot}$  from measured flux density of 1.41 Jy. The flux densities for IRS1 and IRS2 are 1.28 Jy and 0.12 Jy, respectively, leading to respective dust+gas masses of 1.13 and 0.11  $M_{\odot}$  if the same temperature is assumed for both sources. Since IRS1 is a factor of ∼8.6 more luminous than IRS1, its average dust temperature should be a factor of ∼1.7 higher, (34 K) than IRS2 assuming that  $T \propto L^{0.25}$ . With an assumed dust temperature of 34 K for IRS1, the estimated mass becomes 0.59  $M_{\odot}$ , making the combined mass 0.7  $M_{\odot}$  considering the likely higher temperature for IRS1.

These masses are for the inner envelope and disk around the protostars, the larger-scale envelope traced by Herschel (Figure 2) is resolved-out and not detected in the ALMA observations. Thus, these mass estimates should be considered lower limits.

The continuum and photometric data enable us to classify the two protostars and measure the amount of mass surrounding the protostars; however, these data do not specifically reveal what led to the formation of the binary system other than there being a large mass reservoir in the envelope surrounding BHR71. To better characterize the formation of the binary system, we must turn to the molecular line data to trace the kinematics of the gas surrounding the protostars. But first, we will briefly examine the outflows of BHR71 observed by ALMA.

## 3.3. Outflow Morphology from ALMA

The outflows from BHR71 IRS1 and IRS2 are quite prominent in the scattered light images from the near-infrared and *Spitzer*. However, we also traced the outflow with <sup>12</sup>CO and <sup>13</sup>CO observations with ALMA (Figure [6\)](#page-32-0). The integrated intensity maps of the CO emission from IRS1 and IRS2 (Figure [6\)](#page-32-0) clearly show the red and blue-shifted sides of both outflows. The <sup>12</sup>CO maps indicate that two protostars have their outflows inclined in opposite directions. The blue-shifted side of the outflow from IRS1 is on the south and the red-shifted side is to the north, while this is opposite in the case of IRS2. This orientation of the outflows had been previously indicated from the near-infrared and IRAC imaging as well as from single-dish observations [\(Bourke](#page-26-2) [2001;](#page-26-2) [Parise et al.](#page-27-25) [2006\)](#page-27-25). We also find that the  $C^{18}O$  traces the edge of the outflow cavity walls toward IRS1 in Figure [6.](#page-32-0) From these data, we measure outflow position angles of  $174°$  and  $-31°$  for IRS1 and IRS2, respectively. The angles are measured east of north and are found by drawing a line that bisects the outflow cavity and passes through the continuum position.

The outflow from IRS2 has a more narrow opening angle than the outflow of IRS2. The outflow opening angles of IRS1 and IRS2 are 55<sup>°</sup> and ~47<sup>°</sup>, respectively, measured at a distance of 10<sup>"</sup> from the protostars. Note that these are the full opening angles of the outflow cavities, radiative transfer

modeling studies typically quote the half-opening angles [\(Whitney et al.](#page-28-12) [2003\)](#page-28-12). The <sup>13</sup>CO emission from IRS1 traces lower-velocity emission better than <sup>12</sup>CO and we measure a slightly larger opening angle of ∼63◦ . The outflow from IRS2 is not well-detected in <sup>13</sup>CO and we are unable to make a similar measurement.

The red-shifted side of the outflow toward IRS2 shows peculiar structure at distances greater than  $10<sup>''</sup>$  from the protostar. Along the line of sight, the blue-shifted outflow of IRS1 and red-shifted outflow of IRS2 overlap. The northern side of the red-shifted outflow from IRS2 fans out to a much larger opening angle at this point and this continues past the point where the outflow cavities overlap. This wider feature also appears in the <sup>13</sup>CO integrated intensity map at lower signal to noise.

A more in depth study of the outflow system will be presented by Bourke et al. (in prep.). However, we can make some estimates of the source inclinations using the outflow geometry in the absence of detailed modeling. This is possible because there is very limited overlap between the blue- and red-shifted sides of both outflows. Assuming that the outflows are conical, the lack of significant overlap of blue and red-shifted sides, both along the outflow and toward the protostars, means that the IRS1 outflow must have an inclination between 35 and 63<sup>°</sup>, because its opening angle is  $\sim 55^{\circ}$ . The outflow is likely closer to  $63^{\circ}$  than the other extreme because of how extended it is in the plane of the sky. The IRS2 outflow with a more narrow opening angle of ∼47◦ must have an inclination between 43 and 67◦ . We define 90◦ as viewing the system edge-on where the outflow is totally in the plane of the sky, and  $0°$  is edge-on where the outflow is completely along the line of sight.

## 4. DENSE MOLECULAR LINE KINEMATICS

The datasets from Parkes, ATCA, and ALMA (Table 1) combined trace the kinematics of the dense gas from the scale of the entire BHR71 core to the envelopes surrounding the individual protostars. It is instructive to start from the largest scales and move inward, thus we will begin with a discussion of the Parkes data. To introduce the data, we show the spectrum of each line toward each source (both sources for Parkes) and plot them in Figure [7.](#page-33-0) The spectra are extracted from a circular region matching the size of the beam in each observation (see Table 1), except for the ALMA data where we extract the spectra within a  $5''$  diameter circle. The <sup>12</sup>CO and <sup>13</sup>CO have broad linewidths tracing the outflow and  $C^{18}O$  mainly traces the inner envelope with some outflow contamination. N<sub>2</sub>D<sup>+</sup> traces the cold envelope and this molecule is mostly destroyed in the  $5''$  region around of IRS1, and NH<sub>3</sub> also traces the cold gas but on the larger scales probed by ATCA. Given that  $NH_3$  and  $N_2D^+$  have lower intensities at the protostar positions, we also extracted spectra from the position of the  $N_2D^+$ emission peaks east of IRS1 and east of IRS2. These positions better show the hyperfine structure of  $N_2D^+$  and NH<sub>3</sub>.

The largely distinct regions traced by CO and its isotopologues and NH<sub>3</sub>, N<sub>2</sub>D<sup>+</sup>, and N<sub>2</sub>H<sup>+</sup> are the result of both the thermal structure of the infalling envelopes and chemical processes. CO is frozen-out onto dust grains at temperatures below ∼20 K and at typical densities of star forming cores n >10<sup>5</sup> cm<sup>-3</sup> [\(Frerking et al.](#page-27-26) [1982;](#page-27-26) [Benson & Myers](#page-26-16) [1989\)](#page-26-16). Observations and astrochemical models show that  $N_2H^+$ ,  $N_2D^+$ , and  $NH_3$  form abundantly in these cold, dense regions where CO is frozen-out (e.g., [Caselli et al.](#page-26-17) [2002;](#page-26-17) [Lee et al.](#page-27-27) [2004;](#page-27-27) [Bergin et al.](#page-26-18) [2002\)](#page-26-18). However, in the central regions of the protostellar envelopes, the temperature increases due to heating from the protostar and accretion which evaporates CO from the dust grains inside R∼1000 AU (the actual radius depends luminosity). In this region, there is an extremely efficient reaction of CO with  $N_2H^+$  and  $N_2D^+$ that destroys these molecules rapidly. Also, N<sub>2</sub>D<sup>+</sup> formation is shut off at T∼20 K because the key

molecule in its formation pathway,  $H_2D^+$  is converted back into  $H_3^+$  and HD (e.g., [Herbst](#page-27-28) [1982\)](#page-27-28). Then there is also a destruction pathway for  $NH_3$  involving reactions with  $HCO^+$ , which will form in the presence of CO [\(Lee et al.](#page-27-27) [2004\)](#page-27-27); there is also the possibility that  $NH<sub>3</sub>$  will freeze-out onto dust grains before being completely destroyed by  $HCO^+$  at n >10<sup>6</sup> cm<sup>-3</sup> [\(Visser et al.](#page-28-15) [2011\)](#page-28-15). This leads to the onion-like chemical structure of protostellar envelopes where  $N_2H^+$ ,  $N_2D^+$ , and  $NH_3$  exist in gas-phase in the outer regions not in the inner regions, and CO exists most prevalently in the inner regions without  $N_2H^+$ ,  $N_2D^+$ , and  $NH_3$ . We note, however, that these are not the only molecules in the gas phase in these regions.

## 4.1. Parkes NH<sub>3</sub> Kinematics

The Parkes data traced the NH<sub>3</sub> (1,1) emission line with 60<sup> $\prime\prime$ </sup> (12000 AU) resolution out to 8<sup> $\prime$ </sup>  $(\sim 0.5 \text{ pc})$  scales, probing the largest scales available to us in molecular gas. NH<sub>3</sub> emission is detected from the entire region where the envelope is seen in absorption at  $8 \mu m$  (Figure 2), and extending a bit beyond the edge of the map to the southeast as shown in Figure [8.](#page-34-0) The emission does not likely extend beyond the northern end of the map; the intensity appears to rise here due to increased noise at the map edge.

Using the  $NH<sub>3</sub>$  fitting routines in CLASS, we fit the hyperfine lines in each pixel of the map where emission was detected above the  $5\sigma$  level and the line-center velocity and linewidth maps that are also shown in Figure [8.](#page-34-0) The line-center velocity map shows a slight velocity gradient along the long axis of the NH<sub>3</sub> emission from southeast to northwest, in the same direction as the extended  $8 \mu m$ extinction. The total velocity change is about  $0.15$  -  $0.2 \text{ km s}^{-1}$ , corresponding to a gradient of about 1 km s<sup>−</sup><sup>1</sup> pc<sup>−</sup><sup>1</sup> . We measure the velocity gradient more quantitatively by fitting the slope of a 1D cut along the envelope orthogonal to the outflow of IRS1 and by fitting a 2D plane to the line-center velocity maps. The resulting velocity gradient orthogonal to the outflow direction is 1.2 km s<sup>-1</sup> pc<sup>-1</sup> and the 2D velocity gradient fit finds a gradient of 1.15 km s<sup>-1</sup> pc<sup>-1</sup> with a position angle of 101<sup>°</sup> east of north. This velocity gradient could be interpreted as core rotation, as is often done (e.g., [Goodman](#page-27-29) [et al.](#page-27-29) [1993\)](#page-27-29), but it was shown in [Tobin et al.](#page-28-4) [\(2011,](#page-28-4) [2012b\)](#page-28-16) that infalling flows along asymmetric envelopes could also produce velocity gradients. If the velocity gradient of the core reflects rotation, then IRS2 should orbit around IRS1 in the same direction as the core rotation. Furthermore, if IRS1 and IRS2 fragmented within this rotating core, the individual envelopes should have rotation in the same direction in the case of angular momentum conservation.

The linewidth map in Figure [8](#page-34-0) shows some variation across the source. The southern-most tip of the map has a very broad linewidth, this appears real and could be in an area affected by the two outflows. The eastern side of the linewidth map is relatively constant at 0.5 to 0.6 km s<sup> $-1$ </sup> and on the western side the linewidth dips down to ~0.3 km s<sup>-1</sup>. These linewidths are in excess of the expected linewidth of NH<sub>3</sub> at 20 K of 0.13 km s<sup>-1</sup>, indicative of an additional non-thermal component. Some of this non-thermal component could come from unresolved velocity gradients in the envelope within the  $60^{\prime\prime}$  beam.

While the Parkes data are quite informative of the large-scale velocity structure, they also demonstrate a clear need to follow the kinematics to smaller scales with interferometric data to fully characterize the formation pathway of IRS1 and IRS2.

4.2. ATCA NH<sup>3</sup> Kinematics

The ATCA observations detected significant  $NH<sub>3</sub>$  emission near the protostar as shown in Figures [9](#page-35-0)  $\&$  [10.](#page-36-0) The emission morphology is quite filamentary, also concentrated along the extended envelope seen in extinction at 8  $\mu$ m. The strongest emission is found southeast of IRS1 with another subpeak located east of IRS2, between the sources. The emission west of IRS2 is more diffuse with lower surface brightness. We also note that the NH<sub>3</sub> emission does not peak on the locations of the protostars themselves, and the morphology of the emission from the main NH<sup>3</sup> lines is quite similar to that of the satellite lines. We overlay the  ${}^{12}CO$  integrated intensity maps in Figure [9](#page-35-0) to highlight the regions that the outflows may influence. At first glance, it does not appear likely that the  $NH<sub>3</sub>$ emission east and west of IRS1 is significantly affected by the outflows.

The kinematic structure derived from the ATCA observations (Figure [10\)](#page-36-0) is also fit using the  $NH<sub>3</sub>$ fitting routines within CLASS. This was done for each pixel of the map where emission was detected above  $5\sigma$ , see Appendix A. The ATCA kinematic structure bears some similarity to the Parkes observations, but also has some significant departures. The line-center velocity map has features that extend to both more red-shifted and blue-shifted velocities than were evident in the Parkes map. North of IRS1, there is a red-shifted feature with velocities greater than -4.2 km s<sup>-1</sup> and there is a red-shifted feature directly southeast of IRS1. These features may reflect outflow interaction with the envelope as evidenced by the shape of the conical shape of the red-shifted feature north of IRS1, and the position of the red-shifted southeast feature along the edge of the  $NH<sub>3</sub>$  emission, bordering the cavity wall of the southern outflow lobe. The linewidth map also shows broad linewidth south of IRS1 and to the southeast; however, the linewidth is very narrow to the north of IRS1. Outflow interactions can produce increased linewidth and/or changes in the line-center velocity.

Between IRS1 and IRS2 there is a steep velocity gradient from red-shifted to blue-shifted in the line-center map that continues past the position of IRS2. In the region with the steep velocity gradient, there is an accompanying increase in the linewidth map that results from the velocity gradient being unresolved; such a feature showed up several times for other sources in [Tobin et al.](#page-28-4) [\(2011\)](#page-28-4). This gradient from red to blue is in the same direction as the large-scale gradient observed in the Parkes NH<sub>3</sub> map. Beyond  $\sim \pm 40''$  from IRS1 along the direction of the extended envelope, the line center velocities are about equal in both the southeast and northwest sides of the envelope. The linewidth northwest of IRS2 is narrow, between 0.2 and 0.3 km s<sup>-1</sup>. To the southeast, the linewidth remains above 0.3 km s<sup>−</sup><sup>1</sup> and may be the result of outflow-envelope interaction, as suggested already. However, the increased linewidth feature extends beyond the edge of the outflow cavity, yet there is unknown 3D structure along the line of sight. Depending on the configuration of material, the outflow could affect a larger area beyond its boundaries, and increased linewidth propagates into the envelope, or if the envelope is filamentary and curved along the line of sight gas falling-in toward the protstar could yield an increased linewidth. To the west of IRS2, the linewidth is quite narrow, except for one area that has a linewidth of  $0.5 \text{ km s}^{-1}$ .

We zoom-in on the immediate region around IRS1 and IRS2, the inner  $60''$  of the source, in Figure [11,](#page-37-0) showing the line-center velocity in greater detail. From east-to-west, crossing the source, the velocity goes from -4.55 km s<sup>-1</sup> in the eastern-most part of the map to -4.25 km s<sup>-1</sup> at IRS1, -4.6 km s<sup>-1</sup> at IRS2, and down to -4.85 km s<sup>-1</sup> west of IRS2. Qualitatively, the gradient goes from blue, to red, and back to blue; thus, there is not a clear velocity gradient that can be attributed to rotation around IRS1 and IRS2. Furthermore, the strip of broad linewidth directly north of IRS2 can be indicative of a rapid velocity change that can result from two velocity components along the

line of sight. This was observed toward several protostars in [Tobin et al.](#page-28-4) [\(2011\)](#page-28-4), and the higher resolution ALMA  $N_2D^+$  data will be used to investigate this further.

The velocity profiles from the ATCA and Parkes velocity maps are extracted along lines orthogonal to the outflow directions and shown in Figure [14.](#page-40-0) The Parkes data show a velocity gradient comparable to the two-dimensional fit. The ATCA profile shows significant structure near the protostar and then the velocities move back toward agreement with the cloud for IRS1. For IRS2, a very steep velocity gradient is derived from the ATCA data. The fits to the Parkes velocity gradients also find that the core velocity at the position of IRS1 is -4.45 km s<sup>-1</sup>.

The velocity and linewidth maps derived from hyperfine fitting and the profiles extracted are useful for providing a global view of the envelope kinematics. However, they can abstract some more detailed information provided by the line profiles. We show position-velocity (PV) diagrams for both the main  $NH_3$  lines and satellite lines in Figure [12.](#page-38-0) These are extracted from a 10" strip in the east-west direction (position angle =  $90^{\circ}$ ), centered on IRS1. PV diagrams extracted from NH<sub>3</sub> (1,1) datacubes are complex because each hyperfine component is a pair of closely spaced lines, so caution must be exercised in their interpretation. Figure [12](#page-38-0) shows that the western portion of the envelope (containing IRS2) is best described by a single velocity component (-4.6 km s<sup>−</sup><sup>1</sup>and offsets between  $-25$ <sup>n</sup> to  $-5$ <sup>n</sup>), and this component shows a slight gradient to more blue-shifted velocities. However, another distinct velocity component is present on the east side of the envelope (-4.0 km s<sup>−</sup><sup>1</sup>and offsets between  $-5$ <sup>n</sup> to  $25$ <sup>n</sup>), red-shifted with respect to far west side of the envelope. This is most evident in the PV diagram of the satellite lines where there appear to be three peaks; each velocity component of NH<sup>3</sup> has two hyperfine lines and the velocity shift is large enough that one hyperfine from the blue and red component are overlapping. This large shift appears in the velocity map as the redshifted component that is coincident with the red-shifted outflow from IRS1, but this red-shifted component extends beyond the bounds of the outflow. This could indicate that there is a second velocity component to the core, or that the outflow is imparting a bulk redshifted velocity on dense molecular gas in the envelope.

In addition to the PV diagrams in the equatorial cuts, we also show a PV diagram extracted from a larger region along the long-axis of the envelope (a position angle of 123<sup>°</sup>) in Figure [13.](#page-39-0) This PV diagram shows that the features observed in the smaller region are present across much of the envelope with the red-shifted velocity component appearing throughout most of the East side of the envelope. The red-shifted velocity component (-4.0 km s<sup>-1</sup>) begins to appear just before the PV cut crosses IRS1 (offsets from  $-5''$  to 70"), indicating that it could be related to the red-shifted outflow from IRS1. However, the red-shifted velocities continue as the the cut follows the envelope southeast, well-beyond the region of influence of the red-shifted outflow from IRS1. The red-shifted outflow of IRS2 is directed in this region and has substantial <sup>12</sup>CO and <sup>13</sup>CO emission that fans out toward the east side of the envelope. Thus, it is clear that the eastern side of the envelope has a second, redshifted velocity component, but it is unclear if this is outflow-related or part of the cloud structure itself.

## 4.3. ALMA  $N_2D^+$  Kinematics

With the ambiguity of the NH<sub>3</sub> kinematics near IRS1 and IRS2, higher resolution is imperative to disentangle the kinematic structure of gas surrounding the binary system. The ALMA observations include  $N_2D^+$  ( $J = 3 \rightarrow 2$ ), the deuterated form of  $N_2H^+$ .  $N_2H^+$ ,  $N_2D^+$ , and  $NH_3$  have been shown in multiple studies to trace the same (or very similar) kinematic and physical structures in both

starless and protostellar cores in single-dish and interferometric studies [\(Johnstone et al.](#page-27-30) [2010;](#page-27-30) [Tobin](#page-28-4) [et al.](#page-28-4) [2011\)](#page-28-4).  $N_2D^+$  has the caveat that it can be destroyed in the inner regions of the envelope where  $N_2H^+$  and NH<sub>3</sub> can still exist [\(Tobin et al.](#page-28-17) [2013\)](#page-28-17), but in the case of BHR71 the NH<sub>3</sub> and N<sub>2</sub>D<sup>+</sup> peaks align closely with the  $N_2H^+$  peaks found by [Chen et al.](#page-26-3) [\(2008\)](#page-26-3).

We show the  $N_2D^+$  integrated intensity maps overlaid on the *Spitzer* 8  $\mu$ m image, the ATCA  $NH_3$  and the 1.3 mm continuum in Figures [15](#page-41-0) & [16.](#page-42-0) The N<sub>2</sub>D<sup>+</sup> map covers a smaller region than the ATCA NH<sub>3</sub> map; as such, the N<sub>2</sub>D<sup>+</sup> does not map to the 8  $\mu$ m absorption feature as well as  $NH<sub>3</sub>$ . However, there is good correspondence to the east of IRS1. The N<sub>2</sub>D<sup>+</sup> also tends to avoid the region of the outflow toward IRS1, but there is some emission north of IRS1 along the red-shifted outflow. Toward IRS2, the  $N_2D^+$  does not clearly avoid the outflow, which might be due to the three dimensional configuration of the envelope. Finally, the  $N_2D^+$  east of IRS2 corresponds well to the  $NH_3$  map, but east and south of IRS2 the NH<sub>3</sub> intensity is dropping while there is N<sub>2</sub>D<sup>+</sup> emission surrounding IRS2 on three sides. We also notice in Figure [16](#page-42-0) that the extended 1.3 mm continuum around IRS1 seems to avoid the brightest areas of  $N_2D^+$  emission. Specifically, the extended emission west of IRS1 seems to run through the gap in  $N_2D^+$  toward IRS2.

The kinematic structure is derived from the  $N_2D^+$  data using the CLASS hyperfine fitting routines and the known hyperfine line positions and relative line ratios [\(Dore et al.](#page-27-31) [2004\)](#page-27-31); also see [Tobin](#page-28-17) [et al.](#page-28-17) [\(2013\)](#page-28-17); see Appendix A. The overall kinematic structure of  $N_2D^+$  is quite similar to  $NH_3$  in the overlapping regions, except that some of the velocity gradients are better resolved. There are also still indications of outflow-envelope interaction with red shifted line-center velocity southeast of IRS1 and increased linewidth in this region. In the  $N_2D^+$  linewidth map, the linewidth is remarkably low across the entire envelope. The map is a bit noisy due to the smaller beam and lower signal-to-noise in some regions. North of IRS1, there is still the large region of red-shifted velocities that coincide with the likely area of influence from the outflow. Toward IRS2, the line center velocity map is not highly structured, particularly away from the region of red-shifted velocities that seem to be associated with IRS1. The velocity gradient around IRS2 appears to be more along the outflow direction than orthogonal to it. Overall, the  $N_2D^+$  line-center velocity near IRS2 is very close the line-center velocity to the east of IRS1. Thus, it is not clear that the kinematics strongly indicate rotation across the core on scales comparable to the protostar separation, or even around the protostars themselves. The area of large linewidth observed north of IRS2 in NH<sub>3</sub> with ATCA is not reflected in the  $N_2D^+$  map, likely because this was due to a velocity gradient that is resolved in the ALMA map. Given that the velocities of the gas toward IRS2 and the gas east of IRS1 have comparable velocities, the kinematics surrounding IRS1 and IRS2 do not strongly indicate that they are forming out of kinematically distinct cores. The only firm conclusion that we can make is that the outflow from IRS1 appears to be significantly affecting the kinematic structure to the north.

The velocity profiles from the ALMA  $N_2D^+$  velocity maps are also extracted along lines orthogonal to the outflow directions and shown in Figure [14.](#page-40-0) Compared to the ATCA NH<sub>3</sub> velocity profiles, the  $N_2D^+$  profiles are significantly more structured and the velocity away from the protostar position for IRS1 does not as closely agree with the Parkes velocities. This could result from chemical differences between  $N_2D^+$  and  $NH_3$ , or the higher resolution of the  $N_2D^+$  data could be resolving structure that is smoothed out in the lower resolution ATCA data.

In addition to the line center velocity maps and profiles, we also extracted a PV diagram from the  $N_2D^+$  emission toward BHR71. This PV diagram is shown in Figure [17](#page-43-0) and is taken in an

east-west cut 10<sup>n</sup> in width, centered on IRS1. The angular resolution of the N<sub>2</sub>D<sup>+</sup> maps is about 5 times finer than the  $NH_3$  map, so some features appear different, but the major features are similar. The PV cut of  $N_2D^+$  shows very little evidence for a systematic velocity gradient across the position of IRS1. There is also still a second velocity component evident  $(-4.0 \text{ km s}^{-1})$ , but this red-shifted component is most evident near the protostar position at an offset position of  $\pm 5$ ". The red-shifted velocity component is made less obvious due to the brightest hyperfine lines being spread over  $\sim 0.5$  km s<sup>-1</sup> [\(Dore et al.](#page-27-31) [2004\)](#page-27-31), in contrast to the pairs of ammonia lines that are less blended. The lines also appear slightly more red-shifted than  $NH<sub>3</sub>$  because we assigned the rest frequency to match the brightest  $N_2D^+$  hyperfine line, which is at a higher frequency than the other hyperfine lines of comparable relative intensity.

## 4.4. ALMA C <sup>18</sup>O Kinematics

Inside the  $N_2D^+$  and  $NH_3$  emitting regions, we find that  $C^{18}O$  emission peaks directly on the continuum sources as shown in Figure [6.](#page-32-0) There is also extended emission around IRS1 and IRS2, but with most emission concentrated at the location of the continuum sources. Moreover, toward IRS1, the C<sup>18</sup>O appears extended along the walls of the outflow cavities. There is also a thin ridge of increased C<sup>18</sup>O brightness extended between IRS1 and IRS2.

We examined the  $C^{18}O$  line kinematics for signs of rotation, infall, and/or outflow entrainment by first examining the integrated intensity, line center velocity, and linewidth maps shown in Figure [18.](#page-44-0) These are similar to the maps constructed for the  $NH_3$  and  $N_2D^+$  data, but instead of fitting the hyperfine lines, we calculate the standard 1st (centroid velocity) and 2nd (linewidth) moment maps because  $C^{18}O$  is a single emission line. The centroid velocity map shows the signature of the outflow from IRS1, but there is a very slight 'twist' in the velocity map across the source indicating that there is more going on that just the outflow. Also, the linewidth map shows a broader linewidth near the position of IRS1, indicative of more rapid motion along the line of sight. There is not evidence of enhanced linewidth toward IRS2, but moment maps are most often dominated by the low-velocity emission that is strongest, especially in the case of a map with total power emission included (Figure [7\)](#page-33-0)

In order to better examine the motions of the gas near the protostar positions, we examine integrated intensity maps constructed using only the higher velocity blue- and red-shifted emission independently. Toward IRS1, integrated intensity maps are shown in Figure [19](#page-45-0) for the red and blueshifted  $\rm C^{18}O$  in velocity ranges from low (-5.8 to -4.4 km s<sup>-1</sup> and -4.4 to -3.2 km s<sup>-1</sup>), medium (-6.2 to -5.6 km s<sup>-1</sup> and -3.6 to -2.8 km s<sup>-1</sup>), and high (-6.3 to -5.8 km s<sup>-1</sup> and -3.0 to -2.5 km s<sup>-1</sup>). At lower velocities the red and blue-shifted emission are clearly offset along the outflow direction. At medium velocities, the emission becomes more compact and the red and blue-shifted emission are oriented diagonally across the continuum source, possibly tracing both outflow and motion in the equatorial plane. We also note that velocity gradients in the direction of the outflow may result from infall motion [\(Yen et al.](#page-28-18) [2013\)](#page-28-18), but may also be related to outflow-envelope interaction [\(Arce](#page-26-5) [& Sargent](#page-26-5) [2006\)](#page-26-5).

The C<sup>18</sup>O at the highest velocities has blue and red-shifted emission peaks that are oriented in the east-west direction (blue on the east, red on the west), with some low-level emission along the outflow direction. Thus, this velocity gradient is oriented perpendicular to the outflow, as expected for rotational motion on <1000 AU scales in the inner envelope. Note that the direction of this velocity gradient is the opposite of what is observed on large-scales with Parkes and ATCA.

The  $C^{18}O$  emission toward IRS2, on the other hand, is found to have the red-shifted peak on the east and the blue-shifted peak is on the west (Figure [20\)](#page-45-1). The position angle is not exactly orthogonal to the outflow, it is not along the outflow either. Thus, the small-scale velocity gradients in  $C^{18}O$ appear to be in the opposite directions for IRS1 and IRS2.

To examine the kinematics in more detail, we show PV diagrams of the C<sup>18</sup>O emission toward IRS1 and IRS2 in Figure [21.](#page-46-0) These PV cuts are taken orthogonal to the outflow directions of each source, centered on the continuum source. For both sources, positive offsets are to the east and negative offsets are to the west. The PV diagrams in Figure [21](#page-46-0) are generated using only the data from the ALMA 12m array to reduce confusion with the extended emission picked up by the ACA and Total Power observations.

The PV diagram for IRS1 shows that at high-velocities near the continuum position, the blue-shifted side is located to the east and the red-shifted side is located to the west. Toward lower velocities, the C<sup>18</sup>O emission is more symmetrically distributed in both position and velocity. IRS2 shows a considerably different appearance. First, the  $C^{18}O$  is not as spatially extended, likely because IRS2 is ten times less luminous than IRS1, and the emitting region is expected to be smaller. Second, the region with the brightest blue-shifted C<sup>18</sup>O is still on the west side of the envelope, while the brightest red-shifted  $C^{18}O$  is on the east side of the envelope. Thus, both in the integrated intensity maps and the PV diagrams, the velocity gradients of IRS1 and IRS2 are in the opposite direction.

The PV diagrams also show that the  $C^{18}O$  emission of the protostars is slightly different from the velocity of the core. The C<sup>18</sup>O emission is best described with a central velocity of -4.6 km s<sup>-1</sup> in contrast to the average core velocity of -4.45 km s<sup>-1</sup> from Parkes  $NH_3$  emission. Parts of the core also show velocities that are even more different, Figure [16](#page-42-0) shows that the core near IRS2 and east of IRS1 has velocities up to -4.9 km s<sup>-1</sup>, and near IRS1 the velocity can be -4.2 km s<sup>-1</sup>.

To compare the PV diagram of the  $C^{18}O$  emission to that of the NH<sub>3</sub> and N<sub>2</sub>D<sup>+</sup>, we use of the 12m+ACA+TP data to show the PV diagram on comparable spatial scales across both sources and show this larger scale  $C^{18}O$  PV diagram in Figure [22.](#page-47-0) The PV diagram shows that there is extended structure from the surrounding core toward both IRS1 and IRS2. Similar to  $NH_3$  and  $N_2D^+$ , the  $C^{18}O$  shows an apparent second velocity component around -4.0 km s<sup>-1</sup> and present from -25" to beyond  $30''$ . However, the protostars themselves seem to be more closely associated with the blueshifted velocity component. The blueshifted component is at about -4.9 km s<sup>-1</sup> and the protostars are between -4.7 and -4.6 km s<sup>-1</sup>. We note, however, that the large-scale  $C^{18}O$  emision likely reflects the velocity of the gas in the exterior regions of the could because  $C^{18}O$  is likely frozen-out onto dusgrams in the higher density interior of the core/envelope where  $N_2D^+$  and  $NH_3$  are present [\(Frerking et al.](#page-27-26) [1982;](#page-27-26) [Benson & Myers](#page-26-16) [1989\)](#page-26-16).

#### 5. DISCUSSION

As an isolated core harboring a wide binary system, BHR71 should lend itself to being an ideal testing ground for wide binary formation. The formation of such systems have long been thought to be related to the rotation of the core (e.g., [Larson](#page-27-6) [1972;](#page-27-6) [Boss & Bodenheimer](#page-26-19) [1979;](#page-26-19) [Burkert](#page-26-6) [& Bodenheimer](#page-26-6) [1993;](#page-26-6) [Tohline](#page-28-2) [2002\)](#page-28-2). Thus, our initial expectation was to find a classic scenario of clear core rotation, with conserved angular momentum leading to increased rotation velocity at progressively smaller radii. We also expected that a difference in radial velocity of IRS2 relative to IRS1 due to orbital motion could be measured because its separation, lower overall luminosity, and estimated dust+gas mass.

#### 5.1. Kinematic Analysis

The kinematic structure observed in  $NH<sub>3</sub>$  from Parkes and ATCA alone, appeared remarkably consistent with rotation-induced fragmentation. The Parkes data showed a modest velocity gradient on arcminute scales, and the ATCA data showed evidence of a larger velocity gradient across the two protostars. Furthermore, previous ATCA  $N_2H^+$  observations toward IRS1 detected a velocity gradient across the main protostar, but did not detect emission around IRS2 due to the smaller primary beam at 3 mm [\(Chen et al.](#page-26-3) [2008\)](#page-26-3). The velocity gradient observed by Parkes along the major axis of the core as viewed in NH<sub>3</sub> is 1.2 km s<sup>-1</sup> pc<sup>-1</sup>, measured across a diameter of 150″ (30000 AU). The low-resolution of the single-dish observations smooth out any velocity structure at small scales and provides a measure of the velocity gradient on the largest scale, which for the sake of this discussion, we interpret as solid-body rotation. But once material begins free-fall, rotation is no longer solid-body and conserves angular momentum (in the absence of magnetic fields). This gradient would correspond to a rotation velocity of 0.087 km s<sup>-1</sup> at the core radius of 15000 AU and an angular velocity ( $\Omega$ ) of 3.9 ×10<sup>-14</sup> s<sup>-1</sup>. If we assume that the entire core at radii less than 15000 AU is in freefall, the infalling gas should conserve angular momentum. Then, the rotation velocity should be  $\sim 0.33$  km s<sup>-1</sup> at a radius of 4000 AU. At the separation of IRS2, 16" (3200) AU) the rotation rate would be even higher, 0.4 km s<sup>-1</sup>. The ALMA  $N_2D^+$  map shows very little evidence for a velocity gradient of this magnitude from the area directly east of IRS1 to the area surrounding IRS2. Furthermore, the  $C^{18}O$  velocities of IRS1 and IRS2 are consistent with being at the same velocity. The lack of observed rotation velocity on 3200 AU scales, however, could be due to the apparent influence of the outflow on the ambient envelope material, masking hints of rotational velocity increases toward the envelope center. But, the largest scales examined by the ALMA  $N_2D^+$ do not show ordered rotation surrounding either IRS1 or IRS2.

In the preceding paragraph, we assumed that the entire core was falling in and that only the velocity gradient on the largest scales reflected solid-body rotation. If we instead assume that the entire core is not yet collapsing and the 1.2 km s<sup>-1</sup> pc<sup>-1</sup> velocity gradient reflects solid body rotation out to a certain radius, then we can calculate different rotation velocities. Assuming inside-out collapse [\(Shu](#page-28-19) [1977;](#page-28-19) [Terebey et al.](#page-28-20) [1984\)](#page-28-20), the infall radius must be larger than the companion separation. If we assume an infall radius of 6400 AU, twice the companion separation, the solid body rotation velocity at this radius would be 0.037 km s<sup>-1</sup> with the 1.2 km s<sup>-1</sup> pc<sup>-1</sup> gradient ( $\Omega = 3.9 \times 10^{-14}$  s<sup>-1</sup>), below our ability to detect. Furthermore, if angular momentum was conserved from a hypothetical infall radius of 6400 AU, then the rotation velocity at a radius of 3200 AU would only be 0.074 km s<sup>-1</sup>. Both the solid-body rotation rate at 6400 AU and the estimated rotation velocity from conserved angular momentum at 3200 AU would be below our ability to detect. Nevertheless, the core is observed to have velocity structure with significantly larger amplitudes than these extrapolations from the large-scale core rotation. These inconsistencies can be taken as evidence against ordered collapse of the system with rotationally induced fragmentation.

Using these observationally measured quantities of the core velocity structure, we can calculate the estimated stability of the core from the ratio of rotational energy to gravitational potential energy  $(\beta_{\rm rot})$  on the scale of 3200 AU. While this diagnostic assumes solid-body rotation and we use a rotation velocity inferred from the assumption of constant angular momentum, this analysis enables comparisons to other studies that determine  $\beta_{\rm rot}$  as their key diagnostic derived from the observations.

We follow the method outlined by [Chen et al.](#page-26-20) [\(2007\)](#page-26-20) which calculates the rotational energy as

$$
E_{rot} = \frac{1}{2}I\Omega^2 = \frac{1}{2}\alpha_{rot}MR^2\Omega^2
$$
\n(2)

and  $\alpha_{rot} = \frac{2}{3}$  $\frac{2}{3}(3-p)/(5-p)$ ,  $\Omega$  is the angular velocity derived from the observed velocity gradient, R is the radius of the envelope, and M is the mass of the envelope. We adopt  $p=1.5$  for a spherical envelope in free-fall. Note that this is not exact because the envelope around BHR71 IRS1 and IRS2 is non-spherical. Then the gravitational binding energy is defined from the virial theorem to be

$$
E_{grav} = \frac{3GM^2}{5R} \tag{3}
$$

where G is the gravitation constant; M and R are the same as defined for the previous equation. Knowing these two equations,  $\beta_{\rm rot}$  can be calculated

$$
\beta_{\rm rot} = \frac{E_{rot}}{E_{grav}} = \frac{5R^3 \Omega^2 \alpha_{rot}}{6GM}.
$$
\n(4)

We can simplify this relationship by multiplying constants through and converting the more natural observed units such as solar masses, parsecs, and km s<sup>−</sup><sup>1</sup> pc<sup>−</sup><sup>1</sup> , yielding

$$
\beta_{\rm rot} = 55.2 \left(\frac{R}{\rm pc}\right)^3 \left(\frac{M_{\rm env}}{1.0 \text{ M}_{\odot}}\right)^{-1} \left(\frac{v_{\rm grad}}{\rm km s^{-1} pc^{-1}}\right)^2 \tag{5}
$$

The terms of the equation are now defined as R being the core radius in pc,  $v_{\text{grad}}$  is the velocity gradient in terms of km s<sup>-1</sup> pc<sup>-1</sup> across this scale, and M<sub>env</sub> is the core/envelope mass in solar masses. We are interested in determining the level of rotational support both at a radius of 0.05 pc (10000AU) to assess the level of rotational support in the envelope as a whole and at a radius of 3200 AU (0.016 pc), the separation of IRS1 from IRS2. We first calculate  $\beta_{\rm rot}$  for the larger scale using the velocity gradient measured across the BHR71 core of  $\sim$ 1.2 km s<sup>-1</sup> pc<sup>-1</sup> and an envelope mass of  $\sim$ 4.6 M<sub>☉</sub> determined for the radius of 0.05 pc from 8  $\mu$ m extinction from [Tobin et al.](#page-28-6) [\(2010b\)](#page-28-6). With these values, we find  $\beta_{\rm rot} = 0.002$  at a radius of 0.05 pc.

To calculate an upper limit of rotational support on the scale of the companion separation (3200 AU; 0.016 pc), we assume that angular momentum is conserved and the velocity gradient is more rapid at a radius 0.016 pc. Scaling the velocity gradient from the ratio of core to separation/inner envelope radius, the velocity gradient at a radius of 0.016 pc (3200 AU) is 4.83 km s<sup>-1</sup> pc<sup>-1</sup>. We then use the mass measured from the ALMA continuum data for both IRS1 and IRS2, which totals 0.7  $M_{\odot}$ , factoring in the likely higher dust temperature toward IRS1. With these numbers, we calculate an upper limit to  $\beta_{\rm rot} = 0.006$ . Note that we call this an upper limit because we assumed that angular momentum was conserved from larger scales, but the velocity gradient across the envelope from ALMA  $N_2D^+$  is consistent with zero. Furthermore, [Tobin et al.](#page-28-4) [\(2011,](#page-28-4) [2012b\)](#page-28-16) argued that for elongated envelopes such as BHR71 projected infall motions can contribute to the observed velocity gradients. Lastly, if the motions do correspond to rotation, inclination is not taken into account, therefore rotation could be higher if that is the origin of the velocity gradient. However, as outlined in Section 3.3 the inclination of the outflows is very likely to be greater than 35◦ therefore the correction to the rotation velocity is likely less than a factor of 2. The  $\beta_{\rm rot} = 0.006$  measurement for

BHR71 places this system toward the top end of the distribution of single systems shown in [Chen](#page-26-10) [et al.](#page-26-10) [\(2012\)](#page-26-10). However, this might not be an entirely fair comparison because those  $\beta_{\rm rot}$  values are calculated at a variety of core radii, some single-dish and some interferometric. We further emphasize that if we evaluate the velocity gradient from the ALMA  $N_2D^+$ , excluding the regions that are likely outflow influenced,  $\beta_{\rm rot}$  could be much lower. Our calculated value for  $\beta_{\rm rot}$  is lower than the one determined by by [Chen et al.](#page-26-3) [\(2008\)](#page-26-3), but they probed a different scale focused on IRS1 using  $N_2H^+$ .

In addition to the apparent velocity gradients, the observations also revealed the presence of two velocity components of the dense gas to the east of IRS1. The NH<sub>3</sub> and  $N_2D^+$  line center velocities are distinctly red-shifted in the regions that overlap with the red-shifted outflow lobe of IRS1. Furthermore, the PV diagrams shown in Figures [12,](#page-38-0) [17,](#page-43-0) and [22](#page-47-0) show that this red-shifted component is also present in equatorial plane of the envelope. Its presence in the equatorial plane is diminished in the velocity maps that are derived from hyperfine fitting of the  $NH_3$  and  $N_2D^+$  in Figures [10,](#page-36-0) [11,](#page-37-0) and [16](#page-42-0) because the blue-shifted component has greater line strength.

This additional line component must not be very broad since it can be observed as a distinct component in the PV diagrams in Figures [12,](#page-38-0) [17,](#page-43-0) and [22.](#page-47-0) The outflow from IRS1 could be inducing bulk motion in the surrounding molecular gas, similar to the simulations of [Offner & Arce](#page-27-32) [\(2014\)](#page-27-32). Furthermore, the red-shifted lobe of the outflow from IRS2 could also be contributing to the redshifted velocities in  $NH_3$  and  $N_2D^+$  on the southeast side of the envelope. The overlap of IRS2 outflow contours in Figures [10,](#page-36-0) [11,](#page-37-0) and [16](#page-42-0) highlight this possibility, despite the region also being coincident with the edge of the blue-shifted outflow cavity from IRS1. Alternatively, the two velocity components could also result from velocity shear that might have been present during the formation of a core. However, the ambiguity of the formation conditions of the BHR71 core and its structure along the line of sight makes the origin of the second velocity component and the interaction of the outflows and dense molecular gas difficult to confidently ascertain.

#### 5.2. Formation of the Binary System

Models that consider the formation of multiple star systems from rotating, collapsing envelopes have difficulty fragmenting for parameters similar to the observed quantities of  $\beta_{\rm rot}$  and  $\Omega$  for BHR71. One of the classic studies of [Burkert & Bodenheimer](#page-26-6) [\(1993\)](#page-26-6) used a rotating collapsing envelope with an m=2 perturbation to break the symmetry. However, they still needed a fairly rapid rotation rate for fragmentation on >1000 AU scales, using  $\Omega$ =7.2×10<sup>-13</sup> s<sup>-1</sup> resulting in  $\beta_{\rm rot} = 0.16$ . Thus, these models had over an order of magnitude more rotation in order to produce fragmentation on the desired scale.

Recent work by [Boss & Keiser](#page-26-21) [\(2014\)](#page-26-21) explored fragmentation with rotating cloud cores including magnetic fields. This work did not find fragmentation on scales >1000 AU for systems with rotation rates  $\langle 10^{-13} \text{ s}^{-1} \rangle$  nor  $\beta_{\text{rot}} < 0.01$ . The systems that most frequently produced fragments on scales >1000 AU in those simulations had the highest rotation rates  $(\Omega = 3 \times 10^{-13} \text{ s}^{-1})$ . Systems still fragmented in the presence of magnetic fields, but the models with the strongest magnetic fields did inhibit fragmentation. Many other models considering core rotation also had difficulty in forming multiple systems with >1000 AU for rotation rates comparable to or exceeding those found in our observations (e.g. [Price & Bate](#page-27-33) [2007;](#page-27-33) [Machida et al.](#page-27-34) [2008\)](#page-27-34).

Putting all this information on the kinematics of BHR71 together and recent numerical studies of multiple star formation with rotation, we find it unlikely that core rotation could have resulted in the formation of the binary system. This is based on the reasonable upper limit to the value

of  $\beta_{\rm rot} = 0.006$ , but in reality the core rotation is nearly zero across the inner envelope, as probed by  $N_2D^+$  and toward the protostars as measured with  $C^{18}O$ . More to the point, the overall inner envelope kinematics are not consistent with ordered rotation when viewed with  $NH_3$  and  $N_2D^+$  at high enough resolution to both resolve the companion and recover the extended emission of the envelope. However, regardless of the level of true rotation in BHR71, the evidence for rotation in the opposite direction on scales < 1000 AU cannot be easily reconciled. On the scales of disks, the Hall effect could theoretically reverse the direction of rotation in the disk [\(Tsukamoto et al.](#page-28-21) [2015;](#page-28-21) [Krasnopolsky et al.](#page-27-35) [2010\)](#page-27-35), but the motions we observe are on >100s of AU scales where the densities should not be high enough for non-ideal MHD effects to operate efficiently [\(Krasnopolsky et al.](#page-27-35) [2010\)](#page-27-35).

Due to the observational inconsistencies with angular momentum conservation and the difficulty of simulations with similar initial conditions to result in wide companion formation, we must consider alternative scenarios for the formation of this binary system. One of the leading alternatives is turbulent fragmentation of the core [\(Padoan & Nordlund](#page-27-8) [2002;](#page-27-8) [Goodwin et al.](#page-27-36) [2004;](#page-27-36) [Offner et al.](#page-27-9) [2010\)](#page-27-9). The turbulent velocity structure of the simulated molecular clouds will create density perturbations throughout the cloud. This creation of over-densities can be efficient enough such that these regions of locally enhanced density exceed the Jeans mass and collapse to become a protostar. Thus, a protostellar core or two adjacent cores formed in the presence of turbulence could lack an ordered rotation pattern and still form a binary system. Simulations find that the initial separations of these cores are typically >500 AU to 1000s of AU [\(Offner et al.](#page-27-9) [2010;](#page-27-9) [Bate](#page-26-22) [2012\)](#page-26-22).

The apparent rotation in opposite directions on small-scales for IRS1 and IRS2 can also result from turbulent fragmentation. Fragmentation can happen in a turbulent cloud without rotation of the envelope/core, and the angular momenta of the collapsing regions will be derived from the net angular momenta in the turbulent velocities [\(Offner et al.](#page-27-9) [2010;](#page-27-9) [Walch et al.](#page-28-22) [2010;](#page-28-22) [Offner et al.](#page-27-37) [2016\)](#page-27-37). The net angular momenta of the density enhancements that formed IRS1 and IRS2 could have been anti-aligned, leading to the opposite rotation directions observed on <1000 AU scales toward IRS1 and IRS2. Turbulent fragmentation is found to be quite likely for other widely separated protostellar multiple systems as well [\(Lee et al.](#page-27-38) [2017;](#page-27-38) [Pineda et al.](#page-27-39) [2015\)](#page-27-39).

The disordered velocity structure of the envelope as viewed in  $N_2D^+$  and  $NH_3$  could also be the result of turbulence in the core, where the bulk motion of turbulent gas could manifest itself as a disordered velocity structure. We suggest this because the outflow is unlikely to produce all the disordered velocity structure that is observed in the core. Specifically, we are referring to the area east of IRS1 as shown in Figure [16,](#page-42-0) where the velocity along the north-south direction goes from red-shifted to blue-shifted and back to red-shifted.

The misaligned angular momentum vectors are not unique to the BHR71 system. The Class 0 protomultiple system IRAS 16293-2422 may have misaligned kinematics in its binary system separated by only ∼600 AU (e.g., [Zapata et al.](#page-28-23) [2013\)](#page-28-23), the IRAS 04191+1523 system with a separation of ∼860 AU has projected angular momentum vectors that differ by  $\sim 90^{\circ}$  [\(Lee et al.](#page-27-38) [2017\)](#page-27-38), and there are more evolved proto-planetary disk systems that also show similar misalignment [\(Stapelfeldt et al.](#page-28-24) [2003;](#page-28-24) [Jensen & Akeson](#page-27-40) [2014;](#page-27-40) [Williams et al.](#page-28-25) [2014;](#page-28-25) [Brinch et al.](#page-26-23) [2016\)](#page-26-23). Thus, the formation of multiple systems with misaligned or even anti-aligned angular momentum vectors may be common [\(Lee et al.](#page-27-41) [2016\)](#page-27-41).

Despite the seeming consistency with turbulent fragmentation, many of the simulations are focused on fragmentation within a parsec-scale molecular cloud. However, BHR71 is an isolated core and

is not located within a larger cloud [\(Bourke et al.](#page-26-24) [1995b\)](#page-26-24). It is therefore unclear if the results from turbulent fragmentation are directly applicable. However, [Walch et al.](#page-28-22) [\(2010\)](#page-28-22) examined the effects of turbulence within isolated cores that are similar in size and mass to BHR71. They found that the turbulence within the core could lead to non-axisymmetric structure and misaligned net angular momenta. But, these particular simulations used very high levels of initial turbulence, the ratio of turbulent to gravitational binding energy in the simulated cores was between ∼0.3 to ∼0.6. Following [Walch et al.](#page-28-26) [\(2012\)](#page-28-26), BHR71 has a ratio of turbulent to gravitational energy of  $\sim 0.04$ , consistent with other star forming cores. Therefore, the levels of internal turbulence do not appear sufficient to fragment the core in the way that it occurs in the study of [Walch et al.](#page-28-22) [\(2010\)](#page-28-22). Offner  $\&$ [Arce](#page-27-32) [\(2014\)](#page-27-32) also examined isolated, turbulent cores in their outflow study; however, fragmentation in those simulations was not witnessed on scales comparable to BHR71 IRS1 and IRS2, but that was also not the focus of their study.

A variation on the theme of turbulent fragmentation is that the shape of the core itself could result in multiple protostars forming. [Bonnell & Bastien](#page-26-25) [\(1993\)](#page-26-25) considered the collapse of cylindrical cloud cores, and due to the shape the ends of the cylinder collapsed first forming two protostars. These protostars later moved closer together, under their mutual gravitational attraction, forming a bound system. The fact that the BHR71 core appears elongated and asymmetric is similar to such a scenario. Moreover, molecular clouds themselves and even individual protostellar envelopes are often filamentary (e.g., André et al. [2010;](#page-26-11) [Tobin et al.](#page-28-8) [2010a;](#page-28-8) [Looney et al.](#page-27-42) [2007\)](#page-27-42). Therefore it is possible that cloud shape played a role in the formation of BHR71 IRS1 and IRS2, but an asymmetric cloud (or portions of it) can only collapse if they are Jeans unstable.

While turbulent fragmentation can explain many of the features we observe in the velocity field, thermal Jeans fragmentation is another possible route for the formation of the binary system. If the gas is cold and dense, it can collapse without rotation or turbulent motions. Such a scenario is consistent with the lack of radial velocity shift between the sources. However, we might still expect the individual protostars to have a common angular momentum vector because of the overall velocity gradient observed in BHR71.

Using our knowledge of the envelope mass, we can calculate the Jeans length for BHR71 to determine if this is a feasible fragmentation mechanism. The Jeans length is approximately

$$
\lambda_J = \frac{c_s}{\sqrt{G\rho}}\tag{6}
$$

where c<sub>s</sub> is the sound speed ( $\sim$ 0.25 km s<sup>-1</sup> for gas at 20 K), G is the gravitation constant, and  $\rho$  is the average density. We calculate the average density from the mass of 4.6  $M_{\odot}$  that is estimated to be enclosed by a radius of 0.05 pc which is  $\sim 6 \times 10^{-19}$  g cm<sup>-3</sup> or n $\sim 10^5$  cm<sup>-3</sup>. With these values, we estimate the Jeans length to be ∼8400 AU, which is much larger than the projected separation of IRS1 and IRS2. Lower temperatures were certainly possible prior to protostar formation (e.g., [Stutz](#page-28-27) [et al.](#page-28-27) [2010;](#page-28-27) [Sadavoy et al.](#page-28-14) [2018\)](#page-28-14) and higher densities as well, but an average density  $10\times$  higher is necessary to make the Jeans length comparable to the current protostar separation. Thus, thermal Jeans fragmentation is not clearly a favored mechanism on its own to form the binary system, and turbulent fragmentation remains the most likely candidate.

## 5.3. Binary Arrangement Along Line of Sight

The structure of the core and relative positions of IRS1 and IRS2 along the line of sight is a source of uncertainty in the interpretation of the kinematics and companion formation. IRS2 appears to be either behind or at the same distance with respect to IRS1 because the outflows do not appear to strongly interact. Such an interaction is expected to be accompanied by a shock, but  $Spitzer H<sub>2</sub>$  line mapping [\(Giannini et al.](#page-27-43) [2011\)](#page-27-43) and Herschel [O I] mapping [\(Nisini et al.](#page-27-44) [2015\)](#page-27-44) do not show strong emission at the location where the IRS2 outflow fans out. Moreover, the blue-shifted lobe of IRS1 exits the globule, as seen in the infrared and optical imaging (e.g., Figure 1), so it may be near the front edge of the globule (at least, not exactly in the center). Therefore IRS2, if it were even closer to us than IRS1, would be really close to the edge, but we see that it is very embedded and its blue lobe is not prominent in the near-infrared like that of IRS1.

The red-shifted outflow from IRS2 does fan out on the southeast side of it, but this is not what would be expected if the outflow was interacting with the blue-shifted outflow from IRS1. If the two outflows were interacting, the red-shifted outflow from IRS2 would be expected to be deflected to the south, the direction of the blue-shifted flow from IRS1. Since such a deflection is not observed, the arrangement of the sources makes the possibility of IRS2 being located much in front of IRS1 unlikely. The extension of the southeastern side of the red-shifted outflow from IRS2 may more likely be related to how it is interacting with and exiting the core; outflows can also change their orientations over time as they accrete material with different angular momenta [\(Offner et al.](#page-27-37) [2016\)](#page-27-37).

The  $C^{18}O$  velocities of IRS1 and IRS2 are approximately the same, as are the velocities of the  $NH<sub>3</sub>$  and  $N<sub>2</sub>D<sup>+</sup>$  gas closely associated with these protostars. Thus, it is unlikely that they are within separate, but nearby cores along the line of sight, despite the second velocity component. With such an arrangement of IRS1 and IRS2, one could formulate a scenario where the actual separation of IRS1 and IRS2 is 2-3× larger (if e.g., IRS2 is located ∼6,000 AU behind IRS1), and assuming that IRS2 still orbits IRS1, the relative velocity of the protostars as measured by  $C^{18}O$  would be mostly tangential to the line of sight making their relative velocities appear smaller. We note, however, that regardless how IRS1 and IRS2 are arranged in three dimensions, the opposite rotation directions for IRS1 and IRS2 are still inconsistent with rotation induced fragmentation.

#### 5.4. Future Observations to Characterize BHR71

While it is has become somewhat cliche to suggest that further observations are needed to understand a particular system, we outline here a few specific missing pieces of the puzzle that would enable our ideas regarding fragmentation in BHR71 to be confirmed or ruled-out.

First, the evidence for opposite rotation directions toward IRS1 and IRS2 in  $C^{18}O$ , which makes rotational fragmentation seem less likely, could be improved. A factor of 2-4 increases in both sensitivity and resolution would solidify this result. The present observations only have a resolution of ∼1".5, and the effective time on source for the protostars was only ∼62 seconds on each mosaic point.

Second, the envelope kinematics on scales between the  $N_2D^+$  and  $C^{18}O$  need improved resolution to better connect the kinematics from  $>1000$  AU scales to  $<1000$  AU scales where  $C^{18}O$  is the best tracer. [Chen et al.](#page-26-3) [\(2008\)](#page-26-3) published ATCA  $N_2H^+$  observations that had severe limitations of sensitivity and spatial filtering. However, the regions traced in  $N_2H^+$  are coincident with part of the  $N_2D^+$  emission. Further observations of  $N_2H^+$  with ALMA at high-sensitivity and with coverage of both large and small spatial scales would further illuminate the likely opposite rotation of gas near the protostars. While the NH<sub>3</sub> traces down to about the same scale as  $N_2H^+$  and  $N_2D^+$ , observations

of NH<sup>3</sup> cannot currently be obtained at significantly higher resolution due to spatial filtering and lower sensitivity of ATCA. Thus,  $N_2H^+$  with ALMA is the best option going forward.

Third, higher sensitivity observations that could detect the  $NH<sub>3</sub>$  (2,2) transition would be helpful in order to determine the gas kinetic temperature. However, higher sensitivity and resolution  $NH_3$ observations are not trivial because ATCA is currently the only interferometer in the southern hemisphere that can observe the  $NH<sub>3</sub>$  inversion transitions. Knowledge of the gas kinetic temperature could enable the susceptibility of the gas around the protostars to thermal Jeans fragmentation to be assessed. But, such measurements of kinetic temperature may also be possible using  $H_2CO$  [\(Mangum](#page-27-45) [& Wootten](#page-27-45) [1993\)](#page-27-45), if present in the envelope gas surrounding the protostars.

## 6. SUMMARY AND CONCLUSIONS

We have conducted a multi-wavelength and multi-line study of the isolated proto-binary system located within BHR71, unleashing a large battery of ground-based southern and space-based observatories on this system. We have resolved the two protostars IRS1 and IRS2 in the infrared out to 160  $\mu$ m using *Herschel* PACS photometry and conducted the most complete assessment of the SED toward IRS2. We also fully mapped the BHR71 core using Herschel SPIRE mapping. We have further characterized the continuum and kinematic properties of the envelope surrounding both sources, in addition to the individual envelopes around each of IRS1 and IRS2. This was done using ATCA and Parkes to observe NH<sub>3</sub> (1,1) emission from the cold gas, and ALMA to image  $N_2D^+$  and C <sup>18</sup>O in addition to outflow tracers.

Our main results are as follows:

- With observations that resolve the protostars out to 160  $\mu$ m and PSF photometry at 250  $\mu$ m, we find that both are Class 0 protostars, as expected. IRS1 has a bolometric luminosity and temperature of 14.7  $L_{\odot}$  and 68 K, respectively, while IRS2 has values of 1.7  $L_{\odot}$  and 38 K. Thus, IRS2 could be less evolved than IRS1, but there are systematics related to source inclination that add uncertainty to  $T_{bol}$ .
- We resolve the continuum emission toward each protostar at 1.3 mm and 1.3 cm. We find that the radio spectra can be consistent with dust-only emission down to 1.3 cm for both IRS1 and IRS2, even with the 3 mm point from [\(Chen et al.](#page-26-3) [2008\)](#page-26-3) included, but we cannot exclude some free-free emission at 1.3 cm. The total mass measured around the two protostars is 1.25  $M_{\odot}$ , assuming a dust temperature of 20 K for both protostars, and the masses of IRS1 and IRS2 individually are 1.13  $M_{\odot}$  and 0.11  $M_{\odot}$ , respectively with an assumed dust to gas mass ratio of 1:100 and [Ossenkopf & Henning](#page-27-24) [\(1994\)](#page-27-24) dust. If the dust temperature around IRS1 is 34 K, accounting for the higher luminosity, then the mass of IRS1 would be 0.59  $M_{\odot}$ , making the combined mass 0.7  $M_{\odot}$ . This continuum mass is a lower limit since there is substantially more cold dust surrounding two protostars as viewed by *Herschel*. ALMA simply picks up the brightest emission close to each protostar. There is also weak evidence for a bridge of material between IRS1 and IRS2, perhaps similar to what is observed in IRAS 16293-2422 [\(Jacobsen](#page-27-22) [et al.](#page-27-22) [2018\)](#page-27-22).
- The Parkes and ATCA observations seem to indicate a smooth velocity gradient across the core that could be interpreted as rotation. However, the ALMA  $N_2D^+$  observations at higher resolution reveal a very complex velocity field that is not smooth. Position-velocity diagrams

from both ALMA  $N_2D^+$  and ATCA  $NH_3$  find a second velocity component on the envelope east of IRS1. Some of this second component is likely to result from the outflow influence on the envelope, but we also find that there is no clear velocity gradient across the envelope midplane from IRS1 to IRS2.

- The ALMA observations also clearly detect <sup>12</sup>CO, <sup>13</sup>CO, and  $C^{18}O$ . <sup>12</sup>CO and <sup>13</sup>CO mainly trace the outflow emission from the protostars, while  $C^{18}O$  traces some emission along the outflow, but also the kinematics of the envelopes surrounding IRS1 and IRS2 on <1000 AU scales. The C<sup>18</sup>O velocities indicate that IRS1 and IRS2 are at the same velocity along the line of sight, so there is no evidence of orbital motion in their respective radial velocities. Moreover, the apparent rotation of the inner envelopes around IRS1 and IRS2 appear to be in the *opposite* directions!
- We conclude that the binary formation in BHR71 is unlikely to have resulted from rotationallyinduced fragmentation of the core. The upper limit on the ratio of rotational energy to gravitational potential energy is low ( $\beta_{\rm rot} = 0.006$ ). This value is among the low end of the distribution presented in [Chen et al.](#page-26-10) [\(2012\)](#page-26-10). Furthermore, the  $N_2D^+$  velocity field across IRS1 and IRS2, in the equatorial plane of the envelope is consistent with no velocity gradient beyond the separation of the pair. There is no evidence of orbital motion given that the sources have the same systemic velocities along the line of sight, and the apparent rotation of the inner envelopes on scales less than 1000 AU is in the opposite direction for IRS1 and IRS2. Thus, with all the evidence against ordered rotation leading to the formation of the binary system, we conclude that turbulent fragmentation might be the most likely formation scenario, given that the envelope has a velocity field that is not well-ordered.

## Facility: ATCA, Parkes, ALMA, Herschel, Spitzer, CTIO(ISPI), Magellan (PANIC)

Software: Astropy(http://www.astropy.org;[AstropyCollaborationetal.2018;](#page-26-26)[Greenfieldetal.2013\)](#page-27-46), APLpy (http://aplpy.github.com; [Robitaille &Bressert](#page-28-28) [2012\)](#page-28-28), CASA (http://casa.nrao.edu; [McMullin](#page-27-47) [et al.](#page-27-47) [2007\)](#page-27-47), The IDL Astronomy User's Library (https://idlastro.gsfc.nasa.gov/)

We acknowledge the constructive report from the anonymous referee which improved the quality of the manuscript. We also acknowledge useful discussions regarding BHR71 with M. Dunham and C. Hull, as well as help with Starfinder by N. Murillo. J.J.T. acknowledges support from NSF grant AST-1814762, the Homer L. Dodge Endowed Chair, and grant 639.041.439 from the Netherlands Organisation for Scientific Research (NWO). H.G.A. acknowledges the support of NSF grant AST-1714710. X.C. acknowledges support by the NSFC through grant 11473069. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2013.1.00518.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), NSC and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. This work is based in part on observations made with Herschel, a European Space Agency Cornerstone Mission with significant participation by NASA. Support for this work was provided by NASA through an award issued by JPL/Caltech. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This research made use of APLpy, an open-source plotting package for Python hosted at http://aplpy.github.com.

This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration, 2013) http://www.astropy.org.

## APPENDIX

## A. HYPERFINE FITS

We show examples of the hyperfine fitting from CLASS for  $NH_3$  and  $N_2D^+$  in Figure [23](#page-48-0) and [24.](#page-49-0) The fits show that there is evidence of a second component in some positions of the NH<sub>3</sub> and N<sub>2</sub>D<sup>+</sup> velocity maps shown in Figures [10](#page-36-0) and [16,](#page-42-0) respectively. The east side of the envelope in both  $NH<sub>3</sub>$ and  $N_2D^+$  shows two components in Figures [12](#page-38-0) and [17,](#page-43-0) respectively. These appear in the hyperfine fits as broader linewidth because a single velocity component is adopted in the fitting process.

We further examined the issue of the second velocity component for the fit to the east side of the envelope, where the second component causes the most visible change to the linewidth (Figure [25\)](#page-50-0). When we include a second component to the fit, we do find that the linewidths are indeed lower for each velocity component. We do not adopt a two component fit for the entire envelope because the two component fit often requires fine tuning to obtain acceptable results. Thus a two-component fit is not feasible to use over the entire spectral cube.

#### REFERENCES

- <span id="page-26-4"></span>André, P., Ward-Thompson, D., & Barsony, M. 1993, ApJ, 406, 122
- <span id="page-26-11"></span>André, P., Men'shchikov, A., Bontemps, S., et al. 2010, A&A, 518, L102
- <span id="page-26-5"></span>Arce, H. G., & Sargent, A. I. 2006, ApJ, 646, 1070
- <span id="page-26-26"></span>Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package, , , arXiv:1801.02634
- <span id="page-26-22"></span>Bate, M. R. 2012, MNRAS, 419, 3115
- <span id="page-26-16"></span>Benson, P. J., & Myers, P. C. 1989, ApJS, 71, 89
- <span id="page-26-18"></span>Bergin, E. A., Alves, J., Huard, T., & Lada, C. J. 2002, ApJL, 570, L101
- <span id="page-26-15"></span>Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, ApJ, 224, 132
- <span id="page-26-25"></span>Bonnell, I., & Bastien, P. 1993, ApJ, 406, 614
- <span id="page-26-7"></span>Boss, A. P. 1995, in Revista Mexicana de Astronomia y Astrofisica Conference Series, Vol. 1, Revista Mexicana de Astronomia y Astrofisica Conference Series, ed. S. Lizano & J. M. Torrelles, 165
- <span id="page-26-8"></span>Boss, A. P. 2002, ApJ, 568, 743
- <span id="page-26-19"></span>Boss, A. P., & Bodenheimer, P. 1979, ApJ, 234, 289
- <span id="page-26-9"></span>Boss, A. P., & Keiser, S. A. 2013, ApJ, 764, 136
- <span id="page-26-21"></span>—. 2014, ArXiv e-prints, arXiv:1408.2479
- <span id="page-26-2"></span>Bourke, T. L. 2001, ApJL, 554, L91
- <span id="page-26-1"></span>Bourke, T. L., Hyland, A. R., & Robinson, G. 1995a, MNRAS, 276, 1052
- <span id="page-26-24"></span>Bourke, T. L., Hyland, A. R., Robinson, G., James, S. D., & Wright, C. M. 1995b, MNRAS, 276, 1067
- <span id="page-26-23"></span>Brinch, C., Jørgensen, J. K., Hogerheijde, M. R., Nelson, R. P., & Gressel, O. 2016, ApJL, 830, L16
- <span id="page-26-6"></span>Burkert, A., & Bodenheimer, P. 1993, MNRAS, 264, 798
- <span id="page-26-17"></span>Caselli, P., Benson, P. J., Myers, P. C., & Tafalla, M. 2002, ApJ, 572, 238
- <span id="page-26-13"></span>Chen, H., Myers, P. C., Ladd, E. F., & Wood, D. O. S. 1995, ApJ, 445, 377
- <span id="page-26-10"></span>Chen, X., Arce, H. G., Dunham, M. M., & Zhang, Q. 2012, ApJL, 747, L43
- <span id="page-26-3"></span>Chen, X., Launhardt, R., Bourke, T. L., Henning, T., & Barnes, P. J. 2008, ApJ, 683, 862
- <span id="page-26-20"></span>Chen, X., Launhardt, R., & Henning, T. 2007, ApJ, 669, 1058
- <span id="page-26-0"></span>Chen, X., Arce, H. G., Zhang, Q., et al. 2013, ApJ, 768, 110
- <span id="page-26-14"></span>Chiang, H., Looney, L. W., & Tobin, J. J. 2012, ApJ, 709, 470
- <span id="page-26-12"></span>Diolaiti, E., Bendinelli, O., Bonaccini, D., et al. 2000, in Proc. SPIE, Vol. 4007, Adaptive Optical Systems Technology, ed. P. L. Wizinowich, 879–888
- <span id="page-27-31"></span>Dore, L., Caselli, P., Beninati, S., et al. 2004, A&A, 413, 1177
- <span id="page-27-18"></span>Drabek, E., Hatchell, J., Friberg, P., et al. 2012, MNRAS, 426, 23
- <span id="page-27-1"></span>Duchêne, G., & Kraus, A. 2013, ARA&A, 51, 269
- <span id="page-27-5"></span>Dunham, M. M., Stutz, A. M., Allen, L. E., et al. 2014, Protostars and Planets VI, 195
- <span id="page-27-11"></span>Fazio, G. G., Hora, J. L., Allen, L. E., et al. 2004, ApJS, 154, 10
- <span id="page-27-26"></span>Frerking, M. A., Langer, W. D., & Wilson, R. W. 1982, ApJ, 262, 590
- <span id="page-27-16"></span>Furlan, E., Fischer, W. J., Ali, B., et al. 2016, ApJS, 224, 5
- <span id="page-27-43"></span>Giannini, T., Nisini, B., Neufeld, D., et al. 2011, ApJ, 738, 80
- <span id="page-27-29"></span>Goodman, A. A., Benson, P. J., Fuller, G. A., & Myers, P. C. 1993, ApJ, 406, 528
- <span id="page-27-36"></span>Goodwin, S. P., Whitworth, A. P., & Ward-Thompson, D. 2004, A&A, 414, 633
- <span id="page-27-46"></span>Greenfield, P., Robitaille, T., Tollerud, E., et al. 2013, ascl:1304.002
- <span id="page-27-15"></span>Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, A&A, 518, L3
- <span id="page-27-7"></span>Hacar, A., & Tafalla, M. 2011, A&A, 533, A34+
- <span id="page-27-28"></span>Herbst, E. 1982, A&A, 111, 76
- <span id="page-27-13"></span>Houck, J. R., Roellig, T. L., van Cleve, J., et al. 2004, ApJS, 154, 18
- <span id="page-27-22"></span>Jacobsen, S. K., Jørgensen, J. K., van der Wiel, M. H. D., et al. 2018, A&A, 612, A72
- <span id="page-27-40"></span>Jensen, E. L. N., & Akeson, R. 2014, Nature, 511, 567
- <span id="page-27-30"></span>Johnstone, D., Rosolowsky, E., Tafalla, M., & Kirk, H. 2010, ApJ, 711, 655
- <span id="page-27-35"></span>Krasnopolsky, R., Li, Z.-Y., & Shang, H. 2010, ApJ, 716, 1541
- <span id="page-27-2"></span>Kraus, A. L., Ireland, M. J., Martinache, F., & Hillenbrand, L. A. 2011, ApJ, 731, 8
- <span id="page-27-23"></span>Kwon, W., Looney, L. W., Mundy, L. G., Chiang, H.-F., & Kemball, A. J. 2009, ApJ, 696, 841
- <span id="page-27-3"></span>Lada, C. J., & Lada, E. A. 2003, ARA&A, 41, 57
- <span id="page-27-20"></span>Ladd, E. F., Adams, F. C., Fuller, G. A., et al. 1991, ApJ, 382, 555
- <span id="page-27-6"></span>Larson, R. B. 1972, MNRAS, 156, 437
- <span id="page-27-48"></span>Launhardt, R., Nutter, D., Ward-Thompson, D., et al. 2010, ApJS, 188, 139
- <span id="page-27-27"></span>Lee, J., Bergin, E. A., & Evans, II, N. J. 2004, ApJ, 617, 360
- <span id="page-27-38"></span>Lee, J.-E., Lee, S., Dunham, M. M., et al. 2017, Nature Astronomy, 1, 0172
- <span id="page-27-41"></span>Lee, K. I., Dunham, M. M., Myers, P. C., et al. 2016, ApJL, 820, L2
- <span id="page-27-42"></span>Looney, L. W., Tobin, J. J., & Kwon, W. 2007, ApJL, 670, L131
- <span id="page-27-34"></span>Machida, M. N., Tomisaka, K., Matsumoto, T., & Inutsuka, S. 2008, ApJ, 677, 327
- <span id="page-27-17"></span>Magnelli, B., Popesso, P., Berta, S., et al. 2013, A&A, 553, A132
- <span id="page-27-45"></span>Mangum, J. G., & Wootten, A. 1993, ApJS, 89, 123
- <span id="page-27-10"></span>Martini, P., Persson, S. E., Murphy, D. C., et al. 2004, in Proc. SPIE, Vol. 5492, Ground-based Instrumentation for Astronomy, ed. A. F. M. Moorwood & M. Iye, 1653–1660
- <span id="page-27-47"></span>McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, in Astronomical Society of the Pacific Conference Series, Vol. 376, Astronomical Data Analysis Software and Systems XVI, ed. R. A. Shaw, F. Hill, & D. J. Bell, 127
- <span id="page-27-4"></span>Megeath, S. T., Gutermuth, R., Muzerolle, J., et al. 2012, AJ, 144, 192
- <span id="page-27-21"></span>Myers, P. C., Adams, F. C., Chen, H., & Schaff, E. 1998, ApJ, 492, 703
- <span id="page-27-19"></span>Myers, P. C., & Ladd, E. F. 1993, ApJL, 413, L47
- <span id="page-27-44"></span>Nisini, B., Santangelo, G., Giannini, T., et al. 2015, ApJ, 801, 121
- <span id="page-27-32"></span>Offner, S. S. R., & Arce, H. G. 2014, ApJ, 784, 61
- <span id="page-27-37"></span>Offner, S. S. R., Dunham, M. M., Lee, K. I., Arce, H. G., & Fielding, D. B. 2016, ApJL, 827, L11
- <span id="page-27-9"></span>Offner, S. S. R., Kratter, K. M., Matzner, C. D., Krumholz, M. R., & Klein, R. I. 2010, ApJ, 725, 1485
- <span id="page-27-24"></span>Ossenkopf, V., & Henning, T. 1994, A&A, 291, 943
- <span id="page-27-8"></span>Padoan, P., & Nordlund, Å. 2002, ApJ, 576, 870
- <span id="page-27-25"></span>Parise, B., Belloche, A., Leurini, S., et al. 2006, A&A, 454, L79
- <span id="page-27-39"></span>Pineda, J. E., Offner, S. S. R., Parker, R. J., et al. 2015, Nature, 518, 213
- <span id="page-27-14"></span>Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, A&A, 518, L2
- <span id="page-27-33"></span>Price, D. J., & Bate, M. R. 2007, MNRAS, 377, 77
- <span id="page-27-0"></span>Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, ApJS, 190, 1
- <span id="page-27-12"></span>Rieke, G. H., Young, E. T., Engelbracht, C. W., et al. 2004, ApJS, 154, 25
- <span id="page-28-28"></span>Robitaille, T., & Bressert, E. 2012, APLpy: Astronomical Plotting Library in Python, Astrophysics Source Code Library, , , ascl:1208.017
- <span id="page-28-13"></span>Roussel, H. 2013, PASP, 125, 1126
- <span id="page-28-14"></span>Sadavoy, S. I., Keto, E., Bourke, T. L., et al. 2018, ApJ, 852, 102
- <span id="page-28-7"></span>Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in Astronomical Society of the Pacific Conference Series, Vol. 77, Astronomical Data Analysis Software and Systems IV, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 433
- <span id="page-28-19"></span>Shu, F. H. 1977, ApJ, 214, 488
- <span id="page-28-10"></span>Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163
- <span id="page-28-24"></span>Stapelfeldt, K. R., Ménard, F., Watson, A. M., et al. 2003, ApJ, 589, 410
- <span id="page-28-27"></span>Stutz, A., Launhardt, R., Linz, H., et al. 2010, A&A, 518, L87
- <span id="page-28-20"></span>Terebey, S., Shu, F. H., & Cassen, P. 1984, ApJ, 286, 529
- <span id="page-28-11"></span>Tobin, J. J., Bos, S. P., Dunham, M. M., Bourke, T. L., & van der Marel, N. 2018, ApJ, 856, 164
- <span id="page-28-5"></span>Tobin, J. J., Hartmann, L., Bergin, E., et al. 2012a, ApJ, 748, 16
- <span id="page-28-16"></span>Tobin, J. J., Hartmann, L., Chiang, H.-F., et al. 2012b, Nature, 492, 83
- <span id="page-28-8"></span>Tobin, J. J., Hartmann, L., Looney, L. W., & Chiang, H. 2010a, ApJ, 712, 1010
- <span id="page-28-6"></span>Tobin, J. J., Hartmann, L., Looney, L. W., & Chiang, H.-F. 2010b, ApJ, 712, 1010
- <span id="page-28-4"></span>Tobin, J. J., Hartmann, L., Chiang, H.-F., et al. 2011, ApJ, 740, 45
- <span id="page-28-17"></span>Tobin, J. J., Bergin, E. A., Hartmann, L., et al. 2013, ApJ, 765, 18
- <span id="page-28-1"></span>Tobin, J. J., Looney, L. W., Li, Z.-Y., et al. 2016, ApJ, 818, 73
- <span id="page-28-2"></span>Tohline, J. E. 2002, ARA&A, 40, 349
- <span id="page-28-21"></span>Tsukamoto, Y., Iwasaki, K., Okuzumi, S., Machida, M. N., & Inutsuka, S. 2015, ApJL, 810, L26
- <span id="page-28-9"></span>van der Bliek, N. S., Norman, D., Blum, R. D., et al. 2004, in Proc. SPIE, Vol. 5492, Ground-based Instrumentation for Astronomy, ed. A. F. M. Moorwood & M. Iye, 1582–1589
- <span id="page-28-15"></span>Visser, R., Doty, S. D., & van Dishoeck, E. F. 2011, A&A, 534, A132
- <span id="page-28-22"></span>Walch, S., Naab, T., Whitworth, A., Burkert, A., & Gritschneder, M. 2010, MNRAS, 402, 2253
- <span id="page-28-26"></span>Walch, S., Whitworth, A. P., & Girichidis, P. 2012, MNRAS, 419, 760
- <span id="page-28-0"></span>Ward-Duong, K., Patience, J., De Rosa, R. J., et al. 2015, MNRAS, 449, 2618
- <span id="page-28-12"></span>Whitney, B. A., Wood, K., Bjorkman, J. E., & Wolff, M. J. 2003, ApJ, 591, 1049
- <span id="page-28-25"></span>Williams, J. P., Mann, R. K., Di Francesco, J., et al. 2014, ApJ, arXiv:1410.3570
- <span id="page-28-3"></span>Yang, Y.-L., Evans, II, N. J., Green, J. D., Dunham, M. M., & Jørgensen, J. K. 2017, ApJ, 835, 259
- <span id="page-28-18"></span>Yen, H.-W., Takakuwa, S., Ohashi, N., & Ho, P. T. P. 2013, ApJ, 772, 22
- <span id="page-28-23"></span>Zapata, L. A., Loinard, L., Rodríguez, L. F., et al. 2013, ApJL, 764, L14



Figure 1. False color images of BHR71 at multiple wavelengths. The left panel shows BHR71 as viewed in the near-infrared in Ks (PANIC), H (PANIC), and J-bands (ISPI). The outflow cavity from the blue-shifted outflow of IRS1 is quite prominent and the outflow cavity from IRS2 is not distinct, but several knots are apparent in the northwest portion of the image. The middle panel shows the Spitzer 3.6, 4.5, and 8.0  $\mu$ m images on the same scale. IRS1 is still dominant, but IRS2 is much more apparent. The right panel shows the Spitzer 24  $\mu$ m, along with Herschel 70 and 100  $\mu$ m imaging that clearly resolve the mid and far-infrared emission directly associated with both IRS1 and IRS2. IRS1 is quite prominent at all three wavelengths, but IRS2 is not strongly detected until 70 and 100  $\mu$ m due to its deeply embedded nature. All images map the longest wavelength to red, the intermediate wavelength to green, and the shortest wavelength to blue.



**Figure 2.** False-color images of BHR71 showing a larger-scale view from *Spitzer* at 3.6, 4.5, and 8.0  $\mu$ m (left panel). The extended envelope is apparent in absorption against the 8.0  $\mu$ m infrared background. The right panel shows the submillimeter view of BHR71 at 250, 350, and 500  $\mu$ m covering a field of view twice as large as in the left image. There is a red feature at 500  $\mu$ m in the outflow direction that is possibly contamination from CO  $(J = 5 \rightarrow 4)$  emission. The core surrounding the protostars is quite extended and well-resolved even at the longest wavelengths.



<span id="page-30-0"></span>Figure 3. Spectral energy distribution (SED) from the near-infrared to the millimeter constructed from photometry presented in this work, as well as photometry and spectroscopy from the literature. From the SEDs, IRS1 and IRS2 are measured to have  $L_{bol} = 14.7, 1.7 L_{\odot}$  and  $T_{bol} = 68, 38 K$ , respectively. The two protostars are well-resolved shortward of  $70 \mu m$  and at wavelengths where they can be observed with interferometry. The fluxes toward IRS1 and IRS2 are measured with PSF photometry between 70 to 250  $\mu$ m. IRS1 is plotted with square symbols, IRS2 is plotted with triangles, and the combined SED is plotted with the diamonds. IRS1 clearly dominates the SED, but IRS2 has a lower  $T_{\text{bol}}$ , and the SED is seen to peak at slightly longer wavelengths than IRS1 due to its more deeply embedded nature. The solid line is the Spitzer IRS spectrum from [Yang et al.](#page-28-3) [\(2017\)](#page-28-3).



Figure 4. ALMA 1.3 mm and ATCA 1.3 cm contours overlaid on the *Spitzer* 8  $\mu$ m image (grayscale) are shown in the left and right panels, respectively. The ALMA image at higher resolution shows the dust emission corresponding to the location of the brightest 8  $\mu$ m emission toward BHR71 IRS1, but toward IRS2, the 1.3 mm emission is located at the base of the 8  $\mu$ m emission. The same behavior is also shown by the ATCA 1.3 cm image. The ATCA image might be tracing a combination of both free-free and dust emission. The ALMA 1.3 mm emission shows extended features; IRS1 shows features extended apparently along the outflow cavity walls, and there is an extension of emission from IRS1 toward the location of IRS2. The contours in both panels start at  $3\sigma$  and increase on  $3\sigma$  intervals, where  $\sigma_{1.3mm} = 0.5$  mJy beam<sup>-1</sup> and  $\sigma_{1.3cm} = 0.1 \text{ mJy beam}^{-1}.$ 



<span id="page-31-0"></span>Figure 5. Millimeter to centimeter spectrum of BHR71 IRS1 and IRS2. Both are compatible with dust emission down to 1.3 cm, but we cannot rule out a contribution of free-free emission at 1.3 cm toward both protostars. The 3.1 mm data point toward IRS2 seems quite low compared to the others, given that it would indicate an unphysically steep spectral slope, but this is also the lowest S/N data point.



<span id="page-32-0"></span>**Figure 6.** Integrated intensity maps of <sup>12</sup>CO, <sup>13</sup>CO and C<sup>18</sup>O ( $J = 2 \rightarrow 1$ ) overlaid on the ALMA 1.3 mm continuum image (grayscale). The  ${}^{12}CO$  and  ${}^{13}CO$  are divided into red and blue-shifted components, with corresponding contour colors; the  $C^{18}O$  includes both the blue and red components. The <sup>12</sup>CO and <sup>13</sup>CO clearly show the outflow emission from the protostars in both lines, with the red-shifted lobe of the IRS2 outflow being quite broad. The outflow cavities are clearly not axisymmetric, but the IRS1 cavity especially appears to show reflection symmetry. The C18O is peaked toward both protostars, tracing the warm inner envelopes. The <sup>12</sup>CO is integrated between -15 to -5.6 km s<sup>-1</sup> and -3 to 4 km s<sup>-1</sup>, and  $\sigma_{blue, red} = 0.46$ , 0.4 K km s<sup>-1</sup>, respectively. The blue and red contours start  $20\sigma$  and increase at this same interval. The <sup>13</sup>CO is integrated between -7 to -4.7 km s<sup>-1</sup> and -3.2 to -1.5 km s<sup>-1</sup>, and  $\sigma_{blue, red} = 0.13, 0.15$  K km s<sup>-1</sup>, respectively, with the blue contours starting at  $50\sigma$  and increasing on  $20\sigma$  intervals, while the red contours start 20 $\sigma$  and increase at this same interval. The C<sup>18</sup>O is integrated between -5.8 to -3.2 km s<sup>-1</sup>, and  $\sigma$  = 0.13 K km s<sup>-1</sup> and the contours start at  $15\sigma$  and increase on  $10\sigma$  intervals.



<span id="page-33-0"></span>Figure 7. Spectra toward the positions of BHR71 IRS1 and IRS2 for each molecule observed. The ALMA  $spectra are extracted from a 5'' circle centered on the protostars, the ATCA spectra are extracted from a circle$ matching the synthesized beam  $(\sim 10'')$  centered on the protostars, and the Parkes spectrum is extracted from a 60" circle centered on IRS1. ALMA and ATCA  $N_2H^+$  and  $N_2D^+$  spectra are also extracted from the emission peaks closest to IRS1 and IRS2 in the panels containing the text 'Peak Positions.' Some spectra are shown in different velocity ranges to highlight different features of the spectra.



<span id="page-34-0"></span>Figure 8. Parkes single-dish  $NH_3$  (1,1) maps. The upper left panel shows the integrated intensity of the main NH<sub>3</sub> (1,1) hyperfine lines overlaid on the *Spitzer* 8  $\mu$ m image (color scale). The upper right panel shows just the integrated intensity of the main NH<sup>3</sup> (1,1) hyperfine lines (color scale and contours). The lower left panel shows the line center velocity derived from a fit to all the  $NH<sub>3</sub>(1,1)$  hyperfine lines, and the lower right panel shows the linewidth also derived from fitting all hyperfine components. The integrated intensity peaks toward the position of IRS2, and the velocity map shows a very small gradient across the source, only about a 0.15 km s−<sup>1</sup> differential. The linewidth map shows that the lines are narrow, at most only about 0.6 km s−<sup>1</sup> wide. The positions of the IRS1 and IRS2 are marked with X's. The intensity levels in the top panels are from the integrated intensity map of the main hyperfine lines between -6.5 to -2.2 km s<sup>-1</sup>; the intensity levels start at and increase on  $3\sigma$  intervals, where  $\sigma$ =0.045 K km s<sup>-1</sup>.



<span id="page-35-0"></span>Figure 9. The integrated intensity contours from the ATCA  $NH<sub>3</sub>(1,1)$  observations (green) are overlaid on the Spitzer 8  $\mu$ m image (inverse grayscale). The integrated intensity of the main NH<sub>3</sub> lines is shown in the left panel and the satellite lines in the right panel (between -12.6 to -11.4 km s<sup>-1</sup>). We also overlay the  $12$ CO red and blue-shifted integrated intensity contours (with corresponding colors) from Figure [6,](#page-32-0) showing the relationship of the outflow with the dense gas. The  $NH<sub>3</sub>(1,1)$  emission at this higher resolution shows excellent correspondence with the  $8 \mu m$  absorption feature of the envelope on the background, and the NH<sub>3</sub> maps exhibit a cavity and depressions toward the location of IRS1. IRS2 is located directly beside a NH<sup>3</sup> peak. The NH<sub>3</sub> main and (satellite) contours start at  $5\sigma$  ( $3\sigma$ ) and increase on  $5\sigma$  ( $3\sigma$ ) intervals, where  $\sigma$ =0.075 K km s<sup>-1</sup> (0.072 K km s<sup>-1</sup>), a -5 $\sigma$  (-3 $\sigma$ ) contour is also drawn.



<span id="page-36-0"></span>Figure 10. Same as for Figure [8,](#page-34-0) but for the ATCA NH<sub>3</sub> observations. The spatial extents of the outflows probed by ALMA in <sup>12</sup>CO are marked with the gray contours in all panels but the top left. The velocity map shows considerably more structure than the Parkes map. There is a red-shifted component that appears to reflect the red-shifted outflow cavity of IRS1, then there is an abrupt velocity gradient between IRS1 and IRS2. Some of the velocity gradient is likely related to the outflow influence, but even without the additional velocity structure of the outflow, the gradient is quite strong. Furthermore, the gradient also appears as a region of large linewidth which results from the line velocity changing on the scale of one beamwidth. There is also some evidence of outflow influence on the south side where there is increased linewidth where the eastern edge of the IRS1 outflow cavity is located, in addition to the red-shifted side of the IRS2 outflow. However, there is no increased linewidth toward the red-shifted side of the outflow where there appears to be a shift in the line-center map. The intensity levels in the top panels are from the integrated intensity map of the main hyperfine lines between -5.0 to -3.5 km s<sup>-1</sup>; the intensity levels start at  $3\sigma$  and increase on  $6\sigma$  intervals, where  $\sigma$ =0.07 K km s<sup>-1</sup>. The dashed lines in the upper right panel marks the region of position-velocity (PV) extraction.



<span id="page-37-0"></span>Figure 11. Same as Figure [10](#page-36-0) but zooming in. The dashed circles in the upper right panel correspond to regions of spectral extraction to examine the quality of hyperfine fitting, see Appendix A.



<span id="page-38-0"></span>Figure 12. Position-velocity diagrams of  $NH_3$  taken in a 10" wide strip in the east-west direction, centered on IRS1. The region extracted is bounded by the dashed lines in the upper right panel of Figure [11.](#page-37-0) The top panel shows the PV diagram of the main NH<sup>3</sup> hyperfine lines, and the bottom panel shows a PV diagram of a set of NH<sup>3</sup> satellite hyperfine lines that are not as closely spaced as the main lines.



<span id="page-39-0"></span>Figure 13. Position-velocity diagrams of NH<sub>3</sub> taken in a 10<sup> $\prime\prime$ </sup> wide strip along a position angle of 123 $\degree$  east of north, centered on IRS1. The region extracted is bounded by the dashed lines in the upper right panel of Figure [10.](#page-36-0) The top panel shows the PV diagram of the main NH<sup>3</sup> hyperfine lines, and the bottom panel shows a PV diagram of a set of NH<sub>3</sub> satellite hyperfine lines that are not as closely spaced as the main lines.



<span id="page-40-0"></span>Figure 14. Velocity profiles extracted orthogonal to the outflow directions for IRS1 (top row) and IRS2 (bottom row). Velocity profiles are shown for both NH<sub>3</sub> (1,1) (left panels) and N<sub>2</sub>D<sup>+</sup> (right panels). All panels also show the profile extracted from the Parkes NH<sub>3</sub> data (diamonds) which is fit with a line (solid line). The single-dish data show a gradual velocity gradient and the interferometer data (plus signs) show more structure. The  $N_2D^+$  data may have different centroid velocities relative to the NH<sub>3</sub>. This could be due to  $N_2D^+$  tracing a different region of the cloud due to chemistry. The velocity gradients in the interferometer data are fit (dashed lines), but their structure limits the utility of the values derived.



<span id="page-41-0"></span>**Figure 15.** The integrated intensity contours from the ALMA  $N_2D^+$  ( $J = 3 \rightarrow 2$ ) observations are overlaid (green) on the *Spitzer* 8.0  $\mu$ m image (inverse grayscale) in the left panel and the N<sub>2</sub>D<sup>+</sup> is overlaid (green) on the ATCA  $NH_3$  (1,1) integrated intensity map (inverse grayscale) in the right panel. We also overlaid the <sup>12</sup>CO blue- and red-shifted integrated intensity contours (with corresponding colors) from Figure [6.](#page-32-0) The extent of  $C^{18}O$  shown in Figure [6](#page-32-0) is consistent with the size of the  $N_2D^+$  deficit regions. There is some correspondence on the east side of the  $N_2D^+$  and the 8  $\mu$ m absorption of the envelope, but less than the ATCA NH<sub>3</sub> due to the smaller area mapped by ALMA. The ALMA  $N_2D^+$  corresponds well with the NH<sub>3</sub> on the eastern side to the edge of the map and the emission toward IRS1 is lower in both  $NH_3$  and  $N_2D^+$ . Near IRS2, the correspondence of  $N_2D^+$  and  $NH_3$  is not as close, the peak of NH<sub>3</sub> east of IRS2 corresponds with the  $N_2D^+$  peak, but the  $N_2D^+$  east and south of IRS2 has lower NH<sub>3</sub> intensity. The  $N_2D^+$  contours start at  $5\sigma$  and increase on  $10\sigma$  intervals, where  $\sigma$ =0.098 K km s<sup>-1</sup>.



N2D+ (J=3-2) Integrated Intensity K-km/s N2D+ (J=3-2) Integrated Intensity K-km/s + ALMA 1.3 mm image

<span id="page-42-0"></span>**Figure 16.** Same as for Figure [10](#page-36-0) but for the ALMA  $N_2D^+$  ( $J = 3 \rightarrow 2$ ) transition and the data are overlaid on the ALMA 1.3 mm image instead of the Spitzer 8  $\mu$ m. The spatial extents of the outflows probed by ALMA in  ${}^{12}CO$  are marked with the gray contours in all panels but the top left. The integrated intensity maps show clear deficits of  $N_2D^+$  toward the protostar locations, as expected from protostellar heating. However, toward IRS2, the  $N_2D^+$  is found in a horseshoe-shaped feature. The velocity structure is markedly different, going to lower velocities on the eastern side of the envelope in  $N_2D^+$  vs. NH<sub>3</sub>. Some of this is likely due to the higher resolution of the ALMA data and there is not as much blending of line components. Here the more red-shifted features to the north appear to occupy more area than the outflow region of influence delineated by the  ${}^{12}CO$  contours. Ignoring the features on the north side of the envelope, the east and west side of the envelope are at practically the same velocity. The linewidth map is very noisy due to the low S/N in some parts of the data, but linewidths are all quite narrow, much more narrow than NH<sup>3</sup> in most places. The intensity levels in the top panels are from the integrated intensity map of the main hyperfine lines between -5.5 to -3.7 km s<sup>-1</sup>; the intensity levels start at  $3\sigma$  and increase on  $5\sigma$  intervals, where  $\sigma$ =0.1 K km s<sup>-1</sup>. The dashed lines in the upper right panel denote the region of PV extraction, and the dashed circles in the upper right panel correspond to regions of spectral extraction to examine the quality of hyperfine fitting, see Appendix A.



<span id="page-43-0"></span>Figure 17. Position-velocity diagram of  $N_2D^+$  emission toward IRS1 and IRS2 taken in a 10<sup>n</sup> cut across the equatorial plane of the envelope  $(PA = 90^{\circ})$ , centered on IRS1. The region extracted is bounded by the dashed lines in the upper right panel of Figure [16.](#page-42-0) The second velocity component shown in Figure [12](#page-38-0) is difficult to pick out because the hyperfine structure of  $N_2D^+$  ( $J=3\rightarrow 2$ ) is comprised of the closely spaced lines with comparable relative intensities. But, the overall emission is more broad on the east side of the envelope than on the west due to the additional velocity component. The red-shifted velocities related to the outflow are clear at the position of IRS1 where the only detectable emission is red-shifted. The overall line profiles are more shifted toward the red because the rest frequency of our observation was centered on the brightest hyperfine line of  $N_2D^+$  ( $J = 3 \rightarrow 2$ ), which located at a slightly higher frequency that most of the other lines having comparable relative intensity.



C180 (J=2-1) Integrated Intensity K-km/s C180 (J=2-1) Integrated Intensity K-km/s + ALMA 1.3 mm image 1.3 3.9 6.5 9.1 11.7 14.3 16.9 19.5 22.1 24.7 27.3 29.9

<span id="page-44-0"></span>**Figure 18.** Same as for Figure [16](#page-42-0) but for the ALMA C<sup>18</sup>O ( $J = 2 \rightarrow 1$ ) and zoomed-in on a smaller area centered on the two protostars. The line center velocity map in the lower left shows the clear influence of the outflow on the  $C^{18}O$  kinematics, but there is a twist near the location of the protostar that is inconsistent with the outflow. The linewidth map for  $C^{18}O$  shows enhanced linewidth toward IRS1, indicative of more rapid motion along the line of sight; IRS2 does not show an increased line width. The intensity levels shown are the same as in Figure [6,](#page-32-0) using the 12m+ACA+TP data.



<span id="page-45-0"></span>Figure 19.  $C^{18}O$  red and blue-shifted integrated intensity maps toward IRS1 (using  $12m+ACA+TP$  data) shown in different velocity ranges and overlaid on the 1.3 mm continuum image. The low-velocity  $C^{18}O$ clearly is influenced by the outflow, but at moderate and higher velocities, the  $C^{18}O$  begins to trace a velocity gradient orthogonal to the outflow, the expected velocity gradient direction for rotation. The lowvelocity emission is integrated between -5.8 to -4.4 km s<sup>-1</sup> and -4.4 to -3.2 km s<sup>-1</sup>, and  $\sigma_{red,blue} = 0.09$ , 0.1 K km s<sup>-1</sup> with the contours are drawn starting at  $30\sigma$  and increasing on intervals of  $30\sigma$ . The moderatevelocity emission is integrated between -6.2 to -5.6 km s<sup>-1</sup> and -3.6 to -2.8 km s<sup>-1</sup>, and  $\sigma_{red,blue} = 0.08$ , 0.07 K km s<sup>-1</sup> with the contours are drawn starting at  $10\sigma$  and increasing on intervals of  $10\sigma$ . The highvelocity emission is integrated between -6.3 to -5.8 km s<sup>-1</sup> and -3.0 to -2.5 km s<sup>-1</sup>, and  $\sigma_{red,blue} = 0.06$ , 0.06 K km s<sup>-1</sup> with the contours are drawn starting at  $3\sigma$  ( $5\sigma$ ) and increasing on intervals of  $3\sigma$  ( $5\sigma$ ); the values in parentheses correspond to the blue contours. The  $12m+ACA+TP$  data were used to generate this Figure.



<span id="page-45-1"></span>Figure 20.  $\rm C^{18}O$  red and blue-shifted integrated intensity maps toward IRS2 (using  $\rm 12m+ACA+TP$  data). A velocity gradient is evident, in a direction that is near orthogonal to the outflow. However, the velocity gradient direction is in *opposite* sense with respect to the gradient direction in IRS1. The  $C^{18}O$  emission is integrated between -5.1 to -4.4 km s<sup>-1</sup> and -4.4 to -3.5 km s<sup>-1</sup>, and  $\sigma_{red,blue} = 0.08, 0.074$  K km s<sup>-1</sup>. The red and (blue) contours are drawn starting at  $15\sigma$  (35.0 $\sigma$ ) and increasing on intervals of  $5\sigma$  (10 $\sigma$ ).



<span id="page-46-0"></span>Figure 21. Position-velocity diagrams of  $C^{18}O$  emission toward IRS1 (top) and IRS2 (bottom) taken in cuts orthogonal to the outflow directions, 6. 2 in width. The velocity gradient in IRS1 appears to go from the upper left quadrant to the lower right quadrant at the highest velocities, and the velocity gradient in IRS2 appears to go from the lower left quadrant to the upper right quadrant. The two PV diagrams are generated using only the data from the ALMA 12m array, excluding the ACA and Total Power. This limits confusion from the inclusion of the extended  $C^{18}O$  emission.



<span id="page-47-0"></span>Figure 22. Position-velocity diagrams of  $C^{18}O$  emission encompassing both IRS1 and IRS2 taken in the same cut as the  $NH_3$  and  $N_2D^+$  PV diagrams shown in Figures [12](#page-38-0) and [17.](#page-43-0) This PV diagram includes the 12m, ACA, and Total Power data to recover the extended structure from the core. Similar to the NH<sub>3</sub> and  $N_2D^+$ , the C<sup>18</sup>O also shows two velocity components, one at about -4.9 km s<sup>-1</sup> and the other at about -4.0 km s−<sup>1</sup> . The C18O emission concetrated around the protostars themselves is a bit between the two velocity components, but more closely associated with the blue-shifted velocity component, having their velocity centers at about  $-4.6 \text{ km s}^{-1}$ .



<span id="page-48-0"></span>Figure 23. ATCA NH<sub>3</sub> spectra taken toward three positions, showing the spectrum and hyperfine fit in one panel and the residual of the fit below. The coordinates denoted for East, West, and Outflow are relative to the position of IRS1 in Figure [10,](#page-36-0) and the extraction regions are also drawn as circles in the upper right panel of Figure [11.](#page-37-0) The East spectrum has the largest residual which is likely due to the two velocity components that are apparent in Figure [12.](#page-38-0) The two blended components in the East spectrum results in the fit having a broader linewidth. The West and Outflow positions are dominated by a single velocity component; however, the Outflow position has a small blue-shifted residual.



<span id="page-49-0"></span>Figure 24. ALMA  $N_2D^+$  spectra taken toward three positions, showing the spectrum and hyperfine fit in one panel and the residual of the fit below. The coordinates denoted for East, West, and Outflow are relative to the position of IRS1 in Figure [16,](#page-42-0) and the extraction regions are also drawn as circles in the upper right panel of that Figure. The East spectrum has the largest residual which is likely due to the two velocity components that are apparent in Figure [17.](#page-43-0) The two blended components in the East spectrum results in the fit having a broader linewidth. The West position is dominated by a single velocity component, and the Outflow position has a small residual.



<span id="page-50-0"></span>**Figure 25.** ALMA NH<sub>3</sub> and N<sub>2</sub>D<sup>+</sup> spectra taken at the same East envelope position as in Figures [23](#page-48-0) and [24.](#page-49-0) However, instead of a single component fit, we attempt a two-component fit at this position to disentangle and characterize the two velocity components. The two-component fit is visually and statistically better than a single component fit. The vertical dashed lines in both the upper and lower panels correspond to the central velocities of each component fit.

Table 1. Molecular Line Setups

Telescope	Molecule	Frequency	Raw Spectral Resolution	Map Spectral resolution	Map RMS	Beam	Detected?
		(GHz)	$(kHz, km s^{-1})$	$(km s^{-1})$	$(\mathrm{Jy}\ \mathrm{beam}^{-1}\ \mathrm{or}\ \mathrm{K})$	$($ ")	
Parkes	$NH_3(1,1)$	23.694495	1.0, 0.012	0.05	$0.1\mathrm{~K}$	60	Y
<b>ATCA</b>	$NH_3(1,1)$	23.694495	0.5, 0.006	0.1	0.015 Jy beam <sup><math>-1</math></sup> , 0.19 K	$9.1\times7.8$	$\mathbf{Y}$
	$NH_3(2,2)$	23.722633	0.5, 0.006	$0.2\,$	$0.012$ Jy beam <sup>-1</sup>	$10.8 \times 9.2$ M	
	$NH_3(3,3)$	23.870129	0.5, 0.006	0.2	$0.011$ Jy beam <sup>-1</sup>	$10.8 \times 9.2$ N	
	Continuum	23.160	1.0, 13.0	$\cdots$	$0.000048$ Jy beam <sup>-1</sup>	$9.4 \times 8.1$	$\mathbf{Y}$
ALMA	$C^{18}O$ $(J = 2 \rightarrow 1)$	219.560354	61.035, 0.083	0.083	$0.029$ Jy beam <sup>-1</sup>	$1.7\times1.5$	$\mathbf{Y}$
	<sup>13</sup> CO $(J = 2 \rightarrow 1)$	220.398684	61.035, 0.083	0.083	$0.035$ Jy beam <sup>-1</sup>	$1.7\times1.4$	$\mathbf{Y}$
	<sup>12</sup> CO $(J = 2 \rightarrow 1)$	230.538	61.035, 0.079	0.166	$0.029$ Jy beam <sup>-1</sup>	$1.5 \times 1.3$	$\mathbf{Y}$
	$N_2D^+$ $(J = 3 \rightarrow 2)$	231.321828	61.035, 0.079	0.083	0.034 Jy beam <sup><math>-1</math></sup> , 0.23 K	$2.0\times1.8$	$\mathbf{Y}$
	Continuum	232.5	14.6, 18.8	$\cdots$	$0.00087$ Jy beam <sup>-1</sup>	$1.3\times1.2$	$\mathbf{Y}$

NOTE— The final column, with the label 'Detected?' denotes whether the molecule or continuum was detected with high-confidence (Y), a marginal detection (M), or a non-detection (N).

Table 2. Photometry

Wavelength	IRS1 Flux Density	IRS2 Flux Density	Total Flux Density	Instrument	Resolution	Aperture Radius <sup>a</sup>	Reference
$(\mu m)$	$(\mathrm{Jy})$	$(\mathrm{Jy})$	$(\mathrm{Jy})$		('')	('')	
1.25	$0.0048 \pm 0.0005$	$< 1.4e-4$	$0.017 \pm 0.0017$	CTIO/ISPI	0.9	30, 5, 50	$\mathbf{1}$
1.66	$0.025 \pm 0.006$	$< 2.6e-4$	$0.072 \pm 0.007$	CTIO/ISPI	0.9	30, 5, 50	1
2.15	$0.14 \pm 0.014$	$< 2.5e-3$	$0.33 \pm 0.03$	CTIO/ISPI	0.9	30, 5, 50	1.
3.6	$0.073 \pm 0.07$	$0.0045 \pm 0.0004$	$0.15 \pm 0.02$	<b>IRAC</b>	1.7	30, 5, 50	1.
4.5	$0.15 \pm 0.02$	$0.012 \pm 0.007$	$0.27 \pm 0.015$	<b>IRAC</b>	1.7	30, 5, 50	1
5.8	$0.17 \pm 0.02$	$0.015 \pm 0.007$	$0.27 \pm 0.004$	<b>IRAC</b>	1.8	30, 5, 50	1.
8.0	$0.14 \pm 0.03$	$0.009 \pm 0.0006$	$0.17 \pm 0.08$	<b>IRAC</b>	1.9	30, 5, 50	$\mathbf 1$
24.0	$5.0 + 0.3$	$0.09 \pm 0.003$	$5.1 \pm 0.3$	<b>MIPS</b>	6	PSF, PSF, 50	$\overline{2}$
60.0	$\cdots$	$\cdots$	$77.4 \pm 15.5$	<b>IRAS</b>	72	300	3
70.0	$127.2 \pm 12.7$	$8.8 \pm 1.0$	$145.7 \pm 14.5$	<b>PACS</b>	5.6	PSF, PSF, 50	$\mathbf{1}$
100.0	$230.9 \pm 23.0$	$22.8 \pm 4.1$	$292.7 \pm 30.0$	<b>PACS</b>	6.8	PSF, PSF, 50	1.
160.0	$248.4 \pm 25.0$	$37.6 \pm 8.0$	$353.8 \pm 35.0$	<b>PACS</b>	10.7	PSF, PSF, 50	$\mathbf{1}$
250.0	$157.5 \pm 30.0$	$36.1 \pm 6.0$	$213.4 \pm 22.0$	<b>SPIRE</b>	18	PSF, PSF, 22	$\mathbf{1}$
350.0	$\cdots$	$\cdots$	$127.0 \pm 9.1$	<b>SPIRE</b>	24	30	
500.0	$\cdots$	$\cdots$	$64.8 \pm 4.5$	<b>SPIRE</b>	35	40	
1300.0	$\cdots$	$\cdots$	$3.8 \pm 0.72$	<b>SEST</b>	23	60	$\overline{4}$
1300.0	$1.28 \pm 0.01$	$0.12 \pm 0.005$	$1.41 \pm 0.016$	<b>ALMA</b>	1.25	12.5, 5, 17.5	1
3200.0	$0.140 \pm 0.028$	$0.0028 \pm 0.0021$	$0.143 \pm 0.028$	<b>ATCA</b>	3.5	Gaussian	$\overline{2}$
12960.0	$0.0027 \pm 0.0002$	$0.00022 \pm 0.0001$	$0.0029 \pm 0.0002$	<b>ATCA</b>	9	Gaussian	

NOTE— The flux density for IRS1 includes a minor contribution from IRS2 due to the 30" (6000 AU) radius aperture coming close to encompassing the system. The total flux density refers to the flux measured in the third aperture listed in the Aperture column. References: (1) This work; (2) [Chen et al.](#page-26-3) [\(2008\)](#page-26-3); (3) IRAS Photometry, (4) [\(Launhardt et al.](#page-27-48) [2010\)](#page-27-48).

 $a$  The comma separated values in this column represent the aperture used for IRS1, IRS2, and the their combined flux densities, respectively. The aperture for the combined flux densities of IRS1 and IRS2 is chosen to encompass the emission of both sources. The numerical sizes denote aperture photometry, PSF denotes photometry by point spread function fitting and Gaussian denotes photometry by Gaussian fitting.