



Universiteit
Leiden
The Netherlands

Ensemble nonequivalence in random graphs with modular structure

Garlaschelli, D.; Hollander, W.T.F.; Roccaverde, A.

Citation

Garlaschelli, D., Hollander, W. T. F., & Roccaverde, A. (2016). Ensemble nonequivalence in random graphs with modular structure. *Journal Of Physics A: Mathematical And Theoretical*, 50(1), 015001. doi:10.1088/1751-8113/50/1/015001

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/47736>

Note: To cite this publication please use the final published version (if applicable).

Ensemble nonequivalence in random graphs with modular structure

Diego Garlaschelli ¹
Frank den Hollander ²
Andrea Roccaverde ^{1,2}

November 1, 2016

Abstract

Breaking of equivalence between the microcanonical ensemble and the canonical ensemble, describing a large system subject to hard and soft constraints, respectively, was recently shown to occur in large random graphs. Hard constraints must be met by every graph, soft constraints must be met only on average, subject to maximal entropy. In Squartini, de Mol, den Hollander and Garlaschelli (2015) it was shown that ensembles of random graphs are nonequivalent when the degrees of the nodes are constrained, in the sense of a non-zero limiting specific relative entropy as the number of nodes diverges. In that paper, the nodes were placed either on a single layer (uni-partite graphs) or on two layers (bi-partite graphs). In the present paper we consider an arbitrary number of intra-connected and inter-connected layers, thus allowing for modular graphs with a multi-partite, multiplex, time-varying, block-model or community structure. We give a full classification of ensemble equivalence in the sparse regime, proving that breakdown occurs as soon as the number of local constraints (i.e., the number of constrained degrees) is extensive in the number of nodes, irrespective of the layer structure. In addition, we derive an explicit formula for the specific relative entropy and provide an interpretation of this formula in terms of Poissonisation of the degrees.

MSC 2010. 60C05, 60K35, 82B20.

Key words and phrases. Random graph, community structure, multiplex network, multilayer network, stochastic block-model, constraints, microcanonical ensemble, canonical ensemble, relative entropy, equivalence vs. nonequivalence.

Acknowledgment. DG and AR are supported by EU-project 317532-MULTIPLEX. FdH and AR are supported by NWO Gravitation Grant 024.002.003-NETWORKS.

¹Lorentz Institute for Theoretical Physics, Leiden University, P.O. Box 9504, 2300 RA Leiden, The Netherlands

²Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands

1 Introduction and main results

1.1 Background and outline

For systems with many interacting components a detailed microscopic description is infeasible and must be replaced by a probabilistic description, where the system is assumed to be a random sample drawn from a set of allowed microscopic configurations that are consistent with a set of known macroscopic properties, referred to as *constraints*. Statistical physics deals with the definition of the appropriate probability distribution over the set of microscopic configurations and with the calculation of the resulting macroscopic properties of the system. The three main choices of probability distribution are: (1) the *microcanonical ensemble*, where the constraints are *hard* (i.e., are satisfied by each individual configuration); (2) the *canonical ensemble*, where the constraints are *soft* (i.e., hold as ensemble averages, while individual configurations may violate the constraints); (3) the *grandcanonical ensemble*, where also the number of components is considered as a soft constraint.

For systems that are large but finite, the three ensembles are obviously different and, in fact, represent different physical situations: (1) the microcanonical ensemble models completely isolated systems (where both the energy and the number of particles are “hard”); (2) the canonical ensemble models closed systems in thermal equilibrium with a heat bath (where the energy is “soft” and the number of particles is “hard”); (3) the grandcanonical ensemble models open systems in thermal and chemical equilibrium (where both the energy and the number of particles are “soft”). However, in the limit as the number of particles diverges, the three ensembles are traditionally *assumed* to become equivalent as a result of the expected vanishing of the fluctuations of the soft constraints, i.e., the soft constraints are expected to become asymptotically hard. This assumption of *ensemble equivalence*, which dates back to Gibbs [33], has been verified in traditional models of physical systems with short-range interactions and a finite number of constraints, but it does *not* hold in general. Nonetheless, equivalence is considered to be one of the pillars of statistical physics and underlies many of the results that contribute to our current understanding of large real-world systems.

Despite the fact that many textbooks still convey the message that ensemble equivalence holds for all systems, as some sort of universal asymptotic property, over the last decades various examples have been found for which it breaks down. These examples range from astrophysical processes [41], [56], [35], [40], [19], quantum phase separation [10], [5], [25], nuclear fragmentation [21], and fluid turbulence [23], [24]. Across these examples, the signatures of ensemble nonequivalence differ, which calls for a rigorous mathematical definition of ensemble (non)equivalence: (i) *thermodynamic equivalence* refers to the existence of an invertible Legendre transform between the microcanonical entropy and canonical free energy [25]; (ii) *macrostate equivalence* refers to the equivalence of the canonical and microcanonical sets of equilibrium values of macroscopic properties [58]; (iii) *measure equivalence* refers to the asymptotic equivalence of the microcanonical and canonical probability distributions in the thermodynamic limit, i.e., the vanishing of their specific relative entropy [57]. The latter reference reviews the three definitions and shows that, under certain hypotheses, they are identical.

In the present paper we focus on the equivalence between microcanonical and canonical ensembles, although nonequivalence can in general involve the grandcanonical ensemble as well [59]. While there is consensus that nonequivalence occurs when the microcanonical specific entropy is non-concave as a function of the energy density in the thermodynamic limit, the classification of the physical mechanisms at the *origin* of nonequivalence is still open. In most

of the models studied in the literature, nonequivalence appears to be associated with the non-additivity of the energy of the subparts of the system or with phase transitions [15], [16], [57]. A possible and natural mechanism for non-additivity is the presence of *long-range interactions*. Similarly, phase transitions are naturally associated with long-range order. These “standard mechanisms” for ensemble nonequivalence have been documented also in the study of random graphs. In [4], a Potts model on a random regular graph is studied in both the microcanonical and canonical ensemble, where the microscopic configurations are the spin configurations (not the configurations of the network itself). It is found that the long-range nature of random connections, which makes the model non-additive and the microcanonical entropy non-concave, ultimately results in ensemble nonequivalence. In [50], [51], [52] and [18], random networks with given densities of edges and triangles are considered, and phase transitions characterised by jumps in these densities are found, with an associated breaking of ensemble equivalence (where the microscopic configurations are network configurations).

Recently, the study of certain classes of uni-partite and bi-partite random graphs [55], [30] has shown that ensemble nonequivalence can manifest itself via an additional, novel mechanism, unrelated to non-additivity or phase transitions: namely, the presence of an *extensive* number of local topological constraints, i.e., the degrees and/or the strengths (for weighted graphs) of all nodes.¹ This finding explains previously documented signatures of nonequivalence in random graphs with local constraints, such as a finite difference between the microcanonical and canonical entropy densities [1] and the non-vanishing of the relative fluctuations of the constraints [54]. How generally this result holds beyond the specific uni-partite and bi-partite cases considered so far remains an open question, on which we focus in the present paper. By considering a much more general class of random graphs with a variable number of constraints, we confirm that the presence of an extensive number of local topological constraints breaks ensemble equivalence, even in the absence of phase transitions or non-additivity.

The remainder of our paper is organised as follows. In Section 1.2 we give the definition of measure equivalence and, following [55], show that it translates into a simple pointwise criterion for the large deviation properties of the microcanonical and canonical probabilities. In Section 1.3 we introduce our main theorems in pedagogical order, starting from the characterisation of nonequivalence in the simple cases of uni-partite and bi-partite graphs already explored in [55], and subsequently moving on to a very general class of graphs with arbitrary multilayer structure and tunable intra-layer and inter-layer connectivity. Our main theorems, which (mostly) concern the *sparse regime*, not only characterise nonequivalence *qualitatively*, they also provide a *quantitative* formula for the specific relative entropy. In Section 2 we discuss various important implications of our results, describing properties that are fully general but also focussing on several special cases of empirical relevance. In addition, we provide an interpretation of the specific relative entropy formula in terms of Poissonisation of the degrees. We also discuss the implications of our results for the study of several empirically relevant classes of “modular” networks that have recently attracted interest in the literature, such as networks with a so-called multi-partite, multiplex [11], time-varying [38], block-model [37], [39] or community structure [26], [49]. In Section 3, finally, we provide the proofs of our theorems.

In future work we will address the *dense regime*, which requires the use of *graphons*. In that regime we expect nonequivalence to persist, and in some cases become even more pronounced.

¹While in binary (i.e., simple) graphs the *degree* of a node is defined as the number of edges incident to that node, in weighted graphs (i.e., graphs where edges can carry weights) the *strength* of a node is defined as the total weight of all edges incident to that node. In this paper, we focus on binary graphs only.

1.2 Microcanonical ensemble, canonical ensemble, relative entropy

For $n \in \mathbb{N}$, let \mathcal{G}_n denote the set of all simple undirected graphs with n nodes. Let $\mathcal{G}_n^\sharp \subseteq \mathcal{G}_n$ be some non-empty subset of \mathcal{G}_n , to be specified later. Informally, the restriction from \mathcal{G}_n to \mathcal{G}_n^\sharp allows us to forbid the presence of certain links, in such a way that the n nodes are effectively partitioned into $M \in \mathbb{N}$ groups of nodes (or “layers”) of sizes n_1, \dots, n_M with $\sum_{i=1}^M n_i = n$. This restriction can be made explicit and rigorous through the definition of a superstructure, which we call the *master graph*, that will be introduced later. A given choice of \mathcal{G}_n^\sharp corresponds to the selection of a specific class of *multilayer* graphs with desired intra-layer and inter-layer connectivity, such as graphs with a multipartite, multiplex, time-varying, block-model or community structure. In the simplest case, $\mathcal{G}_n^\sharp = \mathcal{G}_n$, which reduces to the ordinary choice of uni-partite (single-layer) graphs. This example, along with various more complicated examples, is considered explicitly later on.

In general, any graph $\mathbf{G} \in \mathcal{G}_n^\sharp$ can be represented as an $n \times n$ matrix with elements

$$g_{i,j}(\mathbf{G}) = \begin{cases} 1 & \text{if there is a link between node } i \text{ and node } j, \\ 0 & \text{otherwise.} \end{cases} \quad (1.1)$$

Let \vec{C} denote a vector-valued function on \mathcal{G}_n^\sharp . Given a specific value \vec{C}^* , which we assume to be *graphic*, i.e., realisable by at least one graph in \mathcal{G}_n^\sharp , the *microcanonical probability distribution* on \mathcal{G}_n^\sharp with *hard constraint* \vec{C}^* is defined as

$$P_{\text{mic}}(\mathbf{G}) = \begin{cases} 1/\Omega_{\vec{C}^*}, & \text{if } \vec{C}(\mathbf{G}) = \vec{C}^*, \\ 0, & \text{else,} \end{cases} \quad (1.2)$$

where

$$\Omega_{\vec{C}^*} = |\{\mathbf{G} \in \mathcal{G}_n^\sharp : \vec{C}(\mathbf{G}) = \vec{C}^*\}| > 0 \quad (1.3)$$

is the number of graphs that realise \vec{C}^* . The *canonical probability distribution* $P_{\text{can}}(\mathbf{G})$ on \mathcal{G}_n^\sharp is defined as the solution of the maximisation of the *entropy*

$$S_n(P_{\text{can}}) = - \sum_{\mathbf{G} \in \mathcal{G}_n^\sharp} P_{\text{can}}(\mathbf{G}) \ln P_{\text{can}}(\mathbf{G}) \quad (1.4)$$

subject to the *soft constraint* $\langle \vec{C} \rangle = \vec{C}^*$, where $\langle \cdot \rangle$ denotes the average w.r.t. P_{can} , and subject to the normalisation condition $\sum_{\mathbf{G} \in \mathcal{G}_n^\sharp} P_{\text{can}}(\mathbf{G}) = 1$. This gives

$$P_{\text{can}}(\mathbf{G}) = \frac{\exp[-H(\mathbf{G}, \vec{\theta}^*)]}{Z(\vec{\theta}^*)}, \quad (1.5)$$

where

$$H(\mathbf{G}, \vec{\theta}) = \vec{\theta} \cdot \vec{C}(\mathbf{G}) \quad (1.6)$$

is the *Hamiltonian* (or *energy*) and

$$Z(\vec{\theta}) = \sum_{\mathbf{G} \in \mathcal{G}_n^\sharp} \exp[-H(\mathbf{G}, \vec{\theta})] \quad (1.7)$$

is the *partition function*. Note that in (1.5) the parameter $\vec{\theta}$ must be set to the particular value $\vec{\theta}^*$ that realises $\langle \vec{C} \rangle = \vec{C}^*$. This value also maximises the likelihood of the model, given the data [31].

It is worth mentioning that, in the social network analysis literature [17], maximum-entropy canonical ensembles of graphs are traditionally known under the name of Exponential Random Graphs (ERGs). Indeed, many of the examples of canonical graph ensembles that we will consider in this paper, or variants thereof, have been studied previously as ERG models of social networks. Recently, ERGs have also entered the physics literature [1], [2], [9], [47], [53], [54], [42], [28], [29], [39], [27], [48], [8] because of the wide applicability of techniques from statistical physics for the calculation of canonical partition functions. We will refer more extensively to these models, and to the empirical situations for which they have been proposed, in Section 2.2. Apart for a few exceptions [1], [48], [55], these previous studies have not addressed the problem of ensemble (non)equivalence of ERGs. The aim of the present paper is to do so exhaustively, and in a mathematically rigorous way, via the following definitions.

The *relative entropy* of P_{mic} w.r.t. P_{can} is

$$S_n(P_{\text{mic}} | P_{\text{can}}) = \sum_{\mathbf{G} \in \mathcal{G}_n^\sharp} P_{\text{mic}}(\mathbf{G}) \ln \frac{P_{\text{mic}}(\mathbf{G})}{P_{\text{can}}(\mathbf{G})}, \quad (1.8)$$

and the *specific relative entropy* is

$$s_n = n^{-1} S_n(P_{\text{mic}} | P_{\text{can}}). \quad (1.9)$$

Following [57], [55], we say that the two ensembles are measure equivalent if and only if their specific relative entropy vanishes in the *thermodynamic limit* $n \rightarrow \infty$, i.e.,

$$s_\infty = \lim_{n \rightarrow \infty} n^{-1} S_n(P_{\text{mic}} | P_{\text{can}}) = 0. \quad (1.10)$$

It should be noted that, for a given choice of \mathcal{G}_n^\sharp and \vec{C} , there may be different ways to realise the thermodynamic limit, corresponding to different ways in which the numbers $\{n_i\}_{i=1}^M$ of nodes inside the M layers grow relatively to each other. So, (1.10) implicitly requires an underlying *specific definition of the thermodynamic limit*. Explicit examples will be considered in each case separately, and certain different realisations of the thermodynamic limit will indeed be seen to lead to different results. With this in mind, we suppress the n -dependence from our notation of quantities like \mathbf{G} , \vec{C} , \vec{C}^* , P_{mic} , P_{can} , H , Z . When letting $n \rightarrow \infty$ it will be understood that $\mathbf{G} \in \mathcal{G}_n^\sharp$ always.

Before considering specific cases, we recall an important observation made in [55]. The definition of $H(\mathbf{G}, \vec{\theta})$ ensures that, for any $\mathbf{G}_1, \mathbf{G}_2 \in \mathcal{G}_n^\sharp$, $P_{\text{can}}(\mathbf{G}_1) = P_{\text{can}}(\mathbf{G}_2)$ whenever $\vec{C}(\mathbf{G}_1) = \vec{C}(\mathbf{G}_2)$ (i.e., the canonical probability is the same for all graphs having the same value of the constraint). We may therefore rewrite (1.8) as

$$S_n(P_{\text{mic}} | P_{\text{can}}) = \ln \frac{P_{\text{mic}}(\mathbf{G}^*)}{P_{\text{can}}(\mathbf{G}^*)}, \quad (1.11)$$

where \mathbf{G}^* is *any* graph in \mathcal{G}_n^\sharp such that $\vec{C}(\mathbf{G}^*) = \vec{C}^*$ (recall that we have assumed that \vec{C}^* is realisable by at least one graph in \mathcal{G}_n^\sharp). The condition for equivalence in (1.10) then becomes

$$\lim_{n \rightarrow \infty} n^{-1} [\ln P_{\text{mic}}(\mathbf{G}^*) - \ln P_{\text{can}}(\mathbf{G}^*)] = 0, \quad (1.12)$$

which shows that the breaking of ensemble equivalence coincides with $P_{\text{mic}}(\mathbf{G}^*)$ and $P_{\text{can}}(\mathbf{G}^*)$ having different large deviation behaviour. Importantly, this condition is entirely local, i.e.,

it involves the microcanonical and canonical probabilities of a *single* configuration \mathbf{G}^* realising the hard constraint. Apart from its theoretical importance, this fact greatly simplifies mathematical calculations. Note that (1.12), like (1.10), implicitly requires a specific definition of the thermodynamic limit. For a given choice of \mathcal{G}_n^\sharp and \vec{C} , different definitions of the thermodynamic limit may result either in ensemble equivalence or in ensemble nonequivalence.

1.3 Main Theorems

Most of the constraints that will be considered below are *extensive* in the number of nodes.

1.3.1 Single layer: uni-partite graphs

The first class of random graphs we consider is specified by $M = 1$ and $\mathcal{G}_n^\sharp = \mathcal{G}_n$. This choice corresponds to the class of (simple and undirected) *uni-partite graphs*, where links are allowed between each pair of nodes. We can think of these graphs as consisting of a single layer of nodes, inside which all links are allowed. Note that in this simple case the thermodynamic limit $n \rightarrow \infty$ can be realised in a unique way, which makes (1.10) and (1.12) already well-defined.

Constraints on the degree sequence. For a uni-partite graph $\mathbf{G} \in \mathcal{G}_n$, the degree sequence is defined as $\vec{k}(\mathbf{G}) = (k_i(\mathbf{G}))_{i=1}^n$ with $k_i(\mathbf{G}) = \sum_{j \neq i} g_{i,j}(\mathbf{G})$. In what follows we constrain the degree sequence to a *specific value* \vec{k}^* , which (in accordance with our aforementioned general prescription for \vec{C}^*) we assume to be *graphical*, i.e., there is at least one graph with degree sequence \vec{k}^* . The constraints are therefore

$$\vec{C}^* = \vec{k}^* = (k_i^*)_{i=1}^n \in \mathbb{N}_0^n, \quad (1.13)$$

where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ with $\mathbb{N} = \{1, 2, \dots\}$. This class is also known as the *configuration model* ([7], [13], [44], [45], [20], [54]; see also [36, Chapter 7]). In [55] the breaking of ensemble equivalence was studied in the *sparse regime* defined by the condition

$$m^* = \max_{1 \leq i \leq n} k_i^* = o(\sqrt{n}). \quad (1.14)$$

Let $\mathcal{P}(\mathbb{N}_0)$ denote the set of probability distributions on \mathbb{N}_0 . Let

$$f_n = n^{-1} \sum_{i=1}^n \delta_{k_i^*} \in \mathcal{P}(\mathbb{N}_0), \quad (1.15)$$

be the *empirical degree distribution*, where δ_k denotes the point measure at k . Suppose that there exists a degree distribution $f \in \mathcal{P}(\mathbb{N}_0)$ such that

$$\lim_{n \rightarrow \infty} \|f_n - f\|_{\ell^1(g)} = 0, \quad (1.16)$$

where $g: \mathbb{N}_0 \rightarrow [0, \infty)$ is given by

$$g(k) = \log \left(\frac{k!}{k^k e^{-k}} \right), \quad k \in \mathbb{N}_0, \quad (1.17)$$

and $\ell^1(g)$ is the vector space of functions $h: \mathbb{Z} \rightarrow \mathbb{R}$ with $\|h\|_{\ell^1(g)} = \sum_{k \in \mathbb{N}_0} |h(k)|g(k) < \infty$. For later use we note that

$$g(0) = 0, \quad k \mapsto g(k) \text{ is strictly increasing}, \quad g(k) = \frac{1}{2} \log(2\pi k) + O(k^{-1}), \quad k \rightarrow \infty. \quad (1.18)$$

Theorem 1.1. *Subject to (1.13)–(1.14) and (1.16), the specific relative entropy equals*

$$s_\infty = \|f\|_{\ell^1(g)} > 0. \quad (1.19)$$

Thus, when we constrain the degrees we break the ensemble equivalence.

Remark 1.2. It is known that \vec{k}^* is graphical if and only if $\sum_{i=1}^n k_i^*$ is even and

$$\sum_{i=1}^j k_i^* \leq j(j-1) + \sum_{i=j+1}^n \min(j, k_i^*), \quad j = 1, \dots, n-1. \quad (1.20)$$

In [3], the case where k_i^* , $i \in \mathbb{N}$, are i.i.d. with probability distribution f is considered, and it is shown that

$$\lim_{n \rightarrow \infty} f^{\otimes n} \left((k_1^*, \dots, k_n^*) \text{ is graphical} \mid \sum_{i=1}^n k_i^* \text{ is even} \right) = 1 \quad (1.21)$$

as soon as f satisfies $0 < \sum_{k \text{ even}} f(k) < 1$ and $\lim_{n \rightarrow \infty} n \sum_{k \geq n} f(k) = 0$. (The latter condition is slightly weaker than the condition $\sum_{k \in \mathbb{N}_0} k f(k) < \infty$.) In what follows we do *not* require the degrees to be drawn in this manner, but when we let $n \rightarrow \infty$ we always implicitly assume that the limit is taken *within the class of graphical degree sequences*.

Remark 1.3. A different yet similar definition of sparse regime, replacing (1.14), is given in van der Hofstad [36, Chapter 7]. This condition is formulated in terms of bounded second moment of the empirical degree distribution f_n in the limit as $n \rightarrow \infty$. Theorem 1.1 carries over.

Constraints on the total number of links only. We now relax the constraints, and fix only the total number of links $L(\mathbf{G}) = \frac{1}{2} \sum_{i=1}^n k_i(\mathbf{G})$. The constraint therefore becomes

$$\vec{C}^* = L^*. \quad (1.22)$$

It should be noted that in this case, the canonical ensemble coincides with the Erdős-Rényi random graph model, where each pair of nodes is independently connected with the same probability. As shown in [1], [55], in this case the usual result that the ensembles are asymptotically equivalent holds.

Theorem 1.4. *Subject to (1.22), the specific relative entropy equals $s_\infty = 0$.*

1.3.2 Two layers: bi-partite graphs

The second class of random graphs we consider are *bi-partite graphs*. Here $M = 2$ and nodes are placed on two (non-overlapping) layers (say, top and bottom), and only links *across* layers are allowed. Let Λ_1 and Λ_2 denote the sets of nodes in the top and bottom layer, respectively. The set of all bi-partite graphs consisting of $n_1 = |\Lambda_1|$ nodes in the top layer and $n_2 = |\Lambda_2|$ nodes in the bottom layer is denoted by $\mathcal{G}_n^\# = \mathcal{G}_{n_1, n_2} \subset \mathcal{G}_n$. Bi-partiteness means that, for all $\mathbf{G} \in \mathcal{G}_{n_1, n_2}$, we have $g_{i,j}(\mathbf{G}) = 0$ if $i, j \in \Lambda_1$ or $i, j \in \Lambda_2$.

In a bipartite graph $\mathbf{G} \in \mathcal{G}_{n_1, n_2}$, we define the degree sequence of the top layer as $\vec{k}_{1 \rightarrow 2}(\mathbf{G}) = (k_i(\mathbf{G}))_{i \in \Lambda_1}$, where $k_i(\mathbf{G}) = \sum_{j \in \Lambda_2} g_{i,j}(\mathbf{G})$. Similarly, we define the degree sequence of the bottom layer as $\vec{k}_{2 \rightarrow 1}(\mathbf{G}) = (k'_i(\mathbf{G}))_{i \in \Lambda_2}$, where $k'_i(\mathbf{G}) = \sum_{j \in \Lambda_1} g_{i,j}(\mathbf{G})$. The symbol $s \rightarrow t$ highlights the fact that the degree sequence of layer s is built from links pointing

from Λ_s to Λ_t ($s, t = 1, 2$). The degree sequences $\vec{k}_{1 \rightarrow 2}(\mathbf{G})$ and $\vec{k}_{2 \rightarrow 1}(\mathbf{G})$ are related by the condition that they both add up to the total number of links $L(\mathbf{G})$:

$$L(\mathbf{G}) = \sum_{i \in \Lambda_1} k_i(\mathbf{G}) = \sum_{j \in \Lambda_2} k'_j(\mathbf{G}). \quad (1.23)$$

Constraints on the top and the bottom layer. We first fix the degree sequence on both layers, i.e., we constrain $\vec{k}_{1 \rightarrow 2}(\mathbf{G})$ and $\vec{k}_{2 \rightarrow 1}(\mathbf{G})$ to the values $\vec{k}_{1 \rightarrow 2}^* = (k_i^*)_{i \in \Lambda_1}$ and $\vec{k}_{2 \rightarrow 1}^* = (k'_j)^*_{j \in \Lambda_2}$ respectively. The constraints are therefore

$$\vec{C}^* = \{\vec{k}_{1 \rightarrow 2}^*, \vec{k}_{2 \rightarrow 1}^*\}. \quad (1.24)$$

As mentioned before, we allow n_1 and n_2 to depend on n , i.e., $n_1 = n_1(n)$ and $n_2 = n_2(n)$. In order not to overburden the notation, we suppress the dependence on n from the notation.

We abbreviate

$$\begin{aligned} m^* &= \max_{i \in \Lambda_1} k_i^*, & m'^* &= \max_{j \in \Lambda_2} k'_j^*, \\ f_{1 \rightarrow 2}^{(n_1)} &= n_1^{-1} \sum_{i \in \Lambda_1} \delta_{k_i^*}, & f_{2 \rightarrow 1}^{(n_2)} &= n_2^{-1} \sum_{j \in \Lambda_2} \delta_{k'_j^*}, \end{aligned} \quad (1.25)$$

and assume the existence of

$$A_1 = \lim_{n \rightarrow \infty} \frac{n_1}{n_1 + n_2}, \quad A_2 = \lim_{n \rightarrow \infty} \frac{n_2}{n_1 + n_2}. \quad (1.26)$$

(This assumption is to be read as follows: choose $n_1 = n_1(n)$ and $n_2 = n_2(n)$ in such a way that the limiting fractions A_1 and A_2 exist.) The *sparse regime* corresponds to

$$m^* m'^* = o(L^{*2/3}), \quad n \rightarrow \infty. \quad (1.27)$$

We further assume that there exist $f_{1 \rightarrow 2}, f_{2 \rightarrow 1} \in \mathcal{P}(\mathbb{N}_0)$ such that

$$\lim_{n \rightarrow \infty} \|f_{1 \rightarrow 2}^{(n_1)} - f_{1 \rightarrow 2}\|_{\ell^1(g)} = 0, \quad \lim_{n \rightarrow \infty} \|f_{2 \rightarrow 1}^{(n_2)} - f_{2 \rightarrow 1}\|_{\ell^1(g)} = 0. \quad (1.28)$$

The specific relative entropy is

$$s_{n_1+n_2} = \frac{S_{n_1+n_2}(P_{\text{mic}} | P_{\text{can}})}{n_1 + n_2}. \quad (1.29)$$

Theorem 1.5. *Subject to (1.24) and (1.26)–(1.28),*

$$s_\infty = \lim_{n \rightarrow \infty} \frac{S_{n_1+n_2}(P_{\text{mic}} | P_{\text{can}})}{n_1 + n_2} = A_1 \|f_{1 \rightarrow 2}\|_{\ell^1(g)} + A_2 \|f_{2 \rightarrow 1}\|_{\ell^1(g)}. \quad (1.30)$$

Since $A_1 + A_2 = 1$, it follows that $s_\infty > 0$, so in this case ensemble equivalence never holds.

Constraints on the top layer only. We now partly relax the constraints and only fix the degree sequence $\vec{k}_{1 \rightarrow 2}(\mathbf{G})$ to the value

$$\vec{C}^* = \vec{k}_{1 \rightarrow 2}^* = (k_i^*)_{i \in \Lambda_1}, \quad (1.31)$$

while leaving $\vec{k}_{2 \rightarrow 1}(\mathbf{G})$ unspecified (apart for the condition (1.23)). The microcanonical number of graphs satisfying the constraint is

$$\Omega_{\vec{k}_{1 \rightarrow 2}^*} = \prod_{i \in \Lambda_1} \binom{n_2}{k_i^*}. \quad (1.32)$$

The canonical ensemble can be obtained from (1.5) by setting

$$H(\mathbf{G}, \vec{\theta}) = \vec{\theta} \cdot \vec{k}_{1 \rightarrow 2}(\mathbf{G}). \quad (1.33)$$

Setting $\vec{\theta} = \vec{\theta}^*$ in order that equation (1.5) is satisfied, we can write the canonical probability as

$$P_{\text{can}}(\mathbf{G}) = \prod_{i \in \Lambda_1} (p_i^*)^{k_i(\mathbf{G})} (1 - p_i^*)^{n_2 - k_i(\mathbf{G})} \quad (1.34)$$

with $p_i^* = \frac{k_i^*}{n_2}$. Let

$$f_{n_1} = n_2^{-1} \sum_{i \in \Lambda_2} \delta_{k_i^*} \in \mathcal{P}(\mathbb{N}_0). \quad (1.35)$$

Suppose that there exists an $f \in \mathcal{P}(\mathbb{N}_0)$ such that

$$\lim_{n \rightarrow \infty} \|f_{n_1} - f\|_{\ell^1(g)} = 0. \quad (1.36)$$

The relative entropy per node can be written as

$$s_{n_1+n_2} = \frac{S_{n_1+n_2}(P_{\text{mic}} | P_{\text{can}})}{n_1 + n_2} = \frac{n_1}{n_1 + n_2} \|f_{n_1}\|_{\ell^1(g_{n_2})}, \quad (1.37)$$

with

$$g_{n_2}(k) = -\log \left[\text{Bin} \left(n_2, \frac{k}{n_2} \right) (k) \right] \mathbb{I}_{0 \leq k \leq n_2}, \quad k \in \mathbb{N}_0, \quad (1.38)$$

and $\text{Bin}(n_2, \frac{k}{n_2})(k) = \binom{n_2}{k} \left(\frac{k}{n_2}\right)^k \left(\frac{n_2-k}{n_2}\right)^{n_2-k}$ for $k = 0, \dots, n_2$ and equals to 0 for $k > n_2$. We follow the convention $0 \log(0) = 0$.

In this partly relaxed case, different scenarios are possible depending on the specific realisation of the thermodynamic limit, i.e., on how n_1, n_2 tend to infinity. The ratio between the sizes of the two layers $c = \lim_{n \rightarrow \infty} \frac{n_2}{n_1} = \frac{A_2}{A_1}$ plays an important role.

Theorem 1.6. *Subject to (1.31) and (1.36):*

- (1) *If $n_2 \rightarrow^{n \rightarrow \infty} \infty$ with n_1 fixed ($c = \infty$), then $s_\infty = \lim_{n \rightarrow \infty} s_{n_1+n_2} = 0$.*
- (2) *If $n_1, n_2 \rightarrow^{n \rightarrow \infty} \infty$ with $c = \infty$, then $s_\infty = \lim_{n \rightarrow \infty} s_{n_1+n_2} = 0$.*
- (3) *If $n_1 \rightarrow^{n \rightarrow \infty} \infty$ with n_2 fixed ($c = 0$), then*

$$s_\infty = \lim_{n \rightarrow \infty} s_{n_1+n_2} = \|f\|_{\ell^1(g_{n_2})}. \quad (1.39)$$

- (4) *If $n_1, n_2 \rightarrow^{n \rightarrow \infty} \infty$ with $c \in [0, \infty)$, then*

$$s_\infty = \frac{1}{1+c} \|f\|_{\ell^1(g)}. \quad (1.40)$$

Constraints on the total number of links only. We now fully relax the constraints and only fix the total number of links, i.e.,

$$\vec{C}^* = L^*. \quad (1.41)$$

In analogy with the corresponding result for the uni-partite case (Theorem 1.4), in this case ensemble equivalence is restored.

Theorem 1.7. *Subject to (1.41), the specific relative entropy equals $s_\infty = 0$.*

1.3.3 Multiple layers

We now come to our most general setting where we fix a finite number $M \in \mathbb{N}$ of layers. Each layer s has n_s nodes, with $\sum_{s=1}^M n_s = n$. Let $v_i^{(s)}$ denote the i -th node of layer s , and $\Lambda_s = \{v_1^{(s)}, \dots, v_{n_s}^{(s)}\}$ denote the set of nodes in layer s . We may allow links *both within and across* layers, while constraining the numbers of links among different layers separately. But we may as well switch off links inside or between (some of the) layers. The actual choice can be specified by a superstructure, which we denote as the *master graph* $\mathbf{\Gamma}$, in which self-loops are allowed but multi-links are not. The nodes set of $\mathbf{\Gamma}$ is $\{1, \dots, M\}$ and the associated adjacency matrix has entries

$$\gamma_{s,t}(\mathbf{\Gamma}) = \begin{cases} 1 & \text{if a link between layers } s \text{ and } t \text{ exists} \\ 0 & \text{otherwise.} \end{cases} \quad (1.42)$$

The chosen set of all multi-layer graphs with given numbers of nodes, layers, and admissible edges (we admit edges only between layers connected in the *master graph*) is $\mathcal{G}_n^\# = \mathcal{G}_{n_1, \dots, n_M}(\mathbf{\Gamma}) \subseteq \mathcal{G}_n$. In 2.2 we discuss various empirically relevant choices of $\mathbf{\Gamma}$ explicitly, while here we keep our discussion entirely general.

Given a graph \mathbf{G} , for each pair of layers s and t (including $s = t$) we define the *t-targeted degree sequence* of layer s as $\vec{k}_{s \rightarrow t}(\mathbf{G}) = (k_i^t(\mathbf{G}))_{i \in \Lambda_s}$, where $k_i^t(\mathbf{G}) = \sum_{j \in \Lambda_t} g_{i,j}(\mathbf{G})$ is the number of links connecting node i to all other nodes in layer t . For each pair of layers s and t such that $\gamma_{s,t}(\mathbf{\Gamma}) = 1$, we enforce the value $\vec{k}_{s \rightarrow t}^* = (k_i^{*t})_{i \in \Lambda_s}$ as a constraint for the t -targeted degree sequence of layer s . For $\gamma_{s,t}(\mathbf{\Gamma}) = 0$ we have $\vec{k}_{s \rightarrow t}^* = \vec{0}$, but this constraint is automatically enforced by the master graph. Thus, the relevant constraints are

$$\vec{C}^* = \left\{ \vec{k}_{s \rightarrow t}^* : s, t = 1, \dots, M \quad \gamma_{s,t}(\mathbf{\Gamma}) = 1 \right\}. \quad (1.43)$$

We abbreviate

$$L_{s,t}^* = \sum_{i \in \Lambda_s} k_i^{*t} = \sum_{j \in \Lambda_t} k_j^{*s}, \quad m_{s \rightarrow t}^* = \max_{i \in \Lambda_s} k_i^{*t}, \quad f_{s \rightarrow t}^{(n_s)} = n_s^{-1} \sum_{i \in \Lambda_s} \delta_{k_i^{*t}}, \quad (1.44)$$

where $L_{s,t}^*$ is the number of links between layers s and t (note that $L_{s,s}^*$ is *twice* the number of links inside layer s), and assume the existence of

$$A_s = \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{n_s}{n} \quad \forall s, \quad (1.45)$$

where $\sum_{s=1}^M A_s = 1$. (As before, this assumption is to be read as follows: choose $n_s = n_s(n)$, $1 \leq s \leq M$, in such a way that the limiting fractions $A_s, 1 \leq s \leq M$, exist.) The *sparse regime* corresponds to

$$\begin{aligned} m_{s \rightarrow t}^* m_{t \rightarrow s}^* &= o(L_{s,t}^{*2/3}), & n_s, n_t &\rightarrow \infty \quad \forall s \neq t, \\ m_{s \rightarrow s}^* &= o(n_s^{1/2}), & n_s &\rightarrow \infty \quad \forall s. \end{aligned} \quad (1.46)$$

We further assume that there exists $f_{s \rightarrow t} \in \mathcal{P}(\mathbb{N}_0)$ such that

$$\lim_{n_s \rightarrow \infty} \|f_{s \rightarrow t}^{(n_s)} - f_{s \rightarrow t}\|_{\ell^1(g)}, \quad \lim_{n_s \rightarrow \infty} \|f_{s \rightarrow s}^{(n_s)} - f_{s \rightarrow s}\|_{\ell^1(g)} = 0. \quad (1.47)$$

Theorem 1.8. *Subject to (1.43) and (1.45)–(1.47),*

$$s_\infty = \sum_{\substack{s,t=1 \\ \gamma_{s,t}(\Gamma)=1}}^M A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \quad (1.48)$$

The above result shows that, unless $A_s = 0$ whenever $\gamma_{s,t}(\Gamma) = 1$ (i.e., unless only the nodes of the master graph that have no links or self-loops contribute a finite fraction of nodes in the corresponding layers), ensemble equivalence does not hold.

1.3.4 Relaxing constraints in the multilayer case

We next study the effects of relaxing constraints. This deserves a separate discussion, since in the multi-partite setting there are more possible ways of relaxing the constraints than in the uni-partite and bi-partite settings.

One class of layers. We first fix *two kinds of constraints*: (1) the total number of links between some pairs of layers; (2) the degree sequence between some other pairs of layers. We define the set of the edges of the *master graph* as $\mathcal{E} = \{(s, t) \in (M \times M) : \gamma_{s,t}(\Gamma) = 1\}$. Then, we partition \mathcal{E} into two parts, namely $\mathcal{D}, \mathcal{L} \subseteq \mathcal{E}$, with $\mathcal{D} \cap \mathcal{L} = \emptyset$, \mathcal{D} and \mathcal{L} symmetric, by requiring that $(s, t) \in \mathcal{D}$ ($\in \mathcal{L}$) when $(t, s) \in \mathcal{D}$ ($\in \mathcal{L}$). For each pair of layers $(s, t) \in \mathcal{D}$ we fix the degree sequence $\vec{k}_{s \rightarrow t}^*$ of every node of Λ_s linking to Λ_t . As before, we impose that $\sum_{i \in \Lambda_s} k_i^{*t} = \sum_{j \in \Lambda_t} k_j^{*s}$. For each pair of layers $(s, t) \in \mathcal{L}$ we fix the total number of links $L_{s,t}^*$ ($L_{s,t}^* = L_{t,s}^*$).

The effect of relaxing some constraints affects the specific relative entropy: this will decrease because the pairs of layers with relaxed constraints (i.e., the pairs in \mathcal{L}) no longer contribute.

Theorem 1.9. *Subject to the above relaxation,*

$$s_\infty = \sum_{(s,t) \in \mathcal{D}} A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \quad (1.49)$$

In particular, equivalence holds if and only if $\mathcal{D} = \emptyset$ or $A_s = 0$ for all s endpoints of elements in \mathcal{E} . Note that, if $\mathcal{D} = \emptyset$, then we have a finite number of constraints (at most M^2), and this implies equivalence of the ensembles.

Two classes of layers. We may further generalise Theorem 1.8 as follows. Suppose that we have two classes of layers, \mathcal{M}_1 and \mathcal{M}_2 . For every pair of layers $s, t \in \mathcal{M}_1$ such that $\gamma_{s,t}(\Gamma) = 1$, we fix the degree sequences $\vec{k}_{s \rightarrow t}^*$ and $\vec{k}_{t \rightarrow s}^*$. For every pair of layers $s \in \mathcal{M}_1$, $t \in \mathcal{M}_2$, $\gamma_{s,t}(\Gamma) = 1$ we fix the degree sequence $\vec{k}_{s \rightarrow t}^*$ from the layer in \mathcal{M}_1 to the layer in \mathcal{M}_2 (but not vice versa). We show that the resulting specific relative entropy is a mixture of the one in Theorem 1.8 and the one in Theorem 1.6. For $s = 1, \dots, M$ we set $A_s = \lim_{n_1, n_2, \dots, n_M \rightarrow \infty} \frac{n_s}{n}$.

Theorem 1.10. *Subject to the above relaxation,*

$$s_\infty = \sum_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_1 \cup \mathcal{M}_2 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \quad (1.50)$$

In particular,

$$s_\infty = 0 \iff A_s = 0 \forall s \in \{u \in \mathcal{M}_1 : \exists t \in \mathcal{M}_1 \cup \mathcal{M}_2 \text{ with } \gamma_{u,t}(\mathbf{\Gamma}) = 1\}. \quad (1.51)$$

Another way for relaxing constraints. We may think about another way for relaxing the constraints. We assume that $\gamma_{s,t}(\mathbf{\Gamma}) = 1$ for all $s, t = 1, 2, \dots, M$ and we fix $\vec{k}_s^* = \sum_{t=1}^M \vec{k}_{s \rightarrow t}^*$ for each $s = 1, 2, \dots, M$. This means that for each node we fix its degree sequence (no matter to which target layer, possibly its own layer). In this case we lose the multi-layer structure: constraints are no longer involving pairs of layers and the graphs are effectively uni-partite. This is the same case described in the configuration model of Theorem 1.1. There are still an extensive number of local constraints, and the ensembles are nonequivalent.

2 Discussion

In this section we discuss various important implications of our results. We first consider properties that are fully general, and afterwards focus on several special cases of empirical relevance.

2.1 General considerations

Poissonisation. The function g in (1.17) has an interesting interpretation, namely,

$$g(k) = S(\delta[k] \mid \text{Poisson}[k]) \quad (2.1)$$

is the relative entropy of the Poisson distribution with average k w.r.t. the Dirac distribution with average k . The specific relative entropy in (1.1) for the uni-partite setting can therefore be seen as a sum over k of contributions coming from the nodes with fixed, respectively, average degree k . The microcanonical ensemble forces the degree of these nodes to be exactly k (which corresponds to $\delta[k]$), while the canonical ensemble, under the sparseness condition in (1.14), forces their degree to be Poisson distributed with average k . The same condition ensures that in the limit as $n \rightarrow \infty$ the constraints act on the nodes essentially independently.

The same interpretation applies to Theorems 1.5–1.6 and 1.8–1.10. The result in Theorem 1.6(3) shows that in the bi-partite setting, when one of the layers tends to infinity while the other layer does not, Poissonisation does not set in fully. Namely, we have

$$s_n = \sum_{k=1}^n f(k) g_n(k), \quad g_n(k) = S(\delta[k] \mid \text{Bin}(n, \frac{k}{n})). \quad (2.2)$$

In words, the canonical ensemble forces the nodes in the infinite layer with average degree k to draw their degrees towards the n nodes in the finite layer essentially independently, giving rise to a binomial distribution. Only in the limit as $n \rightarrow \infty$ does this distribution converge to the Poisson distribution with average k .

Additivity vs. non-additivity. In all the other examples known so far in the literature, the generally accepted explanation for the breaking of ensemble equivalence is the presence of a non-additive energy, induced e.g. by long-range interactions [15], [16]. However, in the examples considered in the present paper, nonequivalence has a different origin, namely, the presence of an extensive number of local constraints. As we now show, this mechanism is completely unrelated to non-additivity and is therefore a novel mechanism for ensemble nonequivalence.

Intuitively, the energy of a system is additive when, upon partitioning the units of the system into non-overlapping subunits, the ‘interaction’ energy between these subunits is negligible with respect to the internal energy of the subunits themselves. The ‘physical’ size of the systems considered in this paper is given by the number n of nodes, i.e., we are defining the network to become ‘twice as large’ when the number of nodes is doubled. Think, for instance, of a population of n individuals and the corresponding social network connecting these individuals: we say that the size of the network doubles when the population doubles. Consistently, in (1.9) we have defined the specific relative entropy s_n by dividing S_n by n . In accordance with this reasoning, in order to establish whether in our systems ensemble equivalence has anything to do with energy additivity, we need to define the latter *node-wise*, i.e., with respect to partitioning the set of nodes into nonoverlapping subsets. Note that, in the presence of more than one layer, we have allowed for the number of nodes in some layer(s) to remain finite (in general, to grow subextensively) as the total number of nodes goes to infinity (see for instance Theorem 1.6). In such a situation it makes sense to study additivity only with respect to the nodes in those layers that are allowed to grow extensively in the thermodynamic limit.

Formally, if we let \mathcal{I} denote the union of all layers for which $A_s > 0$ (see (1.45)), then we say that the energy is *node-additive* if the Hamiltonian (1.6) can be written as

$$H(\mathbf{G}, \vec{\theta}) = \sum_{i \in \mathcal{I}} H_i(\mathbf{G}, \vec{\theta}) \quad \forall \mathbf{G} \in \mathcal{G}_n^\#, \quad (2.3)$$

where the $\{H_i\}_{i \in \mathcal{I}}$ do not depend on common subgraphs of \mathbf{G} (i.e., each of them can be restricted to a distinct subgraph of \mathbf{G}), and are therefore independent random variables.

The case of uni-partite graphs with fixed degree sequence (Theorem 1.1) is an example of ensemble *nonequivalence* with *non-additive* Hamiltonian, because the latter is defined as $H(\mathbf{G}, \vec{\theta}) = \sum_{i=1}^n \theta_i k_i(\mathbf{G})$ and cannot be rewritten in the form of (2.3) with independent $\{H_i(\mathbf{G}, \vec{\theta})\}$: the degrees $k_i(\mathbf{G})$ and $k_j(\mathbf{G})$ of any two distinct nodes i and j depend on a common subgraph of \mathbf{G} , i.e., the dyad $g_{i,j}(\mathbf{G})$. In the example of uni-partite graphs with a fixed total number of links (see (1.22)), the energy has the form $H(\mathbf{G}, \vec{\theta}) = \theta L(\mathbf{G}) = \frac{1}{2} \theta \sum_{i=1}^n k_i(\mathbf{G})$, which is still *non-additive*. However, the ensembles are in this case *equivalent* (see Theorem 1.4).

By contrast, the case of bi-partite graphs with fixed degree sequence on the top layer and the nodes in the other layer growing subextensively (case (3) of Theorem 1.6) is an example of ensemble *nonequivalence* with an *additive* Hamiltonian. Indeed, from (1.33) we see that $H(\mathbf{G}, \vec{\theta})$ is now a linear combination of the n_1 degrees of the nodes in layer Λ_1 , each of which depends only on the (bi-partite) subgraph obtained from the corresponding node of the top layer and all the nodes of the bottom layer. Here, unlike the uni-partite case, all these subgraphs are disjoint. Despite being node-additive, when $A_1 = 1$ ($c = 0$) this Hamiltonian leads to nonequivalence, as established in (1.39). Similar examples can be engineered using some of the relaxations in Section 1.3.4. Finally, the case of bi-partite graphs with fixed

total number of links (Theorem 1.7) is an example of ensemble *equivalence* with an *additive* Hamiltonian.

The four examples above show that additivity or non-additivity of the Hamiltonian does *not* influence the breaking of ensembles equivalence in the examples considered here. What matters is the *extensiveness* of the number of constraints. This observation was already made in [55], and is confirmed in full generality for the multi-layer setting treated in the present paper. Indeed, our results indicate that, whenever the number κ of constraints on the degrees is *subextensive*, i.e., $\kappa = o(n)$ where n is the number of nodes, ensemble equivalence is restored.

Note that the above notion of *node additivity* should not be confused with that of *edge additivity*, i.e., the fact that the Hamiltonian can be written as a sum over independent pairs of nodes. Due to the linearity of the chosen (local) constraints on the entries $\{g_{i,j}\}_{i,j=1}^n$ of the adjacency matrix of the graph \mathbf{G} , our examples are always edge-additive (irrespective of whether they are ensemble-equivalent), while they may or may not be node-equivalent, as we have seen. In either case, there is no relation between additivity and equivalence.

We stress again that the extensivity of the (local) constraints is, with respect to the mechanisms for nonequivalence already explored in the literature so far, an additional (and previously unrecognised) *sufficient* mechanism. It is obviously not the only one, and definitely *not a necessary one*, as exemplified by the fact that, in dense networks, nonequivalence has been found even in the presence of only two constraints, such as the total numbers of edges and triangles [50, 51, 52, 18]. However, while in the previous examples the breaking of equivalence arises from the nonlinearity (with respect to $\{g_{i,j}\}$) of some constraint and is typically found in a specific (usually critical) region of the parameter space separating phases where ensemble equivalence still applies, in our setting ensemble nonequivalence arises from the extensiveness of the number of (linear) constraints and extends to the entire space of parameters of the models. In this sense it is a stronger form of nonequivalence. Moreover, while the nonequivalence of network ensembles with a finite number of constraints was previously reported only for dense graphs, we are documenting it for the unexplored regime of sparse graphs.

A principled choice of ensembles. Ensembles of random graphs with constraints are used for many practical purposes. Two important examples are *pattern detection* and *network reconstruction*. For concreteness, we briefly illustrate these examples before we emphasize the implications that our results have for these and other applications.

Pattern detection is the identification of nontrivial structural properties in a real-world network, through the comparison of such network with a suitable null model [53]. For instance, *community detection* is the identification of groups of nodes that are more densely connected with each other than expected under a null model [26], [49] (in Section 2.2 we discuss the relation between our models and community detection in more detail). A null model is a random graph model that preserves some simple topological properties of the real network (typically local, like the degree sequence) and is otherwise completely random. So, maximum-entropy ensembles of graphs with given degrees are a key tool for pattern detection.

Network reconstruction employs purely local topological information to infer the higher-order structural properties of a real-world network [42]. This problem arises whenever the complete structure of a network is not known (for instance, due to confidentiality or privacy issues), but local properties are. An example relevant for the epidemiology of sexually transmitted diseases is the network of sexual contacts among people, for which only aggregate information (the total number of contacts with different partners) can be typically surveyed in

a population. In such cases, optimal inference about the network can be achieved by maximising the entropy subject to the known (local) constraints, which again leads to the ensembles with fixed degrees considered here.

The aforementioned applications, along with similar ones, make use of random graphs with local constraints. Our proof of nonequivalence of the corresponding ensembles have the following important implication. While for ensemble-equivalent models it makes practically no difference whether a microcanonical or canonical implementation is applied to large networks, for nonequivalent models different choices of the ensemble lead to asymptotically different results. As a consequence, while for applications based on ensemble-equivalent models the choice of the working ensemble can be arbitrary or be done on mathematical convenience (as usually done), for those based on nonequivalent models the choice should be principled, i.e., dictated by a theoretical criterion that indicates *a priori* which ensemble is the appropriate one.

Among the possible criteria, we suggest one that we believe appropriate whenever the available data are subject to (even small) errors, i.e., when the measured value \vec{C}^* entering as input in the construction of the random graph ensemble is, strictly speaking, the best available estimate for some unknown ‘true’ (error-free) value \vec{C}^\times . In this situation, we want that possible small deviations of \vec{C}^* from \vec{C}^\times result in small deviations of P_{mic}^* and P_{can}^* from the corresponding P_{mic}^\times and P_{can}^\times . Now, if $\vec{C}^* \neq \vec{C}^\times$ (no matter how “small” and in which norm this difference is taken), then P_{mic}^* will attach zero probability to any graph \mathbf{G}^\times that realises the ‘true’ constraint \vec{C}^\times : $P_{\text{mic}}^*(\mathbf{G}^\times) = 0$, while $P_{\text{mic}}^\times(\mathbf{G}^\times) \neq 0$. Indeed, P_{mic}^* and P_{mic}^\times will have non-overlapping supports, so they will sample distinct sets of graphs. This means that even small initial errors in the knowledge of the constraints will be severely propagated to the entire microcanonical ensemble, and inference based on the latter will be highly biased. In particular, the ‘true’ network will never be sampled by P_{mic}^* . On the other hand, if the difference between \vec{C}^* and \vec{C}^\times is small, then the difference between P_{can}^* and P_{can}^\times will also be small. So, even though \vec{C}^\times is unknown, any graph \mathbf{G}^\times that realises this value will be given a probability $P_{\text{can}}^*(\mathbf{G}^\times)$ that is nonzero and not very different from the probability $P_{\text{can}}^\times(\mathbf{G}^\times)$ that would be obtained by knowing the true value \vec{C}^\times . In general, small deviations of \vec{C}^* from \vec{C}^\times imply that $P_{\text{can}}^*(\mathbf{G})$ is not very different from $P_{\text{can}}^\times(\mathbf{G})$ for any graph \mathbf{G} , as desired. This implies that *even if \vec{C}^* is affected by small errors, then a principled choice of ensembles is the canonical one*. So, besides being the mathematically simpler option, we argue that canonical ensembles are also the most appropriate choice in the presence of ‘noise’. A similar claim was already made in [54], and is here strengthened by our proof of nonequivalence.

2.2 Special cases of empirical relevance

Different choices of the master graph Γ induce different structural features in the graphs of the ensemble $\mathcal{G}_n^\#$. Convenient choices allow us to consider certain classes of graphs that have been introduced recently to study appropriate types of real-world networks of empirical relevance. We discuss some of these choices below. The full generality of our results in Section 1.3.3 allows us to immediately draw conclusions about the (non)equivalence of the corresponding ensembles in each case of interest. As an important outcome of this discussion, all the empirically relevant ensembles of graphs turn out to be nonequivalent. In line with our general observation at the end of the previous section, this implies that a principled choice of ensembles is needed in all practical applications.

Scale-free uni-partite networks. Clearly, the trivial case when the master graph has a single node ($M = 1$) with a self-loop, i.e., $\gamma_{11}(\mathbf{\Gamma}) = 1$, corresponds to the class of uni-partite graphs we considered in Section 1.3.1. Many real-world networks, at least at a certain level of aggregation, admit such uni-partite representation. Examples include the Internet, the World Wide Web and many biological, social and economic networks. A common property displayed by most of these real-world networks is the presence of a “broad” empirical degree distribution, often consistent with a power-law distribution with an upper cut-off [12]. Networks with a power-law degree distribution are said to be *scale-free* [14]. This empirical observation implies that real-world networks are very different from Erdős-Rényi random graphs (which have a much narrower degree distribution) and are more closely reproduced by a configuration model with a truncated power-law degree distribution f_n (see (1.15)) of the form $f_n(k) = A_{\gamma,n} k^{-\gamma} \mathbb{I}_{1 \leq k \leq k_c(n)}$ with $\gamma > 1$, $A_{\gamma,n}$ the normalisation constant, and $\lim_{n \rightarrow \infty} k_c(n) = \infty$ and $k_c(n) = o(\sqrt{n})$. The so-called *structural cut-off* $k_c(n)$ makes the networks sparse, as in condition (1.14) [12]. Since $\lim_{n \rightarrow \infty} \|f_n - f\|_{\ell^1(g)} = 0$ with $f(k) = k^{-\gamma}/\zeta(\gamma)$ for $k \geq 1$ and 0 elsewhere, where ζ is the Riemann zeta-function, our result in (1.19) tells us that

$$s_\infty = \sum_{k \in \mathbb{N}} g(k) f(k) = \frac{1}{\zeta(\gamma)} \sum_{k \in \mathbb{N}} g(k) k^{-\gamma}. \quad (2.4)$$

Since $g(k) = \frac{1}{2} \log(2\pi k) + O(k^{-1})$ as $k \rightarrow \infty$, we find that s_∞ tends to 1 as $\gamma \rightarrow \infty$ and diverges like $\sim 1/2(\gamma-1)$ as $\gamma \downarrow 1$. This result shows that the simplest random graph ensemble consistent with the scale-free character of real-world networks is nonequivalent. Interestingly, as the tail exponent γ decreases, the degree distribution becomes broader and the degree of nonequivalence increases. A similar conclusion was drawn in [55].

Remark 2.1. Suppose that for each $n \in \mathbb{N}$ the degrees are drawn in an i.i.d. manner from the truncated degree distribution f_n . Suppose further that $\sum_{k \in \mathbb{N}_0} k f(k) < \infty$, i.e., $\gamma > 2$. Then, because $\sup_{n \in \mathbb{N}} \sum_{k \in \mathbb{N}_0} k f_n(k) = \sum_{k \in \mathbb{N}_0} k f(k) < \infty$, conditional on the sum of the degrees being even, the degree sequence is graphical with a probability tending to one as $n \rightarrow \infty$. This fact is the analogue of the result in [3] mentioned in Remark 1.2, and its proof is a straightforward extension of the argument in [3]. Truncation improves the chance of being graphical.

Multipartite networks. The case when the master graph has only $M = 2$ interconnected nodes *and no self-loops*, i.e., $\gamma_{1,2}(\mathbf{\Gamma}) = \gamma_{2,1}(\mathbf{\Gamma}) = 1$ and $\gamma_{1,1}(\mathbf{\Gamma}) = \gamma_{2,2}(\mathbf{\Gamma}) = 0$, coincides with the class of bi-partite graphs discussed in Section 1.3.2. Popular real-world examples relevant to economics, ecology and scientometrics are bank-firm, plant-pollinator and author-paper networks, respectively. In this case as well, empirical evidence shows that real-world bi-partite networks have broad degree distributions (at least on one of the two layers, and typically on both). Random graph models with only a global constraint on the total number of links (as in Theorem 1.7) are therefore unrealistic. The minimal ensemble that is consistent with the properties of most real-world bi-partite networks requires the specification of the degree sequence(s) as constraint(s) and is therefore nonequivalent.

A direct generalisation of the bi-partite case is when $\mathbf{\Gamma}$ is an M -dimensional matrix with zeroes along the diagonal and ones off the diagonal: $\gamma_{s,s}(\mathbf{\Gamma}) = 0 \ \forall s$ and $\gamma_{s,t}(\mathbf{\Gamma}) = 1$ for all $s \neq t$. The induced graphs in \mathcal{G}_n^\sharp have an “all-to-all” multipartite structure (i.e., links are allowed between all pairs of distinct layers, but not inside layers). From our Theorem 1.8 it follows that if the t -targeted degree sequences are specified as a constraint, then the relative

entropy in the all-to-all multipartite case is

$$s_\infty = \sum_{\substack{s,t=1 \\ s \neq t}}^M A_s \|f_{s \rightarrow t}\|_{\ell^1(g)} > 0, \quad (2.5)$$

which proves again ensemble nonequivalence.

Stochastic block-models. Another important example is when the master graph is a complete graph with all self-loops realised, i.e., $\gamma_{s,t}(\mathbf{\Gamma}) = 1$ for all s, t . This prescription generates the class of so-called *stochastic block-models*, which are very popular in the social network analysis literature [37], [39], [27]. The earliest and simplest stochastic block-model [37] is one where only the total numbers of links between all pairs of blocks (including within each block) are specified. When we identify blocks with layers, this model coincides with our relaxed model considered in Theorem 1.9, with $\mathcal{D} = \emptyset$. It follows as a corollary that this model is ensemble equivalent:

$$s_\infty = 0. \quad (2.6)$$

However, this model predicts that, within each block, the expected topological properties of the network are those of an Erdős-Rényi random graph, a property that is contradicted by empirical evidence. So, unless the number of blocks is chosen to be comparable with the number of nodes (which in our case is contradicted by the requirement that M is finite), the traditional block-model is not a good model of real-world networks.

More recently, emphasis has been put on the more realistic *degree-corrected stochastic block-model* [39], where an additional constraint is put on the degree of all nodes. An even more constrained variant of this model has been proposed in [27], where the constraints coincide with the t -targeted degree sequences $\{\vec{k}_{s \rightarrow t}\}_{s,t}$ among all pairs of blocks. To distinguish this model from the “generic” degree-corrected block-model, we call it the *targeted degree-corrected block-model*. This coincides with our model in Section 1.3.3, with the block structure given by the (complete) master graph. From Theorem 1.8 we calculate the relative entropy as

$$s_\infty = \sum_{s,t=1}^M A_s \|f_{s \rightarrow t}\|_{\ell^1(g)} > 0. \quad (2.7)$$

We can therefore conclude that, unlike the traditional block-model considered above, the targeted degree-corrected model is ensemble nonequivalent. We also note that, unlike stated in [27], the targeted degree-corrected block-model is not just a reparametrisation of the untargeted degree-corrected model. While fixing the targeted degree sequences automatically realises the constraints of the untargeted model, the converse is not true. Being a relaxation of the targeted model, we expect the untargeted model to have a relative entropy smaller than in (2.7), further illustrating the difference between the two models. Yet, we expect the relative entropy in the untargeted model to be strictly positive for, every choice of the degree sequence, since there is still an extensive number of active constraints. This would support the claim made in [48] that, *for small values of the degrees*, the degree-corrected block-models with soft and hard constraints are not equivalent in the thermodynamic limit. At the same time, it would contradict the claim made in the same reference that, *if all degrees become large* (but still in the sparse regime), the two ensembles become equivalent. Indeed, from the behaviour of $g(k)$ for large k (see (1.18)) and the normalisation by n in (1.9), we expect a finite specific relative entropy in that case as well.

Networks with community structure. Another very important class of graphs that are studied intensively in the literature are graphs with community structure [26], [49]. This class is related to the block-models described above, but is in general different. Community structure is loosely defined as the presence of groups of nodes that are more densely interconnected internally than with each other. One of the possible ways to quantitatively define the presence of communities in a real-world network is in terms of a positive difference between the realised number of intra-community links and the corresponding expected number calculated under a certain null hypothesis. This definition can be made more explicit by introducing the concept of *modularity* [26], [49]. For a graph with n nodes, a non-overlapping partition of nodes into M communities can be specified by the n -dimensional vector $\vec{\sigma}$, where the i -th entry $\sigma_i \in \{1, \dots, M\}$ is an integer number labelling the community to which node i is assigned by that particular partition. For a given real-world graph \mathbf{G}^* , the modularity is a function on the space of possible partitions, defined as

$$Q_{\mathbf{G}^*}(\vec{\sigma}) = K_{\mathbf{G}^*} \sum_{1 \leq i < j \leq n} (g_{ij}(\mathbf{G}^*) - \langle g_{ij} \rangle) \delta_{\sigma_i, \sigma_j}, \quad (2.8)$$

where $K_{\mathbf{G}^*}$ is an (inessential) normalisation constant (independent of the partition $\vec{\sigma}$) intended to have the property $Q_{\mathbf{G}^*} \in [-1, +1]$, and $\langle g_{ij} \rangle$ is the expected value of $g_{ij}(\mathbf{G})$ under the null hypothesis. The null hypothesis leads to a *null model* for the real-world network \mathbf{G}^* . The most popular choice for this null model is the canonical configuration model in the sparse regime, which gives $\langle g_{ij} \rangle = k_i^* k_j^* / 2L^*$ for $i \neq j$ and $\langle g_{ii} \rangle = 0$, where k_i^* , k_j^* and L^* are all calculated on \mathbf{G}^* (see (3.7) in the proof of Theorem 1.1).

Now, if the real-world network \mathbf{G}^* is indeed composed of communities, then the partition $\vec{\sigma}^\dagger$ that encodes these communities will be such that $Q_{\mathbf{G}^*}(\vec{\sigma}^\dagger) > 0$, i.e., the total number of links inside communities will be larger than the expected number under the null model. More stringently, the ‘optimal’ partition into communities can be defined as the one that maximises $Q_{\mathbf{G}^*}(\vec{\sigma})$, provided that the corresponding value $\max_{\vec{\sigma}} Q_{\mathbf{G}^*}(\vec{\sigma})$ is positive. Indeed, one of the most popular ways in which communities are looked for in real-world networks is through the process of modularity maximisation. The higher the value of the maximised modularity, the sharper the community structure. In practice, the problem of community detection is complicated by the possible existence of many local minima of $Q_{\mathbf{G}^*}(\vec{\sigma})$ and by the fact that $Q_{\mathbf{G}^*}(\vec{\sigma}^\dagger)$ may be positive even for “noisy communities”, i.e., communities induced by chance only out of randomness in the data.

In our setting, community structure can be easily induced in the multilayer graph ensemble $\mathcal{G}_n^\# = \mathcal{G}_{n_1, \dots, n_M}(\mathbf{\Gamma})$ through a convenient choice of the master graph $\mathbf{\Gamma}$ and of the constrained t -targeted degree sequences $\{\vec{k}_{s \rightarrow t}^*\}$. First, we identify the M layers $\{\Lambda_s\}$ with the desired communities and define the corresponding partition $\vec{\sigma}^\dagger$ through $\sigma_i^\dagger = \Lambda_s$ if $i \in \Lambda_s$. Next, we require that the master graph $\mathbf{\Gamma}$ has all possible self-loops, plus a desired number of additional edges that need not be maximal (pairs of distinct communities are not necessarily connected in real-world networks). Finally, we need to require that the t -targeted degree sequences induce an excess of intra-community links with respect to the null model, so that the modularity is at least positive, i.e., $Q_{\mathbf{G}^*}(\vec{\sigma}^\dagger) > 0$, and at best maximised by the desired partition, i.e.,

$\vec{\sigma}^\dagger = \operatorname{argmax}_{\vec{\sigma}} Q_{\mathbf{G}^*}(\vec{\sigma})$. To this end, we rewrite

$$\begin{aligned}
Q_{\mathbf{G}^*}(\vec{\sigma}^\dagger) &= K_{\mathbf{G}^*} \sum_{1 \leq i < j \leq n} (g_{ij}(\mathbf{G}^*) - \langle g_{ij} \rangle) \delta_{\sigma_i^\dagger, \sigma_j^\dagger} \\
&= \frac{K_{\mathbf{G}^*}}{2} \sum_{1 \leq i, j \leq n} (g_{ij}(\mathbf{G}^*) - \langle g_{ij} \rangle) \delta_{\sigma_i^\dagger, \sigma_j^\dagger} \\
&= \frac{K_{\mathbf{G}^*}}{2} \sum_{s=1}^M \sum_{i, j \in \Lambda_s} \left(g_{ij}(\mathbf{G}^*) - \frac{k_i^* k_j^*}{2L^*} \right) \\
&= \frac{K_{\mathbf{G}^*}}{2} \sum_{s=1}^M \left(L_{s,s}^* - \frac{1}{2L^*} \left(\sum_{i \in \Lambda_s} k_i^* \right)^2 \right) \\
&= \frac{K_{\mathbf{G}^*}}{2} \sum_{s=1}^M \left(L_{s,s}^* - \frac{1}{\sum_{s,t=1}^M L_{s,t}^*} \left(\sum_{t=1}^M L_{s,t}^* \right)^2 \right),
\end{aligned} \tag{2.9}$$

where we use $g_{ii}(\mathbf{G}^*) = \langle g_{ii} \rangle = 0$, $k_i^* = \sum_{t=1}^M k_i^{*t}$ and $2L^* = \sum_{s,t=1}^M L_{s,t}^*$. So, the weaker condition $Q_{\mathbf{G}^*}(\vec{\sigma}^\dagger) > 0$ is realised by requiring that $\{\vec{k}_{s \rightarrow t}^*\}$ satisfies the inequality

$$\sum_{s=1}^M L_{s,s}^* > \frac{\sum_{s=1}^M \left(\sum_{t=1}^M L_{s,t}^* \right)^2}{\sum_{s,t=1}^M L_{s,t}^*}, \tag{2.10}$$

where $L_{s,t}^* = \sum_{i \in \Lambda_s} k_i^{*t}$. The above inequality explicitly states that the number of realised intra-community edges counted in the left-hand side should be larger than the expected number calculated in the right-hand side. The stronger condition $\vec{\sigma}^\dagger = \operatorname{argmax}_{\vec{\sigma}} Q_{\mathbf{G}^*}(\vec{\sigma})$ should instead be enforced by looking for the specific $\{\vec{k}_{s \rightarrow t}^*\}$ that maximises (2.9).

Independently of how communities are induced in our framework, our results show that *ensembles of random graphs with community structure* (according to the definition above) are *nonequivalent*, with a relative entropy given by (1.48) where the degree distributions $\{f_{s \rightarrow t}\}$ are induced by suitable t -targeted degree sequences that realise (2.10) and possibly also $\vec{\sigma}^\dagger = \operatorname{argmax}_{\vec{\sigma}} Q_{\mathbf{G}^*}(\vec{\sigma})$.

Multiplex networks and time-varying graphs. Two other important classes of graphs that have recently gained attention are those of *multiplex networks* [11] and *time-varying graphs* [38].

Multiplex networks are networks where the same set of nodes can be connected by M different types of links [11]. Two examples, both studies in [32], are the multiplex of international trade in different products (where nodes are world countries and links of different type represent international trade in different products) and the multiplex of flights by different airlines (where nodes are airports and links of different type represent flights operated by different companies). An equivalent and widely used representation for a multiplex is one where a number M of layers is introduced, the same nodes are replicated in each layer, and inside each layer an ordinary graph is constructed, specified by all links of a single type. Links only exist within layers, and not across layers. Indeed, what ‘couples’ the different layers and makes a real-world multiplex different from a collection of independent layers is the empirical fact that the topological properties of the layer-specific networks are typically strongly (either positively or negatively) correlated. For instance, networks of trade in different products have

a similar structure, and most notably countries that are ‘hubs’ in one layer are likely to be hubs in other layers as well. By contrast, airports that are hubs for a domestic airline are likely not to be hubs for other domestic airlines [32]. This means that, for each node i in real-world networks, the M numbers of intra-layer links (i.e., the *intra-layer degrees*) are in general (anti)correlated.

Time-varying graphs are collections of temporal snapshots of the same network [38]. If the set of nodes in the network does not change with time, then a time-varying graph can be represented as a multiplex where each temporal snapshot is a single layer. (Note that multiplex networks themselves can vary over time [46].) Again, while not interacting directly via links, the different layers are mutually dependent because of empirical correlations between the properties of the same physical network across its temporal snapshots. Therefore this type of time-varying graphs can be treated in a way formally similar to that used for multiplex networks, the only difference being that a natural temporal ordering can be defined for the snapshots of time-varying graphs, while this is in general not true for the layers of a multiplex.

In our framework, a multiplex or time-varying network can be introduced by identifying each link type with a layer Λ_s and by requiring that the only edges of the master graph Γ are self-loops, i.e., $\gamma_{s,s}(\Gamma) = 1$ for $s = 1, M$ and $\gamma_{s,t}(\Gamma) = 0$ for $s \neq t$. Note that this specification, which implies $\vec{k}_{s \rightarrow t}^* = \vec{0}$ for $s \neq t$, is somehow ‘dual’ to the one defining all-to-all multipartite networks (see above). The fact that nodes in different layers are replicas of the same set of n nodes implies that $|\Lambda_s|$ is the same for all s , i.e., $n_s = n/M$. Finally, the ‘coupling’ between the topological properties of different layers can be introduced by assigning (anti)correlated t -targeted degree sequences, i.e., by choosing (anti)correlated entries for every pair of vectors $\vec{k}_{s \rightarrow s}^*$ and $\vec{k}_{t \rightarrow t}^*$, $s \neq t$. Real-world multiplexes, including the two examples made above, are well reproduced by such a model [32]. Our results imply that the relevant ensembles are nonequivalent. In particular, as a corollary of Theorem 1.8 we have

$$s_\infty = \frac{1}{M} \sum_{s=1}^M \|f_{s \rightarrow s}\|_{\ell^1(g)}. \quad (2.11)$$

So, the relative entropy between the microcanonical and canonical distributions is the average of the relative entropy of all layers, where for each layer s the relative entropy is the same as that obtained for a uni-partite network with n/M nodes and limiting degree distribution $f_{s \rightarrow s}$ (see Theorem 1.1). Moreover, the presence of correlations between $\vec{k}_{s \rightarrow s}^*$ and $\vec{k}_{t \rightarrow t}^*$ translate into dependencies between $\|f_{s \rightarrow s}\|_{\ell^1(g)}$ and $\|f_{t \rightarrow t}\|_{\ell^1(g)}$. In particular, in case of perfect correlation ($\vec{k}_{s \rightarrow s}^* = \vec{k}_{t \rightarrow t}^*$ for all s, t), all the degree distributions are equal to a common one $f_{s \rightarrow s} = f \forall s$, and we get

$$s_\infty = \|f\|_{\ell^1(g)}. \quad (2.12)$$

In this case, the degree of nonequivalence is the same as that obtained for a single uni-partite network with n/M nodes and limiting degree distribution f (see Theorem 1.1).

Interdependent multilayer networks. Finally, we discuss the class of *interdependent multilayer networks*, which are multiplex networks with the addition of inter-layer links [11]. Nodes in different layers are still replicas of the same set of nodes, so we still have $n_s = n/M$ for all s . Similarly, the topological properties of different intra-layer networks are still (anti)correlated, which can be again realised by choosing (anti)correlated entries for every pair of vectors $\vec{k}_{s \rightarrow s}^*$ and $\vec{k}_{t \rightarrow t}^*$, $s \neq t$. However, while we still require $\gamma_{s,s}(\Gamma) = 1$ for $s = 1, M$,

now we no longer require $\gamma_{s,t}(\mathbf{\Gamma}) = 0$ for $s \neq t$. Therefore the degree of nonequivalence can only increase with respect to (2.11). Indeed, Theorem 1.8 now leads to

$$s_\infty = \frac{1}{M} \sum_{\substack{s,t=1 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}}^M \|f_{s \rightarrow t}\|_{\ell^1(g)}, \quad (2.13)$$

which shows that the relative entropy is no longer only an average over the layer-specific relative entropies, since inter-layer relative entropies give additional contributions.

Networks of networks. A final class of graphs worth mentioning is the so-called *networks of networks*, sometimes constructed by different ‘micro-networks’ that are coupled together into a ‘macro-network’ where each node is a micro-network itself [22]. This class is similar to the interdependent multilayer networks considered above, but here there is no identification of the nodes in different layers to the same physical entity. An example is provided by multi-scale transport networks, where different cities are internally characterised by their local urban transport networks and at the same time are coupled through a long-distance inter-city transport network (like highways or flights). In our framework, this class of network can be induced by identifying the master graph $\mathbf{\Gamma}$ with the macro-network, and the M intra-layer subgraphs with the micro-networks. To have all micro-networks non-empty, the master graph must have all self-loops realised. This case is similar to the block-model mentioned above, but now the master graph itself can be chosen to have nontrivial structural properties, such as community structure, to resemble the specific properties of real-world networks of networks.

If the t -targeted degree sequences $\{\vec{k}_{s \rightarrow t}^*\}$ ($s, t = 1, M$) are all enforced as constraints, then the relative entropy is given by (1.48) with $\gamma_{s,s}(\mathbf{\Gamma}) = 1$ for all s . However, in this class of models it is often more natural to assume that the internal degree sequence $\vec{k}_{s \rightarrow s}^*$ of each micro-network (layer) s is enforced (in order to get realistic micro-network topologies), while between every pair s, t ($s \neq t$) of micro-networks only the number of links $L_{s,t}^*$ is fixed (because the topology of the master graph is already chosen in order to replicate the empirical macro-network). This leads to the relaxed model in Theorem 1.9 with $\mathcal{D} = \{(s, s) : s = 1, M\}$. The relative entropy is therefore

$$s_\infty = \sum_{s=1}^M A_s \|f_{s \rightarrow s}\|_{\ell^1(g)} \quad (2.14)$$

and is still positive, even though the links among micro-networks do not contribute to it.

3 Proof of the theorems

3.1 Proof of Theorem 1.1

Proof. The microcanonical number $\Omega_{\vec{k}^*}$ is not known in general, but asymptotic results exist in the *sparse regime* defined by the condition (1.14). For this regime it was shown in [6], [43] that

$$\Omega_{\vec{k}^*} = \frac{\sqrt{2} \left(\frac{2L^*}{e}\right)^{L^*}}{\prod_{i=1}^n k_i^*!} e^{-\left(\overline{k^{*2}}/2\overline{k^*}\right)^2 + \frac{1}{4} + o(n^{-1}\overline{k^*}^3)}, \quad (3.1)$$

where $\overline{k^*} = n^{-1} \sum_{i=1}^n k_i^*$ (average degree), $L^* = n\overline{k^*}/2$ (number of links), $\overline{k^{*2}} = n^{-1} \sum_{i=1}^n k_i^{*2}$ (average square degree). The canonical ensemble has Hamiltonian $H(\mathbf{G}, \vec{\theta}) = \sum_{i=1}^n \theta_i k_i(\mathbf{G})$,

where \mathbf{G} is a graph belonging to \mathcal{G}_n , and $k_i(\mathbf{G}) = \sum_{j \neq i} g_{i,j}(\mathbf{G})$ is the degree of the node i . The partition function equals

$$\begin{aligned} Z(\theta) &= \sum_{\mathbf{G} \in \mathcal{G}_n} e^{-H(\mathbf{G}, \vec{\theta})} = \sum_{\mathbf{G} \in \mathcal{G}_n} \prod_{1 \leq i < j \leq n} e^{-\theta_i g_{i,j}(\mathbf{G})} \\ &= \sum_{\mathbf{G} \in \mathcal{G}_n} \prod_{1 \leq i < j \leq n} e^{-(\theta_i + \theta_j) g_{i,j}(\mathbf{G})} = \prod_{1 \leq i < j \leq n} (1 + e^{-(\theta_i + \theta_j)}). \end{aligned} \quad (3.2)$$

The canonical probability equals

$$P_{\text{can}}(\mathbf{G} \mid \vec{\theta}) = \frac{\prod_{1 \leq i < j \leq n} e^{-(\theta_i + \theta_j) g_{i,j}(\mathbf{G})}}{Z(\vec{\theta})} = \prod_{1 \leq i < j \leq n} \frac{e^{-(\theta_i + \theta_j) g_{i,j}(\mathbf{G})}}{1 + e^{-(\theta_i + \theta_j)}}. \quad (3.3)$$

Setting $p_{ij}^* \equiv e^{-\theta_i^* - \theta_j^*} / (1 + e^{-\theta_i^* - \theta_j^*})$, and $\vec{\theta}^*$ such that

$$\sum_{j \neq i} \frac{e^{-\theta_i^* - \theta_j^*}}{1 + e^{-\theta_i^* - \theta_j^*}} = k_i^* \quad \forall i \quad (3.4)$$

we have

$$P_{\text{can}}(\mathbf{G}) = \prod_{1 \leq i < j \leq n} (p_{ij}^*)^{g_{ij}} (1 - p_{ij}^*)^{1 - g_{ij}}. \quad (3.5)$$

It is ensured by (1.14) that $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{1 \leq i < j \leq n} \widehat{p}_{ij}^{*2} = 0$, a condition under which we can show that (3.5) has the same asymptotic behaviour as

$$\widehat{P}_{\text{can}}(\mathbf{G}) = \prod_{1 \leq i < j \leq n} (\widehat{p}_{ij}^*)^{g_{ij}} (1 - \widehat{p}_{ij}^*)^{1 - g_{ij}}, \quad (3.6)$$

with

$$\widehat{p}_{ij}^* = e^{-\theta_i^* - \theta_j^*} = \frac{k_i^* k_j^*}{2L^*}. \quad (3.7)$$

Indeed,

$$\frac{1}{n} \log \left(\frac{\widehat{P}_{\text{can}}(\mathbf{G})}{P_{\text{can}}(\mathbf{G})} \right) = \frac{1}{n} \sum_{1 \leq i < j \leq n} g_{i,j} \log(1 - \widehat{p}_{ij}^*) - \frac{1}{n} \sum_{1 \leq i < j \leq n} \log(1 - \widehat{p}_{ij}^{*2}) \rightarrow 0, \quad n \rightarrow \infty, \quad (3.8)$$

because

$$\sum_{1 \leq i < j \leq n} g_{i,j} \log(1 - \widehat{p}_{ij}^*) \leq (m^*)^2 + O(\widehat{p}_{ij}^{*2}) \quad (3.9)$$

and

$$0 \leq \frac{1}{n} \sum_{1 \leq i < j \leq n} \widehat{p}_{ij}^{*2} = \frac{1}{2} \left[\frac{\sum_{i=1}^n k_i^2}{\sqrt{n} \sum_{i=1}^n k_i} \right]^2 \leq \frac{1}{2} \frac{(m^*)^2}{n} \rightarrow 0, \quad n \rightarrow \infty. \quad (3.10)$$

This implies $\sum_{1 \leq i < j \leq n} \ln(1 - p_{ij}^*) = -\sum_{1 \leq i < j \leq n} k_i^* k_j^* / 2L^* + o(n)$. Thus,

$$\ln P_{\text{can}}(\mathbf{G}^*) = \sum_{i=1}^n k_i^* \ln k_i^* - L^* \ln(2L^*) - L^* + o(n). \quad (3.11)$$

Combining (3.1) and (3.11), we obtain (recall (1.17))

$$S_n(P_{\text{mic}} | P_{\text{can}}) = \sum_{i=1}^n g(k_i^*) + o(n), \quad n \rightarrow \infty, \quad (3.12)$$

where $g(k) = \log\left(\frac{k!}{k^k e^{-k}}\right)$, as defined in (1.17). With the help of (1.15) this reads

$$n^{-1} S_n(P_{\text{mic}} | P_{\text{can}}) = \sum_{k \in \mathbb{N}_0} f_n(k) g(k) + o(1) = \|f_n\|_{\ell^1(g)} + o(1), \quad (3.13)$$

which together with (1.16) yields the claim. \square

3.2 Proof of Theorem 1.4

Proof. The microcanonical ensemble is easy: the number of graphs with a fixed fraction $\lambda \in (0, 1)$ of links is

$$\Omega_{L^*} = \binom{\binom{n}{2}}{L^*} = \binom{K}{\lambda K}, \quad K = \binom{n}{2}. \quad (3.14)$$

The canonical ensemble has the Hamiltonian $H(\mathbf{G}, \theta) = \theta L(\mathbf{G})$, where \mathbf{G} is a graph belonging to \mathcal{G}_n , and $L(\mathbf{G}) = \sum_{1 \leq i < j \leq n} g_{i,j}(\mathbf{G})$ is the number of links in \mathbf{G} . The partition function equals

$$Z(\theta) = \sum_{\mathbf{G} \in \mathcal{G}_n} e^{-H(\mathbf{G}, \theta)} = \sum_{\mathbf{G} \in \mathcal{G}_n} \prod_{1 \leq i < j \leq n} e^{-\theta g_{i,j}(\mathbf{G})} = \prod_{1 \leq i < j \leq n} (1 + e^{-\theta}). \quad (3.15)$$

The canonical probability equals

$$P_{\text{can}}(\mathbf{G} | \theta) = \frac{e^{-\sum_{1 \leq i < j \leq n} \theta g_{i,j}(\mathbf{G})}}{Z(\theta)} = \prod_{1 \leq i < j \leq n} \frac{e^{-\theta g_{i,j}(\mathbf{G})}}{1 + e^{-\theta}} = \prod_{1 \leq i < j \leq n} p^{g_{i,j}(\mathbf{G})} (1-p)^{1-g_{i,j}(\mathbf{G})} \quad (3.16)$$

with $p = \frac{e^{-\theta}}{1+e^{-\theta}}$. We search for θ^* such that

$$L^* = \sum_{1 \leq i < j \leq n} p^*, \quad p^* = \frac{e^{-\theta^*}}{1 + e^{-\theta^*}}. \quad (3.17)$$

It follows that $p^* = \lambda$. Thus,

$$\begin{aligned} \log P_{\text{mic}}(\mathbf{G}^*) &= -\log(K!) + \log(\lambda K)! + \log((1-\lambda)K)! \\ &= -K[\log K - 1] + \lambda K[\log \lambda K - 1] + [(1-\lambda)K][\log((1-\lambda)K) - 1] + o(n) \\ &= K \log(1-\lambda) + \lambda K \log\left(\frac{\lambda}{1-\lambda}\right) + o(n), \\ \log P_{\text{can}}(\mathbf{G}^*) &= \lambda K \log(\lambda) + (1-\lambda)K \log(1-\lambda). \end{aligned} \quad (3.18)$$

This in turn implies that

$$\lim_{n \rightarrow \infty} \frac{S_n(P_{\text{mic}} | P_{\text{can}})}{n} = 0. \quad (3.19)$$

\square

3.3 Proof of Theorem 1.5

Proof. We start by describing the canonical ensemble. The Hamiltonian is

$$\begin{aligned} H(\mathbf{G} | \vec{\theta}, \vec{\phi}) &= \sum_{i \in \Lambda_1} k_i(\mathbf{G}) \theta_i + \sum_{j \in \Lambda_2} k'_j(\mathbf{G}) \phi_j \\ &= \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \theta_i g_{i,j}(\mathbf{G}) + \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \phi_j g_{i,j}(\mathbf{G}) = \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} (\theta_i + \phi_j) g_{i,j}(\mathbf{G}). \end{aligned} \quad (3.20)$$

The partition function is

$$Z(\vec{\theta}, \vec{\phi}) = \sum_{\mathbf{G} \in \mathcal{G}_{n_1, n_2}} e^{-\sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} (\theta_i + \phi_j) g_{i,j}(\mathbf{G})} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} (1 + e^{-(\theta_i + \phi_j)}). \quad (3.21)$$

The canonical probability becomes

$$\begin{aligned} P_{\text{can}}(\mathbf{G} | \vec{\theta}, \vec{\phi}) &= \frac{e^{-\sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} (\theta_i + \phi_j) g_{i,j}(\mathbf{G})}}{Z(\vec{\theta}, \vec{\phi})} \\ &= \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} \frac{e^{-(\theta_i + \phi_j) g_{i,j}(\mathbf{G})}}{1 + e^{-(\theta_i + \phi_j)}} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} p_{i,j}^{g_{i,j}(\mathbf{G})} (1 - p_{i,j})^{1 - g_{i,j}(\mathbf{G})}, \end{aligned} \quad (3.22)$$

where $p_{i,j} = \frac{e^{-(\theta_i + \phi_j)}}{1 + e^{-(\theta_i + \phi_j)}}$. We search for $(\vec{\theta}^*, \vec{\phi}^*)$ that solves the system of equations

$$\begin{cases} \sum_{j \in \Lambda_2} p_{i,j}^* = k_i^*, \\ \sum_{i \in \Lambda_1} p_{i,j}^* = k'_j, \end{cases} \quad (3.23)$$

where $p_{i,j}^* = \frac{e^{-(\theta_i^* + \phi_j^*)}}{1 + e^{-(\theta_i^* + \phi_j^*)}}$. If \mathbf{G}^* is any graph in \mathcal{G}_{n_1, n_2} such that $k_i(\mathbf{G}^*) = k_i^*$ and $k'_j(\mathbf{G}^*) = k'_j$, then

$$P_{\text{can}}(\mathbf{G}^*) = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} p_{i,j}^{* g_{i,j}(\mathbf{G}^*)} (1 - p_{i,j}^*)^{1 - g_{i,j}(\mathbf{G}^*)}. \quad (3.24)$$

Under the sparseness condition (1.27), we can replace $p_{i,j}^*$ with the following quantity. Define $\hat{p}_{i,j}^* = e^{-(\theta_i^* + \phi_j^*)}$ and consider the system of equations

$$\begin{cases} \sum_{j \in \Lambda_2} \hat{p}_{i,j}^* = k_i^*, \\ \sum_{i \in \Lambda_1} \hat{p}_{i,j}^* = k'_j. \end{cases} \quad (3.25)$$

This has solution

$$\hat{p}_{i,j}^* = \frac{k_i^* k'_j}{L^*}, \quad L^* = \sum_{i \in \Lambda_1} k_i^* = \sum_{j \in \Lambda_2} k'_j. \quad (3.26)$$

We define

$$\hat{P}_{\text{can}}(\mathbf{G}) = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} (\hat{p}_{i,j}^*)^{g_{i,j}(\mathbf{G})} (1 - \hat{p}_{i,j}^*)^{1 - g_{i,j}(\mathbf{G})}, \quad (3.27)$$

and note that

$$\frac{1}{n_1 + n_2} \log \left(\frac{\hat{P}_{\text{can}}(\mathbf{G})}{P_{\text{can}}(\mathbf{G})} \right) \rightarrow 0, \quad n_1, n_2 \rightarrow \infty. \quad (3.28)$$

The crucial point is to prove that $\frac{1}{n_1+n_2} \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \widehat{p}_{i,j}^{*2} \rightarrow 0$. This allows us to write

$$\sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \log(1 - p_{i,j}^*) = - \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \frac{k_i^* k_j'^*}{L^*} + o(n_1 + n_2), \quad n_1, n_2 \rightarrow \infty. \quad (3.29)$$

Indeed,

$$0 \leq \frac{1}{n_1 + n_2} \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \widehat{p}_{i,j}^{*2} = \frac{1}{n_1 + n_2} \frac{\sum_{i \in \Lambda_1} k_i^{*2} \sum_{j \in \Lambda_2} k_j'^{*2}}{\sum_{i \in \Lambda_1} k_i^* \sum_{j \in \Lambda_2} k_j'^*} \leq \frac{m^* m'^* \sqrt{n_1 n_2}}{\sqrt{n_1 n_2} (n_1 + n_2)} \rightarrow 0, \quad (3.30)$$

because $m^* m'^* = o(L^{*2/3})$ implies $m^* m'^* = o(\sqrt{n_1 n_2})$.

Combining (3.24) and (3.29), we have

$$\begin{aligned} \log P_{\text{can}}(\mathbf{G}^*) &= \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} g_{i,j}(\mathbf{G}^*) \log \left(\frac{k_i^* k_j'^*}{L^*} \right) - \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \frac{k_i^* k_j'^*}{L^*} + o(n_1 + n_2) \\ &= \sum_{i \in \Lambda_1} k_i^* \log(k_i^*) + \sum_{j \in \Lambda_2} k_j'^* \log(k_j'^*) - L^* \log L^* - L^* + o(n_1 + n_2), \end{aligned} \quad (3.31)$$

which concludes our computation for the canonical ensemble.

Microcanonical probabilities come from the results in [34], where it is shown that, as $n \rightarrow \infty$, the number of bi-partite graphs with degree sequences \vec{k}^*, \vec{k}'^* on the two layers is given by

$$\Omega_{\vec{k}^*, \vec{k}'^*} = \frac{L^*!}{\prod_{i \in \Lambda_1} k_i^*! \prod_{j \in \Lambda_2} k_j'^*!} e^{o(n_1+n_2)}. \quad (3.32)$$

Hence

$$\log P_{\text{mic}}(\mathbf{G}^*) = -\log \Omega_{\vec{k}^*, \vec{k}'^*} = \sum_{i \in \Lambda_1} k_i^*! + \sum_{j \in \Lambda_2} k_j'^*! - \log(L^*!) + o(n_1 + n_2). \quad (3.33)$$

From (3.31) and (3.33) we get

$$\begin{aligned} S_{n_1+n_2}(P_{\text{can}} | P_{\text{mic}}) &= \log P_{\text{mic}}(\mathbf{G}^*) - \log P_{\text{can}}(\mathbf{G}^*) \\ &= \sum_{i \in \Lambda_1} \log \left(\frac{k_i^*!}{k_i^* k_i^*} \right) + \sum_{j \in \Lambda_2} \log \left(\frac{k_j'^*!}{k_j'^* k_j'^*} \right) \\ &\quad + L^* \log L^* + L^* - \log(L^*!) + o(n_1 + n_2) \\ &= \sum_{i \in \Lambda_1} g(k_i^*) + \sum_{j \in \Lambda_2} g(k_j'^*) + o(n_1 + n_2), \end{aligned} \quad (3.34)$$

where in the last line we use $L^* = \sum_{i \in \Lambda_1} k_i^* = \sum_{j \in \Lambda_2} k_j'^*$ and Stirling's approximation for $\log(L^*!)$. Since

$$\begin{aligned} n_1^{-1} \sum_{i \in \Lambda_1} g(k_i^*) &= \sum_{k \in \mathbb{N}_0}^{n_2} f_{1 \rightarrow 2}^{(n_1)}(k) g(k) = \|f_{1 \rightarrow 2}^{(n_1)}\|_{\ell^1(g)}, \\ n_2^{-1} \sum_{j \in \Lambda_2} g(k_j'^*) &= \sum_{k \in \mathbb{N}_0}^{n_1} f_{2 \rightarrow 1}^{(n_2)}(k) g(k) = \|f_{2 \rightarrow 1}^{(n_2)}\|_{\ell^1(g)}, \end{aligned} \quad (3.35)$$

we get, with the help of (1.28),

$$\lim_{n \rightarrow \infty} \frac{S_{n_1+n_2}(P_{\text{can}} | P_{\text{mic}})}{n_1 + n_2} = A_1 \|f_{1 \rightarrow 2}\|_{\ell^1(g)} + A_2 \|f_{2 \rightarrow 1}\|_{\ell^1(g)}, \quad (3.36)$$

which proves the claim. \square

3.4 Proof of Theorem 1.6

Proof. The number of bi-partite graphs with constraint \vec{k}^* on the top layer is

$$\Omega_{\vec{k}^*} = \prod_{i \in \Lambda_1} \binom{n_2}{k_i^*}. \quad (3.37)$$

In order to calculate the canonical probability, we calculate the partition function:

$$Z(\vec{\theta}) = \sum_{\mathbf{G} \in \mathcal{G}_{n_1, n_2}} e^{-\sum_{i \in \Lambda_1} \theta_i \sum_{j \in \Lambda_2} g_{i,j}(\mathbf{G})} = \sum_{\mathbf{G} \in \mathcal{G}_{n_1, n_2}} \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} e^{-\theta_i g_{i,j}(\mathbf{G})} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} [1 + e^{-\theta_i}]. \quad (3.38)$$

The canonical probability becomes

$$P_{\text{can}}(\mathbf{G} | \vec{\theta}) = \frac{e^{-\sum_{i \in \Lambda_1} \theta_i \sum_{j \in \Lambda_2} g_{i,j}(\mathbf{G})}}{Z(\vec{\theta})} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} \frac{e^{-\theta_i g_{i,j}(\mathbf{G})}}{1 + e^{-\theta_i}} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} p_i^{g_{i,j}(\mathbf{G})} (1 - p_i)^{1 - g_{i,j}(\mathbf{G})} \quad (3.39)$$

with $p_i = \frac{e^{-\theta_i}}{1 + e^{-\theta_i}}$. We search for θ_i^* such that

$$k_i^* = \sum_{j \in \Lambda_2} p_i^* = n_2 p_i^*, \quad p_i^* = \frac{e^{-\theta_i^*}}{1 + e^{-\theta_i^*}}. \quad (3.40)$$

It follows that $p_i = \frac{k_i^*}{n_2}$ (recall (1.34)). According to (1.11) we have

$$\begin{aligned} S_{n_1+n_2}(P_{\text{mic}} | P_{\text{can}}) &= \ln \frac{P_{\text{mic}}(\mathbf{G}^*)}{P_{\text{can}}(\mathbf{G}^*)} \\ &= - \sum_{i \in \Lambda_1} \log \binom{n_2}{k_i^*} - \sum_{i \in \Lambda_1} k_i^* \log \left(\frac{k_i^*}{n_2} \right) - \sum_{i \in \Lambda_1} (n_2 - k_i^*) \log \left(1 - \frac{k_i^*}{n_2} \right) \\ &= n_1 n_2 \log n_2 - \sum_{i \in \Lambda_1} \log \left[\binom{n_2}{k_i^*} k_i^{*k_i^*} (n_2 - k_i^*)^{(n_2 - k_i^*)} \right]. \end{aligned} \quad (3.41)$$

Abbreviate $U_a(x) \equiv \log \left[\binom{a}{x} x^x (a-x)^{a-x} \right]$ and write

$$S_{n_1+n_2}(P_{\text{mic}} | P_{\text{can}}) = n_1 n_2 \log n_2 - \sum_{i \in \Lambda_1} U_{n_2}(k_i^*) = n_1 n_2 \log n_2 - n_1 \sum_{k=0}^{n_2} f_{n_1}(k) U_{n_2}(k). \quad (3.42)$$

For the relative entropy per node this gives

$$\begin{aligned} s_{n_1+n_2} &= \frac{n_1}{n_1 + n_2} \sum_{k=0}^{n_2} f_{n_1}(k) n_2 \log n_2 - \frac{n_1}{n_1 + n_2} \sum_{k=0}^{n_2} f_{n_1}(k) U_{n_2}(k) \\ &= - \frac{n_1}{n_1 + n_2} \sum_{k=0}^{n_2} f_{n_1}(k) \log \text{Bin} \left(n_2, \frac{k}{n_2} \right) (k) = \frac{n_1}{n_1 + n_2} \|f_{n_1}\|_{\ell^1(g_{n_2})}. \end{aligned} \quad (3.43)$$

Case (1). Recall (1.17). Note that $x \mapsto z(x) = e^{g(x)}$ is non-decreasing:

$$\frac{z(x-1)}{z(x)} = \left(\frac{x}{x-1} \right)^{x-1} \frac{1}{e} \leq 1. \quad (3.44)$$

It therefore follows that

$$\begin{aligned}
\|f_{n_1}\|_{\ell^1(g_{n_2})} &= -\sum_{k=0}^{n_2} f_{n_1}(k) \log \text{Bin}\left(n_2, \frac{k}{n_2}\right)(k) = \sum_{k=0}^{n_2} f_{n_1}(k) \log \left(\frac{z(k)z(n_2-k)}{z(n_2)} \right) \\
&= \sum_{k \in \mathbb{N}_0} f_{n_1}(k) \log \left(\frac{z(k)z(n_2-k)}{z(n_2)} \right) \mathbb{I}_{k \leq n_2} \leq \sum_{k \in \mathbb{N}_0} \mathbb{I}_{0 \leq k \leq n_2} f_{n_1}(k) \log z(k) \\
&\leq \sum_{k \in \mathbb{N}_0} f_{n_1}(k) \log z(k) = \|f_{n_1}\|_{\ell^1(g)} < \infty.
\end{aligned} \tag{3.45}$$

By (1.36) and dominated convergence, we may exchange limit and sum to obtain

$$\lim_{n \rightarrow \infty} s_{n_1, n_2} = \lim_{n_2 \rightarrow \infty} \frac{n_1}{n_1 + n_2} \sum_{k \in \mathbb{N}_0} f_{n_1}(k) \lim_{n_2 \rightarrow \infty} \log \left(\frac{z(k)z(n_2-k)}{z(n_2)} \right) \mathbb{I}_{0 \leq k \leq n_2} = 0, \tag{3.46}$$

where we use that $\lim_{n \rightarrow \infty} \frac{n_1}{n_1 + n_2} = 0$ and $\lim_{n \rightarrow \infty} \frac{z(n_2-k)}{z(n_2)} = 1$ for all $k \in \mathbb{N}_0$.

Case (2). Using (3.45) and (1.36), we get

$$0 \leq s_{n_1, n_2} = \frac{n_1}{n_1 + n_2} \|f_{n_1}\|_{\ell^1(g)} \xrightarrow{n \rightarrow \infty} \frac{1}{1+c} \|f\|_{\ell^1(g)} = 0. \tag{3.47}$$

Case (3). Estimate

$$0 \leq \left| \|f_{n_1}\|_{\ell^1(g_{n_2})} - \|f_{n_1}\|_{\ell^1(g)} \right| \leq \|f_{n_1} - f\|_{\ell^1(g_{n_2})} \leq \|f_{n_1} - f\|_{\ell^1(g)} \xrightarrow{n \rightarrow \infty} 0. \tag{3.48}$$

Case (4).

$$0 \leq \left| \|f_{n_1}\|_{\ell^1(g_{n_2})} - \|f\|_{\ell^1(g)} \right| \leq \sum_{k \in \mathbb{N}_0} |f_{n_1}(k) - f(k)| |g_{n_2}(k) \mathbb{I}_{k \leq n_2} - g(k)| \leq 2 \|f_{n_1} - f\|_{\ell^1(g)}. \tag{3.49}$$

Since $\frac{n_1}{n_1 + n_2} = \frac{1}{1 + \frac{n_2}{n_1}} \rightarrow \frac{1}{1+c}$, the claim follows. \square

3.5 Proof of Theorem 1.7

Proof. The microcanonical ensemble is easy: the number of bi-partite graphs with a fixed fraction $\lambda \in (0, 1)$ of links is

$$\Omega_{L^*} = \binom{n_1 n_2}{L^*} = \binom{n_1 n_2}{\lambda n_1 n_2}. \tag{3.50}$$

The canonical ensemble has the Hamiltonian $H(\mathbf{G}, \theta) = \theta L(\mathbf{G})$, where \mathbf{G} is a bi-partite graph belonging to \mathcal{G}_{n_1, n_2} , and $L(\mathbf{G}) = \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} g_{i, j}(\mathbf{G})$ is the number of links in \mathbf{G} . The partition function equals

$$Z(\theta) = \sum_{\mathbf{G} \in \mathcal{G}_{n_1, n_2}} e^{-H(\mathbf{G}, \theta)} = \sum_{\mathbf{G} \in \mathcal{G}_{n_1, n_2}} \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} e^{-\theta g_{i, j}(\mathbf{G})} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} (1 + e^{-\theta}). \tag{3.51}$$

The canonical probability equals

$$P_{\text{can}}(\mathbf{G} \mid \vec{\theta}) = \frac{e^{-\sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} \theta g_{i,j}(\mathbf{G})}}{Z(\vec{\theta})} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} \frac{e^{-\theta g_{i,j}(\mathbf{G})}}{1 + e^{-\theta}} = \prod_{i \in \Lambda_1} \prod_{j \in \Lambda_2} p^{g_{i,j}(\mathbf{G})} (1-p)^{1-g_{i,j}(\mathbf{G})} \quad (3.52)$$

with $p = \frac{e^{-\theta}}{1+e^{-\theta}}$. We search for θ^* such that

$$L^* = \sum_{i \in \Lambda_1} \sum_{j \in \Lambda_2} p^*, \quad p^* = \frac{e^{-\theta^*}}{1 + e^{-\theta^*}}. \quad (3.53)$$

It follows that $p^* = \lambda$. Thus,

$$\begin{aligned} \log P_{\text{mic}}(\mathbf{G}^*) &= -\log(n_1 n_2)! + \log(\lambda n_1 n_2)! + \log((1-\lambda)n_1 n_2)! \\ &= -n_1 n_2 [\log n_1 n_2 - 1] + \lambda n_1 n_2 [\log \lambda n_1 n_2 - 1] \\ &\quad + [1 - \lambda n_1 n_2] [\log((1-\lambda)n_1 n_2) - 1] + o(n_1 + n_2) \\ &= n_1 n_2 \log(1-\lambda) + \lambda n_1 n_2 \log\left(\frac{\lambda}{1-\lambda}\right) + o(n_1 + n_2), \\ \log P_{\text{can}}(\mathbf{G}^*) &= n_1 n_2 \log(1-\lambda) + \lambda n_1 n_2 \log\left(\frac{\lambda}{1-\lambda}\right). \end{aligned} \quad (3.54)$$

This in turn implies that

$$\lim_{n_1, n_2 \rightarrow \infty} \frac{S_{n_1+n_2}(P_{\text{mic}} \mid P_{\text{can}})}{n_1 + n_2} = 0. \quad (3.55)$$

□

3.6 Proof of Theorem 1.8

Proof. The proof is based on the previous theorems. We start by looking at the Hamiltonian of the system. For each admitted pair of layers ($\gamma_{s,t}(\mathbf{\Gamma}) = 1$) we define Lagrange multipliers $\vec{\theta}_{s \rightarrow t} = (\theta_1^{(t)}, \dots, \theta_{n_s}^{(t)})$. The Hamiltonian equals

$$\begin{aligned} H(\mathbf{G} \mid \vec{\theta}_{s \rightarrow t}; s, t = 1, \dots, M, \gamma_{s,t}(\mathbf{\Gamma}) = 1) \\ &= \sum_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma}) = 1}} \sum_{\substack{i \in \Lambda_s \\ j \in \Lambda_t}} (\theta_i^t + \theta_j^s) g_{i,j}(\mathbf{G}) + \sum_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma}) = 1}}^M \sum_{\substack{i,j \in \Lambda_s \\ i < j}} (\theta_i^s + \theta_j^s) g_{i,j}(\mathbf{G}) \\ &= \sum_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma}) = 1}} \sum_{\substack{i \in \Lambda_s \\ j \in \Lambda_t}} H_{s,t}(\mathbf{G}^{(st)} \mid \vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s}) + \sum_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma}) = 1}}^M \sum_{\substack{i,j \in \Lambda_s \\ i < j}} H_{s,s}(\mathbf{G}^{(ss)} \mid \vec{\theta}_{s \rightarrow s}), \end{aligned} \quad (3.56)$$

where

$$\begin{aligned} H_{s,t}(\mathbf{G}^{(st)} \mid \vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s}) &= \sum_{\substack{i \in \Lambda_s \\ j \in \Lambda_t \\ i < j}} (\theta_i^t + \theta_j^s) g_{i,j}(\mathbf{G}^{(st)}), \\ H_{s,s}(\mathbf{G}^{(ss)} \mid \vec{\theta}_{s \rightarrow s}) &= \sum_{\substack{i,j \in \Lambda_s \\ i < j}} (\theta_i^s + \theta_j^s) g_{i,j}(\mathbf{G}^{(ss)}), \end{aligned} \quad (3.57)$$

and $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) is the bi-partite (uni-partite) graph between layers s and t (inside layer s) obtained from the multi-partite graph \mathbf{G} . The $n_s \times n_t$ matrix representing the bi-partite graph has, for each $i \in \Lambda_s$ and $j \in \Lambda_t$, elements $g_{i,j}(\mathbf{G}^{(st)}) = g_{i,j}(\mathbf{G})$. Note that $H_{s,t}(\mathbf{G}^{(st)} | \vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s})$ is the Hamiltonian of the bi-partite graph $\mathbf{G}^{(st)}$ between layers s and t with constraints $\vec{k}_{s \rightarrow t}^*$, and $H_{s,s}(\mathbf{G}^{(ss)} | \vec{\theta}_{s \rightarrow s})$ is the Hamiltonian of the uni-partite graph $\mathbf{G}^{(ss)}$ of the layer s with constraints $\vec{k}_{s \rightarrow s}^*$.

The partition function of the canonical ensemble equals

$$\begin{aligned}
Z(\vec{\theta}_{s \rightarrow t}; s, t = 1, \dots, M, \gamma_{s,t}(\Gamma) = 1) &= \sum_{\mathbf{G} \in \mathcal{G}_{n_1, \dots, n_M}(\Gamma)} e^{-H(\mathbf{G} | \vec{\theta}_{s \rightarrow t}; s, t = 1, 2, \dots, M: \gamma_{s,t}(\Gamma) = 1)} \\
&= \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\Gamma) = 1}} \sum_{\mathbf{G}^{(st)} \in \mathcal{G}_{n_s, n_t}} e^{-H_{s,t}(\mathbf{G}^{(st)} | \vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s})} \prod_{\substack{s=1 \\ \gamma_{s,s}(\Gamma) = 1}}^M \sum_{\mathbf{G}^{(ss)} \in \mathcal{G}_{n_s, n_s}} e^{-H_{s,s}(\mathbf{G}^{(ss)} | \vec{\theta}_{s \rightarrow s})} \\
&= \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\Gamma) = 1}} Z^{(st)}(\vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s}) \prod_{\substack{s=1 \\ \gamma_{s,s}(\Gamma) = 1}}^M Z^{(ss)}(\vec{\theta}_{s \rightarrow s}),
\end{aligned} \tag{3.58}$$

where $Z^{(st)}(\vec{\theta}_{s \rightarrow t}, \vec{\theta}_{t \rightarrow s})$ is the partition function of the set of bi-partite graphs \mathcal{G}_{n_s, n_t} with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer, and $Z^{(ss)}(\vec{\theta}_{s \rightarrow s})$ is the partition function of the set of graph \mathcal{G}_{n_s} with constraint $\vec{k}_{s \rightarrow s}^*$. The canonical ensemble is

$$P_{\text{can}}(\mathbf{G}) = \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\Gamma) = 1}} P_{\text{can}}^{(st)}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ \gamma_{s,s}(\Gamma) = 1}}^M P_{\text{can}}^{(ss)}(\mathbf{G}^{(ss)}), \tag{3.59}$$

where $P_{\text{can}}^{(st)}(\mathbf{G}^{(st)})$ is the canonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer, and $P_{\text{can}}^{(ss)}(\mathbf{G}^{(ss)})$ is the canonical probability of the uni-partite graph $\mathbf{G}^{(ss)}$ with constraint $\vec{k}_{s \rightarrow s}^*$.

We can split the microcanonical probability as products of microcanonical probabilities for simpler cases. The number of graphs with constraints \vec{C}^* is

$$\begin{aligned}
\Omega_{\vec{k}_{s \rightarrow t}^*; s, t \in \{1, \dots, M\}, \gamma_{s,t}(\Gamma) = 1} &= \left| \left\{ \mathbf{G} \in \mathcal{G}_{n_1, \dots, n_M}(\Gamma) : \sum_{j \in \Lambda_t} g_{i,j}(\mathbf{G}) = k_i^{*t} \quad \forall i \in \Lambda_s \quad \forall \gamma_{s,t} = 1 \right\} \right| \\
&= \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\Gamma) = 1}} \left| \left\{ \mathbf{G}^{(st)} \in \mathcal{G}_{n_s, n_t} : \sum_{j \in \Lambda_t} g_{i,j}(\mathbf{G}^{(st)}) = k_i^{*t} \quad \forall i \in \Lambda_s, \sum_{i \in \Lambda_s} g_{i,j}(\mathbf{G}^{(st)}) = k_j^{*s} \quad \forall j \in \Lambda_t \right\} \right| \\
&\quad \prod_{\substack{s=1 \\ \gamma_{s,s}(\Gamma) = 1}}^M \left| \left\{ \mathbf{G}^{(ss)} \in \mathcal{G}_{n_s} : \sum_{j \in \Lambda_s} g_{i,j}(\mathbf{G}^{(ss)}) = s_i^{*h} \quad \forall i \in \Lambda_s \right\} \right| \\
&= \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\Gamma) = 1}} \Omega_{\vec{k}_{s \rightarrow t}^*, \vec{k}_{t \rightarrow s}^*} \prod_{\substack{s=1 \\ \gamma_{s,s}(\Gamma) = 1}}^M \Omega_{\vec{k}_{s \rightarrow s}^*}.
\end{aligned} \tag{3.60}$$

This means the microcanonical probability can be factorised as

$$P_{\text{mic}}(\mathbf{G}) = \prod_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} P_{\text{mic}}^{(st)}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}}^M P_{\text{mic}}^{(ss)}(\mathbf{G}^{(ss)}), \quad (3.61)$$

where $P_{\text{mic}}^{(st)}(\mathbf{G}^{(st)})$ is the microcanonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer, and $P_{\text{mic}}^{(ss)}(\mathbf{G}^{(ss)})$ is the microcanonical probability of the uni-partite graph $\mathbf{G}^{(ss)}$ with constraint $\vec{k}_{s \rightarrow s}^*$.

Equations (3.59) and (3.61) imply that the relative entropy equals the sum

$$S_n(P_{\text{mic}} | P_{\text{can}}) = \sum_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} S_n(P_{\text{mic}}^{(st)} | P_{\text{can}}^{(st)}) + \sum_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}}^M S_n(P_{\text{mic}}^{(ss)} | P_{\text{can}}^{(ss)}). \quad (3.62)$$

We can now apply Theorems 1.1 and 1.5 to get the asymptotic relative entropy per nodes as

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}} | P_{\text{can}})}{n} \\ &= \sum_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{(st)} | P_{\text{can}}^{(st)})}{n} + \sum_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}}^M \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{(ss)} | P_{\text{can}}^{(ss)})}{n} \\ &= \sum_{\substack{1 \leq s < t \leq M \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \{A_s \|f_{s \rightarrow t}\|_{\ell^1(g)} + A_t \|f_{t \rightarrow s}\|_{\ell^1(g)}\} + \sum_{\substack{s=1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}}^M \{A_s \|f_{s \rightarrow s}\|_{\ell^1(g)}\} \\ &= \sum_{\substack{s,t=1 \\ \gamma_{s,t}(\mathbf{\Gamma})}}^M A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \end{aligned} \quad (3.63)$$

□

3.7 Proof of Theorem 1.9

Proof. We start by studying the Hamiltonian. For each pair (s, t) of layers in \mathcal{D} , we define Lagrange multipliers $\vec{\theta}_{s \rightarrow t} = (\theta_1^t, \dots, \theta_{n_s}^t)$. For each pair (s, t) of layers in \mathcal{L} , we define a Lagrange multiplier $\theta_{s,t}$. The Hamiltonian is

$$\begin{aligned} H(\mathbf{G} | \vec{\theta}_{s \rightarrow t}, \theta_{l,m}; (s, t) \in \mathcal{D}, (l, m) \in \mathcal{L}) \\ = H_{\mathcal{D}}(\mathbf{G} | \vec{\theta}_{s \rightarrow t}; (s, t) \in \mathcal{D}) + H_{\mathcal{L}}(\mathbf{G} | \theta_{l,m}; (l, m) \in \mathcal{L}) \end{aligned} \quad (3.64)$$

with

$$\begin{aligned} H_{\mathcal{D}}(\mathbf{G} | \vec{\theta}_{s \rightarrow t}; (s, t) \in \mathcal{D}) &= \sum_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{D}}} \sum_{\substack{i \in \Lambda_s \\ j \in \Lambda_t}} (\theta_i^t + \theta_j^s) g_{i,j}(\mathbf{G}) + \sum_{\substack{s=1 \\ (s,s) \in \mathcal{D}}}^M \sum_{\substack{i,j \in \Lambda_s \\ i < j}} (\theta_i^s + \theta_j^s) g_{i,j}(\mathbf{G}), \\ H_{\mathcal{L}}(\mathbf{G} | \theta_{s,t}; (s, t) \in \mathcal{L}) &= \sum_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{L}}} \sum_{\substack{i \in \Lambda_s \\ j \in \Lambda_t}} (\theta_{s,t}) g_{i,j}(\mathbf{G}) + \sum_{\substack{s=1 \\ (s,s) \in \mathcal{L}}}^M \sum_{\substack{i,j \in \Lambda_s \\ i < j}} (\theta_{s,s}) g_{i,j}(\mathbf{G}). \end{aligned} \quad (3.65)$$

Consequently, the canonical ensemble is

$$P_{\text{can}}(\mathbf{G}) = P_{\text{can}}^{\mathcal{D}}(\mathbf{G})P_{\text{can}}^{\mathcal{L}}(\mathbf{G}) \quad (3.66)$$

with

$$\begin{aligned} P_{\text{can}}^{\mathcal{D}}(\mathbf{G}) &= \prod_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{D}}} P_{\text{can}}^{(st)\mathcal{D}}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ (s,s) \in \mathcal{D}}}^M P_{\text{can}}^{(ss)\mathcal{D}}(\mathbf{G}^{(ss)}), \\ P_{\text{can}}^{\mathcal{L}}(\mathbf{G}) &= \prod_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{L}}} P_{\text{can}}^{(st)\mathcal{L}}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ (s,s) \in \mathcal{L}}}^M P_{\text{can}}^{(ss)\mathcal{L}}(\mathbf{G}^{(ss)}). \end{aligned} \quad (3.67)$$

Here,

- $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) is the bi-partite (uni-partite) graph between layers s and t (and itself) obtained from the multi-partite graph \mathbf{G} . The $n_s \times n_t$ ($n_s \times n_s$) matrix representing this bi-partite (uni-partite) graph has, for each $i \in \Lambda_s$ and $j \in \Lambda_t$ (for each $i, j \in \Lambda_s$), elements $g_{i,j}(\mathbf{G}^{(st)}) = g_{i,j}(\mathbf{G})$ ($g_{i,j}(\mathbf{G}^{(ss)}) = g_{i,j}(\mathbf{G})$).
- $P_{\text{can}}^{(st)\mathcal{D}}(\mathbf{G}^{(st)})$ ($P_{\text{can}}^{(ss)\mathcal{D}}(\mathbf{G}^{(ss)})$) is the canonical probability of the bi-partite (uni-partite) graph $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer (with constraint $\vec{k}_{s \rightarrow s}^*$).
- $P_{\text{can}}^{(st)\mathcal{L}}(\mathbf{G}^{(st)})$ ($P_{\text{can}}^{(ss)\mathcal{L}}(\mathbf{G}^{(ss)})$) is the canonical probability of the bi-partite (uni-partite) graph $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) with constraint $L_{s,t}^*$ ($L_{s,s}^*$).

We can split the microcanonical probability as products of microcanonical probabilities of simpler cases. The number of graphs with such a type of constraints is

$$\Omega_{\vec{k}_{s \rightarrow t}^*, L_{l,m}^*; (s,t) \in \mathcal{D}, (l,m) \in \mathcal{L}} = \Omega_{\vec{k}_{s \rightarrow t}^*; (s,t) \in \mathcal{D}} \Omega_{L_{l,m}^*; (l,m) \in \mathcal{L}}. \quad (3.68)$$

This means that the microcanonical probability can be factorised as

$$P_{\text{mic}}(\mathbf{G}) = P_{\text{mic}}^{\mathcal{D}}(\mathbf{G})P_{\text{mic}}^{\mathcal{L}}(\mathbf{G}) \quad (3.69)$$

with

$$\begin{aligned} P_{\text{mic}}^{\mathcal{D}}(\mathbf{G}) &= \prod_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{D}}} P_{\text{mic}}^{(st)\mathcal{D}}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ (s,s) \in \mathcal{D}}}^M P_{\text{mic}}^{(ss)\mathcal{D}}(\mathbf{G}^{(ss)}), \\ P_{\text{mic}}^{\mathcal{L}}(\mathbf{G}) &= \prod_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{L}}} P_{\text{mic}}^{(st)\mathcal{L}}(\mathbf{G}^{(st)}) \prod_{\substack{s=1 \\ (s,s) \in \mathcal{L}}}^M P_{\text{mic}}^{(ss)\mathcal{L}}(\mathbf{G}^{(ss)}). \end{aligned} \quad (3.70)$$

Here,

- $P_{\text{mic}}^{(st)\mathcal{D}}(\mathbf{G}^{(st)})$ ($P_{\text{mic}}^{(ss)\mathcal{D}}(\mathbf{G}^{(ss)})$) is the microcanonical probability of the bi-partite (uni-partite) graph $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer (with constraint $\vec{k}_{s \rightarrow s}^*$).
- $P_{\text{mic}}^{(st)\mathcal{L}}(\mathbf{G}^{(st)})$ ($P_{\text{mic}}^{(ss)\mathcal{L}}(\mathbf{G}^{(ss)})$) is the microcanonical probability of the bi-partite (uni-partite) graph $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) with constraint $L_{s,t}^*$ ($L_{s,s}^*$).

The relative entropy becomes

$$S_n(P_{\text{mic}} | P_{\text{can}}) = S_n(P_{\text{mic}}^{\mathcal{D}} | P_{\text{can}}^{\mathcal{D}}) + S_n(P_{\text{mic}}^{\mathcal{L}} | P_{\text{can}}^{\mathcal{L}}). \quad (3.71)$$

It follows that

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}} | P_{\text{can}})}{n} \\ &= \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{D}} | P_{\text{can}}^{\mathcal{D}})}{n} + \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{L}} | P_{\text{can}}^{\mathcal{L}})}{n}. \end{aligned} \quad (3.72)$$

Using Theorem 1.8 we get

$$\lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{D}} | P_{\text{can}}^{\mathcal{D}})}{n} = \sum_{(s,t) \in \mathcal{D}} A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \quad (3.73)$$

Moreover,

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{L}} | P_{\text{can}}^{\mathcal{L}})}{n} \\ &= \lim_{n_1, \dots, n_M \rightarrow \infty} \sum_{\substack{1 \leq s < t \leq M \\ (s,t) \in \mathcal{L}}} \frac{S_n(P_{\text{mic}}^{(st)\mathcal{L}} | P_{\text{can}}^{(st)\mathcal{L}})}{n} + \lim_{n_1, \dots, n_M \rightarrow \infty} \sum_{\substack{s=1 \\ (s,s) \in \mathcal{L}}}^M \frac{S_n(P_{\text{mic}}^{(ss)\mathcal{L}} | P_{\text{can}}^{(ss)\mathcal{L}})}{n}. \end{aligned} \quad (3.74)$$

Using Theorems 1.4 and 1.7, we get

$$\lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{(st)\mathcal{L}} | P_{\text{can}}^{(st)\mathcal{L}})}{n} = \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{(ss)\mathcal{L}} | P_{\text{can}}^{(ss)\mathcal{L}})}{n} = 0, \quad (3.75)$$

which proves the claim. \square

3.8 Proof of Theorem 1.10

Proof. The proof is based on the previous theorems. For each pair of layers $s, t \in \mathcal{M}_1$ we define Lagrange multipliers $\vec{\theta}_{s \rightarrow t} = (\theta_1^t, \dots, \theta_{n_s}^t)$ and $\vec{\theta}_{t \rightarrow s} = (\theta_1^s, \dots, \theta_{n_t}^s)$. For each pair of layers $s \in \mathcal{M}_1, t \in \mathcal{M}_2$ we define $\vec{\theta}_{s \rightarrow t} = (\theta_1^t, \dots, \theta_{n_s}^t)$. The Hamiltonian is

$$\begin{aligned} & H(\mathbf{G} | \vec{\theta}_{s \rightarrow t}; s \in \mathcal{M}_1, t \in \mathcal{M}_1 \cup \mathcal{M}_2, \gamma_{s,t}(\mathbf{\Gamma}) = 1) \\ &= \sum_{\substack{s,t \in \mathcal{M}_1 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow t} \vec{s}_{s \rightarrow t}(\mathbf{G}) + \sum_{\substack{s \in \mathcal{M}_1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow s} \vec{s}_{s \rightarrow s}(\mathbf{G}) + \sum_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow t} \vec{s}_{s \rightarrow t}(\mathbf{G}) \\ &= H_{\mathcal{M}_1 \rightarrow \mathcal{M}_1} + H_{\mathcal{M}_1 \rightarrow \mathcal{M}_2}, \end{aligned} \quad (3.76)$$

with

$$\begin{aligned} H_{\mathcal{M}_1 \rightarrow \mathcal{M}_1} &= \sum_{\substack{s,t \in \mathcal{M}_1 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow t} \vec{s}_{s \rightarrow t}(\mathbf{G}) + \sum_{\substack{s \in \mathcal{M}_1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow s} \vec{s}_{s \rightarrow s}(\mathbf{G}), \\ H_{\mathcal{M}_1 \rightarrow \mathcal{M}_2} &= \sum_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} \vec{\theta}_{s \rightarrow t} \vec{s}_{s \rightarrow t}(\mathbf{G}). \end{aligned} \quad (3.77)$$

Consequently, the canonical ensemble is

$$P_{\text{can}}(\mathbf{G}) = P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1}(\mathbf{G}) P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2}(\mathbf{G}) \quad (3.78)$$

with

$$\begin{aligned} P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1}(\mathbf{G}) &= \prod_{\substack{s,t \in \mathcal{M}_1 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} P_{\text{can}}^{(st) \text{top}, \text{bot}}(\mathbf{G}^{(st)}) \prod_{\substack{s \in \mathcal{M}_1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}} P_{\text{can}}^{(ss)}(\mathbf{G}^{(ss)}), \\ P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2}(\mathbf{G}) &= \prod_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} P_{\text{can}}^{(st) \text{top}}(\mathbf{G}^{(st)}). \end{aligned} \quad (3.79)$$

Here,

- $\mathbf{G}^{(st)}$ ($\mathbf{G}^{(ss)}$) is the bi-partite (uni-partite) graph between layers s and t (itself) obtained from the multi-partite graph \mathbf{G} . The $n_s \times n_t$ ($n_s \times n_s$) matrix representing this bi-partite (uni-partite) graph has, for each $i \in \Lambda_s$ and $j \in \Lambda_t$ (for each $i, j \in s$), elements $g_{i,j}(\mathbf{G}^{(st)}) = g_{i,j}(\mathbf{G})$ ($g_{i,j}(\mathbf{G}^{(ss)}) = g_{i,j}(\mathbf{G})$).
- $P_{\text{can}}^{(st) \text{top}, \text{bot}}(\mathbf{G}^{(st)})$ is the canonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer.
- $P_{\text{can}}^{(ss)}(\mathbf{G}^{(ss)})$ is the canonical probability of the uni-partite graph $\mathbf{G}^{(ss)}$ with constraint $\vec{k}_{s \rightarrow s}^*$.
- $P_{\text{can}}^{(st) \text{top}}(\mathbf{G}^{(st)})$ is the canonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraint $\vec{k}_{s \rightarrow t}^*$ on the top layer.

We can split the microcanonical probability as products of microcanonical probabilities for simpler cases. The number of graphs with such a type of constraints is

$$\Omega_{\vec{k}_{s \rightarrow t}^*; s \in \mathcal{M}_1, t \in \mathcal{M}_1 \cup \mathcal{M}_2, \gamma_{s,t}(\mathbf{\Gamma})=1} = \Omega_{\vec{k}_{s \rightarrow t}^*; s, t \in \mathcal{M}_1, \gamma_{s,t}(\mathbf{\Gamma})=1} \Omega_{\vec{k}_{s \rightarrow t}^*; s \in \mathcal{M}_1, t \in \mathcal{M}_2, \gamma_{s,t}(\mathbf{\Gamma})=1}. \quad (3.80)$$

This means that the microcanonical probability can be factorised as

$$P_{\text{mic}}(\mathbf{G}) = P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1}(\mathbf{G}) P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2}(\mathbf{G}) \quad (3.81)$$

with

$$\begin{aligned} P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1}(\mathbf{G}) &= \prod_{\substack{s,t \in \mathcal{M}_1 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} P_{\text{mic}}^{(st) \text{top}, \text{bot}}(\mathbf{G}^{(st)}) \prod_{\substack{s \in \mathcal{M}_1 \\ \gamma_{s,s}(\mathbf{\Gamma})=1}} P_{\text{mic}}^{(ss)}(\mathbf{G}^{(ss)}), \\ P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2}(\mathbf{G}) &= \prod_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s,t}(\mathbf{\Gamma})=1}} P_{\text{mic}}^{(st) \text{top}}(\mathbf{G}^{(st)}). \end{aligned} \quad (3.82)$$

Here,

- $P_{\text{mic}}^{(st) \text{top}, \text{bot}}(\mathbf{G}^{(st)})$ is the microcanonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraints $\vec{k}_{s \rightarrow t}^*$ on the top layer and $\vec{k}_{t \rightarrow s}^*$ on the bottom layer.
- $P_{\text{mic}}^{(ss)}(\mathbf{G}^{(ss)})$ is the microcanonical probability of the uni-partite graph $\mathbf{G}^{(ss)}$ with constraint $\vec{k}_{s \rightarrow s}^*$.

- $P_{\text{mic}}^{(st)top}(\mathbf{G}^{(st)})$ is the microcanonical probability of the bi-partite graph $\mathbf{G}^{(st)}$ with constraint $\vec{k}_{s \rightarrow t}^*$ on the top layer.

The relative entropy becomes

$$S_n(P_{\text{mic}} | P_{\text{can}}) = S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1}) + S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2}). \quad (3.83)$$

It follows that

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{can}} | P_{\text{can}})}{n} \\ &= \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1})}{n} + \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2})}{n}. \end{aligned} \quad (3.84)$$

Using again Theorem 1.8 we get

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_1})}{n} \\ &= \sum_{\substack{s, t \in \mathcal{M}_1 \\ \gamma_{s, t}(\Gamma) = 1}} \{A_s \|f_{s \rightarrow t}\|_{\ell^1(g)} + A_t \|f_{t \rightarrow s}\|_{\ell^1(g)}\} + \sum_{\substack{s \in \mathcal{M}_1 \\ \gamma_{s, s}(\Gamma) = 1}} A_s \|f_{s \rightarrow s}\|_{\ell^1(g)} \\ &= \sum_{\substack{s, t \in \mathcal{M}_1 \\ \gamma_{s, t}(\Gamma) = 1}} A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}. \end{aligned} \quad (3.85)$$

From Theorem 1.6 we get

$$\begin{aligned} & \lim_{n_1, \dots, n_M \rightarrow \infty} \frac{S_n(P_{\text{mic}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2} | P_{\text{can}}^{\mathcal{M}_1 \rightarrow \mathcal{M}_2})}{n} \\ &= \lim_{n_1, \dots, n_M \rightarrow \infty} \sum_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s, t}(\Gamma) = 1}} \frac{S_n(P_{\text{mic}}^{(st)\mathcal{M}_1 \rightarrow \mathcal{M}_2} | P_{\text{can}}^{(st)\mathcal{M}_1 \rightarrow \mathcal{M}_2})}{n} = \sum_{\substack{s \in \mathcal{M}_1, t \in \mathcal{M}_2 \\ \gamma_{s, t}(\Gamma) = 1}} A_s \|f_{s \rightarrow t}\|_{\ell^1(g)}, \end{aligned} \quad (3.86)$$

which concludes the proof. \square

References

- [1] K. Anand and G. Bianconi, *Phys. Rev. E* **80**, 045102 (2009).
- [2] K. Anand and G. Bianconi, *Phys. Rev. E* **82**, 011116 (2010).
- [3] R. Arratia and T.M. Liggett, *Ann. Appl. Probab.* **15**, 652 (2005).
- [4] J. Barré and B. Goncalves, *Physica A* **386**, 212 (2007).
- [5] J. Barré, D. Mukamel, and S. Ruffo, *Phys. Rev. Lett.* **87**, 030601 (2001).
- [6] E.A. Bender, *Discrete Math.* **10**, 217 (1974).
- [7] E.A. Bender and E.R. Canfield, *J. Comb. Theory (A)* **24**, 296 (1978).
- [8] G. Bianconi, *Physical Review E* **87**, 062806 (2013).

- [9] G. Bianconi, A.C.C. Coolen, C.J. Perez Vicente, *Phys. Rev. E* **78**, 016114 (2008).
- [10] M. Blume, V.J. Emery, and R.B. Griffiths, *Phys. Rev. A* **4**, 1071 (1971).
- [11] S. Boccaletti, G. Bianconi, R. Criado, C.I. Del Genio, J. Gómez-Gardeñes, M. Romance, I. Sendiña-Nadalj, Z. Wang, and M. Zanin, *Physics Reports* **544**, 1 (2014).
- [12] M. Boguñá, R. Pastor-Satorras, and A. Vespignani, *The European Physical Journal B* **38**, 205-209 (2004).
- [13] B. Bollobás, *European J. Combin.* **1**, 311 (1980).
- [14] G. Caldarelli, *Scale-Free Networks: Complex Webs in Nature and Technology* (OUP, Oxford, 2007).
- [15] A. Campa, T. Dauxois, D. Fanelli, and S. Ruffo, *Physics of Long-Range Interacting Systems* (OUP, Oxford, 2014).
- [16] A. Campa, T. Dauxois, and S. Ruffo, *Physics Reports* **480**, 57 (2009).
- [17] P.J. Carrington, J. Scott, and S. Wasserman, *Models and Methods in Social Network Analysis*, Vol. 28 (Cambridge University Press, 2005).
- [18] S. Chatterjee, and P. Diaconis, *The Annals of Statistics* **41**, 2428 (2013).
- [19] P.-H. Chavanis, *Astron. & Astrophys.* **401**, 15 (2003).
- [20] F. Chung, and L. Lu, *Proceedings of the National Academy of Sciences* **99**, 15879 (2002).
- [21] M. D'Agostino *et al.*, *Phys. Lett. B* **473**, 219 (2000).
- [22] G. D'Agostino, and A. Scala, *Networks of Networks: The Last Frontier of Complexity*, Vol. 340 (Springer, Berlin, 2014).
- [23] R.S. Ellis, K. Haven, and B. Turkington, *J. Stat. Phys.* **101**, 999 (2000).
- [24] R.S. Ellis, K. Haven, and B. Turkington, *Nonlinearity* **15**, 239 (2002).
- [25] R.S. Ellis, H. Touchette, and B. Turkington, *Physica A* **335**, 518 (2004).
- [26] S. Fortunato, *Physics Reports* **486**, 75 (2010).
- [27] P. Fronczak, A. Fronczak, and M. Bujok, *Physical Review E* **88**, 032810 (2013).
- [28] D. Garlaschelli, *New Journal of Physics* **11**, 073005 (2009).
- [29] D. Garlaschelli, S.E. Ahnert, T. Fink, and G. Caldarelli, *Entropy* **15**, 3148 (2013).
- [30] D. Garlaschelli, F. den Hollander, and A. Roccaverde, *Nieuw Archief voor Wiskunde* **5/16**, 207 (2015).
- [31] D. Garlaschelli and M.I. Loffredo, *Physical Review E* **78**, 015101 (2008).
- [32] V. Gemmetto, and D. Garlaschelli, *Scientific Reports* **5**, 9120 (2015).
- [33] J.W. Gibbs, *Elementary Principles of Statistical Mechanics*, Yale University Press, New Haven, Connecticut (1902).

- [34] C. Greenhill, B.D. McKay, and X. Wang, *J. Comb. Theory, Series A* **113**, 291 (2006).
- [35] P. Hertel and W. Thirring, *Annals of Physics* **63**, 520 (1971).
- [36] R. van der Hofstad, *Random Graphs and Complex Networks, Volume I*, to appear with Cambridge University Press.
- [37] P.W. Holland, K.B. Laskey, and S. Leinhardt, *Social Networks* **5**, 109 (1983).
- [38] P. Holme, and J. Saramäki, *Physics Reports* **519**, 97 (2012).
- [39] B. Karrer, and M.E.J. Newman, *Physical Review E* **83**, 016107 (2011).
- [40] D. Lynden-Bell, *Physica A* **263**, 293 (1999).
- [41] D. Lynden-Bell, R. Wood, *Monthly Notices of the Royal Astronomical Society* **138**, 495 (1968).
- [42] R. Mastrandrea, T. Squartini, G. Fagiolo, and D. Garlaschelli, *New Journal of Physics* **16**, 043022 (2014).
- [43] B.D. McKay and N.C. Wormald, *Combinatorica* **11**, 369 (1991).
- [44] M. Molloy, and B. Reed, *Random Structures & Algorithms* **6**, 161 (1995).
- [45] M.E. Newman, S.H. Strogatz, and D.J. Watts, *Physical review E* **64**, 026118 (2001).
- [46] V. Nicosia, G. Bianconi, V. Latora, and M. Barthelemy, *Phys. Rev. Lett.* **111**, 058701 (2013).
- [47] J. Park, and M.E. Newman, *Physical Review E* **70**, 066117 (2004).
- [48] T.P. Peixoto, *Physical Review E* **85**, 056122 (2012).
- [49] M.A. Porter, J.P. Onnela, and P.J. Mucha, *Notices of the AMS* **56**, 1082-1097.
- [50] C. Radin, and L. Sadun, *J. Phys. A: Math. Theor.* **46**, 305002 (2013).
- [51] C. Radin, and L. Sadun, *Journal of Statistical Physics* **158**, 853 (2015).
- [52] C. Radin and M. Yi, *The Annals of Applied Probability* **23**, 2458-2471 (2013).
- [53] T. Squartini, and D. Garlaschelli, *New Journal of Physics* **13**, 083001 (2011).
- [54] T. Squartini, R. Mastrandrea, and D. Garlaschelli, *New J. Phys.* **17**, 023052 (2015).
- [55] T. Squartini, J. de Mol, F. den Hollander, and D. Garlaschelli, *Phys. Rev. Lett.* **115**, 268701 (2015).
- [56] W. Thirring, *Zeitschrift für Physik* **235**, 339 (1970).
- [57] H. Touchette, *J. Stat. Phys.* **159**, 987 (2015).
- [58] H. Touchette, R.S. Ellis, and B. Turkington, *Physica A* **340**, 138 (2004).
- [59] R.M. Ziff, G.E. Uhlenbeck, and M. Kac, *Physics Reports* **32**, 169 (1977).