



Universiteit
Leiden
The Netherlands

spFRET studies of nucleosome dynamics modulated by histone modifications, histone variants and neighboring nucleosomes

Buning, R.

Citation

Buning, R. (2015, January 15). *spFRET studies of nucleosome dynamics modulated by histone modifications, histone variants and neighboring nucleosomes*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/31477>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/31477>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/31477> holds various files of this Leiden University dissertation.

Author: Buning, Ruth

Title: spFRET studies of nucleosome dynamics modulated by histone modifications, histone variants and neighboring nucleosomes

Issue Date: 2015-01-15

spFRET studies of nucleosome dynamics modulated by histone modifications, histone variants and neighboring nucleosomes

PROEFSCHRIFT

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. C. J. J. M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op 15 januari 2015
klokke 11:15 uur

door


Ruth Buning

geboren te Amsterdam

in 1981

Promotiecommissie

Promotor:	Prof. dr. T. Schmidt	(Universiteit Leiden)
Co-promotor:	Dr. ir. S. J. T. van Noort	(Universiteit Leiden)
Overige leden:	Prof. dr. E. R. Eliel	(Universiteit Leiden)
	Dr. D. M. Heinrich	(Universiteit Leiden)
	Dr. H. van Ingen	(Universiteit Leiden)
	Prof. dr. H. Schiessel	(Universiteit Leiden)
	Prof. dr. C. A. M. Seidel	(Heinrich-Heine Universität Düsseldorf)



Universiteit
Leiden

Casimir PhD Series, Delft-Leiden, 2014-34

ISBN: 978-90-8593-206-2

An electronic version of this thesis can be found at <https://openaccess.leidenuniv.nl>

Dit werk maakt deel uit van het onderzoekprogramma van de Stichting voor Fundamenteel Onderzoek der Materie (FOM), die financieel wordt ondersteund door de Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO).

Aan Oma Fien

Contents

Aim and outline of this thesis	1
1 spFRET on nucleosomes	3
1.1 Introduction	6
1.2 The nucleosome	6
1.2.1 Nucleosome dynamics	8
1.3 FRET	10
1.4 Reconstitution of nucleosomes	11
1.4.1 Relation between FRET efficiency and length of unwrapped DNA	14
1.5 spFRET techniques	14
1.5.1 Confocal microscopy	14
1.5.2 Widefield TIRF	15
1.5.3 Alternating Laser Excitation	16
1.6 DNA breathing studied with spFRET	18
1.7 DNA sequence effects	23
1.8 Histone modifications	23
1.9 Nucleosome remodeling	24
1.10 DNA accessibility in nucleosome arrays	27
1.11 Conclusions	29
2 Making spFRET experiments with nucleosomes succeed	31
2.1 Introduction	32
2.2 Reconstitution of nucleosomes with FRET pairs	32
2.2.1 DNA substrate	32
2.2.2 Nucleosome reconstitution	35
2.3 Nucleosomes in a polyacrylamide gel	36
2.4 Sample preparation for spFRET	37
2.4.1 Dilution to single-molecule concentrations	37
2.4.2 Buffer conditions	39

2.4.3	Surface effects	40
2.5	Single-molecule fluorescence spectroscopy	40
2.5.1	Confocal setup	40
2.5.2	Single-molecule burst selection	42
2.5.3	Caveats in spFRET	42
2.6	Quantitative comparison of multiple FRET techniques	44
2.6.1	Bulk fluorescence spectroscopy	44
2.6.2	PAGE and spFRET	46
2.7	Conclusions	50
3	Genetic acetylation defines H3 K56Ac effects	53
3.1	Introduction	56
3.2	Results	57
3.2.1	Production of site-specifically acetylated histones	57
3.2.2	H3 K56Ac does not affect salt-dependent nucleosome stability	59
3.2.3	H3 K56Ac increases DNA breathing in mononucleosomes	60
3.2.4	Formation of higher-order chromatin structure in nucleosome arrays is unaffected by H3 K56Ac	64
3.2.5	The effect of H3 K56Ac on chromatin remodeling	66
3.3	Discussion	68
3.4	Experimental procedures	70
3.4.1	Expression and purification of acetylated histones	70
3.4.2	Single-molecule FRET	71
4	H2A.Z containing nucleosomes with spFRET	73
4.1	Introduction	76
4.2	Materials and Methods	78
4.2.1	Preparation of DNA constructs	78
4.2.2	Preparation of histone octamers	78
4.2.3	Nucleosome reconstitution	80
4.2.4	Polyacrylamide gel electrophoresis	80
4.2.5	spFRET sample preparation	80
4.2.6	Single-molecule fluorescence microscopy	81
4.2.7	Single-molecule data analysis	81
4.3	Results	82
4.3.1	Gel electrophoresis reveals near 100 % reconstitution yield	82
4.3.2	spFRET in gel shows H2A.Z-nucleosomes are more stable	84
4.3.3	FRET distributions are temperature independent	84
4.3.4	spFRET measurements on nucleosomes depend critically on slide passivation	86

4.3.5	H2A.Z nucleosomes are more stable during extended periods of storage	88
4.3.6	Xenopus H2A.Z nucleosomes also more stable	89
4.4	Discussion	89
4.5	Conclusions	95
5	Breathing in dinucleosomes	97
5.1	Introduction	100
5.2	Materials and Methods	102
5.2.1	Preparation of DNA constructs	102
5.2.2	Nucleosome reconstitution	102
5.2.3	Polyacrylamide gel electrophoresis	102
5.2.4	Sample preparation	104
5.2.5	Single-molecule fluorescence microscopy	104
5.2.6	Single-molecule data analysis	106
5.3	Results	108
5.3.1	Gel electrophoresis of (di)nucleosome reconstitutions	108
5.3.2	FRET distributions of mononucleosomes	110
5.3.3	Linker DNA increases DNA breathing	110
5.3.4	DNA breathing is affected by neighboring nucleosomes	112
5.4	Discussion	115
5.4.1	Sample handling	115
5.4.2	Sample heterogeneity	115
5.4.3	Linker DNA increases breathing in mononucleosomes	117
5.4.4	Dinucleosomes	117
5.5	Conclusions	119
	References	121
	Summary	131
	Samenvatting	135
	List of publications	143
	Curriculum Vitae	145
	Dankwoord	147

Aim and outline of this thesis

The principle of encoding genetic information in DNA is common in all living organisms and is very elegant in its simplicity. Yet the translation from genetic code to a functioning living organism that reacts to its environmental conditions is tremendously complex and far from understood. At the basis of the regulation of the genetic code (DNA) in eukaryotes¹, is its organization into nucleosomes: 10 nm wide structures, in which ~ 150 basepairs (bp) of DNA are wrapped around a disk of proteins, the histone octamer. Arrays of nucleosomes are organized in fiber-like structures called chromatin, ultimately forming the well-known chromosome, that, in its most compact state during cell division, can easily be visualized with a normal light microscope. This thesis focusses on the smallest units in chromatin, the nucleosomes, and how the physical nature of these nucleosomes allows for dynamic accessibility of DNA for cellular processes.

Nucleosomes realize a huge compaction of the DNA. In humans, the 2 m of DNA (6 billion bp) present in every cell, forming a coil of $\sim 100 \mu\text{m}$ in diameter if not condensed [1], fits in the cell nucleus of only $\sim 5 \mu\text{m}$ in diameter. The compaction in the form of nucleosomes provides the essential 'tool' to regulate accessibility of DNA. By thermal fluctuations, part of the nucleosomal DNA transiently unwraps from the histone protein core (DNA breathing), making it accessible for the cellular machinery. Changes in the binding energy between DNA and histones alter the breathing probability, which is realized in the cell by modifications to the histones or by incorporating histone variants. The linking of nucleosomes to their neighbors poses constraints on DNA breathing as well. It is therefore not trivial to understand the regulation of DNA accessibility in a physical manner.

To distinguish open nucleosomes (where part of the nucleosomal DNA is unwrapped) from closed nucleosomes (where the nucleosomal DNA is fully wrapped), the use of Fluorescence (or Förster) Resonance Energy Transfer (FRET) is a logical choice. FRET has been used intensively for nucleosome research. Ensemble measurements record the average FRET efficiency of large nucleosome populations. Infor-

¹to which all animals, plants and fungi belong

mation about the conformational heterogeneity or kinetic processes that occur in the sample can only be revealed with single-pair FRET (spFRET), in which FRET is measured in individual nucleosomes.

In this thesis, **Chapter 1** is a review of experiments that have used spFRET to elucidate single nucleosome dynamics. Such experiments show that nucleosomes are in general open for tens of milliseconds and for 10 % of the time. We investigated the influence of modulations of the histone proteins and the presence of linker DNA and neighboring nucleosomes on DNA breathing with spFRET. Several challenges we experienced in spFRET with nucleosomes are described in **Chapter 2**, along with suggestions how to deal with them. In **Chapter 3** we show that specific acetylation of the histone octamers at H3K56 enhances unwrapping of nucleosomal DNA. In **Chapter 4**, nucleosomes containing either the canonical histone H2A or the variant H2A.Z are investigated. We reveal that nucleosomes containing H2A.Z are more stable than their H2A-containing counterparts and thus can act as an accessibility switch for nucleosomal DNA. Finally, in **Chapter 5** we show that electrostatic interactions between linker DNAs and nucleosomes, and interactions between neighboring nucleosomes, enhance DNA breathing.