



Universiteit
Leiden
The Netherlands

The potential use of dendritic cells in mouse models of atherosclerosis

Habets, K.L.L.

Citation

Habets, K. L. L. (2009, December 8). *The potential use of dendritic cells in mouse models of atherosclerosis*. Retrieved from <https://hdl.handle.net/1887/14484>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14484>

Note: To cite this publication please use the final published version (if applicable).

Chapter 5

Deficiency of TGF- β signaling in CD11c⁺ cells accelerates atherosclerosis by disrupting T-cell homeostasis

-Manuscript submitted-

D. Lievens¹; K.L.L. Habets²; Y. Laouar³; A. Robertson⁴;
L. Beckers¹; E. Wijnands¹; T. Rademakers⁵; M. van
Zandvoort⁵; L. Boon⁶; M. Rudling⁷; R.A. Flavell⁸; M.J.A.P.
Daemen¹; J. Kuiper²; E.A. Biessen¹ and E. Lutgens^{1,9}

Abstract

TGF- β is an anti-inflammatory and pro-fibrotic cytokine that plays a bridging role between the innate and adaptive immune system. However, the exact liaison between TGF- β , the innate and the adaptive immune system in atherosclerosis is still unclear. Here we investigate the effect of deficiency of TGF- β signaling in cells of the innate immune system (ie CD11c⁺ DCs, NK cells and macrophages) in atherosclerosis, by using transgenic mice with a targeted functional inactivation of TGF- β RII signaling in CD11c⁺ cells (ApoE^{-/-}CD11cDNR). ApoE^{-/-}CD11cDNR mice were sacrificed at 20 wks of age and plaques in the aortic root were analyzed. ApoE^{-/-}CD11cDNR mice exhibited a 2-fold increase in plaque area and showed a substantial increase in CD45⁺ leukocytes and CD3⁺, CD4⁺ and CD8⁺ T-cells, while macrophage content was not affected. Moreover, a significant reduction in α -smooth muscle cell (ASMA) and collagen content was observed in plaques of ApoE^{-/-}CD11cDNR mice. FACS analysis of lymphoid organs of ApoE^{-/-}CD11cDNR mice revealed an increase in both CD4⁺ and CD8⁺ CD44^{high}CD62L^{low} effector memory T-cells, whereas the number of CD4⁺CD25⁺FoxP3⁺ regulatory T-cells had increased. In the supernatant of CD3/CD28 stimulated splenocytes from ApoE^{-/-}CD11cDNR mice, elevated levels of Th1 (IFN- γ), Th2 (IL-4, IL-10) and Th17 (IL-17) cytokines were detected. In conclusion, deficient TGF- β signaling in CD11c⁺ cells results in a disturbed T-cell homeostasis, thereby accelerating atherosclerosis and inducing a pro-inflammatory plaque phenotype.

Introduction

Atherosclerosis is a chronic inflammatory disease of the arterial wall.¹⁻³ Immune cells from both the innate and the adaptive immune system orchestrate the formation and progression of atherosclerotic plaques by controlling their structure, composition and expansion.³ This complex interplay between different cells and cell types of the immune system is coordinated by a broad array of pro- and anti-inflammatory chemokines and cytokines. The balance between pro- and anti-inflammatory modulators is decisive in the degree of atherosclerosis progression and the plaque phenotype that develops.⁴⁻⁷

One of the central regulators of the immune response is transforming growth factor beta (TGF- β).^{8, 9} Three highly conserved isoforms of TGF- β have been described: TGF- β 1, TGF- β 2 and TGF- β 3. TGF- β 1 is present at high levels in healthy blood vessels and interestingly, higher levels of TGF- β 1 are present in human asymptomatic lesions than in lesions from symptomatic patients.¹⁰ The closely related TGF- β 2 and TGF- β 3 isoforms are either absent or present at low levels.¹¹ The classical signaling receptors for TGF- β are the TGF- β receptor I (TGF- β RI), also known as activin-like kinase (ALK), and the constitutively active type II receptor (TGF- β RII). Access of the ligands to the type I and type II receptors is regulated by soluble ligand binding proteins and by the accessory type III receptor, also known as betaglycan.^{12, 13} TGF- β signals in most cells via TGF- β type II receptor and ALK5.¹⁴⁻¹⁷

A wide range of cell types, important in atherosclerosis, expresses TGF- β .^{8, 11, 18} TGF- β exerts crucial anti-inflammatory functions by inhibiting the proliferation and differentiation of T-cells and the activation of macrophages.^{11, 18} In endothelial cells, TGF- β inhibits the expression of adhesion molecules and chemokines required for the recruitment of leukocytes.^{19, 20} Moreover, TGF- β is also known to stimulate collagen production in smooth muscle cells (SMC).^{21, 22} In 2002 we showed that treatment of ApoE^{-/-} mice with a soluble TGF- β -receptor II protein (TGF- β RII:Fc) that prevents TGF- β signaling, accelerated atherosclerosis. Plaques exhibited an unstable phenotype that contained low amounts of fibrosis, an increased amount of inflammatory cells, and even intraplaque hemorrhages.²³ Mallat *et al.* who injected neutralizing antibodies against TGF- β 1 into ApoE^{-/-} mice, obtained similar results.²⁴ Recently, Frutkin *et al.* showed that mice with cardiac over-expression of TGF- β reduced atherosclerosis formation in the aortic roots and prevented aortic dilatation.²⁵ These effects could be attributed to TGF- β signaling in T-cells. Mice, with deficient TGF- β signaling in CD4⁺ T-cells (CD4-dnTGF- β RII), also showed accelerated plaque progression, and a shift towards an unstable atherosclerotic plaque phenotype.²⁶ This was accompanied by an increased differentiation of T-cells towards both Th1 and Th2 phenotype. Increased levels of IFN- γ , IL-10 and IL-4 were found in cultured splenocytes.²⁷


Although the effects of TGF- β signaling of the adaptive immune system in atherosclerosis have been described, limited data are available on the effect of TGF- β signaling in the innate immune system. In atherosclerosis, the principal effector cells of the innate immunity are the DCs, monocytes/macrophages, neutrophils and NK cells, which all express CD11c. In the present study, we investigated the effect of TGF- β signaling in CD11c⁺ cells in atherosclerosis using ApoE^{-/-} mice

that contained a dysfunctional TGF- β Receptor II (ApoE^{-/-}CD11cDNR). We found that blockade of TGF- β signaling in CD11c accelerated atherogenesis, enhanced the influx of both CD4⁺ and CD8⁺ T-cells into the plaques and decreased fibrosis. Systemically, an increased differentiation of T-cells towards effector memory T-cells (CD44^{high}CD62L^{low}) was found together with increased levels of Th1, Th2 and Th17 cytokines. Our data illustrate the important role of TGF- β as a regulator between innate and adaptive immunity in atherosclerosis.

Materials and methods

ApoE^{-/-}CD11c-dnTGF- β RII transgenic mice

CD11c-dnTGF- β RII (CD11cDNR) mice were generated as described previously and were backcrossed to ApoE^{-/-} mice (both genotypes were on a C57BL/6 background). In short, the human TGF- β type II receptor sequence between nucleotides -7 and +573 that encodes for the extracellular and transmembrane regions of the TGF- β type II receptor was cloned into the EcoRI site of rabbit β -globin gene exon 3 of plasmid CD11c promoter vector pDOI-5. Expression of the transgene was determined by real-time PCR in freshly isolated cell types from spleen. Smad2 phosphorylation in isolated NK cells and DCs was used as a read-out for disturbed TGF- β signaling (Supplemental Figure 1).²⁸



Supplemental Figure 1: Generation and characterization of CD11c-dnTGF- β RII transgene mice.

Transgene construct with in grey the dnTGF- β RII sequence, in white the rabbit β -globin exon gene 3 and in black the plasmid CD11c promoter (A). Normalized values to HPRT of dnTGF- β RII mRNA in cells isolated from spleen as determined by real-time PCR (B). Smad2 phosphorylation in DCs and NK cells after stimulation in the presence or absence of TGF- β with LPS or IL-12 and IL-18, respectively (C).

At the age of 20 weeks, mice were euthanized. Blood was obtained from the retro-orbital plexus and plasma cholesterol was measured using a colorimetric assay (CHOD-PAP, Roche). Spleen, liver and lymph nodes were harvested after in situ perfusion using PBS. Heart and aorta were isolated after subsequent perfusion using 1% paraformaldehyde. Hearts were frozen in Tissue-Tek (Shandon, Veldhoven, The Netherlands). **Organs for autopsy were collected in 4% paraformaldehyde. All animal experiments were performed under approved Institutional Animal Care and Use Committee protocols (Yale University).**

Histology and morphometry

Plaque area was analyzed using serial sections of 6 μm with 42 μm intervals, beginning from the onset of the aortic valves until the valves had disappeared. For histological analysis of atherosclerosis, sections were stained with hematoxylin and eosin (HE). The extent of atherosclerosis was assessed in aortic roots and plaque area was calculated by using a Leica DM3000 Light microscope (Leica Microsystems) coupled to a computerized morphometry system (Leica Qwin 3.5.1).

Immunohistochemistry

Corresponding sections were immunolabeled with CD45 rat monoclonal antibody (1:5000; Pharmingen) to detect all inflammatory cells, Moma-2 rat monoclonal antibody (1:50; Serotec) to detect macrophages and α SMA monoclonal mouse antibody (1:500; DAKO) as a marker for vascular smooth muscle cells and fibroblasts. To detect T lymphocytes CD3 rabbit monoclonal antibody (1:200; DAKO) was used. CD4 and CD8 rat monoclonal antibodies (undiluted, gift from W. Buurman) were used to distinguish between respectively T-helper cells and cytotoxic T-cells. Sirius red staining was used to detect collagen content. Morphometric analyses were performed using a Leica Quantimet with Qwin3.5.1 software (Leica Microsystems). Fluorescent immunohistochemistry was used to determine the presence of CD11c⁺ cells in the aortic lesions. CD11c, CD11b, CD4 and CD8 antibodies (all BD Biosciences) were conjugated to fluorescein isothiocyanate (FITC), phycoerythrin (PE) or peridinin chlorophyll protein (PerCP). Sections were analyzed with a Leica TCS SP5 multi-photon microscope (Leica, Germany).

Flow Cytometry

Spleen and lymph nodes were harvested from donor mice and single cell suspensions were prepared and stained with anti-DX5, -TCR, -CD3, -CD4, -CD8, CD25, FoxP3, -CD44, -CD62L, -CD11c (BD Biosciences) or isotype control IgG (PharMingen). Antibodies were conjugated to fluorescein isothiocyanate (FITC), phycoerythrin (PE), allophycocyanin (APC) or peridinin chlorophyll protein (PerCP). Cells were analyzed using a FACS-Canto II (BD Biosciences) and data were analyzed with FlowJo software (Tree Star).

Cytokine production

Spleen and lymph nodes from ApoE^{-/-} and DNR mice treated (n=5) or mDCs (n=6) were isolated and gently mashed through a 70 µm nylon cell strainer. Red blood cells were lysed using 0.83% NH₄Cl in 0.01 M tris/HCL, pH 7.2. Splenocytes and lymph node cells were cultured in triplicate at 3x10⁵ cells/well in the presence of soluble αCD3/CD28 (2 µg/ml) in cRPMI. 48 Hours later, supernatants were collected and TGF-β, IFN-γ, IL-10, IL-12p70, IL-4, IL-17 were measured. Cytokine levels were determined in undiluted supernatant following the manufacturer's instructions. (All ELISA's were purchased at Ebioscience, Belgium).

Statistical analysis

Values are expressed as mean ± SEM. Data were analyzed by either a two-tailed Student's T test or a non-parametric Mann-Whitney U-test. Statistical analysis was performed using Prism software. Probability values of P<0.05 were considered significant.

Results

General

At 20 weeks of age, no significant differences were observed between ApoE^{-/-} CD11cDNR mice and ApoE^{-/-} mice with regard to body weight. Decreased levels of total plasma-cholesterol were detected in ApoE^{-/-}CD11cDNR animals compared to ApoE^{-/-} controls (ApoE^{-/-}CD11cDNR 5.97±1.25 mg/dl vs ApoE^{-/-} 10.10±0.78 mg/dl; $P < 0.05$) while serum triglycerides (TG) did not differ between both groups (ApoE^{-/-}CD11cDNR 2.33±0.24 mg/dl vs ApoE^{-/-} 2.57±0.29 mg/dl). In addition, autopsy of > 20 organs revealed no abnormalities or pathologies.

Localisation and subtype of CD11c⁺ cells in atherosclerotic plaques

To detect whether CD11c⁺ cells are present in atherosclerotic plaques, where they were localized and which plaque cell types express CD11c, aortic roots of both ApoE^{-/-}CD11cDNR and ApoE^{-/-} mice were double labelled CD11c and cell-type specific markers. Using 2-photon microscopy we were able to clearly distinguish between the different CD11c⁺ expressing cell types (Figure 1).

CD11c positive cells were commonly present in the shoulder region of the plaque and in the adventitia. Interestingly, only a small percentage of CD11c⁺ cells expressed CD11b. This indicates that the majority of CD11c⁺ cells in the plaque were dendritic cells (DC), which was confirmed by double staining with CD11c, CD4 and/or CD8, or macrophage foam cells which are also known to be CD11c⁺CD11b⁻ but not NK-cells (Figure 1).




Figure 1: Multi-photon immunohistochemistry to determine the presence of CD11c⁺ cells in the aortic lesions.

(a) CD11c-FITC and CD11b-PE. Both cell types are clearly present in both lesion and adventitia. Only scarce double positive cells are present. (b) Detail of adventitia where massive infiltration of CD11c positive cells (FITC) is observed. (c) Detail of the atherosclerotic lesion. CD11c positive cells (FITC) together with CD11b positive cells (PE) are present in the shoulder region of the plaque.

Defective TGF- β signaling in CD11c⁺ cells accelerates atherosclerosis and enhances the inflammatory state of the lesion

ApoE^{-/-} mice with disrupted TGF- β signaling in CD11c⁺ cells (ApoE^{-/-}CD11cDNR) exhibited a 2-fold increase in atherosclerotic plaque area in the aortic root (ApoE^{-/-}CD11cDNR $2.37 \times 10^5 \pm 5.33 \times 10^4 \mu\text{m}^2$ vs ApoE^{-/-} $1.11 \times 10^5 \pm 2.09 \times 10^4 \mu\text{m}^2$, $n=8/\text{group}$, $P<0.05$) (Figure 2). Moreover, the percentage of CD45⁺ leukocytes was drastically increased in plaques of ApoE^{-/-}CD11cDNR animals compared to controls (ApoE^{-/-}CD11cDNR $24.04 \pm 6.63\%$ vs ApoE^{-/-} $9.74 \pm 3.14\%$ CD45⁺ cells, $P<0.05$) (Figure 3 A-C).




Figure 2: Defective TGF- β signaling in CD11c⁺ cells accelerates atherosclerosis. Plaque area in μm^2 (A). Representative HE staining of atherosclerotic lesions of the aortic roots (B and C) (magnification 5x) (* $P<0.05$).

In order to have a more precise view on which leukocyte subsets were increased, we analyzed the lesions for the presence of different T-cell (CD3, CD4, CD8, FoxP3), and macrophage markers (Moma-2). Plaques of ApoE^{-/-}CD11cDNR mice contained significantly more CD3⁺ T-cells than the controls (ApoE^{-/-}CD11cDNR $21.71 \pm 7\%$ vs ApoE^{-/-} $5.16 \pm 1\%$ CD3⁺ cells, $P<0.05$) (Figure 3 D-F). Increased levels of CD4⁺ T-helper cells were also found in ApoE^{-/-}CD11cDNR $12.14 \pm 2.73\%$ vs ApoE^{-/-} $5.73 \pm 1.25\%$ CD4⁺ cells, $P<0.05$) (Figure 4 A-C). In addition, a substantial amount of CD8⁺ cytotoxic T-cells was present in plaques of ApoE^{-/-}CD11cDNR mice while CD8⁺ T-cells were almost absent in the control group (ApoE^{-/-}CD11cDNR $9.06 \pm 3.21\%$ vs ApoE^{-/-} $2.13 \pm 0.34\%$ CD8⁺ cells, $P<0.05$) (Figure 4 D-F).




Figure 3: Disruption of TGF- β signaling in CD11c⁺ cells enhances plaque inflammation.

Number of CD45 positive cells per total cells number in the atherosclerotic plaque (A). Representative CD45 immuno staining (B and C) (magnification 20x). Number of CD3 positive cells per total cells number in the atherosclerotic plaque (D). Representative CD3 immuno staining (E and F) (magnification 20x) (* $P < 0.05$).




Figure 4: Defective TGF- β signaling in CD11c⁺ cells attracts both CD4⁺ and CD8⁺ T-cells into the lesions.

Number of CD4 positive cells per total cells number in the atherosclerotic plaque (A). Representative CD4 immuno staining (B and C) (magnification 20x). Number of CD8 positive cells per total cells number in the atherosclerotic plaque (D). Representative CD8 immuno staining (E and F) (magnification 20x) (* $P < 0.05$).

However, no differences in FoxP3 staining could be observed between both groups (ApoE^{-/-}CD11cDNR $1.01 \times 10^{-2} \pm 0.53 \times 10^{-2}$ % vs ApoE^{-/-} $0.87 \times 10^{-2} \pm 0.37 \times 10^{-2}$ % FoxP3⁺ cells). In addition macrophage (Moma-2) content did not differ between both groups (ApoE^{-/-}CD11cDNR 19.10 ± 4.42 % vs ApoE^{-/-} 21.08 ± 2.41 % Moma-2⁺ cells).

Defective TGF- β signaling in CD11c⁺ cells inhibits plaque fibrosis

Collagen content on Sirius red-stained sections revealed significantly lower amounts of collagen in plaques of ApoE^{-/-}CD11cDNR mice than in ApoE^{-/-} mice (ApoE^{-/-}CD11cDNR 30.21 ± 2.15 % vs ApoE^{-/-} 37.96 ± 2.66 % Sirius Red + area, $P < 0.05$) (Figure 5 A-C). The low plaque collagen content was accompanied by a decrease in α -smooth muscle cell⁺ (ASMA) content (ApoE^{-/-}CD11cDNR 4.02 ± 0.71 % vs ApoE^{-/-} 9.46 ± 1.83 % ASMA⁺ area, $P < 0.05$) (Figure 5 D-F). These results demonstrate that TGF- β receptor deficiency in CD11c⁺ cells reduces plaque fibrosis, indicative of plaque destabilization.




Figure 5: Defective TGF- β signaling in CD11c⁺ cells leads to an unstable plaque phenotype.

Percentage of Sirius red positive staining (A). Representative Sirius red staining (B and C) (magnification 20x). Amount of α SMC content in the atherosclerotic plaque (D). Representative ASMA staining (E and F) (magnification 20x) (* $P < 0.05$).

Defective TGF- β signaling in CD11c⁺ cells alters T-cell homeostasis

To check whether the increased T-cell content in the ApoE^{-/-}CD11cDNR lesions was due to an altered T-cell homeostasis, we used FACS analysis to determine the expression of homing receptors CD44 and CD62L on CD4⁺ and CD8⁺ T-cells in the spleen to distinguish from naïve (CD44^{low}CD62L^{high}) or effector-memory (CD44^{high}CD62L^{low}) phenotype (Figure 6). The CD4⁺ T-cell population of the ApoE^{-/-}CD11cDNR group had a significant decrease in the marker for naïve T-cells (CD4⁺ CD44^{low}CD62L^{high}: ApoE^{-/-} 52.96±4.46 % vs ApoE^{-/-}CD11cDNR 24.39±2.71 %; $P < 0.05$) and an increase in the marker for effector memory T-cells (CD4⁺ CD44^{high}CD62L^{low} population: ApoE^{-/-} 45.85±4.40 % vs ApoE^{-/-}CD11cDNR 73.07±2.22 %; $P < 0.05$). Similar effects were seen in the CD8 population (CD8⁺ CD44^{low}CD62L^{high} population: ApoE^{-/-} 81.66±2.50 % vs ApoE^{-/-}CD11cDNR 59.50±5.98 %; $P < 0.05$; CD8⁺ CD44^{high}CD62L^{low} population: ApoE^{-/-} 17.21±2.43 % vs ApoE^{-/-}CD11cDNR 35.46±4.42 %; $P < 0.05$). CD62L surface expression is lost when cells become activated. This means that both CD4⁺ T-helper cells and CD8⁺ cytotoxic T-cells from ApoE^{-/-}CD11cDNR have characteristics of differentiated T-cells. These effector T-cells, in contrast to naïve T-cells, are potent to enter inflamed tissues.




Figure 6: Loss of TGF- β signaling in CD11c⁺ cells leads to the generation of effector T-cells in spleen.

Percentage of naïve CD4⁺ T-cells (A). Percentage of naïve CD8⁺ T-cells (B). Percentage of effector CD4⁺ T-cells (C). Percentage of effector CD8⁺ T-cells (D) (* $P < 0.05$).

In addition to T-cell differentiation, an enhanced CD4/CD8 ratio was observed in the spleen of CD11cDNR animals (ApoE^{-/-} 1.42±0.09 % vs ApoE^{-/-}CD11cDNR 1.826±0.14 % $P<0.05$) (Figure 7 A-C). Finally, a significant increase was observed in the amount of CD25⁺FoxP3⁺ double positive cells within the CD4⁺ population (Tregs) in the spleen of ApoE^{-/-}CD11cDNR animals (ApoE^{-/-} 9.16±0.65 % vs ApoE^{-/-}CD11cDNR 15.33±1.33 %; $P<0.05$) (Figure 7 D).




Figure 7: Inhibition of TGF- β signaling in CD11c⁺ cells disturbs T-cell homeostasis in spleen.

Percentage of CD3⁺CD4⁺ T-cells (A). Percentage of CD3⁺CD8⁺ T-cells (B). CD4/CD8 ratio T-cells (C). Percentage CD25⁺FoxP3⁺ of CD4⁺T-cells (D) (* $P<0.05$).

TGF- β deficiency in CD11c⁺ cells is associated with increased levels of Th1, Th2 and Th17 cytokines

To determine whether the observed T-cell phenotype correlated with enhanced cytokine levels, T-cell cytokines were determined in the spleen after re-stimulation in vitro with aCD3/CD28. An 8-fold increase of the Th1 cytokine IFN- γ was detected in the ApoE^{-/-}CD11cDNR group (ApoE^{-/-}CD11cDNR 1370.61±534.34 pg/ml vs ApoE^{-/-} 171.17±70.5 pg/ml, $P<0.05$) (Figure 8 A). Furthermore, significant increases were found in Th2 cytokines IL-4 and IL-10 (IL-4: ApoE^{-/-}CD11cDNR 161.00±29.99 pg/ml vs ApoE^{-/-} 71.00±10.53 pg/ml, $P<0.05$; IL-10: ApoE^{-/-}CD11cDNR 500.00±80.26 pg/ml vs ApoE^{-/-} 346.67±13.14 pg/ml, $P<0.05$) (Figure 8 B and C). Finally, a 5-fold increase in IL-17 levels was detected (ApoE^{-/-}CD11cDNR 101.78±22.81 pg/ml vs ApoE^{-/-} 20.93±1.73 pg/ml, $P<0.05$) (Figure 8D). In addition to cytokine levels, T-cell proliferation was measured but no significant differences could be observed in T-cell proliferation (data not shown).




Figure 8: Disruption of TGF- β signaling in CD11c⁺ cells increases cytokine expression.

Concentrations (pg/ml) of IFN- γ (A), IL-10 (B), IL-4 (C), and IL-17 (D) in supernatants of cultured splenocytes, stimulated with aCD3/CD28 (* $P < 0.05$).

Inhibition of TGF- β signaling does not affect DC maturation

Since the net outcome of T-cell immune responses is influenced by the activation stage of antigen-presenting cells,³⁰ we analyzed the maturation status of the CD11c⁺ DCs in spleen. TGF- β is known to inhibit critical co-stimulatory molecules on the surface of DCs³¹ and therefore regulate DC maturation. However, analysis of cell surface markers of dendritic cells showed that DCs from ApoE^{-/-}CD11cDNR animals had a differentiation process similar to ApoE^{-/-} DCs. No differences could be observed in MHC-II, CD86 or CD40 expression (Figure 9).




Figure 9: Disruption of TGF- β signaling in CD11c⁺ cells does not alter DC maturation *in vivo*.

FACS analysis of maturation markers CD40 (B), MHC-II (C) and CD86 (D). Percentage of CD11c positive cells in spleen (A).

Discussion

In the present study, we highlighted the importance of TGF- β signaling in cells of the innate immune system in the progression of atherosclerosis. When TGF- β signaling in CD11c⁺ cells (i.e. monocytes, dendritic cells, macrophages and NK-cells)^{28, 32} was abrogated, atherosclerotic plaque size increased significantly. Moreover, plaques contained increased amounts of CD4⁺ and CD8⁺ T-cells and a low percentage of collagen, reminiscent of a rupture prone plaque in humans.

Several studies have provided evidence for an important role for TGF- β as an immune modulating cytokine in atherosclerosis. In patients, lowered serum levels of TGF- β are detected but can be increased with tamoxifen treatment with a consequent cardioprotective effect.³³ In addition, mutations and reduced expression of the type II receptor gene have both been detected in cells of human atherosclerotic lesions.³⁴⁻³⁶ In mice, systemic inhibition of TGF- β signaling in ApoE^{-/-} mice by using a recombinant soluble TGF- β type II receptor²³ or a blocking TGF- β 1 antibody results in an unstable atherosclerotic plaque phenotype, whereas cardiac overexpression of TGF- β 1 (resulting in increased plasma levels of TGF- β) limits plaque growth and induces plaque stabilization (Table 1).²⁴ However, the immune modulating effects of TGF- β are not restricted to T-cells. Although TGF- β producing regulatory T-cells

are well known to control T-cell activation and differentiation, and to play a crucial role in atherosclerosis,³⁷ the immune-regulatory capacity of other cell-types is also controlled via TGF- β . Interestingly, it has been reported that phagocytosis of apoptotic cells leads to TGF- β secretion, which inhibits the production of inflammatory cytokines and chemokines in macrophages including IL-1 β and TNF- α .¹⁸ In addition, several studies have shown that TGF- β inhibits *in vitro* activation and maturation of DCs.^{38, 39} TGF- β inhibits upregulation of critical co-stimulatory molecules on the DC surface, thereby reducing its antigen-presenting capacity and DC mediated T-cell responses.³⁰ However these studies focused on Langerhans cell development, and might not be applicable in the present study, where we did not observe an effect on DC maturation *in vivo*. In this study we proved the importance of TGF- β signaling in CD11c positive cells in atherosclerosis using the CD11c-DNR mouse. In this mouse model, specific expression of dnTGF β RII was found in CD11c expressing cells, including DC subsets and NK cells, whereas no expression was found in NKT cells, T-cells or B-cells.²⁸

In atherosclerotic plaques, an important local and systemic role for CD11c expressing cells has been reported recently.^{29, 40, 41} In conditions of hypercholesterolemia, monocytes expressed high levels of CD11c, and show increased adherence to the arterial wall.²⁹ In CD11c^{-/-}/ApoE^{-/-} mice, monocyte recruitment was impaired and only small atherosclerotic plaques developed.²⁹ This was also observed in the atherosclerotic plaque, where CD11c⁺ cells with foam cell morphology also were positive for Moma-2, a macrophage marker. Cho *et al.* studied gene expression profiles during foam cell formation and confirmed that foam cells develop a dendritic cell-like phenotype accompanied by CD11c expression.³² Moreover, CD11c plays a crucial, but yet unidentified role in cholesterol homeostasis.⁴¹ When there is an expansion of CD11c⁺ cells, cholesterol levels and the amount of atherosclerosis decreases, while a depletion of CD11c⁺ cells results in hypercholesterolemia and an increase in atherosclerosis.⁴¹ In the present study we observed decreased levels of cholesterol in the plasma of ApoE^{-/-}CD11cDNR mice. Recently it has been shown that depletion of DCs resulted in enhanced cholesterolemia suggesting a close relationship between the DC population and plasma cholesterol levels.⁴¹ Moreover, Frutkin *et al.* described that hypercholesterolemia induces TGF- β expression.²⁵ Since in our study we blocked TGF- β signalling in DCs, our data may suggest an interaction between TGF- β , DCs and the cholesterol metabolism. However, the reason why the reduced cholesterol levels did not result in reduced lesion size remains unclear.

In the present study, we found that TGF- β signaling in CD11c⁺ cells is involved in the recruitment of inflammatory cells into the plaque, as well as in maintaining T-cell homeostasis. Lack of TGF- β signaling in CD11c⁺ cells results in massive infiltration of CD4⁺ and CD8⁺ cells in atherosclerotic plaques and in a systemic expansion of CD44^{high}CD62L^{low} T-cells. These data demonstrated that blockage of TGF- β signaling in CD11c cells lead to a shift from naïve T-cells towards effector (CD44^{high}CD62L^{low}) T-cells. This effector-T-cell phenotype explained the influx of large amounts of CD4⁺ and CD8⁺ T-cells into the lesions, since naïve T-cells are not capable of infiltrating inflamed tissues. In addition, elevated levels of IFN- γ , TNF- α , IL-4, IL-10, IL-17 were detected. This broad array of expressed cytokines refers to both Th1 and Th2 subsets. The effect on atherosclerosis can therefore

not be explained by a shift in the Th1/Th2 balance. In addition, high levels of IL-17 were detected in the CD11cDNR animals, referring to an enhanced Th17 phenotype as well. These results are in concordance with Laouar *et al.* who studied the CD11cDNR mouse in experimental autoimmune encephalomyelitis (EAE) where an increase in inflammation in the CNS is accompanied with increased levels of Th1 and Th17 cytokines.⁴² In addition abrogation of IL-17 receptor signaling in LDLr^{-/-} mice attenuated atherosclerosis.⁴³ Moreover, Th17 cells have been identified in patients with acute coronary syndrome.⁴⁴ Taken together these data suggest a potential proatherogenic role. We therefore hypothesize that interruption of TGF- β signaling in CD11c⁺ cells results in altered T-cell priming by DCs.

In conclusion, we state that TGF- β signaling in CD11c expressing cells is fundamental in the immunomodulation during atherogenesis. We showed that impaired TGF- β signalling in CD11c⁺ cells resulted in an increase of activated effector T-cell which can more easily migrate into the atherosclerotic lesions with a consequential development of plaques, which are high in inflammation and low in fibrosis. We therefore provided a first inhibitory link between the innate and adaptive immune response with an important anti-inflammatory function for TGF- β signaling in CD11c⁺ cells.

Table 1: Overview of studies on the effects of TGF- β signaling in atherosclerosis, with focus on plaque area, plaque phenotype and the induced cytokines.

Group (Year)	Reference	Experiment	Plaque area	Plaque inflammation	Plaque fibrosis	Induced cytokines
Mallat (2001)	24	Abs against TGF- β 1, TGF- β 2 and TGF- β 3 ApoE ^{-/-} acceptor mice	+	+	-	
Lutgens (2002)	23	rs TGF- β RII ApoE ^{-/-} acceptor mice	+	+	-	
Robertson (2003)	26	CD4-dnTGF- β RII tg+ ApoE ^{-/-} background	+	+	-	increased IFN- γ , TNF- α IL-4,5, 10
Frutkin (2009)	25	Cardiac specific TGF- β overexpression ApoE ^{-/-} background	-	-	+	decreased IFN- γ TNF- α , MIP-1 α

(Ref: Reference; BMT: Bone Marrow Transplantation; +: increase; - : decrease)

References

1. Galkina E, Ley K. Immune and inflammatory mechanisms of atherosclerosis. *Annu Rev Immunol.* 2009;27:165-197.
2. Lusis AJ. Atherosclerosis. *Nature.* Sep 14 2000;407(6801):233-241.
3. Weber C, Zernecke A, Libby P. The multifaceted contributions of leukocyte subsets to atherosclerosis: lessons from mouse models. *Nat Rev Immunol.* Oct 2008;8(10):802-815.
4. Charo IF, Ransohoff RM. The many roles of chemokines and chemokine receptors in inflammation. *N Engl J Med.* Feb 9 2006;354(6):610-621.
5. Kleemann R, Zadelaar S, Kooistra T. Cytokines and atherosclerosis: a comprehensive review of studies in mice. *Cardiovasc Res.* Aug 1 2008;79(3):360-376.
6. Tedgui A, Mallat Z. Cytokines in atherosclerosis: pathogenic and regulatory pathways. *Physiol Rev.* Apr 2006;86(2):515-581.
7. Zernecke A, Shagdarsuren E, Weber C. Chemokines in atherosclerosis: an update. *Arterioscler Thromb Vasc Biol.* Nov 2008;28(11):1897-1908.
8. Grainger DJ. TGF-beta and atherosclerosis in man. *Cardiovasc Res.* May 1 2007;74(2):213-222.
9. Mallat Z, Tedgui A. The role of transforming growth factor beta in atherosclerosis: novel insights and future perspectives. *Curr Opin Lipidol.* Oct 2002;13(5):523-529.
10. Cipollone F, Fazia M, Mincione G, et al. Increased expression of transforming growth factor-beta1 as a stabilizing factor in human atherosclerotic plaques. *Stroke.* Oct 2004;35(10):2253-2257.
11. Grainger DJ. Transforming growth factor beta and atherosclerosis: so far, so good for the protective cytokine hypothesis. *Arterioscler Thromb Vasc Biol.* Mar 2004;24(3):399-404.
12. Brown CB, Boyer AS, Runyan RB, et al. Requirement of type III TGF-beta receptor for endocardial cell transformation in the heart. *Science.* Mar 26 1999;283(5410):2080-2082.
13. Massague J. TGF-beta signal transduction. *Annu Rev Biochem.* 1998;67:753-791.
14. Attisano L, Wrana JL. Smads as transcriptional co-modulators. *Curr Opin Cell Biol.* Apr 2000;12(2):235-243.
15. Massague J, Gomis RR. The logic of TGFbeta signaling. *FEBS Lett.* May 22 2006;580(12):2811-2820.
16. Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell.* Jun 13 2003;113(6):685-700.
17. Wahl SM, Wen J, Moutsopoulos N. TGF-beta: a mobile purveyor of immune privilege. *Immunol Rev.* Oct 2006;213:213-227.
18. Li MO, Wan YY, Sanjabi S, et al. Transforming growth factor-beta regulation of immune responses. *Annu Rev Immunol.* 2006;24:99-146.
19. Park SK, Yang WS, Lee SK, et al. TGF-beta(1) down-regulates inflammatory cytokine-induced VCAM-1 expression in cultured human glomerular endothelial cells. *Nephrol Dial Transplant.* May 2000;15(5):596-604.
20. Smith WB, Noack L, Khew-Goodall Y, et al. Transforming growth factor-beta 1 inhibits the production of IL-8 and the transmigration of neutrophils through activated endothelium. *J Immunol.* Jul 1 1996;157(1):360-368.
21. Amento EP, Ehsani N, Palmer H, et al. Cytokines and growth factors positively and negatively regulate interstitial collagen gene expression in human vascular smooth muscle cells. *Arterioscler Thromb.* Sep-Oct 1991;11(5):1223-1230.
22. Ignatz RA, Endo T, Massague J. Regulation of fibronectin and type I collagen mRNA levels by transforming growth factor-beta. *J Biol Chem.* May 15 1987;262(14):6443-6446.
23. Lutgens E, Gijbels M, Smook M, et al. Transforming growth factor-beta mediates balance between inflammation and fibrosis during plaque progression. *Arterioscler Thromb Vasc Biol.* Jun 1 2002;22(6):975-982.

- 24.** Mallat Z, Gojova A, Marchiol-Fournigault C, et al. Inhibition of transforming growth factor-beta signaling accelerates atherosclerosis and induces an unstable plaque phenotype in mice. *Circ Res*. Nov 9 2001;89(10):930-934.
- 25.** Frutkin AD, Otsuka G, Stempien-Otero A, et al. TGF-[beta]1 limits plaque growth, stabilizes plaque structure, and prevents aortic dilation in apolipoprotein E-null mice. *Arterioscler Thromb Vasc Biol*. Sep 2009;29(9):1251-1257.
- 26.** Robertson AK, Rudling M, Zhou X, et al. Disruption of TGF-beta signaling in T cells accelerates atherosclerosis. *J Clin Invest*. Nov 2003;112(9):1342-1350.
- 27.** Ovchinnikova O, Robertson AK, Wagsater D, et al. T-cell activation leads to reduced collagen maturation in atherosclerotic plaques of Apoe(-/-) mice. *Am J Pathol*. Feb 2009;174(2):693-700.
- 28.** Laouar Y, Sutterwala FS, Gorelik L, et al. Transforming growth factor-beta controls T helper type 1 cell development through regulation of natural killer cell interferon-gamma. *Nat Immunol*. Jun 2005;6(6):600-607.
- 29.** Wu H, Gower RM, Wang H, et al. Functional role of CD11c⁺ monocytes in atherogenesis associated with hypercholesterolemia. *Circulation*. May 26 2009;119(20):2708-2717.
- 30.** Strobl H, Knapp W. TGF-beta1 regulation of dendritic cells. *Microbes Infect*. Dec 1999;1(15):1283-1290.
- 31.** Strobl H, Bello-Fernandez C, Riedl E, et al. flt3 ligand in cooperation with transforming growth factor-beta1 potentiates in vitro development of Langerhans-type dendritic cells and allows single-cell dendritic cell cluster formation under serum-free conditions. *Blood*. Aug 15 1997;90(4):1425-1434.
- 32.** Cho HJ, Shashkin P, Gleissner CA, et al. Induction of dendritic cell-like phenotype in macrophages during foam cell formation. *Physiol Genomics*. Apr 24 2007;29(2):149-160.
- 33.** McDonald CC, Alexander FE, Whyte BW, et al. Cardiac and vascular morbidity in women receiving adjuvant tamoxifen for breast cancer in a randomised trial. The Scottish Cancer Trials Breast Group. *Bmj*. Oct 14 1995;311(7011):977-980.
- 34.** McCaffrey TA, Consigli S, Du B, et al. Decreased type II/type I TGF-beta receptor ratio in cells derived from human atherosclerotic lesions. Conversion from an antiproliferative to profibrotic response to TGF-beta1. *J Clin Invest*. Dec 1995;96(6):2667-2675.
- 35.** McCaffrey TA, Du B, Consigli S, et al. Genomic instability in the type II TGF-beta1 receptor gene in atherosclerotic and restenotic vascular cells. *J Clin Invest*. Nov 1 1997;100(9):2182-2188.
- 36.** McCaffrey TA, Du B, Fu C, et al. The expression of TGF-beta receptors in human atherosclerosis: evidence for acquired resistance to apoptosis due to receptor imbalance. *J Mol Cell Cardiol*. Sep 1999;31(9):1627-1642.
- 37.** Ait-Oufella H, Salomon BL, Potteaux S, et al. Natural regulatory T cells control the development of atherosclerosis in mice. *Nat Med*. Feb 2006;12(2):178-180.
- 38.** Geissmann F, Prost C, Monnet JP, et al. Transforming growth factor beta1, in the presence of granulocyte/macrophage colony-stimulating factor and interleukin 4, induces differentiation of human peripheral blood monocytes into dendritic Langerhans cells. *J Exp Med*. Mar 16 1998;187(6):961-966.
- 39.** Strobl H, Riedl E, Scheinecker C, et al. TGF-beta 1 promotes in vitro development of dendritic cells from CD34⁺ hemopoietic progenitors. *J Immunol*. Aug 15 1996;157(4):1499-1507.
- 40.** Choi JH, Do Y, Cheong C, et al. Identification of antigen-presenting dendritic cells in mouse aorta and cardiac valves. *J Exp Med*. Mar 16 2009;206(3):497-505.
- 41.** Gautier EL, Huby T, Saint-Charles F, et al. Conventional dendritic cells at the crossroads between immunity and cholesterol homeostasis in atherosclerosis. *Circulation*. May 5 2009;119(17):2367-2375.
- 42.** Laouar Y, Town T, Jeng D, et al. TGF-beta signaling in dendritic cells is a prerequisite for the control of autoimmune encephalomyelitis. *Proc Natl Acad Sci U S A*. Aug 5 2008;105(31):10865-10870.
- 43.** van Es T, van Puijvelde GH, Ramos OH, et al. Attenuated atherosclerosis upon IL-17R signaling disruption in LDLr deficient mice. *Biochem Biophys Res Commun*. Oct 16 2009;388(2):261-265.
- 44.** Cheng X, Yu X, Ding YJ, et al. The Th17/Treg imbalance in patients with acute coronary syndrome. *Clin Immunol*. Apr 2008;127(1):89-97.

¹ Division of Biopharmaceutics, LACDR, Leiden, The Netherlands

² Dept. of Rheumatology, LUMC, Leiden, The Netherlands