



Universiteit  
Leiden

The Netherlands

## Acute myocardial infarction treatment : from prehospital care to secondary prevention

Atary, J.Z.

### Citation

Atary, J. Z. (2011, September 22). *Acute myocardial infarction treatment : from prehospital care to secondary prevention*. Retrieved from <https://hdl.handle.net/1887/17856>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17856>

**Note:** To cite this publication please use the final published version (if applicable).



## Chapter 6

# Distribution of culprit lesions in patients with ST-segment elevation acute myocardial infarction treated with primary percutaneous coronary intervention.

M. Louisa Antoni, MD, Jael Z. Atary, MD, Kai-Hang Yiu, MD,  
Victoria Delgado, MD, PhD, Eduard R. Holman, MD, PhD,  
Ernst E. van der Wall, MD, PhD, Joanne D. Schuijf, PhD, Martin J. Schalij, MD,  
PhD, Jeroen J. Bax, MD, PhD

From the Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands.

## ABSTRACT

### Aims

Data regarding the distribution of vulnerable lesions in the coronary arteries are scarce. The aim was to evaluate the frequency and distribution of culprit lesions in patients with ST-segment elevation acute myocardial infarction (AMI). In addition, the location of culprit lesions was related to infarct size.

### Methods and results

Consecutive patients (N=1533, mean age  $61 \pm 12$  years) were evaluated. All patients were treated with primary percutaneous coronary intervention and underwent 2-dimensional echocardiography <48 hours of admission. The majority of the culprit lesions were located in the left anterior descending coronary artery (LAD,45%), followed by the right coronary artery (RCA,38%) and left circumflex coronary artery (LCX,14%). Subanalysis demonstrated that patients with a culprit lesion in the LAD and LCX had significantly higher peak cardiac enzymes compared to patients with culprit lesions in the RCA. In addition, patients with proximal LAD and LCX lesions had significantly worse left ventricular function compared to patients with mid or distal lesions.

### Conclusion

Plaque rupture resulting in AMI is more likely to occur in the proximal parts of the LAD and RCA. In addition, the location of culprit lesions was related to infarct size. Therefore, knowledge of the distribution of vulnerable lesions is important for identifying patients at risk for acute coronary events.

## INTRODUCTION

Acute coronary syndromes are primarily due to rupture of an atheromatous plaque with superimposed thrombosis. Therefore, identification of vulnerable lesions which are prone to rupture is important and has been studied extensively. Previous studies have particularly focused on characteristic histomorphologic features of vulnerable lesions.<sup>1,2</sup> Besides in the setting of randomized trials, no data have been reported regarding the distribution of culprit lesions among the different coronary arteries in patients presenting with a ST-segment elevation acute myocardial infarction (AMI).<sup>3-5</sup> Furthermore, data about the distribution of culprit lesions within the different segments of the coronary arteries in patients presenting with AMI are scarce.

Accordingly, the aim of the current study was to evaluate the frequency and distribution of culprit lesions within the 3 coronary arteries and within the different segments of the coronary arteries in a large population of patients presenting with ST-segment elevation AMI. In addition, the location of the culprit lesions was related to infarct size as assessed with peak cardiac enzymes and residual left ventricular (LV) systolic function.

## METHODS

Since February 2004, all patients admitted with ST-segment elevation AMI were identified and included in an ongoing registry (MISSION!).<sup>6</sup> The diagnosis ST-segment elevation AMI was defined based on criteria of typical chest pain, elevated cardiac enzyme levels, and typical changes on the electrocardiogram.<sup>7</sup> All patients underwent immediate coronary angiography to identify the location of the culprit lesion followed by primary percutaneous coronary intervention (PCI). Patient data were prospectively collected in the departmental Cardiology Information System (EPD-Vision®, Leiden University Medical Center, Leiden, the Netherlands) and analyzed retrospectively.<sup>8,9</sup> Standardized angiographic projections were chosen for the visual classification of the coronary artery map into segments according to the guidelines of the American College of Cardiology/American Heart Association.<sup>10,11</sup> The infarct-related vessel was determined on the coronary artery territory subtended by the regions of acute electrocardiographic changes. If the culprit vessel had more than 2 lesions, the most severe proximal stenosis or a stenosis identified with thrombus was considered as the culprit lesion. Patients were not included if no clear culprit lesion could be identified on coronary angiography. Patients were treated according to the institutional AMI protocol, which includes 2-dimensional echocardiography performed within 48 hours of admission to assess residual LV function using LV ejection fraction calculated by the biplane Simpson's technique from the apical 2- and 4-chamber views.<sup>6,12</sup>

Continuous data are presented as mean  $\pm$  standard deviation and categorical data are presented as frequencies and percentages. Differences in baseline characteristics between the 3 coronary arteries or the different segments of the coronary arteries were evaluated with 1-way analysis of variance or chi-square test, as appropriate. Of note, patients with a bypass graft identified as culprit vessel were not included in these analyses. Post-hoc comparisons were performed using the Bonferroni adjustments for multiple comparisons. For all tests, a p value  $<0.05$  was considered statistically significant.

## RESULTS

A total of 1533 consecutive patients were evaluated. Mean age of the patient population was  $61.3 \pm 12.2$  years and mean LV ejection fraction was  $45.8 \pm 8.6\%$  (Table 1). Before primary PCI, mean TIMI flow of all patients was  $0.6 \pm 1.0$ . Among the 1533 culprit lesions studied, the majority of the patients showed a 100% stenosis (1013 patients, 66%) or 99% stenosis (279 patients, 18%). Only 6 (0.4%) patients showed a culprit lesion with 50% stenosis, 78 (5%) patients with 75% stenosis and 157 (10%) patients with 90% stenosis as determined by semiquantitative grading.

Table 2 shows the distribution of the culprit lesions. The majority of the culprit lesions were located in the left anterior descending coronary artery (LAD, 668 patients, 45%) and

**Table 1.** Baseline characteristics of the patient population

	All patients (N = 1533)
Clinical characteristics	
Age (years)	$61.3 \pm 12.2$
Male gender	1158 (76%)
Medical History	
Current smoking	715 (47%)
Diabetes	182 (12%)
Family history of coronary artery disease	618 (40%)
Hyperlipidemia	299 (20%)
Hypertension	533 (35%)
Prior myocardial infarction	140 (9%)
Infarct size	
Peak creatine phosphokinase level (U/l)	$2229 \pm 2890$
Peak cardiac troponin T level ( $\mu\text{g/l}$ )	$5.9 \pm 6.5$
TIMI flow	$2.9 \pm 0.4$
Multivessel disease	718 (54%)
Left ventricular ejection fraction (%)	$45.8 \pm 8.6$

Hyperlipidemia= Total cholesterol  $\geq 190$  mg/dl or previous pharmacological treatment.

Hypertension = Blood pressure  $\geq 140/90$  mm Hg or previous pharmacological treatment.

**Table 2.** Distribution of coronary occlusions

Vessel segment	Number of occlusions	LV ejection fraction (%)
Left main (segment 11)	16 (1%)	37.1 ± 8.1
Ramus intermedius (segment 28)	15 (1%)	47.9 ± 7.5
RCA	588 (38%)	46.8 ± 8.2
Proximal (segment 1)	249 (42%)	46.2 ± 8.4
Mid (segment 2)	218 (37%)	47.2 ± 7.7
Distal (segment 3)	96 (16%)	47.2 ± 8.0
Posterior descending (segment 4)	14 (2%)	48.7 ± 11.2
Posterior lateral (segment 6)	11 (2%)	43.8 ± 9.3
LAD	688 (45%)	44.6 ± 8.9
Proximal (segment 12)	401 (58%)	43.7 ± 8.8
Mid (segment 13)	243 (35%)	45.5 ± 8.9
Distal (segment 14)	20 (3%)	45.6 ± 11.2
Diagonal 1 branch (segment 15)	23 (3%)	50.5 ± 7.3
Diagonal 2 branches (segment 16)	1 (0.1%)	48.0 ± 0
LCX	214 (14%)	47.4 ± 7.8
Proximal (segment 18)	82 (38%)	45.1 ± 8.7
Mid (segment 19)	81 (38%)	48.8 ± 6.9
Distal segment 19a)	3 (1%)	47.7 ± 2.9
Obtuse marginal 1 (segment 20)	41 (19%)	48.9 ± 7.0
Obtuse marginal 2 (segment 21)	7 (3%)	49.4 ± 7.3
Bypass graft	12 (0.8%)	44.3 ± 8.6

LAD: left anterior descending coronary artery; LCX: left circumflex artery; RCA: right coronary artery and LV: left ventricular.

the right coronary artery (RCA, 588 patients, 38%) and only a small number of culprit lesions were located in the left circumflex coronary artery (LCX, 214 patients, 14%). Culprit lesions were not uniformly distributed, but tended to be clustered in the proximal or mid vessel segments. In addition, a relatively high percentage was located in the first obtuse marginal branch (41 patients, 19% of all LCX lesions). No differences in baseline characteristics were observed between lesion localization within the coronary arteries except for prior myocardial infarction. Patients with prior myocardial infarction more often had the culprit lesion located in the distal part of the coronary artery compared to the mid and proximal parts (17% vs. 7% and 8%, respectively,  $p = 0.002$ ).

Infarct size was assessed with peak cardiac enzymes (peak creatine phosphokinase (CPK) level and peak cardiac troponin T (cTnT) level) and residual LV function. Both peak CPK level and peak cTnT level were significantly lower in patients with a RCA culprit lesion compared to patients with a LAD culprit lesion ( $1634 \pm 3424$  U/l vs.  $2674 \pm 2565$  U/l,  $p < 0.001$  and  $4.0 \pm 4.2$  µg/l vs.  $7.2 \pm 7.4$  µg/l,  $p < 0.001$ , respectively) or a LCX culprit lesion ( $1634 \pm 3424$  U/l vs.  $2282 \pm 1830$  U/l,  $p = 0.02$  and  $4.0 \pm 4.2$  µg/l vs.  $6.0 \pm 5.3$  µg/l,  $p < 0.001$ , respectively).

No significant differences in peak cardiac enzymes were observed between patients with a LAD and LCX infarction.

In addition, the level of peak cardiac enzymes was evaluated for the proximal, mid and distal segments of the coronary arteries. The RCA and LCX demonstrated no significant differences for the proximal, mid or distal culprit lesions (RCA:  $p = 0.71$  and  $p = 0.37$  and LCX:  $p = 0.11$  and  $p = 0.07$  for peak CPK level and peak cTnT level, respectively). However, LAD proximal culprit lesions resulted in significant higher peak cardiac enzymes compared to lesions in the mid part ( $3192 \pm 2886$  U/l vs.  $2092 \pm 1885$  U/l,  $p < 0.001$  and  $8.4 \pm 8.0$   $\mu\text{g/l}$  vs.  $5.8 \pm 6.5$   $\mu\text{g/l}$ ,  $p < 0.001$ ) or the distal part of the LAD ( $3192 \pm 2886$  U/l vs.  $1261 \pm 1219$  U/l,  $p = 0.005$  and  $8.4 \pm 8.0$   $\mu\text{g/l}$  vs.  $4.0 \pm 4.2$   $\mu\text{g/l}$ ,  $p = 0.04$ ).

Residual LV function assessed with LV ejection fraction differed significantly between patients with different culprit vessels (ANOVA  $p < 0.001$ ) (Table 1). Post-hoc analysis demonstrated that patients with the LAD as culprit vessel had significantly lower LV ejection fraction as compared to patients with the RCA or LCX as culprit vessel ( $44.6 \pm 8.9\%$  vs.  $46.8 \pm 8.2\%$ ,  $p < 0.001$  and  $44.6 \pm 8.9\%$  vs.  $47.4 \pm 7.8\%$ ,  $p = 0.001$ , respectively). Further subanalysis of the different segments per culprit vessel revealed that patients with proximal culprit lesions in the LAD and LCX had significantly worse LV function compared to patients with mid and distal lesions ( $43.7 \pm 8.8\%$  vs.  $45.5 \pm 8.9\%$  and  $45.6 \pm 11.2\%$ ,  $p = 0.04$  for proximal, mid and distal LAD lesions and  $45.1 \pm 8.7\%$  vs.  $48.8 \pm 6.9\%$  and  $47.7 \pm 2.9\%$ ,  $p = 0.02$  for proximal, mid and distal LCX lesions). However, no differences in LV function were observed for the different locations of culprit lesions in the RCA ( $p = 0.37$ ).

## DISCUSSION

Patients with ST-segment elevation AMI treated with primary PCI were more likely to have a LAD or RCA culprit lesion than a LCX culprit lesion. However, infarct size assessed with peak cardiac enzymes demonstrated no significant differences between LAD and LCX infarctions, whereas RCA infarctions were significantly smaller. In addition, patients with proximal lesions in the LAD or LCX demonstrated worse LV function as compared to patients with lesions in the mid and distal parts, whereas no significant difference was observed between patients with proximal, mid or distal occlusions of the RCA.

The results of the present study provide further evidence for what has been described in smaller populations.<sup>11</sup> Wang et al. determined the location of coronary lesions in 208 consecutive patients with ST-segment elevation AMI.<sup>11,13</sup> The authors showed that culprit lesions tended to cluster within the proximal third of the coronary vessels. However, the distance from the ostium to the lesion depended upon which coronary artery was involved. Gibson et al. described that median distances from the ostium to the culprit lesion differed according to the coronary artery and the distance was the smallest in the LAD, followed by

the LCX and the RCA.<sup>13</sup> Interestingly, the same phenomenon has been observed in a large population of 30,386 patients with non-ST elevation AMI undergoing PCI described by Dixon et al, and thus, the current findings may be generalized for all culprit lesions including those of patients with unstable angina.<sup>14</sup>

Although information about the distribution of culprit lesions is important, understanding why plaque ruptures are less likely to occur in the LCX and why proximal occlusions are more prone to rupture remains challenging. To some extent, these observations may be explained by the fact that ischemic events of the LCX artery are underdiagnosed due to limited sensitivity of the 12-lead ECG for detection of ischemia on the lateral and posterior walls.<sup>15,16</sup> Previous studies have reported that only 33% of patients with a LCX occlusion present with ST-segment elevation on the ECG.<sup>17</sup> Recently, From et al. confirmed this hypothesis by demonstrating that in a group of 1500 patients with ST-segment elevation and non-ST-segment elevation AMI, patients with a LCX occlusion were less likely to present with ST-segment elevation on ECG and were referred less frequently for primary PCI.<sup>15</sup> However, among the group of patients presenting with non-ST-segment elevation AMI, patients with a LCX occlusion had the highest peak enzymes. Another explanation for the lower frequency of culprit lesions observed in the LCX may be the greater variation in anatomy as compared to the LAD and RCA. In a large proportion of the population, the LCX is relatively small with few pronounced side-branches, which may also explain why patients with an occlusion of the LCX may be more likely to present with a non-ST-segment elevation AMI or without any changes on the ECG.<sup>15,17</sup> Moreover, since vessel diameter plays an important role in the development of atherosclerosis, it is conceivable that as a consequence also plaque rupture may differ among the coronary arteries. The anatomy of the LCX may result in lower wall shear stress, whereas proximal segments of the coronary arteries conversely may be areas of high shear stress which determine the risk of plaque rupture.

In conclusion, the present study demonstrates that plaque rupture resulting in ST-segment elevation AMI is more likely to occur in the proximal parts of the LAD and RCA. In addition, the location of the culprit lesions in the different coronary arteries was related to infarct size. Therefore, knowledge of the distribution of vulnerable lesions is important for the identification of patients at risk for acute coronary events.



## REFERENCES

1. Kim SH, Hong MK, Park DW, Lee SW, Kim YH, Lee CW, Kim JJ, Park SW, Park SJ. Impact of plaque characteristics analyzed by intravascular ultrasound on long-term clinical outcomes. *Am J Cardiol*. 2009;103:1221-1226.
2. Alsheikh-Ali AA, Kitsios GD, Balk EM, Lau J, Ip S. The vulnerable atherosclerotic plaque: scope of the literature. *Ann Intern Med*. 2010;153:387-395.
3. Spaulding C, Henry P, Teiger E, Beatt K, Bramucci E, Carrie D, Slama MS, Merkely B, Erglis A, Margheri M, Varenne O, Cebrian A, Stoll HP, Snead DB, Bode C. Sirolimus-eluting versus uncoated stents in acute myocardial infarction. *N Engl J Med*. 2006;355:1093-1104.
4. Mauri L, Silbaugh TS, Garg P, Wolf RE, Zelevinsky K, Lovett A, Varma MR, Zhou Z, Normand SL. Drug-eluting or bare-metal stents for acute myocardial infarction. *N Engl J Med*. 2008;359:1330-1342.
5. Stone GW, Lansky AJ, Pocock SJ, Gersh BJ, Dangas G, Wong SC, Witzenbichler B, Guagliumi G, Peruga JZ, Brodie BR, Dudek D, Mockel M, Ochala A, Kellock A, Parise H, Mehran R. Paclitaxel-eluting stents versus bare-metal stents in acute myocardial infarction. *N Engl J Med*. 2009;360:1946-1959.
6. Liem SS, van der Hoeven BL, Oemrawsingh PV, Bax JJ, van der Bom JG, Bosch J, Viergever EP, van RC, Padmos I, Sedney MI, van Exel HJ, Verwey HF, Atsma DE, van der Velde ET, Jukema JW, van der Wall EE, Schalij MJ. MISSION!: optimization of acute and chronic care for patients with acute myocardial infarction. *Am Heart J*. 2007;153:14.e1-11.
7. Myocardial infarction redefined--a consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *Eur Heart J*. 2000;21:1502-1513.
8. Atary JZ, de VM, van den DR, Bosch J, Liem SS, Antoni ML, Bootsma M, Viergever EP, Kirchhof CJ, Padmos I, Sedney MI, van Exel HJ, Verwey HF, Atsma DE, van der Wal EE, Jukema JW, Schalij MJ. Standardised pre-hospital care of acute myocardial infarction patients: MISSION! guidelines applied in practice. *Neth Heart J*. 2010;18:408-415.
9. Borleffs CJ, van Rees JB, van Welsesens GH, van d, V, van EL, Bax JJ, Schalij MJ. Prognostic importance of atrial fibrillation in implantable cardioverter-defibrillator patients. *J Am Coll Cardiol*. 2010;55:879-885.
10. Scanlon PJ, Faxon DP, Audet AM, Carabello B, Dehmer GJ, Eagle KA, Legako RD, Leon DF, Murray JA, Nissen SE, Pepine CJ, Watson RM, Ritchie JL, Gibbons RJ, Chaitlin MD, Gardner TJ, Garson A, Jr., Russell RO, Jr., Ryan TJ, Smith SC, Jr. ACC/AHA guidelines for coronary angiography. A report of the American College of Cardiology/American Heart Association Task Force on practice guidelines (Committee on Coronary Angiography). Developed in collaboration with the Society for Cardiac Angiography and Interventions. *J Am Coll Cardiol*. 1999;33:1756-1824.
11. Wang JC, Normand SL, Mauri L, Kuntz RE. Coronary artery spatial distribution of acute myocardial infarction occlusions. *Circulation*. 2004;110:278-284.
12. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr*. 2005;18:1440-1463.

13. Gibson CM, Kirtane AJ, Murphy SA, Karha J, Cannon CP, Giugliano RP, Roe MT, Harrington RA, Ohman EM, Antman EM. Distance from the coronary ostium to the culprit lesion in acute ST-elevation myocardial infarction and its implications regarding the potential prevention of proximal plaque rupture. *J Thromb Thrombolysis*. 2003;15:189-196.
14. Dixon WC, Wang TY, Dai D, Shunk KA, Peterson ED, Roe MT. Anatomic distribution of the culprit lesion in patients with non-ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention: findings from the National Cardiovascular Data Registry. *J Am Coll Cardiol*. 2008;52:1347-1348.
15. From AM, Best PJ, Lennon RJ, Rihal CS, Prasad A. Acute myocardial infarction due to left circumflex artery occlusion and significance of ST-segment elevation. *Am J Cardiol*. 2010;106:1081-1085.
16. Schmitt C, Lehmann G, Schmieder S, Karch M, Neumann FJ, Schomig A. Diagnosis of acute myocardial infarction in angiographically documented occluded infarct vessel: limitations of ST-segment elevation in standard and extended ECG leads. *Chest*. 2001;120:1540-1546.
17. Shah A, Wagner GS, Green CL, Crater SW, Sawchak ST, Wildermann NM, Mark DB, Waugh RA, Krucoff MW. Electrocardiographic differentiation of the ST-segment depression of acute myocardial injury due to the left circumflex artery occlusion from that of myocardial ischemia of nonocclusive etiologies. *Am J Cardiol*. 1997;80:512-513.

