

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/32006> holds various files of this Leiden University dissertation

Author: Paltansing, Sunita

Title: Antimicrobial resistance in Enterobacteriaceae : characterization and detection

Issue Date: 2015-02-18

CHAPTER 5

Extended-spectrum β -lactamase producing *Enterobacteriaceae* among travelers from the Netherlands

S.Paltansing,¹ J.A. Vlot,² M.E.M. Kraakman,¹ R. Mesman,¹ M.L. Bruijning,¹
A.T. Bernards,¹ L.G. Visser,^{2*} K.E. Veldkamp^{1*}

¹ Department of Medical Microbiology and ² Department of Infectious Diseases,
Leiden University Medical Center, The Netherlands

* These authors contributed equally to this article

Emerging Infectious Diseases 2013 Aug;19(8):1206-1213

5.1 Abstract

Objectives: A prospective cohort study was performed among travelers from the Netherlands to investigate the acquisition of carbapenemase-producing *Enterobacteriaceae* (CP-E) and extended-spectrum β -lactamase producing *Enterobacteriaceae* (ESBL-E) and associated risk factors.

Methods: Questionnaires were administered and rectal swabs were collected and tested before and after return. ESBLs were characterized using a microarray (Checkpoints). Multilocus sequence typing (MLST) was performed on *Escherichia coli* isolates.

Results: Of 370 travelers, 32 (8.6%) were colonized with ESBL-E before travel, 113 (30.5%) acquired an ESBL-E during travel and 26 were still colonized six months after return. No CPE were found. Independent risk factors for ESBL-E acquisition were travel to South and East Asia. MLST showed extensive genetic diversity among *E. coli*. Predominant ESBLs were CTX-M enzymes.

Conclusion: The acquisition rate of ESBL-E in travelers from the Netherlands was high (30.5%). Active surveillance for ESBL-E and CP-E and contact isolation precautions may be recommended at admission to medical facilities for patients, who traveled to Asia during the previous six months.

5.2 Introduction

The effect of international travel on the spread of multidrug resistant *Enterobacteriaceae* (MDR-E) has become more evident in the past few years. Data obtained in prospective studies among returning travelers from Australia, Canada, Sweden and New York during 2007-2010 revealed high carriage rates of extended-spectrum β -lactamase producing *Enterobacteriaceae* (ESBL-E) varying from 18-25% after foreign travel.¹⁻⁴ Two of these studies also reported a pre-travel ESBL-E carriage rate of 7.8%.

The identification of carbapenemase-producing *Enterobacteriaceae* (CP-E) produced another set of challenges. Carbapenemases, such as *Klebsiella pneumoniae* carbapenemases (KPC), New Delhi metallo- β -lactamase (NDM), OXA-48, VIM and IMP, are plasmid-encoded enzymes, which have emerged worldwide. The rate of acquisition of CP-E after foreign travel is unknown and only reported in case reports, as recently reviewed by Van der Bij and Pitout.⁵ In The Netherlands, CP-E were found for the first time in 2010.⁶ No data were available on the pre-and post-travel carriage rates among travelers from the Netherlands.

Our objective was to investigate whether Dutch travelers are at risk of acquiring MDR-E (ESBL-E and/or CP-E) by use of a prospective cohort study design. Because detailed microbiological data of the isolates and epidemiological data are crucial for assessing the real public health impacts of these organisms, we also investigated the persistence of intestinal colonization and possible spread to household contacts six months after the travelers returned.

5.3 Materials and Methods

5.3.1 Study design

A prospective cohort study was conducted at the travel clinic at the Leiden University Medical Center (LUMC) and at the Hollands Midden Municipal Health Services (MHS) in Leiden, The Netherlands. During March-September 2011, all adults who made an appointment for travel advice and had the intention to travel to areas outside Europe, North America and Australia were invited to participate in the study.

Travelers < 18 years of age and those who traveled > 3 months were excluded. Only one person of a couple or travel group was included. Participants were asked to

complete an electronic questionnaire and to deliver a rectal swab immediately before and immediately after travel. Questionnaires were used to collect demographic data, previous medical history and travel information. Travelers who acquired a MDR-E after foreign travel were asked to fill out a third questionnaire and deliver a third rectal swab six months after return.

If travelers were positive for MDR-E six months after return, their household contacts were also requested to submit a rectal swab and questionnaire. Household contacts were defined as persons who shared the same household with a participant on a regular basis. MDR-E positive participants were asked to deliver a fourth rectal swab at the same time. The study was approved by the Leiden University Medical Center medical ethics committee.

5.3.2 Bacterial isolates

Rectal swabs were collected with the Stuart Agar Gel Medium Transport Swabs (Copan Diagnostics, Corona, CA). The swabs were inoculated in trypticase soy broth (TSB) supplemented with cefotaxime 0.25 mg/L and vancomycin 8 mg/L (MP products, Groningen, The Netherlands) and incubated for 24 hours at 37°C. After overnight incubation the TSB broths were subcultured on chromogenic ESBL screening agar (ESBL-ID, BioMérieux) and sheep blood agar as a growth control. All Gram-negative rods growing on the ESBL-ID were identified by MaldiToF-MS with BioTyper software (version 3.0; Bruker Daltonics, Germany) and antimicrobial susceptibility testing was performed using the VITEK2® system (BioMérieux, Marcy-l'Étoile, France). All isolates underwent ESBL confirmatory disk testing by disk diffusion for ceftazidime and cefotaxime or cefepime (in ceftazidime resistant isolates) with and without clavulanic acid as recommended by the Clinical Laboratory Standards Institute's guidelines.

MICs for meropenem and ertapenem were determined using Etests (AB Biodisk, Solna, Sweden) according to the manufacturer's instructions. MICs were interpreted using EUCAST criteria (http://www.eucast.org/clinical_breakpoints/).

5.3.4 Molecular characterization of β -lactamases

Molecular characterization of the β -lactamase genes in ESBL-E was performed by using Check-MDR CT103 microarray, version 1.1 (Checkpoints B.V., Wageningen, The Netherlands). The principals of the microarray system and interpretation software have been described previously.⁷ Concisely, the system combines ligation-

mediated amplification with the detection of amplified products on a microarray to detect the various carbapenemase genes: (OXA-48, NDM-1, IMP, VIM and KPC), CTX-M groups (CTX-M group 1, 2, 9 or combined 8/25) and the most prevalent ESBL-associated single nucleotide polymorphisms (SNPs) in TEM and SHV-variants. Furthermore, the six plasmid-mediated AmpC β -lactamases can be identified (www.lahey.org/studies).

5.3.5 Molecular typing of *Escherichia coli* isolates

Multilocus sequence typing (MLST) was performed on all *E. coli* isolates using seven housekeeping genes (*adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA* and *recA*) to determine the corresponding sequence type (ST) and to designate the sequence type complex (STC) by using the MLST Databases at the Environmental Research Institute, University College Cork website (<http://mlst.warwick.ac.uk/mlst/>).

5.3.6 Data analysis

A logistic regression model was used to determine risk factors for the acquisition of ESBL-E/CP-E after foreign travel for a total of 338 participants. Associations between acquiring an ESBL-E/CP-E post-travel and different variables are calculated as odds ratios and p-values. Participants who were positive for ESBL-E/CP-E before travel were analyzed separately. Database processing and statistical analyses (univariate and multivariate analysis) were performed by using the SPSS software version 20.0 (SPSS Inc., Chicago, IL, USA). MLST analysis was performed using BioNumerics software v.6.6 (Applied Maths, St-Martens-Lathem, Belgium).

5.4 Results

5.4.1 Study population and travel characteristics

In total, 521 participants were invited to participate in the study. 370 travelers completed two questionnaires and sent in two rectal swabs and were included in the analysis (Figure 1). The median age of the study population was 33 years (range 19-82) and 234 (63.2%) were women. The median length of stay abroad was 21 days (range 6-90 days). The most common reason for travel was vacation (n=277). Of the 370 participants, 113 (30.5%), whose pretravel swabs were negative, acquired MDR-E during foreign travel. Of these 113 participants, 19 (16.8%) still

carried MDR-E six months after return. In 32 of the 370 participants (8.6%), MDR-E was identified before travel. Twenty (62.5%) of these 32 participants returned with MDR-E, 7 of whom were still colonized after six months (35%). No MDR-E was found before or after travel in 225 participants (60.8%).




Figure 1. Flowchart of participants in the study

5.4.2 Travel-associated risk factors for ESBL-acquisition in returning travelers

For the analysis of travel-associated risk factors, data of 338 returning travelers with negative pretravel swabs were used (Table 1). In total, 65 countries were visited; these are subdivided in 10 subcontinents. The most common destinations were Indonesia (n=62), Thailand (n=30), Malaysia (n=27), Cambodia (n=21), China (n=39), Kenya (n=30), Tanzania (n=24), Surinam (n=20) and South Africa (n=19).

The highest ESBL-E acquisition rates were identified among travelers who visited countries in Asia: 73% in South Asia and 67% in East Asia. Univariate and multivariate analysis showed that the travel destinations South and East Asia were significant risk factors for the acquisition of ESBL-E in returning travelers ($p < 0.001$). Participants traveling to Asia (all subcontinents) were more likely to return with an ESBL-E after a self-arranged trip (odds ratio 1.7; $p = 0.07$) and stay in hostels/lodges (odds ratio 1.9; $p = 0.08$), although this was not statistically significant. There were no other risk factors for the acquisition of ESBL-E after foreign travel.

Table 1. Person and travel characteristics and risk factors for ESBL-E acquisition in a cohort of 338 travelers from the Netherlands*

Variable	No.(%) negative pre- and post- travel N=225 (%)	No. (%) positive post- travel only, N=113	Univariate analysis OR (95% CI)	p-value †	Multivariate analysis OR (95% CI)	p-value
Gender, female	144 (64)	69 (61.1)	0.88 (0.55-1.41)	0.60		
Age						
18-25 years	54 (24.0)	28 (24.8)	1.0			
26-33 years	66 (29.3)	24 (21.2)	0.39 (0.17-0.92)	0.03		
34-51 years	56 (24.9)	28 (24.8)	0.67 (0.28-1.61)	0.37		
≥ 52 years	43 (21.8)	33 (29.2)	1.46 (0.60-3.54)	0.41		
Vegetarian	13 (5.8)	6 (5.3)	0.91 (0.34-2.47)	0.86		
Health care worker	59 (26.2)	27 (23.9)	0.88 (0.52-1.49)	0.64		
Daily contact with farm animals	8 (3.6)	4 (3.5)	1.0 (0.29-3.38)	0.99		
Visit to identified risk areas during previous 12 months			1.01 (0.64-1.61)	0.96		
None	138 (61.3)	69 (61.1)				
Africa	26 (11.6)	15 (13.3)				
Asia	21 (9.3)	12 (10.6)				
India	5 (2.2)	4 (3.5)				
Middle East	26 (11.6)	13 (11.5)				
Central America and Caribbean Region	17 (7.6)	7 (6.2)				
South America	6 (2.7)	5 (4.4)				
Medical problem ‡						
None	161 (71.6)	84 (74.3)				
Inflammatory bowel disease	2 (0.9)	1 (0.9)				
Chronic diarrhea	3 (1.3)	0				
Chronic constipation	3 (1.3)	1 (0.9)				
Irritable bowel syndrome	17 (3.1)	7 (6.2)				
Diabetes mellitus	3 (1.3)	1 (0.9)				
Gastroesophageal reflux	12 (5.3)	4 (3.5)				
Recurrent UTIs	4 (1.8)	1 (0.9)				
Autoimmune disease	7 (3.1)	2 (1.8)				
Abdominal pain with unknown origin	5 (2.2)	2 (1.8)				
Gallbladder problems	4 (1.7)	1 (0.9)				
Transplantation	1 (0.4)	0				
Coeliac disease	0	2 (1.8)				
Other	30 (13.3)	18 (15.9)				
Antibiotic use during 12 months before travel	47 (20.9)	17 (15.1)	0.85 (0.56-1.29)	0.45		
Hospitalization during 12 months before travel						
<3 months prior	5 (2.2)	3 (2.7)				
3-6 months prior	2 (0.9)	1 (0.9)				
6-9 months prior	1 (0.4)	1 (0.9)				
9-12 months prior	2 (0.9)	2 (1.8)				
Travel destination, by subcontinents¶						
Southeast Asia	73 (32.4)	37 (32.7)	1.01 (0.63-1.64)	0.96		
East Asia	11 (4.9)	22 (19.5)	4.70 (2.19-10.1)	<0.001	3.95 (1.78-8.73)	0.001
South Asia	7 (3.1)	18 (15.9)	5.90 (2.39-14.60)	<0.001	5.09 (2-12.92)	0.001
Central Asia	2 (0.9)	1 (0.9)	1.0 (0.089-1.11)	0.99		
Middle East	13 (5.8)	2 (1.8)	0.29 (0.07-1.33)	0.11	0.28 (0.06-1.30)	0.103
North Africa	6 (2.7)	4 (3.5)				
Central Africa	39 (17.3)	17 (15)				
Southern Africa	23 (10.2)	3 (2.7)	0.24 (0.07-0.82)	0.02	0.24 (0.07-0.85)	0.027
Central America and the Caribbean	21 (9.3)	7 (6.2)	0.64 (0.26-1.56)	0.33		
South America	30 (13.3)	2 (1.8)	0.12 (0.027-0.50)	0.004	0.14 (0.03-0.59)	0.008

Continued

Table 1. Continued

Variable	No. (%) negative pre- and post-travel N=225 (%)	No. (%) positive post- travel only, N=113	Univariate analysis OR (95% CI)	p-value †	Multivariate analysis OR (95% CI)	p-value
Median duration of stay in days (range)	21 (6-90)	22 (6-89)	0.99 (0.976-1.004)	0.17	1.0 (0.97-1.0)	0.22
Type of travel						
Self-arranged travel	95 (42.2)	52 (46)	1.17 (0.74-1.84)	0.51		
Backpacking	51 (22.7)	25 (22.1)	0.97 (0.56-1.67)	0.91		
Organized group travel	62 (27.6)	27 (23.9)	0.83 (0.49-1.39)	0.47		
Cruise	1 (0.4)	0				
Other	16 (7.1)	9 (8)				
Own (holiday) home	16 (7.1)	3 (2.7)				
Other	3 (1.3)	3 (2.7)				
Reason for travel						
Vacation	166 (73.8)	83 (73.5)				
Visit family/friends	8 (3.6)	8 (7.1)				
Business	15 (6.7)	9 (8.0)				
Study	18 (8)	7 (6.2)				
Volunteer work	10 (4.4)	5 (4.4)				
Travel group composition						
Alone	25 (11.1)	14 (12.4)	1.13 (0.56-2.27)	0.73		
With 1 partner	102 (45.3)	44 (38.9)	0.77 (0.49-1.22)	0.26		
More partners	44 (19.6)	30 (26.5)				
Group travel	54 (24.0)	25 (22.1)	1.23 (0.78-1.93)	0.37		
Accommodation during travel						
Luxury hotels	78 (34.7)	34 (30.1)	0.81 (0.50-1.32)	0.40		
Hostels	50 (22.2)	30 (26.5)	1.27 (0.75-2.13)	0.38		
Budget hotels	49 (21.8)	27 (23.9)	1.13 (0.66-1.93)	0.66		
Own holiday home	16 (7.1)	3 (2.7)				
Camping	10 (4.4)	6 (5.3)				
With family/friends	8 (3.6)	5 (4.4)				
Locals	7 (3.1)	3 (2.7)				
Boat	4 (1.8)	2 (1.8)				
Other	3 (1.3)	3 (2.7)				
Diarrhea during travel	83 (36.9)	45 (39.8)	1.13 (0.71-1.80)	0.60		
Companion travelers with diarrhea	115 (51.1)	61 (54)	1.1 (0.71-1.77)	0.62		
Antibiotic use during travel	10 (4.4)	9 (8.0)	1.86 (0.73-4.72)	0.19	1.98 (0.72-5.47)	0.16

*Data are presented as no. (%), unless stated otherwise. Blank cells indicate no data available for value. OR, odds ratio; UTI, urinary tract infection.

†Variables with $p < 0.2$ in the univariate analysis were included in the multivariate logistic regression model.

‡Participants could report >1 medical problem.

§ Travel destinations visited by the travelers who completed the study were divided in 10 subcontinents (n = no. of travelers per destination). One participant could have visited >1 country: Southeast Asia: Cambodia (n = 21), Philippines (n = 1), Indonesia (n = 62), Laos (n = 9), Malaysia, (n = 27), Singapore (n = 9), Thailand (n = 30) and Vietnam (n = 17); East Asia: People's Republic of China (n = 39), Japan (n = 1), Mongolia (n = 4) and Taiwan (n = 1); South Asia: Bangladesh (n = 1), India, (n = 20) Maldives (n = 2), Nepal (n = 8) and Sri Lanka (n = 5); Central Asia: Kazakhstan, (n = 2), Kyrgyzstan (n = 2) Uzbekistan (n = 2) and Turkmenistan (n = 1); Middle East: Iran (n = 1), Jordan (n = 1), Turkey (n = 14) Emirates (n = 3); North Africa: Egypt (n = 10) and Morocco (n = 5); Central Africa: Benin (n = 1), Cameroon (n = 1), Congo (n = 7), Gambia (n = 2), Ghana (n = 1), Kenya (n = 30), Liberia (n = 1), Rwanda (n = 1), Sierra Leone (n = 1), Tanzania (n = 24) and Uganda (n = 9); Southern Africa: Angola (n = 1), Botswana (n = 5), Lesotho (n = 2), Madagascar (n = 3), Malawi (n = 5), Mauritius (n = 1), Mozambique (n = 2), Namibia (n = 7) South Africa (n = 19), Swaziland (n = 6), Zambia (n = 6) and Zimbabwe. (n = 1); Central America and the Caribbean: Belize (n = 2), Bonaire (n = 1), Costa Rica (n = 9), Cuba (n = 5), Curacao (n = 1), Dominican Republic (n = 4), Grenada (n = 1), Guatemala (n = 4), Honduras (n = 2), Mexico (n = 9), Nicaragua (n = 5) and Panama (n = 3); South-America: Argentina (n = 3), Bolivia (n = 2), Brazil (n = 5), Chile (n = 2), Ecuador (n = 3), Guyana (n = 3), Peru (n = 3), Surinam (n = 20), Trinidad and Tobago (n = 2) and Venezuela (n = 1).

The incidence proportions of ESBL-E after foreign travel are listed in Table 2.

Table 2. Incidence proportions and incidence rates for ESBL-E in 338 Dutch travelers

Destination	No of travelers	No of travelers with ESBL-E (%)	Incidence proportion % (SE)	Person days (all travelers)	Mean duration of travel (all travelers)	ESBL incidence rate per 100 pdt* (SE†)
Southeast Asia	110	37 (34)	34 (4.5)	2980	27	1.24 (0.20)
East Asia	33	22 (67)	67 (8.3)	776	24	2.83 (0.60)
South Asia	25	18 (72)	72 (9.2)	599	24	3.01 (0.70)
Central Asia	3	1 (30)	33 (33.3)	94	31	1.06 (1.06)
North Africa	10	4 (40)	40 (16.3)	112	11.2	3.57 (1.76) ‡
Middle Africa	56	17 (30)	30 (6.2)	1637	29	1.04 (0.25)
Southern Africa	25	3 (12)	12 (6.6)	631	25	0.48 (0.27)
Middle East	15	2 (13)	13 (9.1)	222	14.8	0.90 (0.64)
Central America and the Caribbean	28	7 (25)	25 (8.3)	544	19	1.29 (0.48)
South America	32	2 (6)	6 (4.4)	922	29	0.22 (0.15)
All travelers	338	113 (33)	33 (2.6)	8536	25	1.32 (0.12)

* Pdt: person days of travel;

† SE standard error.

‡ The ESBL incidence rate per 100 person days of travel is represented by 4 ESBL-E carrying returning travelers from North Africa. Three of them had traveled for 7 days and one traveler had a 25-day stay abroad, which explains the high standard error.

5.4.3 Microbiological results and molecular characterization

A total of 133 participants were colonized with an MDR-E after travel. This group consisted of 113 travelers, who had initially negative pretravel swabs. In addition, 20 participants who had positive pretravel swabs also returned colonized with MDR-E. The ESBL-E of these 133 post-travel swabs consisted of 146 *E. coli*, 10 *K. pneumoniae* and 2 *Enterobacter cloacae* isolates.

No CP-E were found among the pre-and post-travel isolates. Molecular characterization of the post-travel isolates demonstrated that CTX-M group 1 ESBL (n=110) predominated (CTX-M-1 like, n=4; CTX-M-3 like, n=1; CTX-M-15 like, n=85; CTX-M-32 like, n=20), followed by CTX-M group 9 ESBL (n=42), CTX-M group 2 (n=2) and CTX-M group 8/25 (n=1). One *E. coli* isolate carried an SHV-ESBL (238S+240K). In addition some isolates coproduced plasmid-mediated AmpC β -lactamase, ACT/MIR (n=1) or CMY-2 (n=2).

Thirty-four ESBL-E were isolated from pretravel rectal swabs from 32 participants. Twenty-nine (85.3%) of them were positive for *E. coli*, 4 *Klebsiella pneumoniae* (11.8%) and 1 for *Citrobacter freundii* (2.9%). The CTX-M group 1 ESBL (n=22) predominated (CTX-M-1 like, n=4; CTX-M-15 like, n=16; CTX-M-32 like, n=2); the remaining ESBL isolates belonged to CTX-M group 9 ESBL (n=8) and CTX-M group 2 (n=1).

Two *E. coli* isolates carried an SHV-ESBL (238S+240K).

Co-resistance to other classes of antimicrobial drugs was common in pre- and post-travel isolates; 67% displayed resistance to cotrimoxazole, 36% to ciprofloxacin, 37% to tobramycin, 35% to gentamicin and 29% to nitrofurantoin. All isolates were susceptible to colistin and carbapenems.

5.4.4 MLST of ESBL-producing *E. coli* isolates

MLST of 146 *E. coli* isolates from the post-travel swabs identified 86 different sequence types (ST); 31 new STs were found. The most prevalent STs were: ST38 (12%; n=17), ST10 (7%; n=10) and ST131 (4%; n=9). The distribution of the CTX-M groups and types and STs is displayed in Figure 2. There was no association between ST and ESBL-type, nor were STs associated with specific travel destinations. Pretravel isolates showed a similar diversity of STs, of which three were ST131.




Figure 2. Multilocus sequence typing of *Escherichia coli* (n=146) from the post-travel isolates of 133 travelers from the Netherlands. The numbers indicate the most prevalent sequence types (STs). Gray shadow indicates that >1 ST belongs to the same complex. The following sequences belong to STC10: ST4,10, 34, 43, 44, 48, 167, 193, 215, 218, 227, and 617. Thick connecting lines indicate single-locus variants; thin connecting lines indicate variants with 2–3 loci differences; dashed connecting lines indicate variants with 4 loci differences; dotted connecting lines indicate 5–7 loci differences.

5.4.5 Prolonged carriage and household contacts

Of the 133 participants whose samples were positive for ESBL-E after return, 127 (95.4%) completed the follow-up survey and provided samples after six months. ESBL-E was isolated from 26 samples (20.4%) (Table 3). None of these participants reported the use of antibiotics, or were hospitalised during the previous six months; none were healthcare workers and none reported contact with farm animals. Seven participants reported diarrhea.

Of 113 participants, who had initially negative pretravel swabs and positive swabs immediately after return, 19 (16.8%) were still colonized after six months. Of these, 7 participants had samples that were positive for *E. coli* with the same ST six months after return. Nine participants carried an *E. coli* with a different ST six months after return. Three were positive for a different species six months after return. Eleven household contacts of 4 MDR-E positive participants agreed to cooperate and submitted a rectal swab. ESBL-producing *E. coli* was isolated from 2 (18.1%) household contacts, each from different households. The first household contact carried a different ESBL-producing *E. coli* than the associated traveler before and after travel. Both isolates carried a CTX-M group 9 enzyme. The second household contact was positive for SHV-ESBL producing *E. coli* ST2599. The associated traveler's sample were positive for *E. coli* ST617 and ST38 immediately after travel, *K. pneumoniae* six months after return and a CTX-M-15 like *E. coli* ST3363 in the fourth rectal swab.

Of 20 participants whose samples were positive before and after return, 7 participants (35.0%) were still colonized six months after return. Of these 7 participants, 5 carried a similar strain: 2 carried a CTX-M- group 9-producing *E. coli* with an identical ST as before travel, 2 carried a similar ST but with a different CTX-M group enzyme as before travel and one participant carried a CTX-M group 1 producing *K. pneumoniae* during the study period. Two participants returned with an *E. coli* with a different ST. No household contacts participated in this subgroup of travelers.

Table 3. Microbiological and molecular characteristics of rectal swabs samples collected from travelers from the Netherlands immediately pre-and post-travel and 6 months after return*

ID	Pretravel swab			Immediate post-travel samples						Post-travel sample 6 months after return†		
	Species	CTX-M group	ST	Isolate 1			Isolate 2			Species	CTX-M group	ST
				Species	CTX-M group	ST	Species	CTX-M group	ST			
25	Neg	NA	NA	<i>E. coli</i>	9	131	Neg	NA	NA	<i>E. coli</i>	9	131
45	Neg	NA	NA	<i>E. coli</i>	1	405	<i>E. coli</i>	9	38	<i>E. coli</i>	1	405
56	Neg	NA	NA	<i>E. coli</i>	1	3036	<i>E. coli</i>	1	517	<i>E. coli</i>	1	3267
60	Neg	NA	NA	<i>E. coli</i>	1	648	<i>K. p.</i>	1	ND	<i>E. coli</i>	1	648
61	Neg	NA	NA	<i>E. coli</i>	1	648	<i>E. coli</i>	9	227	<i>E. coli</i>	1	131
62	Neg	NA	NA	<i>E. coli</i>	9	3037	Neg	NA	NA	<i>E. coli</i>	9	501
80	Neg	NA	NA	<i>E. coli</i>	1	131	Neg	NA	NA	<i>E. coli</i>	9	1177
86	Neg	NA	NA	<i>E. coli</i>	1	93	<i>E. coli</i>	1	2090	<i>E. cloacae</i>	9	ND
137	Neg	NA	NA	<i>E. coli</i>	1	155	<i>E. coli</i>	1	617	<i>E. coli</i>	9	131
204	Neg	NA	NA	<i>E. coli</i>	1	38	Neg	NA	NA	<i>K. p.</i>	1	ND
211	Neg	NA	NA	<i>E. coli</i>	1	3044	Neg	NA	NA	<i>K. p.</i>	1	ND
222	Neg	NA	NA	<i>E. coli</i>	9	2003	Neg	NA	NA	<i>E. coli</i>	9	2003
238	Neg	NA	NA	<i>E. coli</i>	9	414	Neg	NA	NA	<i>E. coli</i>	9	10
251	Neg	NA	NA	<i>E. coli</i>	1	34	Neg	NA	NA	<i>E. coli</i>	1	450
309	Neg	NA	NA	<i>E. coli</i>	1	3045	Neg	NA	NA	<i>E. coli</i>	1	3045
373	Neg	NA	NA	<i>E. coli</i>	1	38	Neg	NA	NA	<i>E. coli</i>	1	3266
387	Neg	NA	NA	<i>E. coli</i>	1	131	Neg	NA	NA	<i>E. coli</i>	1	131
454	Neg	NA	NA	<i>E. coli</i>	9	10	Neg	NA	NA	<i>E. coli</i>	9	10
474	Neg	NA	NA	<i>E. coli</i>	1	154	Neg	NA	NA	<i>E. coli</i>	1	131
12	<i>E. coli</i>	9	38	<i>E. coli</i>	1	3074	Neg	NA	NA	<i>E. coli</i>	1	38
105	<i>E. coli</i>	1	191	<i>E. coli</i>	1	120	<i>E. coli</i>	1	38	<i>E. coli</i>	1	120
255	<i>E. coli</i>	9	131	<i>E. coli</i>	1	617	Neg	NA	NA	<i>E. coli</i>	9	131
269	<i>K. p.</i>	1	ND	<i>K. p.</i>	1	ND	Neg	NA	NA	<i>K. p.</i>	1	ND
283	<i>E. coli</i>	9	131	<i>E. coli</i>	1	46	Neg	NA	NA	<i>E. coli</i>	9	131
505	<i>E. coli</i>	1	1163	<i>E. coli</i>	1	69	Neg	NA	NA	<i>E. coli</i>	9	3268
512	<i>E. coli</i>	9	657	<i>E. coli</i>	9	657	Neg	NA	NA	<i>E. coli</i>	1	657

*ID, participant identification number; CTX-M, extended-spectrum β -lactamase enzyme; ST, sequence type; Neg, no species were isolated from sample; *E.*, *Escherichia*; *K.p.*, *Klebsiella pneumoniae*; NA, not applicable; ND, no sequence type data available.

†None of the participants with a positive rectal swab sample after 6 months reported antimicrobial drug use during the 6 months after return.

5.5 Discussion

The results of this study show a high ESBL-E carriage rate of 30.5% among healthy participating travelers from the Netherlands after return. This finding is worrisome, because this ESBL-E carriage rate is higher compared with those in recent studies that identified international travel as an independent risk factor for ESBL-E colonization.¹⁻⁴ It is striking that none of the potential travel-associated risk factors investigated in this study, other than traveling to South and East Asia, were found to contribute to this high ESBL-E carriage rate. Additional risk factors were not revealed by including the univariate analysis the 13 participants who had a positive pretravel swab, and acquired an ESBL-producing *E. coli* during travel with a different ST than before travel. Gastroenteritis during travel was found to be associated with the risk of ESBL-E acquisition in Swedish travelers.³ That association was not found in this study, which may reflect less fecal-oral contamination while traveling. Baaten et. al. reported that diseases transmitted by the fecal-oral route among travelers to non-industrialized countries have declined because of improved hygiene standards at the destination as measured by the human developmental index, sanitation index and the water source index.⁸ The sanitation index (SI) levels, which represent the proportion of the population that has access to sanitation, were the lowest for Sub-Saharan Africa and the Indian subcontinent. On the basis of these indices, we would expect the incidence of ESBL-E acquisition to be similar among travelers in countries in Asia and Africa. Nonetheless, participating travelers to Asia had the highest post-travel colonization rates. Travelers to Asia most likely differ in their eating habits compared to travelers to African countries, since the former are more likely to eat in individual establishments outside of hotels or from street vendors. Thus, the high incidence rate found in returning travelers from Asia in this study may result from the increased risk for foodborne exposure.

No CP-E were found despite the fact that countries were visited where CP-E are prevalent in hospitals and in the environment.⁵ Other known risk areas besides India, for the acquisition of CP-E, such as the USA, Greece, Italy and the Balkan region were not included in this study, because these travelers do not visit the Travel Clinic of the LUMC. Many citizens from the Netherlands have relatives in North-African countries or Turkey whom they visit frequently. OXA-48 producing bacteria are endemic to these countries.⁹ These travelers do not consult travel clinics and may well return carrying OXA-48 producing isolates unnoticed.

Peirano et. al.² reported that the prevalence of ST131, an uropathogenic *E. coli*, notorious for its worldwide expansion and spread of CTX-M-15, was similar among travelers and non-travelers from the Calgary region. The most prevalent ESBL among the travelers participating in this study was the CTX-M-15 like enzyme. However, this enzyme was found in a plethora of different sequences types of *E. coli*. Participants in the Leiden area not only showed a great heterogeneity of sequence types, but also harboured different CTX-M-types after travel and six months after return. The majority of the *E. coli* strains identified in the participants in this study were of sequence types that clustered around ST10 and belonged to sequence type complex 10 (STC10). STC10 strains essentially belong to the non-virulent, commensal phylogenetic group A.¹⁰ In a recent study based in France, isolates belonging to STC10 were found to be the most prevalent among fecal samples of healthy carriers of nalidixic acid resistant (but ESBL-negative) *E. coli*.¹¹ It is also the most prevalent STC in the MLST database. Data from this study show that transmissible genetic elements containing resistance genes are exchanged with naïve *E. coli* strains of the human intestinal microbiota during foreign travel combined with foodborne exposure.

Although 26 participants had positive results for ESBL-E six months after travel, they were not all positive for the same enterobacterial strain that was identified immediately after travel. In 8 travelers colonized with *E. coli*, an ESBL of the same CTX-M group was found in the immediate post-travel sample as after six months, but *E. coli* with a different ST was detected. In 11 travelers, the strain persisted during the study period. It is possible that more strain types were present in the rectal samples where colony morphology of different strains was not discriminative. However, it is also possible that the transfer of ESBL genes between strains within a host is a frequent occurrence. Or, the acquisition of a new ESBL-E occurs at the expense of the resident strain.

Interhousehold transmission of ESBL-E has been demonstrated in the community setting.^{12,13} Clonally related strains could be found for 66% of the isolates from infected community patients and their corresponding household contacts.¹³ Because of the limited data on household contacts in the present study, the transmission dynamics of ESBL-E in households after foreign travel remain to be discovered.

The high pretravel ESBL-E carriage rate among our study participants (8.6%) was an unexpected finding. Two recent studies on the ESBL-E carriage rate in the community have been conducted in the Amsterdam area. In the first study, 10.1% of

the fecal samples from outpatients with gastrointestinal discomfort being assessed by their general practitioners yielded ESBL-E, predominantly CTX-M-15 producing *E. coli*.¹⁴ In a second study, investigating the prevalence of ESBL-E carriage in the general community, a carriage rate of 8.5% was found (E.A. Reuland, unpublished data). Although no data on travel history were given, the investigators pointed out that foreign travel might be responsible for at least part of ESBL-E carriage rates among outpatients from the Netherlands. This finding is supported by data from our study: 50% of participants who had a positive pretravel swab had traveled during the previous 12 months. This high percentage of carriers identified in this study before travel points towards ongoing importation of ESBL-E. Other potential reservoirs for ESBL-E are poultry and other retail meat, which have been found to be contaminated with ESBL-producing *E. coli* strains harboring the genes on identical plasmids as found in human isolates.^{15, 16}

International travel is growing and the number of intercontinental flights has increased during the past decade. The findings in this study support the role of international travel on the ESBL-E acquisition and carriage rates among travelers from the Netherlands, especially to South and East Asia. The high pre- and post-travel carriage rates among persons traveling from the Netherlands indicate that the consequences of increased foreign travel are already manifest in this country. The lack of apparent travel-associated risk factors, the spread of CTX-M enzymes through a highly diverse population of *E. coli*, the association of ESBL-production with multidrug resistance, and the possible role of other sources make containing the spread difficult. These factors also complicate the implementation of other strategies, such as pretravel advice, and imply that all travelers to Asia should be considered for carriage of ESBL-E. Although CP-E were not found in this study, CP-E have been introduced into the Netherlands by returning travelers,^{6,17-19} and introduction by asymptomatic travelers to the Netherlands from countries where CP-E are endemic, may largely go unnoticed. There is no reason to assume that, after CP-E are introduced, their spread will be less dynamic than of ESBL-E. This inference has serious implications for implementation of screening methods and effective infection control strategies. On the basis of the results of this study, we recommend active surveillance of CP-E and ESBL-E and at least temporary contact isolation precautions for patients being admitted to hospitals after travel to Asia during the previous six months.

Acknowledgements

We thank the technicians from the Department of Medical Microbiology for the cultures performed in this study and Check-Points B.V. for technical support.

This study was partly funded by the European Society of Clinical Microbiology and Infectious Diseases (ESCMID SG Research Grant)

5.6 References

- 1 Kennedy K, Collignon P. Colonisation with *Escherichia coli* resistant to "critically important" antibiotics: a high risk for international travellers. *Eur J Clin Microbiol Infect Dis* 2010; **29**: 1501-6.
- 2 Peirano G, Laupland KB, Gregson DB, et al. Colonization of returning travelers with CTX-M-producing *Escherichia coli*. *J Travel Med* 2011; **18**: 299-303.
- 3 Tangden T. Foreign travel is a major risk factor for colonization with *Escherichia coli* producing CTX-M-type extended-spectrum β -lactamases: a prospective study with Swedish volunteers. *Antimicrob Agents and Chemother* 2010; **54**: 3564-8.
- 4 Weisenberg SA, Mediavilla JR, Chen L, et al. Extended spectrum β -lactamase-producing *Enterobacteriaceae* in international travelers and non-travelers in New York City. *PLoS One* 2012; **7**: e45141.
- 5 van der Bij AK, Pitout JD. The role of international travel in the worldwide spread of multiresistant *Enterobacteriaceae*. *J Antimicrob Chemother* 2012; **67**: 2090-100.
- 6 Leverstein-van Hall MA, Stuart JC, Voets GM, et al. Carbapenem-resistant *Klebsiella pneumoniae* following foreign travel. *Ned Tijdschr Geneesk* 2010; **154**: A2013.
- 7 Cohen SJ, Dierikx C, Al NN, et al. Rapid detection of TEM, SHV and CTX-M extended-spectrum β -lactamases in *Enterobacteriaceae* using ligation-mediated amplification with microarray analysis. *J Antimicrob Chemother* 2010; **65**: 1377-81.
- 8 Baaten GG, Sonder GJ, Van Der Loeff MF, et al. Fecal-orally transmitted diseases among travelers are decreasing due to better hygienic standards at travel destination. *J Travel Med* 2010; **17**: 322-8.
- 9 Nordmann P, Naas T, Poirel L. Global spread of carbapenemase-producing *Enterobacteriaceae* *Emerg Infect Dis* 2011; **17**: 1791-8.
- 10 Bengtsson S, Naseer U, Sundsfjord A, et al. Sequence types and plasmid carriage of uropathogenic *Escherichia coli* devoid of phenotypically detectable resistance. *J Antimicrob Chemother* 2012; **67**: 69-73.
- 11 Leflon-Guibout V, Blanco J, Amaqdouf K, et al. Absence of CTX-M enzymes but high prevalence of clones, including clone ST131, among fecal *Escherichia coli* isolates from healthy subjects living in the area of Paris, France. *J Clin Microbiol* 2008; **46**: 3900-5.
- 12 Lo WU, Ho PL, Chow KH, et al. Fecal carriage of CTX-M type ESBL-producing organisms by children and their household contacts. *J Infect* 2010; **60**: 286-92.
- 13 Valverde A, Grill F, Coque TM, et al. High rate of intestinal colonization with extended-spectrum- β -lactamase-producing organisms in household contacts of infected community patients. *J Clin Microbiol* 2008; **46**: 2796-9.
- 14 Reuland EA, Overdevest IT, Al NN, et al. High prevalence of ESBL-producing *Enterobacteriaceae* carriage in Dutch community patients with gastrointestinal complaints. *Clin Microbiol Infect* 2013; **19**: 542-9.
- 15 Kluytmans JA, Overdevest IT, Willemsen I, et al. ESBL-producing *Escherichia coli* from retail chicken meat and humans: Comparison of strains, plasmids, resistance genes, and virulence factors. *Clin Infect Dis* 2013; **56**: 478-87.
- 16 Overdevest I, Willemsen I, Rijnsburger M, et al. Extended-spectrum β -lactamase genes of *Escherichia coli* in chicken meat and humans, The Netherlands. *Emerg Infect Dis* 2011; **17**: 1216-22.
- 17 Halaby T, Reuland AE, Al NN, et al. A case of New Delhi metallo- β -lactamase 1 (NDM-1)-producing *Klebsiella pneumoniae* with putative secondary transmission from the Balkan region in the Netherlands. *Antimicrob Agents Chemother* 2012; **56**: 2790-1.
- 18 Kalpoe JS, Al NN, Poirel L, et al. Detection of an Ambler class D OXA-48-type β -lactamase in a *Klebsiella pneumoniae* strain in The Netherlands. *J Med Microbiol* 2011; **60**: 677-8.
- 19 Leverstein-van Hall MA, Stuart JC, Voets GM, et al. Global spread of New Delhi metallo- β -lactamase 1. *Lancet Infect Dis* 2010; **10**: 830-1.