

**Evaluating the dietary micro-remain record in dental calculus and its application in deciphering hominin diets in Palaeolithic Eurasia** Power, R.C.F.

# Citation

Power, R. C. F. (2016, November 1). *Evaluating the dietary micro-remain record in dental calculus and its application in deciphering hominin diets in Palaeolithic Eurasia*. Retrieved from https://hdl.handle.net/1887/43970

Version:	Not Applicable (or Unknown)
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/43970

Note: To cite this publication please use the final published version (if applicable).

Cover Page



# Universiteit Leiden



The handle <u>http://hdl.handle.net/1887/43970</u> holds various files of this Leiden University dissertation.

Author: Power, R.C.F. Title: Evaluating the dietary micro-remain record in dental calculus and its application in deciphering hominin diets in Palaeolithic Eurasia Issue Date: 2016-11-01

# Dental calculus evidence of Taï Forest Chimpanzee plant consumption and life history transitions

# Robert C. Power,<sup>1</sup> Domingo C. Salazar-García,<sup>2,3,4</sup> Roman M. Wittig,<sup>5,6</sup> Martin Freiberg,<sup>7</sup> Amanda G. Henry<sup>1</sup>

<sup>1</sup>Max Planck Research Group on Plant Foods in Hominin Dietary Ecology, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany.

<sup>2</sup>Department of Archaeology, University of Cape Town, Cape Town, South Africa.

<sup>3</sup>Departament de Prehistòria y Arqueologia, Universitat de València, València, Spain.

<sup>4</sup>Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, Leipzig, Germany.

<sup>5</sup>Department of Primatology, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6 Leipzig, Germany.

<sup>6</sup>Taï Chimpanzee Project, Centre Suisse de Recherches Scientifiques, Abidjan, Cote d'Ivoire <sup>7</sup>Institute of Botany, University of Leipzig, 04103 Leipzig, Germany.

Published in Scientific Reports (2015, Volume 5, 15161).

## Abstract

Dental calculus (calcified dental plaque) is a source of multiple types of data on life history. Recent research has targeted the plant microremains preserved in this mineralised deposit as a source of dietary and health information for recent and past populations. However, it is unclear to what extent we can interpret behaviour from microremains. Few studies to date have directly compared the microremain record from dental calculus to dietary records and none with long-term observation dietary records thus limiting how we can interpret diet, food acquisition and behaviour. Here we present a high-resolution analysis of calculus microremains from wild chimpanzees (*Pan troglodytes verus*) of Taï National Park, Côte d'Ivoire. We test microremain assemblages against more than two decades of field behavioural observations to establish the ability of calculus to capture the composition of diet. Our results show that some microremain classes accumulate as long-lived dietary markers. Phytolith abundance in calculus can reflect the proportions of plants in the diet, yet this pattern is not true for starches. We also report microremains can record information about other dietary behaviours, such as the age of weaning and learned food processing techniques like nut-cracking.

# 4.1 Introduction

Understanding feeding ecology is crucial for recognising the evolutionary pressures that shaped the great apes and humans. It is long recognised that factors such as dietary specialization, tool-assisted food acquisition and the weaning age of infants are important in great apes and humans, and differ significantly among species (Boesch et al., 1994; Ross, 1998; Boesch and Boesch-Achermann, 2000; Teaford and Ungar, 2000).

However, many approaches to dietary reconstruction leave unanswered specific questions on diet and related life history events, especially for fossil specimens. There is a need for new methods to reconstruct food acquisition from populations that can avoid some of the shortfalls of other techniques like direct observation, stable isotope analysis and microwear studies (Phillips and Lancelotti, 2014; Fiorenza et al., 2015). In some contexts, direct observation is simply not possible, for example with extinct great apes and human groups. Stable isotope analysis and dental microwear studies fail to provide total dietary data, and instead only give a picture of broad dietary patterns such as consumption of particular plant categories or mechanical properties of diet (Grine, 1986; Scott et al., 2012). Furthermore, even where direct observational data on food acquisition is available, data collection is frequently constrained because observation is only feasible over a short window of the lifetime of an individual that may live up to several decades.

Dental calculus sampled from living or dead individuals is rapidly gaining recognition as an invaluable material for the reconstruction of life history. Since Armitage (Armitage, 1975) first recognised plant remains from the teeth of ungulates, studies have reported starch grains, phytoliths, pollen grains, diatoms, mineral particles, proteins and DNA from diverse human and animal populations (Kucera et al., 2011; Adler et al., 2013; Salazar-García et al., 2013; Power et al., 2014b, 2015a; Warinner et al., 2014). Using dental calculus from present day forager-horticulturalists, Leonard (Leonard et al., 2015) showed for the first time that recovered microremains also occur in consumed foods verifying the link between microremains in calculus and diet. As demand grows for dietary history data, analysis of phytoliths and starches in dental calculus is been increasingly used to

reconstruct dietary ecology and ecological niches (Lalueza-Fox et al., 1996; Gobetz and Bozarth, 2001; Boyadjian et al., 2007; Henry et al., 2012; Mickleburgh and Pagán-Jiménez, 2012; Salazar-García et al., 2013; Buckley et al., 2014; Lazzati et al., 2015).

Despite the promise of calculus dietary studies, they are hindered by the lack of research that cross-validates the dietary material recovered in calculus with the organism's actual feeding ecology and life history. Until recently, our understanding of what the plant matter preserved in calculus precisely represents has been speculative. The initial effort to characterise the microremain record by Leonard (Leonard et al., 2015) reported that calculus captured only a limited proportion of dietary breadth. In this study many vegetable foods lacked phytoliths and starches, and cooking may have significantly reduced starch abundance even if present. Dietary patterns were established through interviewing and short-term camp stays by Leonard, and though the recovered microremains corresponded to the average diet of the population, the dietary records lacked insight into the long term life history of individuals. Without dietary records that cross intra- and inter-annual cycles, our knowledge of the nature of the calculus record and its potential for archaeological studies is incomplete. Furthermore, it is unclear if the calculus dietary record has input from non-dietary sources (e.g. preparation of plant-based tools and oral hygiene) as well as the consumption of stomach contents (Buck and Stringer, 2014; Dudgeon and Tromp, 2014; Tromp and Dudgeon, 2015), with bias from diagenetic and taphonomic factors rendering it ultimately purely stochastic.

In our study, we compare the plant microremains from the calculus of the chimpanzees of the Taï Forest to 22 years of group averaged dietary observation data in order to validate the calculus record and explore its potential as a source on information on life histories. For this purpose, the study of chimpanzees provides several strengths as a model. First, the chimpanzee mouth is analogous to humans, in that chimpanzees often accumulate large deposits of calculus unlike some mammals. Secondly, chimpanzees produce salivary amylase unlike some primates (Santos et al., 2012) although it is less abundant than in humans (Perry et al., 2007). Thirdly, Taï chimpanzees have a broad diet that includes nearly all food classes (e.g. fruits, piths, leaves, mammals, birds, invertebrates and honey) and is thus relevant to understanding hominin evolution in the African tropics and dietary ecology of hunter-gatherers living in other tropical regions. Fourthly, chimpanzees only consume raw foods meaning the microremains in their food are unaltered by cooking processes.

We sampled calculus from 24 individual chimpanzees using established methods (Dobney and Brothwell, 1986; Mickleburgh and Pagán-Jiménez, 2012), and built a random forest model in the R software environment (Breiman, 2001) to identify the microremains based on multivariate comparison to reference material (Fenwick et al., 2011; Saul et al., 2012; Out et al., 2014; Coster and Field, 2015; Out and Madella, 2015) (see detailed methods 7.2). We predict that if microremains reflect diet, they are accumulative in calculus and should increase with age of the individual. Chimpanzee sex might also influence the abundance of microremains, since male and female chimpanzee are known to vary in their time allocation to different food resources (Wrangham and Smuts, 1980; Boesch and Boesch-Achermann, 2000; N'guessan et al., 2009; Fahy et al., 2013). We also anticipate that the proportions of microremains from each plant will be determined by the frequency with which that plant is consumed and how abundant the microremains are in the plant tissue. Although we knew the taxonomic identity of the reference plants at the level of species, an important amount of dietary observation data was present only at the genus level. Therefore, we performed our analyses at the genus level in order to have a higher chance of capturing long term dietary averages for the group, and refer only to genera in the text. Except where otherwise noted, our analyses were done at the group level observational data, since the records for individual chimpanzees were not complete enough to provide a detailed overview of life history. We found the phytoliths in dental calculus to be an approximate record of diet, and furthermore that microremains can reflect important behaviour like nut-cracking and episodes of Taï Chimpanzee life history such as the age of weaning.

# 4.2 Materials

# 4.2.1 Taï Forest reference material

A reference collection with 91 genera (113 species) of the most frequently consumed chimpanzee plant foods in the Taï Forest was collected and examined for phytoliths and starch (Table 7; Appendix table 4) starches. Phytoliths and starches were isolated from reference plants using conventional approaches (Piperno, 2006). We selected thirteen starch- and seven phytolith-producing genera from the 91 we analysed for the identification model (Appendix table 4).

Plant category (Genus)	Туре	Plant part	% Microremain/Dry Weight	Minutes eaten
Elaeis	Phytolith	Fruit and leaf	4.81	9379
Eremospatha	Phytolith	Pith	1.72	25,046
Laccosperma	Phytolith	Pith and seed	2.15	5311
Aframomum	Phytolith	Seed and leaf	2.13	1704
Sarcophrynium	Phytolith	Fruit	3.32	1847
Cola	Starch	Seed	40	35,778
Aframomum	Starch	Seed	54.58	1704
Piper	Starch	Seed	39.22	492
Sacoglottis	Starch	Fruit	2.46	258,225
Panda	Starch	Seed	0.85	17,299
Coula	Starch	Seed	31.15	118,095
Napoleona	Starch	Seed	20.79	51
Gilbertiodendron	Starch	Seed	23.87	11,808
Eremospatha	Starch	Pith	2.93	25,046
Calpocalyx	Starch	Fruit	9.1	49,074
Sarcophrynium	Starch	Fruit	23.83	1847
Xylia	Starch	Seed	19.58	46,587
Treculia	Starch	Seed	23.87	58,093

Table 7: Plant genera selected from reference collection species for the predictive identification model with the microremain content of the dried plant material provided as a percent of dried plant material, and the frequency of observed consumption provided as number of minutes eaten. We chose to use genus as the taxonomic rank as some dietary records only identify genus.

# 4.2.2 Calculus sampling

The calculus samples used for our analysis come from permanent and deciduous molars of 24 chimpanzees from the Taï Chimpanzee Osteology Collection at the Max Planck Institute for Evolutionary Anthropology (MPI-EVA) with varied life histories (4.6; Table 8, Appendix table 5). We selected only molars to standardise the sampling, and chose teeth that were encrusted with a prominent band of supragingival calculus (calculus present above the original gumline) on the enamel crown. Deposits of supragingival calculus were present on all individuals  $\geq$  1 years old. Subgingival calculus was also present but was not sampled since it occurs below the former gums and it is unclear if it preserves food remains. Calculus on the teeth was documented with photography before sampling, and the colour noted with how each skeleton as treated before our sampling (Appendix table 5). Packing material was sampled as a control. An unidentified adhesive used in the curation of some specimens was removed before sampling. A dental scalar was then used to remove small areas of calculus. The amount of calculus sampled had no relationship with the

amount of calculus present on the teeth except in the youngest chimpanzees (<5.3 years), where calculus was rare and almost entirely collected. We sampled in clean conditions in a laminar flow cabinet at positive-pressure at the MPI-EVA. We then weighed each of the samples and transferred them to microcentrifuge tubes. After sampling, the teeth and surviving calculus were photographed.

Some studies have highlighted the risks of laboratory contamination from modern plant microremains (Crowther et al., 2014; Barton and Torrence, 2015; Weyrich et al., 2015). To address the possibility of contamination, we conducted a regime of weekly laboratory cleaning to remove contamination. All work surfaces were wiped with hot water, washed with starch-free soap and wiped with 5 % sodium hydroxide (NaOH). We additionally performed wipe tests before and after weekly cleaning to quantify starch contamination and assess contaminating types. Wipe tests retrieved settled particles of the surface area (74 x 43 cm<sup>2</sup>) of the laboratory positive-pressure laminar flow hood used for mounting. Results of these intensive contamination control tests are in Power and colleagues (2015b).

Name	ID	Tooth	Sex	Weight (mg)	Age at death	
					In years	In months
Ophelia	14993	Lower Left DM1	F	0.025	1	12
Leonardo	13432	Upper DM2	М	0.329	1.92	23
Bambou	11777	Lower Left DM1	М	0.135	2.08	25
Piment	11788	Lower Right DM1	F	0.27	3.58	43
Oreste	14995	Lower Left M1	М	0.536	5.25	63
Hector	12175	Upper Right M1	М	0.689	5.67	68
Noah	15011	Lower M1	М	1.165	7	84
Lefkas	13433	Upper Left M2	М	0.595	7.58	91
Tina	11790	Lower Right M1	F	1.36	9.08	109
Dorry	15020	Lower Right M2	F	0.742	11	132
Zerlina	11792	Lower Left M3	F	0.878	12.3	144
Clyde	11779	Lower Right M1	М	1.131	13	156
Agathe	11775	Lower Right M2	F	6.076	16	192
Leo	15012	Lower Right M3	М	1.085	19	228
Bijou	11778	Lower Left M2	F	5.041	19	228
Castor	13439	Lower Left M1	F	6.982	22	264
Kendo	11781	Lower Left M2	М	2.895	25	300
Fanny	11780	Lower Left M3	F	3.915	25	300
Venus	15001	Upper Right M1	F	1.133	27	324
Goma	15004	Upper Right M3	М	13.208	28	336
Rubra	15023	Lower Left M2	F	6.751	38	456
Ondine	11786	Lower Left M1	F	1.529	39	468
Mkubwa	13435	Lower Right M3	М	0.324	40	480
Brutus	15029	Upper Left M3	М	3.246	46	552

Table 8: All chimpanzee dental calculus samples analysed.

#### 4.3 Methods

# 4.3.1 Optical microscopy

Optical microscopy was performed at the Plant Foods in Hominin Dietary Ecology laboratory in the MPI-EVA (for reference collection microscopy see 4.6 and Appendix table 6). We added 150  $\mu$ l of 10 % hydrochloric acid to the calculus sample for one to three hours. The samples were then centrifuged at 1691 x g (Heraeus MEGAFUGE 16 with a microcentrifuge rotor) for 10 minutes and then about 100  $\mu$ l of supernatant was decanted and replaced with distilled water. This was repeated three times to remove the hydrochloric acid. After the second decanting, it was refilled with a 25 % glycerine solution. The sample was then ground in the solution in the 1.5 ml Eppendorf microcentrifuge tube to reduce sample loss due to static

electricity. The samples then were centrifuged again at the same speed, and about 1 ml of supernatant was decanted. We mounted 20  $\mu$ l per slide on as many slides as needed in order to examine the entire sample. Microscopy was used as in conventional phytolith and starch studies (Power et al., 2014a; b). We examined each slide under bright field and cross-polarized light on a Zeiss Axioscope microscope at 400 × magnification. We photographed each microremain and described each with the international microremain nomenclature including the International Code of Phytolith Nomenclature (Madella et al., 2005). In some cases, starch aggregates were identified in calculus. In this case, each component grain of each aggregate was counted as an individual starch.

# 4.3.2 Microremain identification

We identified microremains with a reference collection using multivariate analysis with a random forest algorithm. We collected five general microremain measurements and four specific to phytoliths and six specific to starches from a total of 900 reference microremains (Table 7). With the reference collection generated (Appendix table 6) we generated a certainty score that each matched each reference collection genus. The validity was tested through cross-validation with a subset of reference data. We identified the microremain as coming from the genus with the highest certainty score.

# 4.3.3 Behavioural records

The chimpanzees of the Taï Forest have been studied since the commencement of the Taï Chimpanzee Project in 1979 (Boesch and Boesch-Achermann, 2000). The detailed recorded behaviour of the group included observation of feeding time and food item consumed. The feeding records used in our study span the period between 1992 and 2014. The database includes 1,165,150 million behavioural observations of about 128 chimpanzees, with a total of 417,628 dietary observations (2,380,202 minutes). However, only roughly 30,000 observations come from chimpanzees available for sampling at the osteology collection. Furthermore, most of these chimpanzees have only sporadic coverage of their life history. Therefore, instead of using dietary records of individual chimpanzees or the

collated records of the 24 chimpanzees we sampled, we chose to combine dietary records from all 128 chimpanzees to best represent the average Taï Forest diet.

The feeding record includes the times when a chimpanzee started and finished eating, and the food consumed. We chose only those feeding records where the genus of the plant food eaten was documented, and calculated the total amount of time spent consuming each resource. Behavioural records do not account for variations in the volume of food consumed in given number of minutes. In addition, although some observations record the specific plant part that was eaten, most do not, so we do not include this information.

# 4.3.4 Statistical analysis

To test for the effects of age on microremains we used a negative binomial regression (log link) with a count of each microremain class treated as a response (phytoliths, starches and other unsilicified plant fragments) using a likelihood ratio test in R 3.1.0 (R Core Team, 2014). We ran the regression using the glm.nb function of R package MASS (Venables and Ripley, 2002). The full model included the fixed effects of age and sex (4.6). The mg weight of each calculus sample was used to weigh the model to account for larger samples likely being more representative of overall diet due to the potential of microremains to have a clustered distribution in the calculus matrix. Controlling for weight, heavier samples have less variable microremain counts (Compare Table 8 with Fig. 10). The full model was compared with a null model using an ANOVA. We used likelihood ratio tests comparing the full models with reduced models in which each fixed effect was dropped, one at a time. Model assumptions were met. Collinearity was not an issue (largest Variance Inflation Factor=1.001) and leverage values as well as DFBeta values indicated no obvious highly influential cases.

To explore the relationship between diet and the phytolith microremains found in dental calculus we tested an observational random effect Poisson model with likelihood ratio tests. We used counts of each genus predicted to be present with the total minutes spent consuming each genus. For this, we used the glmer function of the R package lme4 (Bates et al., 2013). If any genus was not predicted to be present in a chimpanzee sample, they were included as a 0 value. Our full model included minutes and chimpanzee age in months as fixed effects, and sex as a control predictor. The model included the weight of each calculus sample and the successful identification rate of each type of genera as model weights, and used microremain content as an offset to factor in significant differences in content between different genera. To prepare the data, we z-transformed the minutes and age variables. The chimpanzee individual was included as a random slope term while year of death, tooth and food type were treated as random intercept terms. An id was assigned to each observation, and this was also included as a random thus reducing overdispersion to  $(x^2=13.369, df=116, df=116,$ intercept, dispersion parameter=0.115) in the phytolith model. To test the significance of the full model it was compared with a null model excluding fixed effects of minutes of observation and age. Variance inflation factors (VIF) were derived to assess collinearity using the function vif of the R-package car, from a standard linear model minus random effects, as offsets and weights (Fox and Weisberg, 2002; Field, 2005). Variance inflation factors indicated collinearity to not be an issue (largest VIF=1.02). We tested model stability by excluding each random effect one by one from the data set, running the full model and comparing the results with those from the original model that suggest no highly influential cases.

To explore the relationship between diet and the starch microremains we could not use the same approach due to high zero inflation present in the starch data. To overcome this, we implemented a mixed effects logistic regression using the same terms, random effects, weighs and offset as the phytolith Poisson model. This required the counts data (the response) to be treated as presence and absence data resulting in some loss of data. Variance inflation factors (Field, 2005) were derived to access collinearity using the function vif of the R-package car, from a standard linear model minus random effects as well as offsets and weights (Fox and Weisberg, 2002). Variance inflation factors indicated collinearity to not be an issue (largest VIF=1.018).

# 4.4 Results

#### 4.4.1 Identification of the microremain assemblages

We were able to examine 91 of the 157 genera (113 of 230 species) of plants that the 128 Taï chimpanzees consumed during the observation period. After assessing these plants, we noted thirteen starch-producing genera that could be included in our identification model. Unlike starches, phytolith were abundant in most plants in many different morphotypes. Not all morphotypes could be included in analysis, so we choose one morphotype (globular and spheroid) and identified the five genera that produces them (Table 7; Fig. 9). Most of these thirteen starch-producing and five-phytolith producing genera are major sources of food for these chimpanzees (Appendix fig. 2).

For each microremain-producing plant genus, we collected data from 50 microremains, to provide a range of measurements within each genus. We collected nine types of measurements for phytoliths and 11 for starches from 900 microremains using Zeiss AxioVision Microscopy Software (Appendix table 6). By using a subset of the reference collection to test the model, we assessed the success rate of identification of each genus with the model (Appendix table 8, Appendix table 9). Some genera were reliably identified, and others were more difficult to identify. For example, *Sarcophrynium* phytoliths were identified successfully 94 % of the time while *Panda* starch was only identified 22 % of the time. Generally, phytoliths were identified more reliably (Appendix table 8, Appendix table 9). Using this random forest model, we were able to proceed with identification of microremains recovered from the calculus.



Fig. 9: Starch and phytolith morphotypes used in the identification model. Each scale bar represents 10 μm. (a) *Aframomum sceptrum* seed phytolith, (b) *Aframomum excarpum* leaf (b) *Aframomum excarpum* leaf phytolith, (c) *Aframomum excarpum* seed starch under bright field (left) and cross polarized light (right), (d) *Laccosperma opacum* pith phytolith, (e) *Laccosperma secondiflorum* seed phytolith, (f) *Calpocalyx* sp. fruit starch under bright field (left) and cross polarized light (right), (g) *Cola nitida* seed starch under bright field (left) and cross polarized light (right), (h) *Coula edulis* seed starch under bright field (left) and cross polarized light (right), (i) *Elaeis guineensis* fruit phytolith, (j) *Elaeis guineensis* leaf phytolith, (k) *Gilbertiodendron splendidum* seed starch under bright field (left) and cross polarized light (right), (m) *Eremospatha macrocarpa* pith starch under bright field (left) and cross polarized light (right), (o) *Panda olesosa* seed starch, (p) *Piper guineense* seed starch under bright field (left) and cross polarized light (right), (right

Of the 24 chimpanzee calculus samples, we found starches in 17 of the samples, and phytoliths in 20 (Fig. 10, Fig. 11; Table 8, Appendix table 10). We also found unidentified phytoliths, unsilicified plant fragments, diatoms, pollen and insects, but these were not identified to taxon (Table 8). In ambiguous cases microremains were classified as possible starches and specifically stated, but were not used for statistical genera identification. Most definite starches and phytoliths that were free from damage (234 starches and 1035 phytoliths) were identified to genus using the random forest model, which assigned each unknown microremain to a genus and provided a certainty score that indicated the confidence with which that assignment was made. A microremain was considered to be damaged if it showed pitting, ruptured surfaces or other major irregularities. The highest certainty score for each individual microremain depended heavily on each genus identification rate (as described above), but generally ranged between 0.25 and 0.95.



Fig. 10: Microremains recovered in calculus samples. Microremains include Unsilicified microremains, starches (definite and possible) and phytoliths recovered with chimpanzee age at death (in years) and approximate age of the cessation of weaning highlighted. a=total counts and b=counts per milligram of calculus. The number of microremains per mg in Ophelia was affected by an unusually small amount of calculus in the sample.



Fig. 11: Microremain assemblages recovered in calculus. (top) Bar chart of the composition of the phytolith assemblage recovered from calculus. (bottom) Bar chart of the composition of the starch assemblage recovered from calculus. The individuals are ordered by age from youngest to oldest.

#### 4.4.2 Assessment of biases in our data

First of all, it was important to ascertain if the treatment of the skeletal material to prevent the spread of disease (including one year of burial, and various chemical treatments) had impacted microremain preservation in the calculus. After 2004 chorine and formalin was used to clean skeletal material. Boiling may have been used on some skeletons to clean them and remove Ebola pathogens between the Autumn of 1994 and the Spring of 1996 (Appendix table 10). To test if the three types of treatments significantly influenced starch preservation we used a Kruskal-Wallis test starch per mg on samples from each period (H=3.7633, df=2, p-value=0.1523). We included microremains classified as possible starches in the starch per mg count (Appendix fig. 1). Due to the small sample size we calculated a Kruskal-Wallis p-value based on 999 random permutations. This indicated no differences between the three groups (Permutation H=7.1215, df=2, p=0.159).

Previous studies of other organic material (bone collagen) in the Taï skeletal collection have indicated no significant post-mortem alteration (Fahy et al., 2013, 2014). While collagen does not necessarily behave in the same manner as plant microremains, it is likely that the comparable hydroxyapatite mineral matrices of bone and calculus have a similar protective effect on the organic materials trapped within them (Nicholson, 1996).

Before comparing the calculus results to the observational records, we wanted to see if there was excessive variation in plant representation among the calculus samples. Phytoliths from four of the five phytolith-producing genera were found on most, but not all, of the calculus samples, suggesting that there is not much variability among these calculus samples (Fig. 11, Appendix table 11). Some genera are found in each sample (Eremospatha and Elaeis) while others, like Sarcophrynium, were rare (Appendix table 12). However, the starch record varies significantly among individuals, with most of the thirteen starch-producing genera seldom found. This probably reflects the far lower numbers of starches compared with phytoliths. Several genera dominate the starch record, namely Gilbertiodendron, Coula, Eremospatha, Treculia and Cola (Fig. 11, Fig. 12). Most microremains were isolated, but three calculus samples had four starch aggregates from Piper; each starch in the aggregate was counted as an individual starch grain and thus constitutes a large proportion of the total number of the recovered starches. This potentially biases the starch assemblage's dietary representativeness (Fig. 11; Appendix table 12). In sum, it seems that there is not much variation in the phytolith record of our chimpanzee samples, but the starch record is less homogeneous. Another potential source of bias comes from the differential preservation of microremains relating to their inherent properties, like size and shape. We noted that our results were biased towards foods with larger sized microremains. Elaeis phytoliths and Cola starches, the largest microremains in the study (Fig. 11, Fig. 12), are disproportionately frequent across the assemblages even after controlling for the high concentration of microremains within these genera. They are frequently found, but are not dominant foods (Appendix fig. 2).



Fig. 12: Plant genera represented by microremain assemblages and Chimpanzees diet. Microremain counts are normalised by dividing counts by the percent content of by dry plant weight of starches and phytoliths among different genera. (a) Phytolith counts compared with feeding records. Outermost ring=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of all 128 chimpanzees, innermost=phytolith counts from the sampled 24 chimpanzees. (b) Starch counts compared with feeding records outermost ring=proportions of minutes spent consuming each genus averaged across the feeding records compared with feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of sampled 24 chimpanzees, middle=proportions of minutes spent consuming each genus averaged across the feeding records of all 128 chimpanzees, innermost=starch counts.

#### 4.4.3 Microremain accumulation, chimpanzee age and sex

If microremains reflect diet they should be more abundant in older chimpanzees, and might vary by sex. We tested this using (see detailed methods below). We ran separate tests for phytoliths, starches and unsilicified remains. For phytoliths, the full model of age and sex significantly influenced the count of phytoliths (x<sup>2</sup>=11.794, df=2, P=0.0003), and the effect of age was also significant by itself (x<sup>2</sup>=12.753, df=1, P=0.0004) (Appendix table 13). Older chimpanzees generally have a higher abundance of phytoliths. However, sex by itself did not explain the abundance of phytoliths we found (x<sup>2</sup>=0.028, df=1, P=0.866). For unsilicified microremains, age and sex as the full model significantly influenced the microremain count (x<sup>2</sup>=10.067, df=1, P=0.015), the effect of age alone was also significant (x<sup>2</sup>=9.202, df=1, P=0.0015), but not sex by itself (x<sup>2</sup>=0.59, df=1, P=0.806). Starch abundance was significantly determined by age and sex together (x<sup>2</sup>=23.994, df=2, p=6.1622e-06). Older chimpanzees generally have a higher abundance of starches (x<sup>2</sup>=3.559, df=1, p=0.0592). Unlike with phytoliths and unsilicified remains, sex strongly influenced the abundance of starch, with female chimpanzees having more starches (x<sup>2</sup>=17.301, df=1, p=3.1897e-05) (Appendix table 13).

#### 4.4.4 Microremain dietary picture and observational feeding records

We predicted that more frequently consumed plants should be highly represented in the chimpanzee calculus. To test this, we used an observational random effect Poisson model (4.3.4). The count of identified classes of microremains (phytoliths and starches) belonging to a particular genus was our response variable, and the fixed predictors were: (a) minutes spent consuming each genus, and (b) chimpanzee age in months. Sex was included as a control predictor, and both calculus sample weight and successful identification rate of each genus were included as weights. We used counts of each genus predicted to be present with the total minutes spent consuming each genus. The chimpanzee individual was included as a random slope term, while year of death, tooth and food type were treated as random intercept terms (see methods below for 7.2).

When comparing the genera proportions present in diet (calculated as the number of minutes spent foraging on a genus) with the recovered phytolith assemblages, we found a clear relationship. The number of minutes spent consuming a given plant genus influences its phytolith count in the calculus assemblage even when accounting for the effects of sex, the tooth we sampled, variation in phytolith production between different plants, and the year the individual died. More specifically, an increased reliance on a genus leads to an increase in its representation in calculus ( $x^2$ =4.048, df=1, P=0.045; Appendix table 13; Fig. 13). The age of the chimpanzees was found to not influence how well it matches group diet ( $x^2$ =0.356, df=1, P=0.55; Appendix table 13).



Fig. 13: Mixed Poisson regression model predicted values. The number of phytoliths from a genus increased as the minutes spent consuming this plant resource increased.

In contrast to phytoliths, there was no significant effect of consumption time on starch numbers ( $x^2$ =1.95, df=2, P=0.376). The number of minutes this group spent eating a specific genus of starchy foods does not influence its frequency in dental calculus. Yet there is an element of uncertainty because starches vary more among individuals than do phytoliths, as described above, and do not seem to be as good a record of dietary behaviour. Fig. 10 and Fig. 11 show the discrepancy between the consistency of starch and phytolith results clearly. These results may be a product of post-mortem diagenesis that influenced our chimpanzee samples, including burial to deflesh the remains (Appendix table 5).

# 4.4.5 Weaning and other behavioural signatures in calculus

The microremains in the Taï calculus record other aspects of their behaviour. First, microremains were strikingly rare in samples from individuals less than 5.3 years old (Fig. 10, Fig. 11; Appendix table 5). The calculus deposits were sparse on these individuals, but despite the small volume of calculus, it was notable that only a single starch and an unsilicified plant fragment were found in these samples. Chimpanzees more than 5.3 years old typically show high numbers of microremains, regardless of the size of the calculus deposit.

Second, the exact plants that were recovered in the calculus provide an interesting view of an important learned behaviour. In our sample, many calculus samples had starches from the *Coula* nut, which is mainly consumed once chimpanzees learn to crack open these nuts. *Coula* nut starches were found in samples from individuals across all age ranges (except those under 5.7 years) (Fig. 14). Although common, *Coula* nut appears to be underrepresented in our sample. It was found only in nine calculus samples, despite this plant being a major food source, comprising of 4.7 % of the total Taï diet.



Fig. 14: Abundance of *Coula* nut starches with chimpanzee age at death. *Coula* nut consumption requires nut cracking and its presence implies nut cracking and tool use or food sharing. The individuals are ordered by age from youngest to oldest.

## 4.5 Discussion

Much of the chimpanzee calculus is relatively rich in plant microremains compared with what has been reported in previous studies of human calculus (Kucera et al., 2011; Mickleburgh and Pagán-Jiménez, 2012; Leonard et al., 2015). This is not entirely unexpected for several reasons. First, our samples are modern, and post-mortem microremain diagenesis is therefore less acute than in ancient remains. Secondly, Taï Chimpanzee diet is uncooked, plant-dominated and voluminous (Appendix fig. 1; Appendix fig. 2). Thirdly, and in contrast to humans, chimpanzees consume a large amount of phytolith-rich material. This richness in microremains is largely confined to phytoliths. Starch abundance falls within ranges observed elsewhere (Mickleburgh and Pagán-Jiménez, 2012; Power et al., 2014b; Leonard et al., 2015). It is evident that starches are underrepresented and in some samples are even totally absent. In addition, phytoliths present a far more uniform picture of diet between different chimpanzees. This may be due to diagenesis occurring during the preparation of the skeletons for the osteology collection. It is known that all skeletons were buried for short periods of time during the defleshing process (Appendix table 5). These processes may preferentially alter or remove starches from the calculus record that are not sufficiently mineralised and sealed, while leaving the phytoliths relatively unaltered. Yet, our Kruskal-Wallis test indicates cleaning processes have not influenced starch numbers.

The comparison of microremains with diet was possible because our methodology generated a deep profile of dietary history. However, our metric of food use (minutes spent consuming a food) is not identical to the total volume of consumed food by each chimpanzee, or a food's total energetic value. Unfortunately, our analysis was not able to account for this.

Additionally, we found that the microremain record was likely biased by the differential survivability of microremain from different plants. The plants with the largest starches and phytoliths were overrepresented in our sample, possibly due to the larger surface area. This ties in with research that shows that phytolith and starch morphology and surface area is linked to long term stability (Haslam, 2004; Cabanes et al., 2011). Larger blocky microremains may be preferentially preserved. This is noteworthy given that their larger surface area would enhance their contact with bacterial and chemical process and alteration.

Overall, our results verify that the calculus record can be accumulative by showing that older individuals present more microremains. Sex may be a factor to take into consideration, and seems to influence the accumulation of starches but not phytoliths or unsilicified remains. It may reflect higher consumption of starches by female chimpanzees, or sex differences in amylase production or calculus formation, as has been suggested for humans (Monteiro da Silva et al., 1998). We do not currently have the ability to distinguish among these possibilities. The increase in microremains with age and possibly sex implies that microremain accumulation is bound up in aspects of diet that regulate calculus formation. Thus, microremain presence and proportions are likely effected or confounded by all the factors that influence plaque and calculus such as the intake of protein, smoking, polysilicic acid and silica (Damen and Ten Cate, 1989; Roberts-Harry and Clerehugh, 2000; Jin and Yip, 2002). Calculus clearly can approach a long-term dietary signal, although the timespan involved is not yet clarified.

Our results strongly suggest that care must be taken when interpreting the microremains record preserved in dental calculus, particularly the starch grain record. However, our results also indicate that microremains in calculus can be used to recover important information about diet, behaviour and life history. For example, we observed a lack of microremains from individuals sampled from deciduous teeth of chimpanzees less than 5.3 years old. The microremain assemblages could indicate a rapid accumulation of microremains as solid food enters the diet (Fig. 10). This pattern matches what is generally reported for age of weaning using other measures. Much information on the age of chimpanzee weaning is estimated from inter-birth interval (Fahy et al., 2014). Inter-Birth Interval estimates of weaning ages vary from 4.5 years at Gombe to 5.75 years at Taï (Boesch, 1997), to 6 years at Mahale (Nishida and Hasegawa, 1992). Yet inter-Birth Interval is an indirect measure as it includes more than simply suckling duration. Isotopic based data indicates weaning at Taï commences at 2 years and ends at 3-4.5 years varying by factors such as sex of the offspring. If we combine this infant microremain signal together with the verification of the accumulative nature of the microremain assemblages, we can conclude calculus reflects information on the weaning transition that may be useful for studying unhabituated populations. Researchers should expand this research to infants from recent foragers and horticulturists to develop its applicability.

Furthermore, though the starch dietary record appears more stochastic than that of phytoliths, starches can still provide useful information about behaviour. Many of our starches come from the *Coula* nut (Fig. 11, Fig. 14). Among chimpanzees, *Coula* consumption requires a learned behaviour: nut cracking with a hammer and anvil. This behaviour is restricted to a limited area of the chimpanzee range in West Africa (Boesch et al., 1994). The presence of *Coula* starches (Fig. 11, Fig. 14) shows calculus can reveal nut-cracking behaviour in a group. The fact that tooluse in a group is discernible is relevant for dental calculus studies in both primatology and hominin evolution. The use of *Coula* nut is influenced by age and sex differences in nut cracking (Boesch and Boesch, 1984; Boesch and Boesch-Achermann, 2000), and, as expected, *Coula* starches are absent in youngest chimpanzees who are not yet weaned. Even after weaning infant nut consumption is low and is derived from nuts cracked by the mother as it takes years to learn how to crack nuts (Boesch and Boesch-Achermann, 2000). Beyond this, we do not have enough calculus samples to examine if there are sex or age differences in the calculus record of nut cracking.

This profile of Taï Chimpanzee diet reflects high amount of contextual information available on a single population. Much research utilizing dental calculus is interested in reconstructing diet of collections of individuals and not contemporaneous populations from archaeological sites. Researchers using archaeological populations with small and even large samples must be aware of the unlikeness of being able to capture the full dietary breadth with dental calculus. The best strategy to account for this issue is to maximize the number of samples in a study.

In summary, the study verifies the relevance of dental calculus for investigations on diet, food acquisition behaviour and life history. It is the first to link dental calculus with foods that entered the oral cavity in quantified abundances but it also identifies clear weaknesses of this method. The data also provide valuable information on the commencement of plant food consumption in wild chimpanzees, and confirms the consumption of solid foodstuffs from at least 5.3 years in life. Our study suggests that calculus analysis provides a rich but wavering insight into complex dietary structure, and that phytoliths, when present in calculus and in diet, may provide a more reliable record of diet than starches.