



**Universiteit  
Leiden**  
The Netherlands

## **Immune modulation by mannosylated peptides**

Kel, J.M.

### **Citation**


Kel, J. M. (2008, April 2). *Immune modulation by mannosylated peptides*. Retrieved from <https://hdl.handle.net/1887/12665>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded  
from: <https://hdl.handle.net/1887/12665>

**Note:** To cite this publication please use the final published version (if applicable).



Immune modulation by mannosylated peptides

# Immune modulation by mannosylated peptides

Junda M. Kel

Junda M. Kel

Figure 4 of Chapter 7 (page 137)

M-PLP<sub>139-151</sub>-treatment is associated with reduced CNS inflammation. Recipient mice were treated with PBS (left panels), PLP<sub>139-151</sub> (middle panels) or M-PLP<sub>139-151</sub> (right panels) on days -1, 1 and 3 and sacrificed on day 16. Step serial, longitudinal sections from the cervical to lumbar part of the spinal cord were collected for immunohistochemistry. Antibodies specific for CD3, IFN-γ, CD11b, MHC II and IL-23 were employed. The presence of infiltrates was not limited to particular spinal cord regions. Instead, infiltrates were scattered over the whole tissue, predominantly located in white matter tissue. Representative regions are depicted. Magnification: 200x