



Universiteit
Leiden

The Netherlands

Glycomics based biomarkers of the rate of aging : development and applications of high-throughput N-glycan analysis

Ruhaak, L.R.

Citation

Ruhaak, L. R. (2011, March 24). *Glycomics based biomarkers of the rate of aging : development and applications of high-throughput N-glycan analysis*. Retrieved from <https://hdl.handle.net/1887/16559>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/16559>

Note: To cite this publication please use the final published version (if applicable).

References

- [1] Shental-Bechor, D & Levy, Y (2009) Folding of glycoproteins: toward understanding the biophysics of the glycosylation code. *Curr. Opin. Struct. Biol.*, 19, 524-533.
- [2] Fukuda, MN, Sasaki, H, Lopez, L, & Fukuda, M (1989) Survival of recombinant erythropoietin in the circulation: the role of carbohydrates. *Blood*, 73, 84-89.
- [3] Gu, J, Sato, Y, Kariya, Y, Isaji, T, Taniguchi, N, & Fukuda, T (2009) A mutual regulation between cell-cell adhesion and N-glycosylation: implication of the bisecting GlcNAc for biological functions. *J. Proteome Res.*, 8, 431-435.
- [4] Takahashi, M, Kuroki, Y, Ohtsubo, K, & Taniguchi, N (2009) Core fucose and bisecting GlcNAc, the direct modifiers of the N-glycan core: their functions and target proteins. *Carbohydr. Res.*, 344, 1387-1390.
- [5] Marth, JD & Grewal, PK (2008) Mammalian glycosylation in immunity. *Nat. Rev. Immunol.*, 8, 874-887.
- [6] Ohtsubo, K & Marth, JD (2006) Glycosylation in cellular mechanisms of health and disease. *Cell*, 126, 855-867.
- [7] Arnold, JN, Saldo, R, Hamid, UM, & Rudd, PM (2008) Evaluation of the serum N-linked glycome for the diagnosis of cancer and chronic inflammation. *Proteomics*, 8, 3284-3293.
- [8] Gornik, O, Wagner, J, Pucic, M, Knezevic, A, Redzic, I, & Lauc, G (2009) Stability of N-glycan profiles in human plasma. *Glycobiology*, 19, 1547-1553.
- [9] Packer, NH, von der Lieth, CW, Aoki-Kinoshita, KF, Lebrilla, CB, Paulson, JC, Raman, R, Rudd, P, Sasisekharan, R, Taniguchi, N, & York, WS (2008) Frontiers in glycomics: bioinformatics and biomarkers in disease. An NIH white paper prepared from discussions by the focus groups at a workshop on the NIH campus, Bethesda MD (September 11-13, 2006). *Proteomics*, 8, 8-20.
- [10] Varki, A., Cummings, R.D., Esko, J.D., Freeze, H.H., Stanley, P., Bertozzi, C.R.,

- Hart,G.W., & Etzler,M.E. (2008) Essentials of glycobiology, 2 edn. Cold Spring Harbor Laboratory Press.
- [11] Kornfeld, R & Kornfeld, S (1985) Assembly of asparagine-linked oligosaccharides. *Annu. Rev. Biochem.*, 54, 631-664.
- [12] Burda, P & Aebi, M (1999) The dolichol pathway of N-linked glycosylation. *Biochim. Biophys. Acta*, 1426, 239-257.
- [13] Lowe, JB & Marth, JD (2003) A genetic approach to Mammalian glycan function. *Annu. Rev. Biochem.*, 72, 643-691.
- [14] Adachi, J, Kumar, C, Zhang, Y, Olsen, JV, & Mann, M (2006) The human urinary proteome contains more than 1500 proteins, including a large proportion of membrane proteins. *Genome Biol.*, 7, R80.
- [15] Jacobs, JM, Adkins, JN, Qian, WJ, Liu, T, Shen, Y, Camp, DG, & Smith, RD (2005) Utilizing human blood plasma for proteomic biomarker discovery. *J. Proteome Res.*, 4, 1073-1085.
- [16] Anderson, NL & Anderson, NG (2002) The human plasma proteome: history, character, and diagnostic prospects. *Mol. Cell. Proteomics*, 1, 845-867.
- [17] An, HJ, Ninonuevo, M, Aguilan, J, Liu, H, Lebrilla, CB, Alvarenga, LS, & Mannis, MJ (2005) Glycomics analyses of tear fluid for the diagnostic detection of ocular rosacea. *J. Proteome Res.*, 4, 1981-1987.
- [18] Botella-Lopez, A, Burgaya, F, Gavin, R, Garcia-Ayllon, MS, Gomez-Tortosa, E, Pena-Casanova, J, Urena, JM, Del Rio, JA, Blesa, R, Soriano, E, & Saez-Valero, J (2006) Reelin expression and glycosylation patterns are altered in Alzheimer's disease. *Proc. Natl. Acad. Sci. U. S. A.*, 103, 5573-5578.
- [19] Blomme, B, Van Steenkiste, C, Callewaert, N, & Van Vlierberghe, H (2009) Alteration of protein glycosylation in liver diseases. *J. Hepatol.*, 50, 592-603.
- [20] Vanhooren, V, Desmyter, L, Liu, XE, Cardelli, M, Franceschi, C, Federico, A, Libert, C, Laroy, W, Dewaele, S, Contreras, R, & Chen, C (2007) N-glycomic changes in serum proteins during human aging. *Rejuvenation Res.*, 10, 521-531a.
- [21] Knezevic, A, Polasek, O, Gornik, O, Rudan, I, Campbell, H, Hayward, C, Wright, A, Kolcic, I, O'Donoghue, N, Bones, J, Rudd, PM, & Lauc, G (2009)

- Variability, Heritability and Environmental Determinants of Human Plasma N-Glycome. *J. Proteome Res.*, 8, 694-701.
- [22] Lauc, G, Rudan, I, Campbell, H, & Rudd, PM (2010) Complex genetic regulation of protein glycosylation. *Mol. Biosyst.*, 6, 329-335.
- [23] Lauc, G, Huffman, J, Hayward, C, Knezevic, A, Polasek, O, Gornik, O, Vitar, V, Kolcic, I, Biloglav, Z, Zgaga, L, Hastie, ND, Wright, AF, Campbell, H, Rudd, PM, & Rudan, I (2009) Genome-wide association study identifies FUT8 and ESR2 as co-regulators of a biantennary N-linked glycan A2 (GlcNAc2Man3GlcNAc2) in human plasma proteins. *Nature precedings*, <http://hdl.handle.net/10101/npre.2009.2864.1>.
- [24] Knezevic, A, Gornik, O, Polasek, O, Pucic, M, Redzic, I, Novokmet, M, Rudd, PM, Wright, AF, Campbell, H, Rudan, I, & Lauc, G (2010) Effects of aging, body mass index, plasma lipid profiles, and smoking on human plasma N-glycans. *Glycobiology*, in press.
- [25] Itoh, N, Sakaue, S, Nakagawa, H, Kuroguchi, M, Ohira, H, Deguchi, K, Nishimura, S, & Nishimura, M (2007) Analysis of N-glycan in serum glycoproteins from db/db mice and humans with type 2 diabetes. *Am. J. Physiol. Endocrinol. Metab.*, 293, E1069-E1077.
- [26] An, HJ, Miyamoto, S, Lancaster, KS, Kirmiz, C, Li, B, Lam, KS, Leiserowitz, GS, & Lebrilla, CB (2006) Profiling of glycans in serum for the discovery of potential biomarkers for ovarian cancer. *J. Proteome Res.*, 5, 1626-1635.
- [27] Barrabes, S, Pages-Pons, L, Radcliffe, CM, Tabares, G, Fort, E, Royle, L, Harvey, DJ, Moenner, M, Dwek, RA, Rudd, PM, De Llorens, R, & Peracaula, R (2007) Glycosylation of serum ribonuclease 1 indicates a major endothelial origin and reveals an increase in core fucosylation in pancreatic cancer. *Glycobiology*, 17, 388-400.
- [28] Callewaert, N, Van, VH, Van, HA, Laroy, W, Delanghe, J, & Contreras, R (2004) Noninvasive diagnosis of liver cirrhosis using DNA sequencer-based total serum protein glycomics. *Nat. Med.*, 10, 429-434.
- [29] de Leoz, ML, An, HJ, Kronewitter, S, Kim, J, Beecroft, S, Vinall, R, Miyamoto, S, de Vere, WR, Lam, KS, & Lebrilla, C (2008) Glycomic approach for poten-

- tial biomarkers on prostate cancer: profiling of N-linked glycans in human sera and pRNS cell lines. *Dis. Markers*, 25, 243-258.
- [30] Kirmiz, C, Li, B, An, HJ, Clowers, BH, Chew, HK, Lam, KS, Ferrige, A, Alecio, R, Borowsky, AD, Sulaimon, S, Lebrilla, CB, & Miyamoto, S (2007) A serum glycomics approach to breast cancer biomarkers. *Mol. Cell. Proteomics*, 6, 43-55.
- [31] Okuyama, N, Ide, Y, Nakano, M, Nakagawa, T, Yamanaka, K, Moriwaki, K, Murata, K, Ohigashi, H, Yokoyama, S, Eguchi, H, Ishikawa, O, Ito, T, Kato, M, Kasahara, A, Kawano, S, Gu, J, Taniguchi, N, & Miyoshi, E (2006) Fucosylated haptoglobin is a novel marker for pancreatic cancer: a detailed analysis of the oligosaccharide structure and a possible mechanism for fucosylation. *Int. J. Cancer*, 118, 2803-2808.
- [32] Peracaula, R, Tabares, G, Royle, L, Harvey, DJ, Dwek, RA, Rudd, PM, & De Llorens, R (2003) Altered glycosylation pattern allows the distinction between prostate-specific antigen (PSA) from normal and tumor origins. *Glycobiology*, 13, 457-470.
- [33] Saldo, R, Royle, L, Radcliffe, CM, Abd Hamid, UM, Evans, R, Arnold, JN, Banks, RE, Hutson, R, Harvey, DJ, Antrobus, R, Petrescu, SM, Dwek, RA, & Rudd, PM (2007) Ovarian cancer is associated with changes in glycosylation in both acute-phase proteins and IgG. *Glycobiology*, 17, 1344-1356.
- [34] Fujimura, T, Shinohara, Y, Tissot, B, Pang, PC, Kurogochi, M, Saito, S, Arai, Y, Sadilek, M, Murayama, K, Dell, A, Nishimura, S, & Hakomori, SI (2008) Glycosylation status of haptoglobin in sera of patients with prostate cancer vs. benign prostate disease or normal subjects. *Int. J. Cancer*, 122, 39-49.
- [35] Hoagland, LF, Campa, MJ, Gottlin, EB, Herndon, JE, & Patz, EF, Jr. (2007) Haptoglobin and posttranslational glycan-modified derivatives as serum biomarkers for the diagnosis of nonsmall cell lung cancer. *Cancer*, 110, 2260-2268.
- [36] Imre, T, Kremmer, T, Heberger, K, Molnar-Szollosi, E, Ludanyi, K, Pocsfalvi, G, Malorni, A, Drahos, L, & Vekey, K (2008) Mass spectrometric and linear discriminant analysis of N-glycans of human serum alpha-1-acid glycoprotein

- in cancer patients and healthy individuals. *J. Proteomics*, 71, 186-197.
- [37] Kanoh, Y, Mashiko, T, Danbara, M, Takayama, Y, Ohtani, S, Imasaki, T, Abe, T, & Akahoshi, T (2004) Analysis of the oligosaccharide chain of human serum immunoglobulin g in patients with localized or metastatic cancer. *Oncology*, 66, 365-370.
- [38] Kyselova, Z, Mechref, Y, Kang, P, Goetz, JA, Dobrolecki, LE, Sledge, GW, Schnaper, L, Hickey, RJ, Malkas, LH, & Novotny, MV (2008) Breast cancer diagnosis and prognosis through quantitative measurements of serum glycan profiles. *Clin. Chem.*, 54, 1166-1175.
- [39] Huhn, C, Selman, MH, Ruhaak, LR, Deelder, AM, & Wuhrer, M (2009) IgG glycosylation analysis. *Proteomics*, 9, 882-913.
- [40] Parekh, R, Roitt, I, Isenberg, D, Dwek, R, & Rademacher, T (1988) Age-related galactosylation of the N-linked oligosaccharides of human serum IgG. *J. Exp. Med.*, 167, 1731-1736.
- [41] Yamada, E, Tsukamoto, Y, Sasaki, R, Yagyu, K, & Takahashi, N (1997) Structural changes of immunoglobulin G oligosaccharides with age in healthy human serum. *Glycoconj. J.*, 14, 401-405.
- [42] Shikata, K, Yasuda, T, Takeuchi, F, Konishi, T, Nakata, M, & Mizuochi, T (1998) Structural changes in the oligosaccharide moiety of human IgG with aging. *Glycoconj. J.*, 15, 683-689.
- [43] Parekh, RB, Dwek, RA, Sutton, BJ, Fernandes, DL, Leung, A, Stanworth, D, Rademacher, TW, Mizuochi, T, Taniguchi, T, Matsuta, K, & . (1985) Association of rheumatoid arthritis and primary osteoarthritis with changes in the glycosylation pattern of total serum IgG. *Nature*, 316, 452-457.
- [44] Parekh, RB, Roitt, IM, Isenberg, DA, Dwek, RA, Ansell, BM, & Rademacher, TW (1988) Galactosylation of IgG associated oligosaccharides: reduction in patients with adult and juvenile onset rheumatoid arthritis and relation to disease activity. *Lancet*, 1, 966-969.
- [45] van Zeben D., Rook, GA, Hazes, JM, Zwinderman, AH, Zhang, Y, Ghelani, S, Rademacher, TW, & Breedveld, FC (1994) Early agalactosylation of IgG is associated with a more progressive disease course in patients with rheumatoid

- arthritis: results of a follow-up study. *Br. J. Rheumatol.*, 33, 36-43.
- [46] Nakagawa, H, Hato, M, Takegawa, Y, Deguchi, K, Ito, H, Takahata, M, Iwasaki, N, Minami, A, & Nishimura, S (2007) Detection of altered N-glycan profiles in whole serum from rheumatoid arthritis patients. *J. Chromatogr. B*, 853, 133-137.
- [47] Mehta, AS, Long, RE, Comunale, MA, Wang, M, Rodemich, L, Krakover, J, Philip, R, Marrero, JA, Dwek, RA, & Block, TM (2008) Increased levels of galactose-deficient anti-Gal immunoglobulin G in the sera of hepatitis C virus-infected individuals with fibrosis and cirrhosis. *J. Virol.*, 82, 1259-1270.
- [48] Poland, DC, Schalkwijk, CG, Stehouwer, CD, Koeleman, CA, van het, HB, & van Dijk, W (2001) Increased alpha3-fucosylation of alpha1-acid glycoprotein in Type I diabetic patients is related to vascular function. *Glycoconj. J.*, 18, 261-268.
- [49] Higai, K, Azuma, Y, Aoki, Y, & Matsumoto, K (2003) Altered glycosylation of alpha1-acid glycoprotein in patients with inflammation and diabetes mellitus. *Clin. Chim. Acta*, 329, 117-125.
- [50] Yoo, EM & Morrison, SL (2005) IgA: an immune glycoprotein. *Clin. Immunol.*, 116, 3-10.
- [51] Karnaukhova, E, Ophir, Y, & Golding, B (2006) Recombinant human alpha-1 proteinase inhibitor: towards therapeutic use. *Amino Acids*, 30, 317-332.
- [52] Wuhrer, M, Stam, JC, van de Geijn, FE, Koeleman, CA, Verrips, CT, Dolhain, RJ, Hokke, CH, & Deelder, AM (2007) Glycosylation profiling of immunoglobulin G (IgG) subclasses from human serum. *Proteomics*, 7, 4070-4081.
- [53] Wada, Y, Tajiri, M, & Yoshida, S (2004) Hydrophilic affinity isolation and MALDI multiple-stage tandem mass spectrometry of glycopeptides for glycoproteomics. *Anal. Chem.*, 76, 6560-6565.
- [54] Lim, A, Reed-Bogan, A, & Harmon, BJ (2008) Glycosylation profiling of a therapeutic recombinant monoclonal antibody with two N-linked glycosylation sites using liquid chromatography coupled to a hybrid quadrupole time-of-flight mass spectrometer. *Anal. Biochem.*, 375, 163-172.
- [55] Zhang, Y, Go, EP, & Desaire, H (2008) Maximizing coverage of glycosylation

- heterogeneity in MALDI-MS analysis of glycoproteins with up to 27 glycosylation sites. *Anal. Chem.*, 80, 3144-3158.
- [56] Yu, YQ, Fournier, J, Gilar, M, & Gebler, JC (2007) Identification of N-linked glycosylation sites using glycoprotein digestion with pronase prior to MALDI tandem time-of-flight mass spectrometry. *Anal. Chem.*, 79, 1731-1738.
- [57] Schahs, M, Strasser, R, Stadlmann, J, Kunert, R, Rademacher, T, & Steinkellner, H (2007) Production of a monoclonal antibody in plants with a humanized N-glycosylation pattern. *Plant Biotechnol. J.*, 5, 657-663.
- [58] Selman, MH, McDonnell, LA, Palmblad, M, Ruhaak, LR, Deelder, AM, & Wuhrer, M (2010) Immunoglobulin G glycopeptide profiling by matrix-assisted laser desorption ionization Fourier transform ion cyclotron resonance mass spectrometry. *Anal. Chem.*, 82, 1073-1081.
- [59] Qian, J, Liu, T, Yang, L, Daus, A, Crowley, R, & Zhou, Q (2007) Structural characterization of N-linked oligosaccharides on monoclonal antibody cetuximab by the combination of orthogonal matrix-assisted laser desorption/ionization hybrid quadrupole-quadrupole time-of-flight tandem mass spectrometry and sequential enzymatic digestion. *Anal. Biochem.*, 364, 8-18.
- [60] Holland, M, Yagi, H, Takahashi, N, Kato, K, Savage, CO, Goodall, DM, & Jefferis, R (2006) Differential glycosylation of polyclonal IgG, IgG-Fc and IgG-Fab isolated from the sera of patients with ANCA-associated systemic vasculitis. *Biochim. Biophys. Acta*, 1760, 669-677.
- [61] Omtvedt, LA, Royle, L, Husby, G, Sletten, K, Radcliffe, CM, Harvey, DJ, Dwek, RA, & Rudd, PM (2006) Glycan analysis of monoclonal antibodies secreted in deposition disorders indicates that subsets of plasma cells differentially process IgG glycans. *Arthritis Rheum.*, 54, 3433-3440.
- [62] Youngs, A, Chang, SC, Dwek, RA, & Scragg, IG (1996) Site-specific glycosylation of human immunoglobulin G is altered in four rheumatoid arthritis patients. *Biochem. J.*, 314 (Pt 2), 621-630.
- [63] Wada, Y, Azadi, P, Costello, CE, Dell, A, Dwek, RA, Geyer, H, Geyer, R, Kakehi, K, Karlsson, NG, Kato, K, Kawasaki, N, Khoo, KH, Kim, S, Kondo, A, Lattova, E, Mechref, Y, Miyoshi, E, Nakamura, K, Narimatsu, H, Novotny, MV,

- Packer, NH, Perreault, H, Peter-Katalinic, J, Pohlentz, G, Reinhold, VN, Rudd, PM, Suzuki, A, & Taniguchi, N (2007) Comparison of the methods for profiling glycoprotein glycans--HUPO Human Disease Glycomics/Proteome Initiative multi-institutional study. *Glycobiology*, 17, 411-422.
- [64] Stadlmann, J, Pabst, M, Kolarich, D, Kunert, R, & Altmann, F (2008) Analysis of immunoglobulin glycosylation by LC-ESI-MS of glycopeptides and oligosaccharides. *Proteomics.*, 8, 2858-2871.
- [65] Wagner-Rousset, E, Bednarczyk, A, Bussat, MC, Colas, O, Corvaia, N, Schaeffer, C, Van, DA, & Beck, A (2008) The way forward, enhanced characterization of therapeutic antibody glycosylation: comparison of three level mass spectrometry-based strategies. *J. Chromatogr. B*, 872, 23-37.
- [66] Takegawa, Y, Deguchi, K, Ito, H, Keira, T, Nakagawa, H, & Nishimura, S (2006) Simple separation of isomeric sialylated N-glycopeptides by a zwitterionic type of hydrophilic interaction chromatography. *J. Sep. Sci.*, 29, 2533-2540.
- [67] Harazono, A, Kawasaki, N, Itoh, S, Hashii, N, Matsuishi-Nakajima, Y, Kawaniishi, T, & Yamaguchi, T (2008) Simultaneous glycosylation analysis of human serum glycoproteins by high-performance liquid chromatography/tandem mass spectrometry. *J. Chromatogr. B*, 869, 20-30.
- [68] Ito, H, Takegawa, Y, Deguchi, K, Nagai, S, Nakagawa, H, Shinohara, Y, & Nishimura, S (2006) Direct structural assignment of neutral and sialylated N-glycans of glycopeptides using collision-induced dissociation MSn spectral matching. *Rapid Commun. Mass Spectrom.*, 20, 3557-3565.
- [69] Wuhler, M, Koeleman, CA, Hokke, CH, & Deelder, AM (2005) Protein glycosylation analyzed by normal-phase nano-liquid chromatography--mass spectrometry of glycopeptides. *Anal. Chem.*, 77, 886-894.
- [70] Geyer, H & Geyer, R (2006) Strategies for analysis of glycoprotein glycosylation. *Biochim. Biophys. Acta*, 1764, 1853-1869.
- [71] Ruhaak, LR, Zauner, G, Huhn, C, Bruggink, C, Deelder, AM, & Wuhler, M (2010) Glycan labeling strategies and their use in identification and quantification. *Anal. Bioanal. Chem.*, in press.

- [72] Huang, Y, Mechref, Y, & Novotny, MV (2001) Microscale nonreductive release of O-linked glycans for subsequent analysis through MALDI mass spectrometry and capillary electrophoresis. *Anal. Chem.*, 73, 6063-6069.
- [73] Duk, M, Ugorski, M, & Lisowska, E (1997) beta-Elimination of O-glycans from glycoproteins transferred to immobilon P membranes: method and some applications. *Anal. Biochem.*, 253, 98-102.
- [74] Karlsson, NG & Packer, NH (2002) Analysis of O-linked reducing oligosaccharides released by an in-line flow system. *Anal. Biochem.*, 305, 173-185.
- [75] Morelle, W & Michalski, JC (2007) Analysis of protein glycosylation by mass spectrometry. *Nat. Protoc.*, 2, 1585-1602.
- [76] Harvey, DJ (2003) Matrix-assisted laser desorption/ionization mass spectrometry of carbohydrates and glycoconjugates. *Int. J. Mass Spectrom.*, 226, 1-35.
- [77] Merry, AH, Neville, DC, Royle, L, Matthews, B, Harvey, DJ, Dwek, RA, & Rudd, PM (2002) Recovery of intact 2-aminobenzamide-labeled O-glycans released from glycoproteins by hydrazinolysis. *Anal. Biochem.*, 304, 91-99.
- [78] Nakakita, S, Sumiyoshi, W, Miyanishi, N, & Hirabayashi, J (2007) A practical approach to N-glycan production by hydrazinolysis using hydrazine monohydrate. *Biochem. Biophys. Res. Commun.*, 362, 639-645.
- [79] Deguchi, K, Takegawa, Y, Ito, H, Miura, N, Yoshioka, S, Nagai, S, Nakagawa, H, & Nishimura, S (2006) Structural assignment of isomeric 2-aminopyridine-derivatized monosialylated biantennary N-linked oligosaccharides using negative-ion multistage tandem mass spectral matching. *Rapid Commun. Mass Spectrom.*, 20, 412-418.
- [80] Royle, L, Campbell, MP, Radcliffe, CM, White, DM, Harvey, DJ, Abrahams, JL, Kim, YG, Henry, GW, Shadick, NA, Weinblatt, ME, Lee, DM, Rudd, PM, & Dwek, RA (2008) HPLC-based analysis of serum N-glycans on a 96-well plate platform with dedicated database software. *Anal. Biochem.*, 376, 1-12.
- [81] Wuhler, M & Deelder, AM (2005) Negative-mode MALDI-TOF/TOF-MS of oligosaccharides labeled with 2-aminobenzamide. *Anal. Chem.*, 77, 6954-6959.
- [82] Kang, P, Mechref, Y, Klouckova, I, & Novotny, MV (2005) Solid-phase perme-

- thylation of glycans for mass spectrometric analysis. *Rapid Commun. Mass Spectrom.*, 19, 3421-3428.
- [83] Costello, CE, Contado-Miller, JM, & Cipollo, JF (2007) A glycomics platform for the analysis of permethylated oligosaccharide alditols. *J. Am. Soc. Mass Spectrom.*, 18, 1799-1812.
- [84] Morelle, W, Flahaut, C, Michalski, JC, Louvet, A, Mathurin, P, & Klein, A (2006) Mass spectrometric approach for screening modifications of total serum N-glycome in human diseases: application to cirrhosis. *Glycobiology*, 16, 281-293.
- [85] You, J, Sheng, X, Ding, C, Sun, Z, Suo, Y, Wang, H, & Li, Y (2008) Detection of carbohydrates using new labeling reagent 1-(2-naphthyl)-3-methyl-5-pyrazolone by capillary zone electrophoresis with absorbance (UV). *Anal. Chim. Acta*, 609, 66-75.
- [86] Lattova, E & Perreault, H (2003) Labelling saccharides with phenylhydrazine for electrospray and matrix-assisted laser desorption-ionization mass spectrometry. *J. Chromatogr. B*, 793, 167-179.
- [87] Bigge, JC, Patel, TP, Bruce, JA, Goulding, PN, Charles, SM, & Parekh, RB (1995) Nonselective and efficient fluorescent labeling of glycans using 2-amino benzamide and anthranilic acid. *Anal. Biochem.*, 230, 229-238.
- [88] Evangelista, RA, Guttman, A, & Chen, FT (1996) Acid-catalyzed reductive amination of aldoses with 8-aminopyrene-1,3,6-trisulfonate. *Electrophoresis*, 17, 347-351.
- [89] Anumula, KR (1994) Quantitative determination of monosaccharides in glycoproteins by high-performance liquid chromatography with highly sensitive fluorescence detection. *Anal. Biochem.*, 220, 275-283.
- [90] Anumula, KR (2006) Advances in fluorescence derivatization methods for high-performance liquid chromatographic analysis of glycoprotein carbohydrates. *Anal. Biochem.*, 350, 1-23.
- [91] Shilova, NV & Bovin, NV (2003) Fluorescent labels for analysis of mono- and oligosaccharides. *Russ. J. Bioorg. Chem.*, 29, 309-324.
- [92] Deguchi, K, Keira, T, Yamada, K, Ito, H, Takegawa, Y, Nakagawa, H, & Ni-

- shimura, S (2008) Two-dimensional hydrophilic interaction chromatography coupling anion-exchange and hydrophilic interaction columns for separation of 2-pyridylamino derivatives of neutral and sialylated N-glycans. *J. Chromatogr. A*, 1189, 169-174.
- [93] Takahashi, N., Yagi, H., & Kato, K. (2008) Release of N-glycans by Enzymatic Methods. In *Experimental glycoscience* (Taniguchi, N., Suzuki, A., Ito, H., Narimatsu, H., Kawasaki, N., & Hase, S., eds), pp. 7-11. Springer Tokyo.
- [94] Anumula, KR & Dhume, ST (1998) High resolution and high sensitivity methods for oligosaccharide mapping and characterization by normal phase high performance liquid chromatography following derivatization with highly fluorescent anthranilic acid. *Glycobiology*, 8, 685-694.
- [95] Yu, YQ, Gilar, M, Kaska, J, & Gebler, JC (2005) A rapid sample preparation method for mass spectrometric characterization of N-linked glycans. *Rapid Commun. Mass Spectrom.*, 19, 2331-2336.
- [96] Kamoda, S, Ishikawa, R, & Takechi, K (2006) Capillary electrophoresis with laser-induced fluorescence detection for detailed studies on N-linked oligosaccharide profile of therapeutic recombinant monoclonal antibodies. *J. Chromatogr. A*, 1133, 332-339.
- [97] Ruhaak, LR, Huhn, C, Waterreus, WJ, de Boer, AR, Neuss, C, Hokke, CH, Deelder, AM, & Wuhrer, M (2008) Hydrophilic interaction chromatography-based high-throughput sample preparation method for N-glycan analysis from total human plasma glycoproteins. *Anal. Chem.*, 80, 6119-6126.
- [98] Callewaert, N, Geysens, S, Molemans, F, & Contreras, R (2001) Ultrasensitive profiling and sequencing of N-linked oligosaccharides using standard DNA-sequencing equipment. *Glycobiology*, 11, 275-281.
- [99] Laroy, W, Contreras, R, & Callewaert, N (2006) Glycome mapping on DNA sequencing equipment. *Nat. Protoc.*, 1, 397-405.
- [100] Suzuki, H, Muller, O, Guttman, A, & Karger, BL (1997) Analysis of 1-aminopyrene-3,6,8-trisulfonate-derivatized oligosaccharides by capillary electrophoresis with matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Anal. Chem.*, 69, 4554-4559.

- [101] Dalpathado, DS, Jiang, H, Kater, MA, & Desaire, H (2005) Reductive amination of carbohydrates using NaBH(OAc)₃. *Anal. Bioanal. Chem.*, 381, 1130-1137.
- [102] Locke, D, Bevans, CG, Wang, LX, Zhang, Y, Harris, AL, & Lee, YC (2004) Neutral, acidic, and basic derivatives of anthranilamide that confer different formal charge to reducing oligosaccharides. *Carbohydr. Res.*, 339, 221-231.
- [103] Ruhaak, LR, Steenvoorden, E, Koeleman, CAM, Deelder, AM, & Wuhrer, M (2010) 2-Picoline-borane: a non-toxic reducing agent for oligosaccharide labeling by reductive amination. *Proteomics*, 10, 2330-2336.
- [104] Evangelista, RA, Chen, FT, & Guttman, A (2006) Reductive amination of N-linked oligosaccharides using organic acid catalysts. *J. Chromatogr. A*, 745, 273-280.
- [105] Chen, FT, Dobashi, TS, & Evangelista, RA (1998) Quantitative analysis of sugar constituents of glycoproteins by capillary electrophoresis. *Glycobiology*, 8, 1045-1052.
- [106] Anumula, KR (2000) High-sensitivity and high-resolution methods for glycoprotein analysis. *Anal. Biochem.*, 283, 17-26.
- [107] Pabst, M, Kolarich, D, Poltl, G, Dalik, T, Lubec, G, Hofinger, A, & Altmann, F (2009) Comparison of fluorescent labels for oligosaccharides and introduction of a new postlabeling purification method. *Anal. Biochem.*, 384, 263-273.
- [108] Ciucanu, I & Kerek, CI (1984) A simple and rapid method for the permethylation of carbohydrates. *Carbohydr. Res.*, 131, 209-217.
- [109] Vanderschaeghe, D, Laroy, W, Sablon, E, Halfon, P, Van Hecke, A, Delanghe, J, & Callewaert, N (2009) GlycoFibroTest: DNA sequencer-based serum protein glycomics yields a highly performant liver fibrosis biomarker. *Mol. Cell. Proteomics*, 5, 986-949.
- [110] Royle, L, Mattu, TS, Hart, E, Langridge, JI, Merry, AH, Murphy, N, Harvey, DJ, Dwek, RA, & Rudd, PM (2002) An analytical and structural database provides a strategy for sequencing O-glycans from microgram quantities of glycoproteins. *Anal. Biochem.*, 304, 70-90.
- [111] Morelle, W, Page, A, & Michalski, JC (2005) Electrospray ionization ion trap

- mass spectrometry for structural characterization of oligosaccharides derivatized with 2-aminobenzamide. *Rapid Commun. Mass Spectrom.*, 19, 1145-1158.
- [112] Xia, B, Kawar, ZS, Ju, T, Alvarez, RA, Sachdev, GP, & Cummings, RD (2005) Versatile fluorescent derivatization of glycans for glycomic analysis. *Nat. Methods*, 2, 845-850.
- [113] Xia, B, Feasley, CL, Sachdev, GP, Smith, DF, & Cummings, RD (2009) Glycan reductive isotope labeling for quantitative glycomics. *Anal. Biochem.*, 387, 162-170.
- [114] Lei, M, Mechref, Y, & Novotny, MV (2009) Structural analysis of sulfated glycans by sequential double-permethylation using methyl iodide and deuteromethyl iodide. *J. Am. Soc. Mass Spectrom.*, 20, 1660-1671.
- [115] Kapkova, P (2009) Mass spectrometric analysis of carbohydrates labeled with a biotinylated tag. *Rapid Commun. Mass Spectrom.*, 23, 2775-2784.
- [116] Wuhrer, M, Koeleman, CA, Deelder, AM, & Hokke, CH (2006) Repeats of LacdiNAc and fucosylated LacdiNAc on N-glycans of the human parasite *Schistosoma mansoni*. *FEBS J.*, 273, 347-361.
- [117] Broberg, A (2007) High-performance liquid chromatography/electrospray ionization ion-trap mass spectrometry for analysis of oligosaccharides derivatized by reductive amination and N,N-dimethylation. *Carbohydr. Res.*, 342, 1462-1469.
- [118] Prater, BD, Anumula, KR, & Hutchins, JT (2007) Automated sample preparation facilitated by PhyNexus MEA purification system for oligosaccharide mapping of glycoproteins. *Anal. Biochem.*, 369, 202-209.
- [119] Olajos, M, Hajos, P, Bonn, GK, & Guttman, A (2008) Sample preparation for the analysis of complex carbohydrates by multicapillary gel electrophoresis with light-emitting diode induced fluorescence detection. *Anal. Chem.*, 80, 4241-4246.
- [120] Schwarzer, J, Rapp, E, & Reichl, U (2008) N-glycan analysis by CGE-LIF: profiling influenza A virus hemagglutinin N-glycosylation during vaccine production. *Electrophoresis*, 29, 4203-4214.

- [121] Yamaguchi, Y, Nishimura, M, Nagano, M, Yagi, H, Sasakawa, H, Uchida, K, Shitara, K, & Kato, K (2006) Glycoform-dependent conformational alteration of the Fc region of human immunoglobulin G1 as revealed by NMR spectroscopy. *Biochim. Biophys. Acta*, 1760, 693-700.
- [122] Kita, Y, Miura, Y, Furukawa, J, Nakano, M, Shinohara, Y, Ohno, M, Takimoto, A, & Nishimura, S (2007) Quantitative glycomics of human whole serum glycoproteins based on the standardized protocol for liberating N-glycans. *Mol. Cell. Proteomics*, 6, 1437-1445.
- [123] Atwood, JA, III, Cheng, L, varez-Manilla, G, Warren, NL, York, WS, & Orlando, R (2008) Quantitation by isobaric labeling: applications to glycomics. *J. Proteome. Res.*, 7, 367-374.
- [124] Guillard, M, Gloerich, J, Wessels, HJ, Morava, E, Wevers, RA, & Lefeber, DJ (2009) Automated measurement of permethylated serum N-glycans by MALDI-linear ion trap mass spectrometry. *Carbohydr. Res.*, 344, 1550-1557.
- [125] Botelho, JC, Atwood, JA, Cheng, L, varez-Manilla, G, York, WS, & Orlando, R (2008) Quantification by isobaric labeling (QUIBL) for the comparative glycomic study of O-linked glycans. *Int. J. Mass Spectrom.*, 278, 137-142.
- [126] Saba, JA, Shen, X, Jamieson, JC, & Perreault, H (2001) Investigation of different combinations of derivatization, separation methods and electrospray ionization mass spectrometry for standard oligosaccharides and glycans from ovalbumin. *J. Mass Spectrom.*, 36, 563-574.
- [127] Packer, NH, Lawson, MA, Jardine, DR, & Redmond, JW (1998) A general approach to desalting oligosaccharides released from glycoproteins. *Glycoconj. J.*, 15, 737-747.
- [128] Ruhaak, LR, Deelder, AM, & Wuhrer, M (2009) Oligosaccharide analysis by graphitized carbon liquid chromatography-mass spectrometry. *Anal. Bioanal. Chem.*, 394, 163-174.
- [129] Royle, L, Radcliffe, CM, Dwek, RA, & Rudd, PM (2006) Detailed structural analysis of N-glycans released from glycoproteins in SDS-PAGE gel bands using HPLC combined with exoglycosidase array digestions. *Methods Mol. Biol.*, 347, 125-143.

- [130] Takahashi, N (1996) Three-dimensional mapping of N-linked oligosaccharides using anion-exchange, hydrophobic and hydrophilic interaction modes of high-performance liquid chromatography. *J. Chromatogr. A*, 720, 217-225.
- [131] Vanderschaeghe, D, Festjens, N, Delanghe, J, & Callewaert, N (2010) Glycome profiling using modern glycomics technology: technical aspects and applications. *Biol. Chem.*, 391, 149-161.
- [132] Zaia, J (2004) Mass spectrometry of oligosaccharides. *Mass Spectrom. Rev.*, 23, 161-227.
- [133] Mechref, Y & Novotny, MV (2002) Structural investigations of glycoconjugates at high sensitivity. *Chem. Rev.*, 102, 321-369.
- [134] Wuhrer, M, de Boer, AR, & Deelder, AM (2009) Structural glycomics using hydrophilic interaction chromatography (HILIC) with mass spectrometry. *Mass Spectrom. Rev.*, 28, 192-206.
- [135] Campbell, MP, Royle, L, Radcliffe, CM, Dwek, RA, & Rudd, PM (2008) GlycoBase and autoGU: tools for HPLC-based glycan analysis. *Bioinformatics*, 24, 1214-1216.
- [136] Thomsson, KA, Karlsson, NG, & Hansson, GC (1999) Liquid chromatography-electrospray mass spectrometry as a tool for the analysis of sulfated oligosaccharides from mucin glycoproteins. *J. Chromatogr. A*, 854, 131-139.
- [137] Wuhrer, M, Koeleman, CA, Deelder, AM, & Hokke, CH (2004) Normal-phase nanoscale liquid chromatography-mass spectrometry of underivatized oligosaccharides at low-femtomole sensitivity. *Anal. Chem.*, 76, 833-838.
- [138] Maslen, SL, Goubet, F, Adam, A, Dupree, P, & Stephens, E (2007) Structure elucidation of arabinoxylan isomers by normal phase HPLC-MALDI-TOF/TOF-MS/MS. *Carbohydr. Res.*, 342, 724-735.
- [139] Wuhrer, M, Deelder, AM, & Hokke, CH (2005) Protein glycosylation analysis by liquid chromatography-mass spectrometry. *J. Chromatogr. B*, 825, 124-133.
- [140] Chen, X & Flynn, GC (2009) Gas-phase oligosaccharide nonreducing end (GONE) sequencing and structural analysis by reversed phase HPLC/mass

- spectrometry with polarity switching. *J. Am. Soc. Mass Spectrom.*, 20, 1821-1833.
- [141] Chu, CS, Ninonuevo, MR, Clowers, BH, Perkins, PD, An, HJ, Yin, H, Killeen, K, Miyamoto, S, Grimm, R, & Lebrilla, CB (2009) Profile of native N-linked glycan structures from human serum using high performance liquid chromatography on a microfluidic chip and time-of-flight mass spectrometry. *Proteomics*, 9, 1939-1951.
- [142] Pabst, M, Bondili, JS, Stadlmann, J, Mach, L, & Altmann, F (2007) Mass + retention time = structure: a strategy for the analysis of N-glycans by carbon LC-ESI-MS and its application to fibrin N-glycans. *Anal. Chem.*, 79, 5051-5057.
- [143] Bruggink, C, Wuhrer, M, Koeleman, CA, Barreto, V, Liu, Y, Pohl, C, Ingendoh, A, Hokke, CH, & Deelder, AM (2005) Oligosaccharide analysis by capillary-scale high-pH anion-exchange chromatography with on-line ion-trap mass spectrometry. *J. Chromatogr. B*, 829, 136-143.
- [144] Scheinthal, BM & Bettelheim, FA (1968) Multiple forms of Sialic acids. *Carbohydr. Res.*, 6, 257-265.
- [145] Zhuang, Z, Starkey, JA, Mechref, Y, Novotny, MV, & Jacobson, SC (2007) Electrophoretic analysis of N-glycans on microfluidic devices. *Anal. Chem.*, 79, 7170-7175.
- [146] Chen, FT & Evangelista, RA (1998) Profiling glycoprotein n-linked oligosaccharide by capillary electrophoresis. *Electrophoresis*, 19, 2639-2644.
- [147] Kamoda, S, Nomura, C, Kinoshita, M, Nishiura, S, Ishikawa, R, Kakehi, K, Kawasaki, N, & Hayakawa, T (2004) Profiling analysis of oligosaccharides in antibody pharmaceuticals by capillary electrophoresis. *J. Chromatogr. A*, 1050, 211-216.
- [148] Vanderschaeghe, D, Debruyne, E, Van Vlierberghe, H, Callewaert, N, & Delanghe, J (2009) Analysis of gamma-globulin mobility on routine clinical CE equipment: Exploring its molecular basis and potential clinical utility. *Electrophoresis*, 30, 2617-2623.
- [149] Callewaert, N, Contreras, R, Mitnik-Gankin, L, Carey, L, Matsudaira, P, & Ehrlich, D (2004) Total serum protein N-glycome profiling on a capillary electrop-

- horesis-microfluidics platform. *Electrophoresis*, 25, 3128-3131.
- [150] Heiskanen, A, Hirvonen, T, Salo, H, Impola, U, Olonen, A, Laitinen, A, Tiitinen, S, Natunen, S, Aitio, O, Miller-Podraza, H, Wuhrer, M, Deelder, AM, Natunen, J, Laine, J, Lehenkari, P, Saarinen, J, Satomaa, T, & Valmu, L (2009) Glycomics of bone marrow-derived mesenchymal stem cells can be used to evaluate their cellular differentiation stage. *Glycoconj. J.*, 26, 367-384.
- [151] Oeppen, J & Vaupel, JW (2002) Demography. Broken limits to life expectancy. *Science*, 296, 1029-1031.
- [152] Bartke, A (2008) Insulin and aging. *Cell Cycle*, 7, 3338-3343.
- [153] Bartke, A (2008) Impact of reduced insulin-like growth factor-1/insulin signaling on aging in mammals: novel findings. *Aging Cell*, 7, 285-290.
- [154] Redman, LM, Martin, CK, Williamson, DA, & Ravussin, E (2008) Effect of caloric restriction in non-obese humans on physiological, psychological and behavioral outcomes. *Physiol. Behav.*, 94, 643-648.
- [155] Atzmon, G, Barzilai, N, Hollowell, JG, Surks, MI, & Gabriely, I (2009) Extreme longevity is associated with increased serum thyrotropin. *J. Clin. Endocrinol. Metab.*, 94, 1251-1254.
- [156] Kapahi, P, Boulton, ME, & Kirkwood, TB (1999) Positive correlation between mammalian life span and cellular resistance to stress. *Free Radic. Biol. Med.*, 26, 495-500.
- [157] Harper, JM, Salmon, AB, Leiser, SF, Galecki, AT, & Miller, RA (2007) Skin-derived fibroblasts from long-lived species are resistant to some, but not all, lethal stresses and to the mitochondrial inhibitor rotenone. *Aging Cell*, 6, 1-13.
- [158] Salmon, AB, Murakami, S, Bartke, A, Kopchick, J, Yasumura, K, & Miller, RA (2005) Fibroblast cell lines from young adult mice of long-lived mutant strains are resistant to multiple forms of stress. *Am. J. Physiol. Endocrinol. Metab.*, 289, E23-E29.
- [159] Westendorp, RG, van Heemst, D, Rozing, MP, Frolich, M, Mooijaart, SP, Blauw, GJ, Beekman, M, Heijmans, BT, de Craen, AJ, & Slagboom, PE (2009) Nonagenarian siblings and their offspring display lower risk of mortality and morbidity than sporadic nonagenarians: The Leiden Longevity Study. *J. Am.*

- Geriatr. Soc., 57, 1634-1637.
- [160] Schoenmaker, M, de Craen, AJ, de Meijer, PH, Beekman, M, Blauw, GJ, Slagboom, PE, & Westendorp, RG (2006) Evidence of genetic enrichment for exceptional survival using a family approach: the Leiden Longevity Study. *Eur. J. Hum. Genet.*, 14, 79-84.
- [161] Rozing, MP, Westendorp, RG, Frolich, M, de Craen, AJ, Beekman, M, Heijmans, BT, Mooijaart, SP, Blauw, GJ, Slagboom, PE, van, HD, & Group, OB (2009) Human insulin/IGF-1 and familial longevity at middle age. *Aging (Albany, NY)*, 1, 714-722.
- [162] Heijmans, BT, Beekman, M, Houwing-Duistermaat, JJ, Cobain, MR, Powell, J, Blauw, GJ, van der, OF, Westendorp, RG, & Slagboom, PE (2006) Lipoprotein particle profiles mark familial and sporadic human longevity. *PLoS. Med.*, 3, e495.
- [163] Rozing, MP, Westendorp, RG, de Craen, AJ, Frolich, M, Heijmans, BT, Beekman, M, Wijsman, C, Mooijaart, SP, Blauw, GJ, Slagboom, PE, & van, HD (2010) Low Serum Free Triiodothyronine Levels Mark Familial Longevity: The Leiden Longevity Study. *J Gerontol. A Biol Sci. Med. Sci.*, 65, 365-368.
- [164] Dekker, P, Maier, AB, van, HD, de Koning-Treurniet, C, Blom, J, Dirks, RW, Tanke, HJ, & Westendorp, RG (2009) Stress-induced responses of human skin fibroblasts in vitro reflect human longevity. *Aging Cell*, 8, 595-603.
- [165] Rudd, PM, Guile, GR, Kuster, B, Harvey, DJ, Opdenakker, G, & Dwek, RA (1997) Oligosaccharide sequencing technology. *Nature*, 388, 205-207.
- [166] Miura, Y, Hato, M, Shinohara, Y, Kuramoto, H, Furukawa, J, Kuroguchi, M, Shimaoka, H, Tada, M, Nakanishi, K, Ozaki, M, Todo, S, & Nishimura, S (2008) BlotGlycoABCTM, an integrated glycoblotting technique for rapid and large scale clinical glycomics. *Mol. Cell. Proteomics*, 7, 370-377.
- [167] Kang, P, Mechref, Y, & Novotny, MV (2008) High-throughput solid-phase permethylation of glycans prior to mass spectrometry. *Rapid Commun. Mass Spectrom.*, 22, 721-734.
- [168] Tajiri, M, Yoshida, S, & Wada, Y (2005) Differential analysis of site-specific glycans on plasma and cellular fibronectins: application of a hydrophilic af-

- finity method for glycopeptide enrichment. *Glycobiology*, 15, 1332-1340.
- [169] Neville, DC, Coquard, V, Priestman, DA, te Vrucchte, DJ, Sillence, DJ, Dwek, RA, Platt, FM, & Butters, TD (2004) Analysis of fluorescently labeled glycosphingolipid-derived oligosaccharides following ceramide glycanase digestion and anthranilic acid labeling. *Anal. Biochem.*, 331, 275-282.
- [170] Skov, T & Bro, R (2008) Solving fundamental problems in chromatographic analysis. *Anal. Bioanal. Chem.*, 390, 281-285.
- [171] Maley, F, Trimble, RB, Tarentino, AL, & Plummer, TH, Jr. (1989) Characterization of glycoproteins and their associated oligosaccharides through the use of endoglycosidases. *Anal. Biochem.*, 180, 195-204.
- [172] Wopereis, S, Abd Hamid, UM, Critchley, A, Royle, L, Dwek, RA, Morava, E, Leroy, JG, Wilcken, B, Lagerwerf, AJ, Huijben, KM, Lefeber, DJ, Rudd, PM, & Wevers, RA (2006) Abnormal glycosylation with hypersialylated O-glycans in patients with Sialuria. *Biochim. Biophys. Acta*, 1762, 598-607.
- [173] Kim, YG, Jang, KS, Joo, HS, Kim, HK, Lee, CS, & Kim, BG (2007) Simultaneous profiling of N-glycans and proteins from human serum using a parallel-column system directly coupled to mass spectrometry. *J. Chromatogr. B*, 850, 109-119.
- [174] Zhou, Y, Aebersold, R, & Zhang, H (2007) Isolation of N-linked glycopeptides from plasma. *Anal. Chem.*, 79, 5826-5837.
- [175] Hase, S (1994) High-performance liquid chromatography of pyridylaminated saccharides. *Methods Enzymol.*, 230, 225-237.
- [176] Ohara, K, Sano, M, Kondo, A, & Kato, I (1991) Two-dimensional mapping by high-performance liquid chromatography of pyridylamino oligosaccharides from various glycosphingolipids. *J. Chromatogr.*, 586, 35-41.
- [177] Wuhrer, M (2007) Glycosylation profiling in clinical proteomics: heading for glycan biomarkers. *Expert. Rev. Proteomics*, 4, 135-136.
- [178] Jang-Lee, J, North, SJ, Sutton-Smith, M, Goldberg, D, Panico, M, Morris, H, Haslam, S, & Dell, A (2006) Glycomic profiling of cells and tissues by mass spectrometry: fingerprinting and sequencing methodologies. *Methods Enzymol.*, 415, 59-86.

- [179] Sato, S, Sakamoto, T, Miyazawa, E, & Kikugawa, Y (2004) One-pot reductive amination of aldehydes and ketones with *N*-picoline-borane in methanol, in water, and in neat conditions. *Tetrahedron*, 60, 7899-7906.
- [180] Domon, B & Costello, CE (1988) A Systematic Nomenclature for Carbohydrate Fragmentations in FAB-MS/MS Spectra of Glycoconjugates. *Glycoconj. J.*, 5, 397-409.
- [181] Apweiler, R, Hermjakob, H, & Sharon, N (1999) On the frequency of protein glycosylation, as deduced from analysis of the SWISS-PROT database. *Biochim. Biophys. Acta*, 1473, 4-8.
- [182] Bowman, MJ & Zaia, J (2010) Comparative glycomics using a tetraplex stable-isotope coded tag. *Anal. Chem.*, 82, 3023-3031.
- [183] van de Geijn, FE, Wuhler, M, Selman, MH, Willemsen, SP, de Man, YA, Deelder, AM, Hazes, JM, & Dolhain, RJ (2009) Immunoglobulin G galactosylation and sialylation are associated with pregnancy-induced improvement of rheumatoid arthritis and the postpartum flare: results from a large prospective cohort study. *Arthritis Res. Ther.*, 11, R193.
- [184] Rook, GAW, Steele, JF, Brealey, R, Whyte, A, Isenberg, D, Sumar, N, Nelson, JL, Bodman, KB, Young, A, Roitt, IM, Williams, P, Scragg, IG, Edge, CI, Arkwright, PD, Ashford, D, Wormald, MR, Rudd, PM, Redman, CWG, Dwek, RA, & Rademacher, TW (1991) Changes in IgG glycoform levels are associated with remission of arthritis during pregnancy. *J. Autoimmun.*, 4, 779-794.
- [185] Ruhaak, LR, Uh, HW, Beekman, M, Koeleman, CAM, Hokke, CH, Westendorp, RG, Wuhler, M, Houwing-Duistermaat, JJ, Slagboom, PE, & Deelder, AM (2010) Decreased levels of bisecting GlcNAc glycoforms of IgG mark human longevity. *PIOS ONE*, submitted.
- [186] Stockert, RJ (1995) The asialoglycoprotein receptor: relationships between structure, function, and expression. *Physiol. Rev.*, 75, 591-609.
- [187] Kaneko, Y, Nimmerjahn, F, & Ravetch, JV (2006) Anti-inflammatory activity of immunoglobulin G resulting from Fc sialylation. *Science*, 313, 670-673.
- [188] Anthony, RM, Wermeling, F, Karlsson, MC, & Ravetch, JV (2008) Identification of a receptor required for the anti-inflammatory activity of IVIG. *Proc. Natl.*

- Acad. Sci. U. S. A., 105, 19571-19578.
- [189] Tolstikov, VV & Fiehn, O (2002) Analysis of highly polar compounds of plant origin: combination of hydrophilic interaction chromatography and electro-spray ion trap mass spectrometry. *Anal. Biochem.*, 301, 298-307.
- [190] Ikegami, T, Horie, K, Saad, N, Hosoya, K, Fiehn, O, & Tanaka, N (2008) Highly efficient analysis of underivatized carbohydrates using monolithic-silica-based capillary hydrophilic interaction (HILIC) HPLC. *Anal. Bioanal. Chem.*, 391, 2533-2542.
- [191] Sarmini, K & Kenndler, E (1999) Ionization constants of weak acids and bases in organic solvents. *J. Biochem. Biophys. Methods*, 38, 123-137.
- [192] Arnold, JN, Wormald, MR, Sim, RB, Rudd, PM, & Dwek, RA (2007) The impact of glycosylation on the biological function and structure of human immunoglobulins. *Annu. Rev. Immunol.*, 25, 21-50.
- [193] Pepys, MB & Hirschfield, GM (2003) C-reactive protein: a critical update. *J. Clin. Invest.*, 111, 1805-1812.
- [194] Al-Regaiey, KA, Masternak, MM, Bonkowski, M, Sun, L, & Bartke, A (2005) Long-lived growth hormone receptor knockout mice: interaction of reduced insulin-like growth factor i/insulin signaling and caloric restriction. *Endocrinology*, 146, 851-860.
- [195] Crimmins, E, Vasunilashorn, S, Kim, JK, & Alley, D (2008) Biomarkers related to aging in human populations. *Adv. Clin. Chem.*, 46, 161-216.
- [196] Johnson, WE, Li, C, & Rabinovic, A (2007) Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics.*, 8, 118-127.
- [197] Narasimhan, S (1982) Control of glycoprotein synthesis. UDP-GlcNAc:glycopeptide beta 4-N-acetylglucosaminyltransferase III, an enzyme in hen oviduct which adds GlcNAc in beta 1-4 linkage to the beta-linked mannose of the trimannosyl core of N-glycosyl oligosaccharides. *J. Biol. Chem.*, 257, 10235-10242.
- [198] Shaper, NL, Shaper, JH, Meuth, JL, Fox, JL, Chang, H, Kirsch, IR, & Hollis, GF (1986) Bovine galactosyltransferase: identification of a clone by direct im-

- munological screening of a cDNA expression library. *Proc. Natl. Acad. Sci. U. S. A.*, 83, 1573-1577.
- [199] Narimatsu, H, Sinha, S, Brew, K, Okayama, H, & Qasba, PK (1986) Cloning and sequencing of cDNA of bovine N-acetylglucosamine (beta 1-4)galactosyl-transferase. *Proc. Natl. Acad. Sci. U. S. A.*, 83, 4720-4724.
- [200] Umana, P, Jean-Mairet, J, Moudry, R, Amstutz, H, & Bailey, JE (1999) Engineered glycoforms of an antineuroblastoma IgG1 with optimized antibody-dependent cellular cytotoxic activity. *Nat. Biotechnol.*, 17, 176-180.
- [201] Iida, S, Kuni-Kamochi, R, Mori, K, Misaka, H, Inoue, M, Okazaki, A, Shitara, K, & Satoh, M (2009) Two mechanisms of the enhanced antibody-dependent cellular cytotoxicity (ADCC) efficacy of non-fucosylated therapeutic antibodies in human blood. *BMC. Cancer*, 9, 58.
- [202] Shibata-Koyama, M, Iida, S, Misaka, H, Mori, K, Yano, K, Shitara, K, & Satoh, M (2009) Nonfucosylated rituximab potentiates human neutrophil phagocytosis through its high binding for FcγRIIIb and MHC class II expression on the phagocytotic neutrophils. *Exp. Hematol.*, 37, 309-321.
- [203] Peipp, M, Lammerts van Bueren, JJ, Schneider-Merck, T, Bleeker, WW, Dechant, M, Beyer, T, Repp, R, van Berkel, PH, Vink, T, van de Winkel, JG, Parren, PW, & Valerius, T (2008) Antibody fucosylation differentially impacts cytotoxicity mediated by NK and PMN effector cells. *Blood*, 112, 2390-2399.
- [204] Skov, T, van den Berg, F, Tomasi, G, & Bro, R (2006) Automated alignment of chromatographic data. *Journal of Chemometrics*, 20, 484-497.
- [205] Yang, Z & Hancock, WS (2004) Approach to the comprehensive analysis of glycoproteins isolated from human serum using a multi-lectin affinity column. *J. Chromatogr. A*, 1053, 79-88.
- [206] Nielsen, N-PV, Carstensen, JM, & Smedsgaard, J (1998) Aligning of single and multiple wavelength chromatographic profiles for chemometric data analysis using correlation optimised warping. *J. Chromatogr. A*, 805, 17-35.
- [207] Kolarich, D, Weber, A, Turecek, PL, Schwarz, HP, & Altmann, F (2006) Comprehensive glyco-proteomic analysis of human alpha1-antitrypsin and its charge isoforms. *Proteomics.*, 6, 3369-3380.

- [208] Mills, K, Mills, PB, Clayton, PT, Johnson, AW, Whitehouse, DB, & Winchester, BG (2001) Identification of alpha(1)-antitrypsin variants in plasma with the use of proteomic technology. *Clin. Chem.*, 47, 2012-2022.
- [209] Mattu, TS, Pleass, RJ, Willis, AC, Kilian, M, Wormald, MR, Lellouch, AC, Rudd, PM, Woof, JM, & Dwek, RA (1998) The glycosylation and structure of human serum IgA1, Fab, and Fc regions and the role of N-glycosylation on Fc alpha receptor interactions. *J. Biol. Chem.*, 273, 2260-2272.
- [210] Gomes, MM, Wall, SB, Takahashi, K, Novak, J, Renfrow, MB, & Herr, AB (2008) Analysis of IgA1 N-glycosylation and its contribution to Fc alphaRI binding. *Biochemistry*, 47, 11285-11299.
- [211] Field, MC, matayakul-Chantler, S, Rademacher, TW, Rudd, PM, & Dwek, RA (1994) Structural analysis of the N-glycans from human immunoglobulin A1: comparison of normal human serum immunoglobulin A1 with that isolated from patients with rheumatoid arthritis. *Biochem. J.*, 299 (Pt 1), 261-275.
- [212] Corthals, GL, Wasinger, VC, Hochstrasser, DF, & Sanchez, JC (2000) The dynamic range of protein expression: a challenge for proteomic research. *Electrophoresis*, 21, 1104-1115.
- [213] Fang, X & Zhang, WW (2008) Affinity separation and enrichment methods in proteomic analysis. *J. Proteomics*, 71, 284-303.
- [214] Boschetti, E & Righetti, PG (2009) The art of observing rare protein species in proteomes with peptide ligand libraries. *Proteomics*, 9, 1492-1510.
- [215] Mairal, T, Ozalp, VC, Lozano, SP, Mir, M, Katakis, I, & O'Sullivan, CK (2008) Aptamers: molecular tools for analytical applications. *Anal. Bioanal. Chem.*, 390, 989-1007.
- [216] Bereman, MS & Muddiman, DC (2010) The effects of abundant plasma protein depletion on global glycan profiling using nanoLC FT-ICR mass spectrometry. *Anal Bioanal. Chem*, 396, 1473-1479.
- [217] Atkinson, AJ, Colburn, WA, DeGruttola, VG, DeMets, DL, Downing, GJ, Hoth, DF, Oates, JA, Peck, CC, Schooley, RT, Spilker, BA, Woodcock, J, & Zeger, SL (2001) Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin. Pharmacol. Ther.*, 69, 89-95.

- [218] Rifai, N, Gillette, MA, & Carr, SA (2006) Protein biomarker discovery and validation: the long and uncertain path to clinical utility. *Nat. Biotechnol.*, 24, 971-983.
- [219] Robling, AG, Niziolek, PJ, Baldrige, LA, Condon, KW, Allen, MR, Alam, I, Mantila, SM, Gluhak-Heinrich, J, Bellido, TM, Harris, SE, & Turner, CH (2008) Mechanical stimulation of bone in vivo reduces osteocyte expression of Sost/sclerostin. *J. Biol. Chem.*, 283, 5866-5875.