



Universiteit
Leiden
The Netherlands

Spreading depolarizations : the missing link between migraine and stroke

Ayata, C.

Citation

Ayata, C. (2016, November 24). *Spreading depolarizations : the missing link between migraine and stroke*. Retrieved from <https://hdl.handle.net/1887/44385>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/44385>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/44385> holds various files of this Leiden University dissertation.

Author: Ayata, C.

Title: Spreading depolarizations : the missing link between migraine and stroke

Issue Date: 2016-11-24

CHAPTER 1

GENERAL INTRODUCTION

1. MIGRAINE

Migraine is a common and disabling episodic neurovascular disease¹ that affects ~15% of the world population; it is three times more common among women of reproductive age than men.^{2,3} Recurrent migraine attacks are characterized by severe headache accompanied to various degrees by nausea, vomiting, and increased sensitivity to light, sound, and smell, lasting anywhere between an hour to days (migraine *without* aura).⁴ Attack frequency differs widely among patients, from one per year to several per week. Every day, millions of people are plagued by blistering migraines, which ranks among the most disabling medical conditions worldwide.^{2,5}

Migraine aura. One-third of migraine patients also experience transient focal neurological symptoms termed *aura* that usually precede or sometimes accompany the headache (migraine *with* aura).^{4,6} Aura symptoms can be visual, sensory or motor, or involve language or brain stem function. Aura often shows a characteristic spread, such as a scintillating scotoma gradually expanding from a point of origin in one hemi visual field (Figure 1), or paresthesias marching up from the fingers and hand to sequentially involve the arm and then lower face. This form of spread pattern suggests an intracortical event propagating in a retinotopic or somatotopic fashion. It is widely accepted^{7,8} that the symptomatology of migraine aura is caused by the electrophysiological event called *spreading depression* (SD), an intense neuronal and glial spreading depolarization wave slowly propagating (~3 mm/min) in cerebral cortical or subcortical grey matter.⁹⁻¹¹ Mechanisms of migraine headache are still debated, although there is general agreement that activation and sensitization of the trigeminovascular nociceptive pathways is a critical step.^{12,13} A large body of experimental evidence strongly suggests that SD is capable of activating the trigeminovascular system, and thus directly relevant for headache mechanisms.¹¹ The




Figure 1. Scintillating scotoma of visual migraine aura. Fovea is indicated by x, and the time between each successive drawing by the numbers in minutes. The zig-zag pattern at the propagating wavefront represents scintillations (i.e., positive symptom at the wavefront) while the gray shaded area represents visual loss (i.e., scotoma, negative symptom).¹⁵ Visual aura is caused by SD in occipital cortex.

trigeminovascular system consists of nociceptive trigeminal afferents surrounding the intracranial vessels. These perivascular trigeminal afferents project through the trigeminal ganglion to neurons in the trigeminocervical complex, and then relayed to the thalamus where all nociceptive inputs are integrated.¹⁴

2. SPREADING DEPRESSION

Definition. Spreading depression (SD) is an intense neuronal and glial depolarization wave that slowly propagates in brain tissue (~3 mm/min) by way of gray matter contiguity irrespective of functional divisions or vascular territories.¹⁶ The near-complete loss of membrane potential is a result of massive transmembrane ionic and water shifts, including K^+ and glutamate efflux, and Na^+ , Ca^{2+} and water influx, which last up to a minute and does not lead to injury in otherwise normal brain tissue. All these ionic and water shifts create a signature extracellular negative slow potential shift (aka DC shift) accompanied by suppression of action potentials and synaptic activity (Figure 2). As a result, electrocorticogram (ECoG) is depressed, hence the historical term “spreading depression” originally coined by Leao.¹⁷ Similar spreading depolarization events also occur in ischemic brain, and are termed peri-infarct depolarizations (PIDs) or injury depolarizations (see below). Therefore, SD is a form of spreading depolarization event that occurs in otherwise normal brain.

Basic electrophysiology. SD is triggered when a sufficiently strong stimulus (e.g., topical application of concentrated KCl solution, direct cathodal electrical stimulation, seizure activity or tissue ischemia) simultaneously depolarizes a minimum critical volume of brain tissue estimated to be ~1 mm³ in rodent cortex, *in vivo*.¹⁸ The depolarizing stimulus overloads the extracellular K^+ ($[K^+]_e$) clearance mechanisms causing $[K^+]_e$ to exceed a critical threshold concentration of ~12 mM.¹⁸⁻²⁴ These thresholds can vary in different species and brain regions depending on neuronal and excitatory synaptic density among other factors.^{25,26} The inciting event causes a sudden drop in membrane resistance via opening of non-selective large conductance cation channels, the presence and identity of which are yet incompletely understood.²⁷ As a result, both intracellular and extracellular ions move along their transmembrane concentration gradients. Massive K^+ efflux raises $[K^+]_e$ from ~3 mM at resting state²⁸⁻³⁰ to about ~30-50 mM, and sometimes as high as 80 mM, in an all-or-none fashion, in most species, tissues and model systems.^{19,22,29,31-34} This large K^+ efflux is reciprocated by Na^+ and Cl^- influx that pulls water, causing cell swelling.^{22,35-40} Extracellular space shrinks by as much as 50%.^{41, 42} Depolarization also triggers Ca^{2+} influx and a more than 10-fold drop in $[Ca^{2+}]_e$,^{22,43,44} which, along with Na^+ and water influx, leads to release of many if not all neurotransmitters and neuromodulators within the depolarized tissue. Extracellular glutamate, aspartate, glycine, GABA and taurine concentrations increase during SD,^{32,45,46} and similar increases have been shown for catecholamine and ascorbate levels.⁴⁷⁻⁴⁹




Figure 2. Representative tracings of electrocorticogram (ECoG) and extracellular DC potential during a cortical SD wave in rat brain, triggered by 1M KCl briefly applied on to the occipital cortex, and recorded by two intracortical glass microelectrodes placed in series with the KCl application site. SD is characterized by transient depression of cortical electrical activity and a slow DC shift 20-30 mV in amplitude lasting less than 1 min. When triggered by an intense stimulus that simultaneously depolarizes a minimum critical volume of brain tissue, SD spreads centrifugally to be detected first at the proximal (E1), and after a latency, at the distal (E2) microelectrode. In otherwise normal tissue, ECoG activity starts to recover a few minutes after the onset of SD, but may take up to 10 min to return to normal. The speed of propagation is calculated from the distance and the latency between the electrodes. Modified from ¹⁰.

Propagation of SD. It is believed that the rise in $[K^+]_e$ and glutamate act as chemical signals diffusing to and depolarizing adjacent cells, and in this way the depolarization slowly spreads. The massive rise in $[K^+]_e$ to levels sufficient to depolarize neighboring cells is the critical factor mediating the contiguous spread of the wave.^{20,22,50} Elevated extracellular levels of the strongly depolarizing excitatory amino acids (glutamate and aspartate) further fuel SD and facilitate its propagation by activating NMDA receptors.^{51,52} For the released K^+ and glutamate to reach the critical depolarization threshold of adjacent cells, high neuronal and synaptic density and low extracellular space volume are required; therefore, white matter is characteristically resistant to SD. However, SD can be triggered in subcortical grey matter structures such as striatum, thalamus and hippocampus, with the exception of brain stem, which is resistant to SD unless tested in immature animals or after pharmacological preconditioning (e.g., K^+ -channel blockade).⁵³⁻⁵⁷ Lastly, cortical SD can propagate into subcortical structures that have direct gray matter contiguity with the cortex.⁵⁸

Recovery of SD. The massive redistribution of ions, water and neurotransmitters is self-limited. A number of mechanisms, including the Na^+/K^+ -ATPase, intracellular buffering of $[Ca^{++}]_i$, reuptake and metabolism or spatial buffering by the astrocytic network, and quite likely vascular clearance, all help restore the homeostasis usually

within a minute. The process is in part energy-dependent and strongly stimulates O_2 and glucose consumption. Therefore, in severely hypoperfused (i.e., ischemic) tissue, restoration of homeostasis is delayed, with deleterious consequences on tissue viability.

SD and migraine. Since the discovery of SD decades ago, similarities between the electrophysiological properties of SD and the neurological signs and symptoms during migraine aura suggested a causative link between the two.⁵⁹⁻⁶¹ Experimental evidence also suggests that SD can trigger headache by activating the trigeminovascular system.^{11,62-67} Although whether an asymptomatic SD triggers migraine headache without a perceived aura is still debated, suppression of SD susceptibility by migraine prophylactic drugs as a class effect supported this notion since these drugs have been equally efficacious in migraine with or without a perceived aura. Therefore, SD is now considered a potential therapeutic target in migraine, and experimental models of SD susceptibility are increasingly being used in migraine drug screening.

SD susceptibility. The ease with which SD can be initiated and sustained is often termed *SD susceptibility*, and can be used as an experimental surrogate for migraine (with aura) susceptibility. Indeed, modulation of SD susceptibility by drugs appears to be the basis for migraine prophylaxis.⁶⁸ There are several experimental models that can be used to measure SD susceptibility.⁶⁹ The most common attributes used to define SD susceptibility are: i) the threshold stimulus intensity that triggers an SD (i.e., electrical charge measured in Coulombs, or KCl concentration threshold) determined by sequential stimuli of stepwise escalating intensity until an SD is triggered, ii) the frequency of SDs triggered during continuous constant suprathreshold stimulus for up to an hour (i.e., topical high concentration of KCl), and iii) the propagation speed of SD.

3. ISCHEMIC STROKE

Definition of the problem. Ischemic stroke is an acute cerebrovascular catastrophe and a leading cause of death and disability worldwide.⁷⁰ It is caused by occlusion of an artery supplying the brain or spinal cord. Despite intense research into the pathophysiology and treatment of ischemic stroke, the only proven and widely adopted therapeutic modality is thrombolysis to achieve reperfusion in a timely manner. Brain tissue is highly sensitive to ischemia. Infarction ensues unless reperfusion is achieved within a few hours after cerebral arterial occlusion. Therefore, the therapeutic efficacy and safety window of thrombolysis is limited to only 4.5 hours after stroke onset.⁷¹

Ischemic core. Cerebral arterial occlusion typically creates a focal perfusion defect within its territory (Figure 3). Some degree of collateral blood flow reaches the focal ischemic tissue via surrounding patent arteries. However, efficacy of collateral flow drops as a function of distance, creating an inverted bell-shaped gradient of blood flow from the relatively well-perfused periphery towards the severely ischemic center. Irreversible ischemic injury (i.e., infarct) starts in the center and gradually expands over minutes to days into the periphery. Energy shortage in the severely ischemic center results in failure of Na^+/K^+ ATPase, gradual increase in extracellular potassium concentrations ($[\text{K}^+]_e$). Within a few minutes after the arterial occlusion, when $[\text{K}^+]_e$ reaches a critical threshold of ~ 12 mM (from a normal resting level of ~ 3 -4 mM), neurons develop a sudden onset catastrophic loss of resting membrane potential (V_m), termed *anoxic depolarization* (AD), which marks the *ischemic core*. The process is analogous to the triggering of SD as described above, with the critical distinction being tissue inability to restore the transmembrane ionic gradients because of ongoing ischemia. In that sense, AD is a persistent form of SD that can become permanent. It is generally believed that AD that continues longer than ~ 10 -15 minutes results in irreversible cell injury and eventual death by necrosis and/or apoptosis.

Ischemic penumbra. Surrounding the core, there is sufficient collateral flow to maintain the resting V_m , but neural activity and synaptic transmission are suppressed via intrinsic protective mechanisms creating a ring of electrical silence (Figure 3). This ‘still-viable tissue at risk for infarction’ is called *ischemic penumbra*. The spatiotemporal evolution of injury in penumbra is complex. Depending on residual tissue perfusion and the intrinsic tissue sensitivity to ischemia, portions of penumbra succumb to AD, and get incorporated into the ischemic core over time. Therefore, ischemic core gradually expands into the penumbra (Figure 4). A critical factor that facilitates ischemic injury, expands the infarct and worsens the neurological outcome is the occurrence of spontaneous spreading depression waves within the peri-infarct tissue, called peri-infarct depolarizations.




Figure 3. The hemodynamic status and membrane potential of acute focal ischemic brain tissue showing the definitions of ischemic penumbra and core. V_m , membrane potential.




Figure 4. Ischemic core gradually expands into the penumbra because of persistent ischemia and the occurrence of peri-infarct depolarizations.

Peri-infarct depolarizations. In addition to AD, ischemic brain also develops spontaneous recurrent *peri-infarct spreading depolarization waves* (PIDs) that originate at the boundary and propagate throughout the ischemic and non-ischemic tissue. When PIDs enter the non-ischemic tissue, they are electrophysiologically indistinguishable from SD, the substrate of migraine aura, suggesting that PIDs are analogous to SDs (Figure 5). The local rise in $[K^+]_e$ and the hypoxic suppression of Na^+/K^+ -ATPase in ischemic brain serve as the strongly depolarizing stimuli that trigger PIDs, analogous to the topical KCl or electrical stimulation to trigger SDs. Importantly, occurrence of PIDs have been shown in human brain as well.⁷² Therefore, SDs and PIDs represent a mechanistic overlap between migraine and stroke.




Figure 5. Recurrent peri-infarct injury depolarizations (PIDs; four negative deflections on the tracing) recorded during filament occlusion of the middle cerebral artery by an intracortical glass micropipette placed outside the ischemic tissue in a representative mouse. These PIDs are indistinguishable from SDs triggered by intense depolarization in otherwise normal cortex. Lower right panel is the summary of 10 experiments, where each horizontal line shows the beginning and end of electrophysiological recordings after the onset of ischemia, and each circle represents a PID. Grey shaded area shows the approximate location of the perfusion defect upon arterial occlusion. Recurrent PIDs appear spontaneously throughout the recording period in all mice.

Acute infarct progression on MRI. Ionic changes and cell swelling in tissue that has undergone AD cause a characteristic decrease in apparent diffusion coefficient (ADC) of water on diffusion-weighted MRI (DWI) as the *MRI signature of core infarction* (Figure 6). Perfusion-weighted MRI (PWI) outlines the total tissue at risk for infarction. Therefore, the mismatch between DWI and PWI lesion volumes is a measure of viable tissue at risk for infarction (i.e., penumbra). Ischemic penumbra is the primary target of all acute stroke rescue interventions. The volume of penumbra present at any given time is a good measure of how much tissue is available that can be rescued from ischemic infarction.




Figure 6. Acute MRI of a typical middle cerebral artery stroke. Diffusion-weighted imaging (DWI) shows the infarcted tissue (yellow), whereas perfusion-weighted imaging (PWI; measured by mean transit time, MTT, blue) delineates the hypoperfused tissue. The DWI/PWI mismatch (blue tissue outside the yellow) is a surrogate MRI measure of ischemic penumbra, defined as ‘still-viable tissue at risk for infarction’. Day 7 T2 MRI shows the final infarct volume in this patient. Courtesy of Dr. Hakan Ay.

4. CLINICAL ASSOCIATION BETWEEN MIGRAINE AND STROKE

Migraine has traditionally been viewed as a benign chronic episodic condition. However, accumulating evidence suggests that migraine with aura, can be associated with increased risk for stroke and white matter lesions.⁷³⁻⁷⁹

Observational data. Abundant data from retrospective and population- or hospital-based case-control studies, as well as small and large population-based prospective studies including tens of thousands of subjects, have firmly established a link between migraine and ischemic stroke.^{77,80,81} The odds ratio (OR) for ischemic stroke is ~2 among migraineurs (95% confidence interval [CI] 1.72-2.43). Important insights are:

- I. The association is explained by migraine with aura alone (OR 2.5, 95% CI 1.5-4.1), and does not reach statistical significance for migraine without aura (OR 1.3, 95% CI 0.8-2.1),

- II. The association is stronger in women (OR 2.9, 95% CI 2.4-3.5),⁷⁷ and in subjects younger than 45 (OR 2.7, 95% CI 1.4-5.0),
- III. The risk is higher in patients who experience active migraine attacks (OR 1.9, 95% CI 1.2-3.1), and not in those with just a history of migraine without recent attacks,⁸²
- IV. The risk is higher in those who experience >12 attacks per year (OR 1.7, 95% CI 1.1-2.8).⁸³

Neuroimaging. A number of neuroimaging studies over the past decade revealed a higher prevalence of subclinical brain lesions in migraineurs, including infarcts and white matter hyperintensities, suggesting acute or chronic ischemic disease. CAMERA (Cerebral Abnormalities in Migraine and Epidemiological Risk Analysis) was a cross-sectional, population-based MRI lesion prevalence study in subjects between ages 30 and 60 (mean age 48; 161 migraine with aura, 134 migraine without aura and 140 matched controls) randomly selected from the Genetic Epidemiology of Migraine study. Data suggested an increased risk of subclinical posterior circulation infarct-like lesions, mostly located in the cerebellum, in migraineurs compared to controls (OR 7.1, 95% CI 0.9-55).^{78,84} Important insights are:

- I. The risk was substantially higher in migraineurs with aura (OR 13.7, 95% CI 1.7-112),
- II. The risk was also higher with frequent migraine attacks (≥ 1 attack/month) (OR 15.8, 95% CI 1.8-140),
- III. The risk was independent of triptan use or vascular risk factors,
- IV. In the 9-year follow up of the same cohort, none of the lesions disappeared, and new posterior circulation lesions were found in 5% of migraineurs compared with none in control subjects.⁸⁵

The conclusions of CAMERA study were later independently confirmed in the Age Gene/Environment Susceptibility Reykjavik study,⁸⁶ and in the population-based Epidemiology of Vascular Aging study (780 subjects, mean age of 69).⁸⁷ Clinical contrasts between migraine and stroke make this association highly intriguing:

- 1. Unlike the perceived benign nature of migraine (i.e., no imminent risk of injury), stroke is: (i) an acute and often catastrophic cerebrovascular event, (ii) the leading cause of acquired physical disability in adults in the US,⁸⁸ and (iii) the second leading cause of mortality worldwide.⁷⁰
- 2. Stroke is predominantly a disease of the elderly, while migraine prevalence peaks around age 40.
- 3. Stroke risk is higher in males than females of reproductive age, whereas migraine is higher in females of reproductive age and increases stroke risk most prominently in this group.

Investigation of the mechanisms underlying the association between migraine and stroke can impact hundreds of millions of people worldwide.

5. TRANSGENIC ANIMAL MODELS OF HUMAN MIGRAINE SYNDROMES

Modeling susceptibility to migraine. In order to elucidate the mechanisms linking migraine with aura and stroke, animal models are needed. While animal models of stroke are plenty, animal models of migraine with aura are relatively scarce. Migraine is an inherited disease.⁸⁹ Genetic determinants of migraine susceptibility, occurrence and severity range from rare monogenic inherited conditions (e.g., a single mutation is sufficient to cause disease in a patient) with large effect sizes to common polygenic influences (e.g., multiple polymorphisms are needed to cause disease in a patient) with small effect sizes. While the latter (i.e., common variants with small effects) is difficult to model experimentally, transgenic mouse models expressing human migraine mutations have been generated that recapitulate the clinical features of monogenic inherited conditions characterized by migraine with aura as well as ischemic stroke (e.g., CADASIL^{90,91}). Among these are mutant mouse models of familial hemiplegic migraine (FHM).

Familial hemiplegic migraine. Hemiplegic migraine is a rare monogenic form of migraine with aura characterized by motor weakness during the attacks sometimes accompanied by sensory, aphasic, visual and basilar symptoms, that can be sporadic, or familial (autosomal dominant), linked to genes involved in ion regulation (Table 1). Auras are often severe and prolonged, and a third of patients can experience a decrease in level of consciousness and even coma, which may be prolonged.⁹² The net result of all FHM mutations identified to date is dysregulation of membrane ionic equilibrium and hyperexcitability, predicting enhanced susceptibility to SD as the overarching theme and the overlapping feature between migraine and stroke. FHM has been used as a model for more common forms of migraine with aura because of shared clinical features and trigger factors, female preponderance, and because two thirds of FHM patients and their first-degree relatives also suffer from attacks of common migraine with and without aura.

Table 1. Familial hemiplegic migraine genes identified to date. Modified from Russell and Ducros, 2011.⁹²

	FHM1	FHM2	FHM3
Chromosome	19p13	1q23	2q24
Gene	CACNA1A	ATP1A2	SCN1A
Protein	Pore-forming $\alpha 1$ subunit of neuronal $\text{Ca}_v2.1$ (P/Q-type) voltage-gated Ca^{2+} channels	Catalytic $\alpha 2$ subunit of a glial and neuronal Na^+/K^+ ATPase	Pore-forming $\alpha 1$ subunit of neuronal $\text{Na}_v1.1$ voltage-gated Na^+ channels

Familial hemiplegic migraine type 1. FHM1 is caused by mutations in the pore-forming α_{1A} subunit of neuronal $\text{Ca}_v2.1$ (P/Q-type) voltage-gated Ca^{2+} channels (Figure 7). Mutations shift the voltage-current relationship of the channel to the left so that channels open at more negative membrane potentials (i.e., upon smaller membrane depolarizations), and stay open longer, increasing the Ca^{2+} influx. $\text{Ca}_v2.1$ channels are major regulators of presynaptic glutamate release. The net result is increased presynaptic Ca^{2+} entry and glutamate release resulting in enhanced cerebral excitability⁹³ that may be shared with more common forms of migraine.¹³ Among the FHM1 mutations identified to date, the S218L missense mutation confers stronger gain of $\text{Ca}_v2.1$ channel function and a more severe clinical phenotype with prolonged auras that can progress to coma.^{94,95} In contrast, the R192Q mutation is associated with modest gain of $\text{Ca}_v2.1$ channel function and pure hemiplegic auras.⁹⁶




Figure 7. FHM1 mutations and the $\text{Ca}_v2.1$ gain-of-function. The two missense mutations studied as part of this thesis are also shown (S218L and R192Q). Blue arrows show the direction of change by the S218L mutation with an allele dosage effect. Modified from Pietrobon, 2005.⁹⁷

6. OUTLINE OF THE THESIS

The overall hypothesis. The central hypothesis states that susceptibility to SD determines both the susceptibility to migraine with aura and the susceptibility to hypoxic/ischemic injury in the same direction. Therefore, factors that enhance the susceptibility to SD increase the likelihood of migraine with aura as well as ischemic stroke. Such factors that can modulate the susceptibility to SD may include genes, hormones and pharmacological agents.




Figure 8. The central hypothesis.

The hypothesis predicts that genetic, hormonal and pharmacological modulators that enhance or suppress SD susceptibility will render the brain more or less susceptible to ischemic injury, respectively (Figure 8). As such, SD is hypothesized to be the missing link between migraine and ischemic stroke. In order to test the hypothesis in a logical sequential manner, it is first necessary to identify and characterize a set of factors that modulate the susceptibility to SD that mimic the clinical observations of susceptibility to migraine aura (Figure 8, step 1), and then study the impact of those factors on outcome of a cerebral ischemic event (Figure 8, step 2).

The aim of this thesis is to investigate how genetic, hormonal and pharmacological modulators of SD susceptibility will influence the susceptible to ischemic injury. To this end we will unravel underlying mechanisms of SD susceptibility and susceptibility to ischemic injury by making use of two transgenic mouse models of migraine that carry migraine-relevant gene mutations in voltage-gated $\text{Ca}_v2.1$ Ca^{2+} channels (**Chapters 2 and 3**) and will review all relevant clinical and experimental data from the literature (**Chapter 4**).

In **Chapter 2**, we describe the genetic and gonadal hormone modulation of cortical SD susceptibility using state-of-the-art *in vivo* electrophysiological studies under full systemic physiological monitoring and maintenance in mice. To characterize the genetic modulation, we utilize two transgenic (knock-in) mouse models of FHM1 expressing the *Cacna1a* R192Q or S218L missense mutation in $\text{Ca}_v2.1$ voltage-gated (P/Q-type) Ca^{2+} channels. As mentioned above, the magnitude of single channel gain-of-function is larger in S218L mutants compared with R192Q, and with homozygous mutants compared with heterozygotes. Therefore, the use of two mutations and three genotypes (wild-type controls, heterozygous and homozygous mutants) allows a graded modulation the SD susceptibility. **Chapter 2A** also dissects the female gonadal hormone modulation of CSD susceptibility by gonadectomy (i.e., ovariectomy) and hormone replacement, and draws parallels between the clinical symptomatology of FHM and the neurological signs displayed by FHM1 mutant mice after an SD. Because higher SD susceptibility in female mutants can in part be due to an inhibitory effect of male gonadal hormones on SD in male mice, in **Chapter 2B** we investigate the effect of

androgens on SD once again using gonadectomy (i.e., orchiectomy). Ischemic events are not limited to cortex; therefore, in **Chapter 2C** we study the SD susceptibility in subcortical structures, including striatum, thalamus and hippocampus, and its genetic and gonadal hormone modulation, with the same graded approach using different mutants, genotypes and sexes as described above. Altogether, studies in **Chapter 2** set the stage to investigate the same genetic and hormonal modulators on the outcome of cerebral ischemic events in **Chapter 3**.

In **Chapter 3**, we build upon the data obtained in **Chapter 2** and investigate how SD modulators impact the outcome of cerebral ischemic events. In **Chapter 3A**, we test the impact of genetic susceptibility to SD on susceptibility to ischemic injury by making use of FHM1 knock-in mouse models. By comparing the R192Q and S218L mutants, different genotypes (heterozygotes vs. homozygotes) and sexes, we get an opportunity to study the correspondence between SD susceptibility and ischemic sensitivity across a wide range of phenotypes. **Chapter 3B** tests the opposite effect, i.e., decreased SD susceptibility, using chronic treatment by migraine prophylactic drugs known to suppress SD. By studying two drugs and comparing the effects of chronic treatment (i.e., mimicking migraine prophylaxis) to acute effects of a single dose, we once again get an opportunity to study a wide range of effect size. And finally in **Chapter 3C**, we take the first step towards clinical translation in a retrospective case-control study, by testing whether a history of migraine, particularly migraine with aura, accelerates infarction of ischemic penumbra in acute human stroke as well.

Lastly, **Chapter 4** puts the data generated in this thesis and all relevant clinical and experimental data from the literature together into a comprehensive review to form a synthesis that can explain the clinical association between migraine and stroke. Novel hypotheses and potential experimental approaches to test them are proposed to elucidate potential mechanisms.

REFERENCES

1. Goadsby PJ, Lipton RB, Ferrari MD. Migraine--current understanding and treatment. *N Engl J Med*. 2002;346:257-270
2. Jensen R, Stovner LJ. Epidemiology and comorbidity of headache. *Lancet Neurol*. 2008;7:354-361
3. Victor TW, Hu X, Campbell JC, Buse DC, Lipton RB. Migraine prevalence by age and sex in the united states: A life-span study. *Cephalalgia*. 2010;30:1065-1072
4. Headache Classification Committee of the International Headache S. The international classification of headache disorders, 3rd edition (beta version). *Cephalalgia*. 2013;33:629-808
5. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (ylds) for 1160 sequelae of 289 diseases and injuries 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012;380:2163-2196
6. Silberstein SD. Migraine. *Lancet*. 2004;363:381-391
7. Leao A. Spreading depression of activity in the cerebral cortex. *J Neurophysiol* 1944;7:359-390
8. Ayata C. Cortical spreading depression triggers migraine attack: Pro. *Headache*. 50:725-730

9. Eikermann-Haerter K, Ayata C. Cortical spreading depression and migraine. *Curr Neurol Neurosci Rep.* 2010;10:167-173
10. Ayata C. Spreading depression: From serendipity to targeted therapy in migraine prophylaxis. *Cephalalgia.* 2009;29:1095-1114
11. Ayata C. Cortical spreading depression triggers migraine attack: Pro. *Headache.* 2010;50:725-730
12. Pietrobon D, Moskowitz MA. Pathophysiology of migraine. *Annu Rev Physiol.* 2012
13. Moskowitz MA, Bolay H, Dalkara T. Deciphering migraine mechanisms: Clues from familial hemiplegic migraine genotypes. *Ann Neurol.* 2004;55:276-280
14. Ferrari MD, Klever RR, Terwindt GM, Ayata C, van den Maagdenberg AM. Migraine pathophysiology: Lessons from mouse models and human genetics. *Lancet Neurol.* 2015;14:65-80
15. Lashley KS. Patterns of cerebral integration indicated by the scotomas of migraine. *Archives of neurology and psychiatry.* 1941;46:331-339
16. Leao AAP. Spreading depression of activity in cerebral cortex. *J Neurophysiol.* 1944;7:359-390
17. Leao AAP. Spreading depression of activity in cerebral cortex. *J Neurophysiol.* 1944;7:359-390
18. Matsuura T, Bures J. The minimum volume of depolarized neural tissue required for triggering cortical spreading depression in rat. *Exp Brain Res.* 1971;12:238-249
19. Lothman E, Lamanna J, Cordingley G, Rosenthal M, Somjen G. Responses of electrical potential, potassium levels, and oxidative metabolic activity of the cerebral neocortex of cats. *Brain Res.* 1975;88:15-36
20. Grafstein B. Mechanism of spreading cortical depression. *J Neurophysiol.* 1956;19:154-171
21. Heinemann U, Lux HD. Ceiling of stimulus induced rises in extracellular potassium concentration in the cerebral cortex of cat. *Brain Res.* 1977;120:231-249
22. Hansen AJ, Zeuthen T. Extracellular ion concentrations during spreading depression and ischemia in the rat brain cortex. *Acta Physiol Scand.* 1981;113:437-445
23. Reid KH, Marrannes R, Wauquier A. Spreading depression and central nervous system pharmacology. *J Pharmacol Methods.* 1988;19:1-21
24. Nicholson C, Kraig RP. The behaviour of extracellular ions during spreading depression. In: Zeuthen T, ed. *The application of ion-selective microelectrodes.* Amsterdam: Elsevier; 1981:217-238.
25. Hansen A, Lauritzen M, Tfelt-Hansen P. Spreading cortical depression and antimigraine drugs. In: Amery WK, Van Nueten JM, Waquier A, eds. *The pharmacological basis of migraine therapy.* London: Putman; 1984:161-170.
26. Bures J, Buresova O, Krivanek J. *The mechanism and applications of leao's spreading depression of electroencephalographic activity.* New York: Academic Press; 1974.
27. Somjen GG. Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol Rev.* 2001;81:1065-1096
28. Katzman R. Maintenance of a constant brain extracellular potassium. *Fed Proc.* 1976;35:1244-1247
29. Vyskocil F, Kritiz N, Bures J. Potassium-selective microelectrodes used for measuring the extracellular brain potassium during spreading depression and anoxic depolarization in rats. *Brain Res.* 1972;39:255-259
30. Prince DA, Lux HD, Neher E. Measurement of extracellular potassium activity in cat cortex. *Brain Res.* 1973;50:489-495
31. Brinley FJ, Jr., Kandel ER, Marshall WH. Potassium outflux from rabbit cortex during spreading depression. *J Neurophysiol.* 1960;23:246-256
32. Kraig RP, Nicholson C. Extracellular ionic variations during spreading depression. *Neuroscience.* 1978;3:1045-1059

33. Nicholson C, ten Bruggencate G, Stockle H, Steinberg R. Calcium and potassium changes in extracellular microenvironment of cat cerebellar cortex. *J Neurophysiol.* 1978;41:1026-1039
34. Hansen AJ, Quistorff B, Gjedde A. Relationship between local changes in cortical blood flow and extracellular k^+ during spreading depression. *Acta Physiol Scand.* 1980;109:1-6
35. Freygang WH, Jr., Landau WM. Some relations between resistivity and electrical activity in the cerebral cortex of the cat. *Journal of cellular physiology.* 1955;45:377-392
36. Van Harrevel A, Ochs S. Electrical and vascular concomitants of spreading depression. *Am J Physiol.* 1957;189:159-166
37. Van Harrevel A. Changes in the diameter of apical dendrites during spreading depression. *Am J Physiol.* 1958;192:457-463
38. Van Harrevel A, Khattab FI. Changes in cortical extracellular space during spreading depression investigated with the electron microscope. *J Neurophysiol.* 1967;30:911-929
39. Nicholson C, Kraig RP. Chloride and potassium changes measured during spreading depression in catfish cerebellum. *Brain Res.* 1975;96:384-389
40. Van Harrevel A, Schade JP. Chloride movements in cerebral cortex after circulatory arrest and during spreading depression. *J Cell Comp Physiol.* 1959;54:65-84
41. Phillips JM, Nicholson C. Anion permeability in spreading depression investigated with ion-sensitive microelectrodes. *Brain Res.* 1979;173:567-571
42. Hansen AJ, Olsen CE. Brain extracellular space during spreading depression and ischemia. *Acta Physiol Scand.* 1980;108:355-365
43. Nicholson C, Bruggencate GT, Steinberg R, Stockle H. Calcium modulation in brain extracellular microenvironment demonstrated with ion-selective micropipette. *Proc Natl Acad Sci U S A.* 1977;74:1287-1290
44. Gido G, Katsura K, Kristian T, Siesjo BK. Influence of plasma glucose concentration on rat brain extracellular calcium transients during spreading depression. *J Cereb Blood Flow Metab.* 1993;13:179-182
45. Molchanova S, Koobi P, Oja SS, Saransaari P. Interstitial concentrations of amino acids in the rat striatum during global forebrain ischemia and potassium-evoked spreading depression. *Neurochem Res.* 2004;29:1519-1527
46. Fabricius M, Jensen LH, Lauritzen M. Microdialysis of interstitial amino acids during spreading depression and anoxic depolarization in rat neocortex. *Brain Res.* 1993;612:61-69
47. Moghaddam B, Schenk JO, Stewart WB, Hansen AJ. Temporal relationship between neurotransmitter release and ion flux during spreading depression and anoxia. *Can J Physiol Pharmacol.* 1987;65:1105-1110
48. Rice ME, Nicholson C. Interstitial ascorbate in turtle brain is modulated by release and extracellular volume change. *J Neurochem.* 1987;49:1096-1104
49. Szerb JC. Glutamate release and spreading depression in the fascia dentata in response to microdialysis with high k^+ : Role of glia. *Brain Res.* 1991;542:259-265
50. Grafstein B. Locus of propagation of spreading cortical depression. *J Neurophysiol.* 1956;19:308-316
51. Mody I, Lambert JD, Heinemann U. Low extracellular magnesium induces epileptiform activity and spreading depression in rat hippocampal slices. *J Neurophysiol.* 1987;57:869-888
52. Marrannes R, Willems R, De Prins E, Wauquier A. Evidence for a role of the n-methyl-d-aspartate (nmda) receptor in cortical spreading depression in the rat. *Brain Res.* 1988;457:226-240
53. Richter F, Bauer R, Lehmenkuhler A, Schaible HG. Spreading depression in the brainstem of the adult rat: Electrophysiological parameters and influences on regional brainstem blood flow. *J Cereb Blood Flow Metab.* 2007

54. Richter F, Rupprecht S, Lehmenkuhler A, Schaible HG. Spreading depression can be elicited in brainstem of immature but not adult rats. *J Neurophysiol.* 2003
55. Fifkova E. Spreading depression in subcortical structures in rabbit. *Brain Res.* 1966;2:61-70
56. Fifkova E, Bures J. Spreading depression in the mammalian striatum. *Arch Int Physiol Biochim.* 1964;72:171-179
57. Fifkova E. Thalamic spreading depression in the rat. *Electroencephalogr Clin Neurophysiol.* 1966;20:68-76
58. Vinogradova LV, Koroleva VI, Bures J. Re-entry waves of leao's spreading depression between neocortex and caudate nucleus. *Brain Res.* 1991;538:161-164
59. Leao AAP, Morison RS. Propagation of spreading cortical depression. *J Neurophysiol.* 1945;8:33-45
60. Milner PM. Note on a possible correspondence between the scotomas of migraine and spreading depression of leao. *Electroencephalogr Clin Neurophysiol.* 1958;10:705
61. Lauritzen M. On the possible relation of spreading cortical depression to classical migraine. *Cephalalgia.* 1985;5 Suppl 2:47-51
62. Moskowitz MA, Nozaki K, Kraig RP. Neocortical spreading depression provokes the expression of c-fos protein-like immunoreactivity within trigeminal nucleus caudalis via trigeminovascular mechanisms. *J Neurosci.* 1993;13:1167-1177
63. Bolay H, Reuter U, Dunn AK, Huang Z, Boas DA, Moskowitz MA. Intrinsic brain activity triggers trigeminal meningeal afferents in a migraine model. *Nat Med.* 2002;8:136-142
64. Buzzi MG, Bonamini M, Moskowitz MA. Neurogenic model of migraine. *Cephalalgia.* 1995;15:277-280
65. Wahl M, Schilling L, Parsons AA, Kaumann A. Involvement of calcitonin gene-related peptide (cgrp) and nitric oxide (no) in the pial artery dilatation elicited by cortical spreading depression. *Brain Res.* 1994;637:204-210
66. Read SJ, Smith MI, Hunter AJ, Parsons AA. The dynamics of nitric oxide release measured directly and in real time following repeated waves of cortical spreading depression in the anaesthetised cat. *Neurosci Lett.* 1997;232:127-130
67. Reuter U, Weber JR, Gold L, Arnold G, Wolf T, Dreier J, et al. Perivascular nerves contribute to cortical spreading depression-associated hyperemia in rats. *Am J Physiol.* 1998;274:H1979-1987
68. Ayata C, Jin H, Kudo C, Dalkara T, Moskowitz MA. Suppression of cortical spreading depression in migraine prophylaxis. *Ann Neurol.* 2006;59:652-661
69. Ayata C. Pearls and pitfalls in experimental models of spreading depression. *Cephalalgia.* 2013;33:604-613
70. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the global burden of disease study 2010. *Lancet.* 2012;380:2095-2128
71. Emberson J, Lees KR, Lyden P, Blackwell L, Albers G, Bluhmki E, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: A meta-analysis of individual patient data from randomised trials. *Lancet.* 2014
72. Dohmen C, Sakowitz OW, Fabricius M, Bosche B, Reithmeier T, Ernestus RI, et al. Spreading depolarizations occur in human ischemic stroke with high incidence. *Ann Neurol.* 2008;63:720-728
73. Kurth T, Diener HC. Migraine and stroke: Perspectives for stroke physicians. *Stroke.* 2012;43:3421-3426
74. Bigal ME, Kurth T, Hu H, Santanello N, Lipton RB. Migraine and cardiovascular disease: Possible mechanisms of interaction. *Neurology.* 2009;72:1864-1871

75. Kurth T, Chabriat H, Boussier MG. Migraine and stroke: A complex association with clinical implications. *Lancet Neurol.* 2012;11:92-100
76. Pezzini A, Del Zotto E, Giossi A, Volonghi I, Costa P, Dalla Volta G, et al. The migraine-ischemic stroke relation in young adults. *Stroke Res Treat.* 2010;2011:304921
77. Spector JT, Kahn SR, Jones MR, Jayakumar M, Dalal D, Nazarian S. Migraine headache and ischemic stroke risk: An updated meta-analysis. *Am J Med.* 2010;123:612-624
78. Kruit MC, van Buchem MA, Hofman PA, Bakkers JT, Terwindt GM, Ferrari MD, et al. Migraine as a risk factor for subclinical brain lesions. *JAMA.* 2004;291:427-434
79. Bashir A, Lipton RB, Ashina S, Ashina M. Migraine and structural changes in the brain: A systematic review and meta-analysis. *Neurology.* 2013;81:1260-1268
80. Etminan M, Takkouche B, Isorna FC, Samii A. Risk of ischaemic stroke in people with migraine: Systematic review and meta-analysis of observational studies. *BMJ.* 2005;330:63
81. Schurks M, Rist PM, Bigal ME, Buring JE, Lipton RB, Kurth T. Migraine and cardiovascular disease: Systematic review and meta-analysis. *BMJ.* 2009;339:b3914
82. Kurth T, Gaziano JM, Cook NR, Logroscino G, Diener HC, Buring JE. Migraine and risk of cardiovascular disease in women. *JAMA.* 2006;296:283-291
83. MacClellan LR, Giles W, Cole J, Wozniak M, Stern B, Mitchell BD, et al. Probable migraine with visual aura and risk of ischemic stroke: The stroke prevention in young women study. *Stroke.* 2007;38:2438-2445
84. Kruit MC, Launer LJ, Ferrari MD, van Buchem MA. Infarcts in the posterior circulation territory in migraine. The population-based mri camera study. *Brain.* 2005;128:2068-2077
85. Palm-Meinders IH, Koppen H, Terwindt GM, Launer LJ, Konishi J, Moonen JM, et al. Structural brain changes in migraine. *JAMA.* 308:1889-1897
86. Scher AI, Gudmundsson LS, Sigurdsson S, Ghambaryan A, Aspelund T, Eiriksdottir G, et al. Migraine headache in middle age and late-life brain infarcts. *JAMA.* 2009;301:2563-2570
87. Kurth T, Mohamed S, Maillard P, Zhu YC, Chabriat H, Mazoyer B, et al. Headache, migraine, and structural brain lesions and function: Population based epidemiology of vascular ageing-mri study. *BMJ.* 2011;342:c7357
88. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart disease and stroke statistics--2014 update: A report from the american heart association. *Circulation.* 2014;129:e28-e292
89. Eising E, de Vries B, Ferrari MD, Terwindt GM, van den Maagdenberg AM. Pearls and pitfalls in genetic studies of migraine. *Cephalalgia.* 2013;33:614-625
90. Arboleda-Velasquez JF, Manent J, Lee JH, Tikka S, Ospina C, Vanderburg CR, et al. Hypomorphic notch 3 alleles link notch signaling to ischemic cerebral small-vessel disease. *Proc Natl Acad Sci U S A.* 2011;108:E128-135
91. Eikermann-Haerter K, Yuzawa I, Dilekoz E, Joutel A, Moskowitz MA, Ayata C. Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy syndrome mutations increase susceptibility to spreading depression. *Ann Neurol.* 2011;69:413-418
92. Russell MB, Ducros A. Sporadic and familial hemiplegic migraine: Pathophysiological mechanisms, clinical characteristics, diagnosis, and management. *Lancet Neurol.* 2011;10:457-470
93. Tottene A, Conti R, Fabbro A, Vecchia D, Shapovalova M, Santello M, et al. Enhanced excitatory transmission at cortical synapses as the basis for facilitated spreading depression in ca(v)2.1 knockin migraine mice. *Neuron.* 2009;61:762-773
94. Kors EE, Terwindt GM, Vermeulen FL, Fitzsimons RB, Jardine PE, Heywood P, et al. Delayed cerebral edema and fatal coma after minor head trauma: Role of the cacna1a calcium channel subunit gene and relationship with familial hemiplegic migraine. *Ann Neurol.* 2001;49:753-760

95. Stam AH, Lujckx CJ, Poll-The BT, Ginjaar IB, Frants RR, Haan J, et al. Early seizures and cerebral oedema after trivial head trauma associated with the cacna1a s218l mutation. *J Neurol Neurosurg Psychiatry*. 2009;80:1125-1129
96. Ophoff RA, Terwindt GM, Vergouwe MN, van Eijk R, Oefner PJ, Hoffman SM, et al. Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the ca²⁺ channel gene cacn1a4. *Cell*. 1996;87:543-552
97. Pietrobon D. Function and dysfunction of synaptic calcium channels: Insights from mouse models. *Current opinion in neurobiology*. 2005;15:257-265

