

Anomalous diffusion of Dirac fermions

Groth, C.W.

Citation

Groth, C. W. (2010, December 8). Anomalous diffusion of Dirac fermions. Casimir PhD Series. Retrieved from https://hdl.handle.net/1887/16222

Version:	Not Applicable (or Unknown)
License:	Leiden University Non-exclusive license
Downloaded from:	https://hdl.handle.net/1887/16222

Note: To cite this publication please use the final published version (if applicable).

5 Switching of electrical current by spin precession in the first Landau level of an inverted-gap semiconductor

5.1 Introduction

A central goal of spin-transport electronics (or spintronics) is the ability to switch current between spin-selective electrodes by means of spin precession [151]. In the original Datta-Das proposal for such a spin-based transistor [38], the current which is switched carries both spin and charge. It has proven difficult to separate the effects of spin precession from purely orbital effects (deflection of electron trajectories), so most succesful implementations use a nonlocal geometry [60] to modulate the spin current at zero charge current [58, 85, 139]. Even in the absence of an orbital effect, the fact that different electrons (moving along different trajectories) experience different amounts of spin precession prevents a complete switching of the current from one electrode to the other.

If the electron motion could somehow be confined to a single spatial dimension, it would be easier to isolate spin effects from orbital effects and to ensure that all electron spins precess by the same amount. Complete switching of the current would then be possible, limited only by spin relaxation processes. Edge state transport in the quantum Hall effect is one-dimensional and spin selective (in sufficiently strong perpendicular magnetic fields B_{\perp}), but spin precession plays no role in the traditional experiments



Figure 5.1: Top panel: Schematic illustration of the onedimensional pathway along which the electron spin is injected, precessed, and detected (filled circles: occupied states; open circles: empty states). Bottom panel: Potential profile of the *p*-*n* junction, shown for $B_{\perp} > B_c$ (for $B_{\perp} < B_c$ the labels E_+ and E_- should be interchanged).

on a two-dimensional electron gas [17]. In this paper we show how the quantum Hall effect in an inverted-gap semiconductor offers the unique possibility to perform a one-dimensional spin precession experiment.

The key idea is to combine the spin-selectivity of edge states with free precession along a *p*-*n* interface. The geometry, shown in Fig. 5.1, has been studied in graphene [146, 2, 105, 142] – but there spin is only weakly coupled to the orbit and plays a minor role [62, 1]. The strong spin-orbit coupling in inverted-gap semiconductors splits the first Landau level into a pair of levels E_{\pm} of opposite magnetic moment [74, 123]. One level E_{+} (say, with spin up) has electron-like character and produces edge states in the

conduction band. The other level E_- (with spin down) has holelike character and produces edge states in the valence band. The edge states from E_+ and E_- have opposite chirality, meaning that one circulates clockwise along the edge while the other circulates counter-clockwise. These spin-selective, chiral edge states provide the spin injection at x = 0 and detection at x = W.

For the spin precession we need to combine states from E_+ and E_- . This is achieved by means of a gate electrode, which creates a smooth potential step (height U_0 , width d) centered at y = 0, such that the Fermi level lies in the conduction band for y < 0 (*n*-doped region) and in the valence band for y > 0 (*p*-doped region). At the *p*-*n* interface states from the first Landau levels E_+ and E_- overlap at the Fermi energy E_F , to form a spin-degenerate one-dimensional state. Spin precession can be realized externally by a parallel magnetic field B_{\parallel} (in the x - y plane) or internally by bulk or structure inversion asymmetry [74].

Good overlap at E_F of the states from E_+ and E_- is crucial for effective spin precession. The requirement is that the spatial separation $\delta y \simeq |E_+ - E_-|d/U_0$ of the states should be small compared to the magnetic length $l_m = (\hbar/eB_\perp)^{1/2}$ (which sets their spatial extent). This is where the inverted gap comes in, as we now explain.

Inversion of the gap means that the first Landau level in the conduction band goes down in energy with increasing magnetic field (because it has hole-like character), while the first Landau level in the valence band goes up in energy (because it has electron-like character). As a consequence, the gap $|E_+ - E_-|$ has a minimal value E_c much less than the cyclotron energy $\hbar\omega_c$ at a crossover magnetic field B_c . Indeed, $E_c = 0$ in the absence of inversion asymmetry [74]. Good overlap can therefore be reached in an inverted-gap semiconductor, simply by tuning the magnetic field. In a normal (non-inverted) semiconductor, such as GaAs, the cyclotron energy difference between E_+ and E_- effectively prevents the overlap of Landau levels from conduction and valence bands.

In the following two sections, we first present a general, model independent analysis and then specialize to the case of a HgTe quantum well (where we test the analytical theory by computer simulation).

5.2 General theory

We introduce a one-dimensional coordinate s_{\pm} along the E_{\pm} edge states, increasing in the direction of the chirality (see Fig. 5.1). The wave amplitudes $\psi_{\pm}(s_{\pm})$ of these two states can be combined into the spinor $\Psi = (\psi_+, \psi_-)$. Far from the *p*-*n* interface, ψ_+ and ψ_- evolve independently with Hamiltonian

$$H_0 = \begin{pmatrix} H_+ & 0\\ 0 & H_- \end{pmatrix}, \quad H_\pm = v_\pm \left(-i\hbar \frac{\partial}{\partial s_\pm} - p_F^\pm \right). \tag{5.1}$$

This is the generic linearized Hamiltonian of a chiral mode, with group velocity $v_{\pm} \equiv v(s_{\pm})$ and Fermi momentum $p_F^{\pm} \equiv p_F(s_{\pm})$. Near the *p*-*n* interface the spin-up and spin-down states are coupled by the generic precession Hamiltonian,

$$H_{\rm prec} = \begin{pmatrix} 0 & \mathcal{M}^* \\ \mathcal{M} & 0 \end{pmatrix}, \qquad (5.2)$$

with a matrix element \mathcal{M} to be specified later.

We seek the transfer matrix *T*, defined by

$$\Psi(s_{+}^{f}, s_{-}^{f}) = T\Psi(s_{+}^{i}, s_{-}^{i}).$$
(5.3)

We take for Ψ a solution of the Schrödinger equation,

$$(H_0 + H_{\rm prec})\Psi = 0,$$
 (5.4)

at zero excitation energy (appropriate for electrical conduction in linear response). The initial and final points s_{\pm}^{i} and s_{\pm}^{f} are taken away from the *p*-*n* interface. The unitary scattering matrix *S* (relating incident and outgoing current amplitudes) is related to *T* by a similarity transformation,

$$S = \begin{pmatrix} v_{+}^{f} & 0\\ 0 & v_{-}^{f} \end{pmatrix}^{1/2} T \begin{pmatrix} v_{+}^{i} & 0\\ 0 & v_{-}^{i} \end{pmatrix}^{-1/2}.$$
 (5.5)

The two-terminal linear-response conductance G of the p-n junction is given by the Landauer formula,

$$G = \frac{e^2}{h} |S_{21}|^2.$$
(5.6)

The transition matrix element $\mathcal{M}(s_+, s_-)$ between $\psi_+(s_+)$ and $\psi_-(s_-)$ vanishes if the separation $|s_+ - s_-|$ of the two states is large compared to the magnetic length l_m . We assume that B_{\perp} is sufficiently close to B_c that $|s_+ - s_-| < l_m$ at the *p*-*n* interface y = 0, 0 < x < W, where we may take $\mathcal{M} = \text{constant}$ (independent of *x*). At the two edges x = 0 and x = W we set $\mathcal{M} = 0$, neglecting the crossover region within l_m of (0,0) and (W,0). (The precession angle there will be small compared to unity for $l_m \ll \hbar v_{\pm}/|\mathcal{M}|$.)

In this "abrupt approximation" we may identify the initial and final coordinates s^i_{\pm} and s^f_{\pm} with the points (0,0) and (W,0), at the two ends of the *p*-*n* interface. Integration of the Schrödinger equation (5.4) along the *p*-*n* interface gives the transfer matrix, and application of Eq. (5.5) then gives the scattering matrix

$$S = \exp\left[-i\frac{W}{\hbar} \begin{pmatrix} p_F^+ & \mathcal{M}^*/\sqrt{v_+v_-}\\ \mathcal{M}/\sqrt{v_+v_-} & p_F^- \end{pmatrix}\right].$$
 (5.7)

(We have assumed that v_{\pm} and p_F^{\pm} , as well as \mathcal{M} , do not vary along the *p*-*n* interface, so we may omit the labels *i*, *f*.) One verifies that *S* is unitary, as it should be.

Evaluation of the matrix exponent in Eq. (5.7) and substitution into Eq. (5.6) gives the conductance,

$$G = \frac{e^2}{h} \sin^2 \left(\frac{|\mathbf{p}_{\text{eff}}|W}{\hbar} \right) \sin^2 \alpha.$$
 (5.8)

The effective precession momentum

$$\boldsymbol{p}_{\text{eff}} = \left(\frac{\operatorname{Re}\mathcal{M}}{\bar{v}}, \frac{\operatorname{Im}\mathcal{M}}{\bar{v}}, \frac{\delta p_F}{2}\right)$$
(5.9)

(with $\delta p_F = p_F^+ - p_F^-$ and $\bar{v} = \sqrt{v_+v_-}$) makes an angle α with the *z*-axis. This is the final result of our general analysis.

5.3 Application to a HgTe quantum well

We now turn to a specific inverted-gap semiconductor, a quantum well consisting of a 7 nm layer of HgTe sandwiched symmetrically between $Hg_{0.3}Cd_{0.7}Te$ [73]. The properties of this socalled topological insulator have been reviewed in [74]. The low-energy excitations are described by a four-orbital tight-binding Hamiltonian [23, 44],

$$H = \sum_{n} c_{n}^{\dagger} \mathcal{E}_{n} c_{n} - \sum_{n,m \text{ (nearest neighb.)}} c_{n}^{\dagger} \mathcal{T}_{nm} c_{m}.$$
 (5.10)

Each site *n* on a square lattice (lattice constant a = 4 nm) has four states $|s, \pm\rangle$, $|p_x \pm ip_y, \pm\rangle$ – two electron-like *s*-orbitals and two hole-like *p*-orbitals of opposite spin $\sigma = \pm$. Annihilation operators $c_{n,\tau\sigma}$ for these four states (with $\tau \in \{s, p\}$) are collected in a vector

$$c_n = (c_{n,s+}, c_{n,p+}, c_{n,s-}, c_{n,p-}).$$

States on the same site are coupled by the 4×4 potential matrix \mathcal{E}_n and states on adjacent sites by the 4×4 hopping matrix \mathcal{T}_{nm} .

In zero magnetic field and without inversion asymmetry H decouples into a spin-up block H_+ and a spin-down block H_- , defined in terms of the 2 × 2 matrices

$$\mathcal{E}_{n}^{+} = \mathcal{E}_{n}^{-} = \operatorname{diag}\left(\varepsilon_{s} - U_{n}, \varepsilon_{p} - U_{n}\right), \tag{5.11}$$

$$\mathcal{T}_{nm}^{+} = \left(\mathcal{T}_{nm}^{-}\right)^{*} = \begin{pmatrix} t_{ss} & t_{sp}e^{i\theta_{nm}} \\ t_{sp}e^{-i\theta_{mn}} & -t_{pp} \end{pmatrix}.$$
 (5.12)

Here U_n is the electrostatic potential and θ_{nm} is the angle between the vector $r_n - r_m$ and the positive *x*-axis (so $\theta_{mn} = \pi - \theta_{nm}$). The orbital effect of a perpendicular magnetic field B_{\perp} is introduced into the hopping matrix elements by means of the Peierls substitution

$$\mathcal{T}_{nm} \mapsto \mathcal{T}_{nm} \exp[i(eB_{\perp}/\hbar)(y_n - y_m)x_n].$$

This breaks the degeneracy of the spin-up and spin-down energy levels, but it does not couple them.

Spin-up and spin-down states are coupled by the Zeeman effect from a parallel magnetic field (with gyromagnetic factor g_{\parallel}) and by spin-orbit interaction without inversion symmetry (parameterized by a vector Δ). In first-order perturbation theory, the correction $\delta \mathcal{E}$ to the on-site potential has the form [74]

$$\delta \mathcal{E} = (\mathbf{\Delta} \cdot \boldsymbol{\sigma}) \otimes \tau_{y} + \frac{1}{2} \mu_{B} g_{\parallel} (\boldsymbol{B}_{\parallel} \cdot \boldsymbol{\sigma}) \otimes (\tau_{0} + \tau_{z}) + \mu_{B} B_{\perp} \sigma_{z} \otimes (\bar{g}_{\perp} \tau_{0} + \delta g_{\perp} \tau_{z}).$$
(5.13)

The Pauli matrices $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ act on the spin-up and spindown blocks, while the Pauli matrices τ_y, τ_z and the unit matrix τ_0 act on the orbital degree of freedom *s*, *p* within each block.

The parameters of the tight-binding model for a 7 nm thick HgTe/Hg_{0.3}Cd_{0.7}Te quantum well (grown in the (001) direction) are as follows [74]: $t_{ss} = 74.9$ meV, $t_{pp} = 10.9$ meV, $t_{sp} = 45.6$ meV, $\varepsilon_s = 289.5$ meV, $\varepsilon_p = -33.5$ meV, $\bar{g}_{\perp} = 10.75$, $\delta g_{\perp} = 11.96$, $g_{\parallel} = -20.5$, $\Delta = (0, 1.6$ meV, 0).

The quantum well is symmetric, so only bulk inversion asymmetry contributes to Δ . The *p*-*n* junction is defined by the potential profile

$$U(x,y) = \frac{1}{2}U_0[1 + \tanh(4y/d)], \quad 0 < x < W,$$
(5.14)

with $U_0 = 32 \text{ meV}$, d = 12 nm, and $W = 0.8 \mu \text{m}$. We fix the Fermi level at $E_F = 25 \text{ meV}$, so that it lies in the conduction band for y < 0 and in the valence band for y > 0. (We have checked that none of the results are sensitive to the choice of potential profile or parameter values.) The scattering matrix of the *p*-*n* junction is calculated with the recursive Green function technique, using the "knitting" algorithm¹ of Ref. [65]. Results for *G* as a function of B_{\parallel} are shown in Figs. 5.2 and 5.3.

The dependence of the conductance on the parallel magnetic field B_{\parallel} shows a striking "bullseye" pattern, which can be understood

¹ The computer code for the knitting algorithm was kindly provided to us by Dr. Waintal.



Figure 5.2: Dependence of the conductance of the HgTe quantum well on the parallel magnetic field B_{\parallel} , calculated from the tight-binding model for $B_{\perp} = B_c = 6.09$ T.

as follows. To first order in B_{\parallel} , the edge state parameters v_{\pm} and p_F^{\pm} are constant, while the precession matrix element

$$\mathcal{M} = \Delta_{\text{eff}} + \mu_B g_{\text{eff}} (B_{\parallel x} + iB_{\parallel y}) \tag{5.15}$$

varies linearly. Substitution into Eqs. (5.8) and (5.9) gives a circularly symmetric dependence of *G* on B_{\parallel} ,

$$G = \frac{e^2}{h} \left(1 + \frac{(\bar{v}\delta p_F)^2}{4|\mu_B g_{\text{eff}}|^2 |\mathbf{B}_{\parallel} - \mathbf{B}_0|^2} \right)^{-1} \times \sin^2 \left[\frac{W}{\hbar \bar{v}} \sqrt{|\mu_B g_{\text{eff}}|^2 |\mathbf{B}_{\parallel} - \mathbf{B}_0|^2 + \frac{1}{4} (\bar{v}\delta p_F)^2} \right], \quad (5.16)$$

$$\boldsymbol{B}_{0} = \boldsymbol{\mu}_{B}^{-1} \big(\operatorname{Re}[\Delta_{\mathrm{eff}}/g_{\mathrm{eff}}], \operatorname{Im}[\Delta_{\mathrm{eff}}/g_{\mathrm{eff}}], 0 \big).$$
(5.17)

The parallel magnetic field B_0 corresponds to the center of the bullseye, at which the coupling between the \pm edge states along



Figure 5.3: Dependence of the conductance on $B_{\parallel y}$ for $B_{\parallel x} = 0$, at three values of the perpendicular magnetic field. The solid curves are calculated numerically from the tight-binding model, the dashed curves are the analytical prediction (5.16). The arrow indicates the value of B_0 from Eq. (5.17). (Only the numerical curve is shown in the upper panel, because the analytical curve is nearly indistinguishable from it.)

the *p*-*n* interface by bulk inversion asymmetry is cancelled by the Zeeman effect.

The Fermi momentum mismatch δp_F vanishes at a perpendicular magnetic field B^* close to, but not equal to, B_c . Then the

magnetoconductance oscillations are purely sinusoidal,

$$G = \frac{e^2}{h} \sin^2[(W/\hbar\bar{v})\mu_B g_{\text{eff}}|\boldsymbol{B}_{\parallel} - \boldsymbol{B}_0|].$$
 (5.18)

For a quantitative comparison between numerics and analytics, we extract the parameters v_{\pm} and p_F^{\pm} from the dispersion relation of the edge states ψ_{\pm} along an infinitely long *p*-*n* interface (calculated for uncoupled blocks H_{\pm}). The overlap of ψ_{\pm} and ψ_{-} determines the coefficients

$$\Delta_{\rm eff} = (\Delta_x + i\Delta_y) \langle \psi_- | \tau_y | \psi_+ \rangle, \qquad (5.19)$$

$$g_{\text{eff}} = \frac{1}{2} g_{\parallel} \langle \psi_{-} | \tau_{0} + \tau_{z} | \psi_{+} \rangle.$$
(5.20)

For $B_{\perp} = B_c = 6.09 \text{ T}$ we find $\bar{v}\delta p_F = 0.86 \text{ meV}$, $\hbar \bar{v}/W = 0.23 \text{ meV}$, $\Delta_{\text{eff}} = -1.59 \text{ meV}$, $g_{\text{eff}} = -4.99$. The Fermi momentum mismatch δp_F vanishes for $B_{\perp} = B^* = 5.77 \text{ T}$. Substitution of the parameters into Eq. (5.16) gives the dashed curves in Fig. 5.3, in reasonable agreement with the numerical results from the tight-binding model (solid curves). In particular, the value of B_0 extracted from the numerics is within a few percent of the analytical prediction (5.17).

Because of the one-dimensionality of the motion along the *p*-*n* interface, electrostatic disorder and thermal averaging have a relatively small perturbing effect on the conductance oscillations. For disorder potentials ΔU and thermal energies $k_B T$ up to 10% of U_0 the perturbation is hardly noticeable (a few percent). As shown in Fig. 5.4, the conductance oscillations remain clearly visible even for ΔU and $k_B T$ comparable to U_0 . In particular, we have found that the center of the bullseye pattern remains within 10% of B_0 even for ΔU as large as the *p*-*n* step height U_0 .

5.4 Conclusion

In conclusion, we have proposed a one-dimensional spin precession experiment at a p-n junction in an inverted-gap semiconductor. The



Figure 5.4: The solid blue curve in both panels is the same as in Fig. 5.3, top panel, calculated for $B_{\perp} = B^*$ from the tight-binding model at zero temperature without any disorder. The dotted black curve in the lower panel shows the effect of raising the temperature to $30 \text{ K} \approx U_0/3k_B$. The dotted red curve and dashed green curve in the upper panel show the effect of disorder at zero temperature. The on-site disorder potential is drawn uniformly from the interval $(-\Delta U_0, \Delta U_0)$, with, respectively, $\Delta U = U_0/4$ and $\Delta U = U_0/2$.

conductance as a function of parallel magnetic field oscillates in a bullseye pattern, centered at a field B_0 proportional to the matrix element Δ_{eff} of the bulk inversion asymmetry. Our numerical and analytical calculations show conductance oscillations of amplitude not far below e^2/h , robust to disorder and thermal averaging. Realization of the proposed experiment in a HgTe quantum well [74] (or in other inverted-gap semiconductors [82]) would provide a

unique demonstration of full-current switching by spin precession.

As directions for future research, we envisage potential applications of this technique as a sensitive measurement of the degree of bulk inversion asymmetry, or as a probe of the effects of interactions on spin precession. It might also be possible to eliminate the external magnetic field and realize electrical switching of the current in our setup: The role of the perpendicular magnetic field in producing spin-selective edge states can be taken over by magnetic impurities or a ferromagnetic layer [83], while the role of the parallel magnetic field in providing controlled spin precession can be taken over by gate-controlled structural inversion asymmetry.