



Universiteit
Leiden
The Netherlands

Physiotherapy in Parkinson's disease. towards evidence-based practice.

Keus, S.H.J.

Citation

Keus, S. H. J. (2010, April 29). *Physiotherapy in Parkinson's disease. towards evidence-based practice*. Retrieved from <https://hdl.handle.net/1887/15341>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/15341>

Note: To cite this publication please use the final published version (if applicable).

R

Reference List

- (1) De Lau LM, Koudstaal PJ, Hofman A, Breteler MM. [Parkinson disease is more prevalent than people think. Research results]. *Ned Tijdschr Geneesk* 2009; 153(3):63-68.
- (2) von Campenhausen S, Bornschein B, Wick R, Botzel K, Sampaio C, Poewe W et al. Prevalence and incidence of Parkinson's disease in Europe. *Eur Neuropsychopharmacol* 2005; 15(4):473-490.
- (3) Dorsey ER, Constantinescu R, Thompson JP, Biglan KM, Holloway RG, Kieburtz K et al. Projected number of people with Parkinson disease in the most populous nations, 2005 through 2030. *Neurology* 2006; 68(5):384-386.
- (4) Rijk MCd, Tzourio C, Breteler MM, Dartigues JF, Amaducci L, Lopez-Pousa S et al. Prevalence of parkinsonism and Parkinson's disease in Europe: the EUROPARKINSON Collaborative Study. European Community Concerted Action on the Epidemiology of Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1997; 62(1):10-15.
- (5) Pressley JC, Louis ED, Tang MX, Cote L, Cohen PD, Glied S et al. The impact of comorbid disease and injuries on resource use and expenditures in parkinsonism. *Neurology* 2003; 60(1):87-93.
- (6) Schrag A, Jahanshahi M, Quinn N. How does Parkinson's disease affect quality of life? A comparison with quality of life in the general population. *Movement Disorders* 2000; 15(6):1112-1118.
- (7) Gage H, Hendricks A, Zhang S, Kazis L. The relative health related quality of life of veterans with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2003; 74(2):163-169.
- (8) Poewe W. The natural history of Parkinson's disease. *J Neurol* 2006; 253:2-6.
- (9) Obeso JA, Rodriguez-Oroz MC, Itez-Temino B, Blesa FJ, Guridi J, Marin C et al. Functional organization of the basal ganglia: therapeutic implications for Parkinson's disease. *Mov Disord* 2008; 23(Suppl 3):S548-S559.
- (10) Braak H, Del TK. Cortico-basal ganglia-cortical circuitry in Parkinson's disease reconsidered. *Exp Neurol* 2008; 212(1):226-229.
- (11) Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992; 55(3):181-184.
- (12) Hughes AJ, Daniel SE, Lees AJ. Improved accuracy of clinical diagnosis of Lewy body Parkinson's disease. *Neurology* 2001; 57(8):1497-1499.
- (13) Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. *Arch Neurol* 1999; 56(1):33-39.
- (14) Jankovic J. Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry* 2008; 79(4):368-376.
- (15) Alves G, Wentzel-Larsen T, Aarsland D, Larsen JP. Progression of motor impairment and disability in Parkinson disease: a population-based study. *Neurology* 2005; 65(9):1436-1441.
- (16) Jankovic J, McDermott M, Carter J, Gauthier S, Goetz C, Golbe L et al. Variable expression of Parkinson's disease: a base-line analysis of the DATATOP cohort. The Parkinson Study Group. *Neurology* 1990; 40(10):1529-1534.
- (17) Muslimovic D, Schmand B, Speelman JD, de Haan RJ. Course of cognitive decline in Parkinson's disease: a meta-analysis. *J Int Neuropsychol Soc* 2007; 13(6):920-932.

Reference List

- (18) Post B, Speelman JD, de Haan RJ. Clinical heterogeneity in newly diagnosed Parkinson's disease. *J Neurol* 2008; 255(5):716-722.
- (19) World Health Organization (WHO). International Classification of Functioning, Disability and Health (ICF). www.who.int/classifications/icf/en/index.html [2007; cited 2009 June 9].
- (20) WHO. International Statistical Classification of Diseases and Related Health Problems - 10th revision. <http://apps.who.int/classifications/apps/icd/icd10online/> [2007; cited 2009 June 9].
- (21) Bloem BR, Bhatia KP. Gait and balance in basal ganglia disorders. In: Bronstein AM, Brandt T, Woollacott MH, Nutt JG, editors. *Clinical disorders of balance, posture and gait*. 2nd ed. London, UK: Arnold; 2004. 173-206.
- (22) Groenewegen HJ. The basal ganglia and motor control. *Neural Plast* 2003; 10(1-2):107-120.
- (23) Morris ME, Iansek R. Characteristics of motor disturbance in Parkinson's disease and strategies for movement rehabilitation. *Human Movement Science* 1996; 15:649-669.
- (24) Hughes AJ, Daniel SE, Blankson S, Lees AJ. A clinicopathologic study of 100 cases of Parkinson's disease. *Arch Neurol* 1993; 50(2):140-148.
- (25) Rajput AH, Rozdilsky B, Rajput A. Accuracy of clinical diagnosis in parkinsonism--a prospective study. *Can J Neurol Sci* 1991; 18(3):275-278.
- (26) Stamey W, Davidson A, Jankovic J. Shoulder pain: a presenting symptom of Parkinson disease. *J Clin Rheumatol* 2008; 14(4):253-254.
- (27) Ziemssen T, Reichmann H. Non-motor dysfunction in Parkinson's disease. *Parkinsonism Relat Disord* 2007; 13(6):323-332.
- (28) Poewe W. Non-motor symptoms in Parkinson's disease. *Eur J Neurol* 2008; 15(Suppl 1):14-20.
- (29) Chaudhuri KR, Healy DG, Schapira AH. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol* 2006; 5(3):235-245.
- (30) Chaudhuri KR, Naidu Y. Early Parkinson's disease and non-motor issues. *J Neurol* 2008; 255(Suppl 5):33-38.
- (31) Naidu Y, Martinez-Martin P, Mitra T, Tluk S, Stocchi F, Macphee G et al. Non declaration of non motor symptoms (NMS) of PD: an international study using the NMSQuest. *Mov Disord* 2009; 24(Suppl 1):S386.
- (32) Keus SHJ, Nijkrake M, Bloem BR, Kwakkel G, Berendse HW, Roos RAC et al. ParkNet implementation: evaluation of health benefits. *Parkinsonism Relat Disord* 2008; 14(Suppl 1):S9.
- (33) Wimmers RH, Kamsma YPT. [A survey to assess activity limitations in patients with Parkinson's disease]. *Ned Tijdschr Fysiother* 1998; 3:54-61.
- (34) Driver JA, Kurth T, Buring JE, Gaziano JM, Logroscino G. Parkinson disease and risk of mortality: a prospective comorbidity-matched cohort study. *Neurology* 2008; 70(16):1423-1430.
- (35) Garrett NA, Brasure M, Schmitz KH, Schultz MM, Huber MR. Physical inactivity: direct cost to a health plan. *Am J Prev Med* 2004; 27(4):304-309.
- (36) Marcus BH, Williams DM, Dubbert PM, Sallis JF, King AC, Yancey AK et al. Physical activity intervention studies: what we know and what we need to know: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity); Council on Cardiovascular Disease in the Young; and the Interdisciplinary Working Group on Quality of Care and Outcomes Research. *Circulation* 2006; 114(24):2739-2752.
- (37) Benecke R, Rothwell JC, Dick JP, Day BL, Marsden CD. Disturbance of sequential movements in patients with Parkinson's disease. *Brain* 1987; 110(2):361-379.
- (38) Kamsma YPT, Brouwer WH, Lakke JPWF. Training of compensatory strategies for impaired gross motor skills in patients with Parkinson's disease. *Physiother Th Pract* 1995; 11:209-229.
- (39) Kamsma Y. Functional reorganisation of basic motor actions in Parkinson's disease. [PhD thesis Groningen University]. Groningen: YPT Kamsma; 2002.
- (40) Morris ME. Movement disorders in people with Parkinson disease: a model for physical therapy. *Phys Ther* 2000; 80(6):578-597.
- (41) Fellows SJ, Noth J, Schwarz M. Precision grip and Parkinson's disease. *Brain* 1998; 121(9):1771-1784.

- (42) Ponsen MM, Daffertshofer A, Wolters EC, Beek PJ, Berendse HW. Impairment of complex upper limb motor function in de novo Parkinson's disease. *Parkinsonism Relat Disord* 2008; 14(3):199-204.
- (43) Giladi N, McDermott MP, Fahn S, Przedborski S, Jankovic J, Stern M et al. Freezing of gait in PD: prospective assessment in the DATATOP cohort. *Neurology* 2001; 56(12):1712-1721.
- (44) Bloem BR, Hausdorff JM, Visser JE, Giladi N. Falls and freezing of gait in Parkinson's disease: A review of two interconnected, episodic phenomena. *Mov Disord* 2004; 19(8):871-884.
- (45) Nieuwboer A, Giladi N. The challenge of evaluating freezing of gait in patients with Parkinson's disease. *Br J Neurosurg* 2008; 22(Suppl 1):S16-S18.
- (46) Snijders AH, Nijkrake MJ, Bakker M, Munneke M, Wind C, Bloem BR. Clinimetrics of freezing of gait. *Mov Disord* 2008; 23(Suppl 2):S468-S474.
- (47) Giladi N, Nieuwboer A. Understanding and treating freezing of gait in parkinsonism, proposed working definition, and setting the stage. *Mov Disord* 2008; 23(Suppl 2):S423-S425.
- (48) Morris ME. Locomotor training in people with Parkinson disease. *Phys Ther* 2006; 86(10):1426-1435.
- (49) Schaafsma JD, Balash Y, Gurevich T, Bartels AL, Hausdorff JM, Giladi N. Characterization of freezing of gait subtypes and the response of each to levodopa in Parkinson's disease. *Eur J Neurol* 2003; 10(4):391-398.
- (50) Snijders AH, van de Warrenburg BP, Giladi N, Bloem BR. Neurological gait disorders in elderly people: clinical approach and classification. *Lancet Neurol* 2007; 6(1):63-74.
- (51) Bloem BR, Valkenburg VV, Slabbekoorn M, Gert vD. The multiple tasks test. Strategies in Parkinson's disease. *Exp Brain Res* 2001; 137(3-4):478-486.
- (52) Bloem BR, Grimbergen YA, van Dijk JG, Munneke M. The "posture second" strategy: a review of wrong priorities in Parkinson's disease. *J Neurol Sci* 2006; 248(1-2):196-204.
- (53) Bond JM, Morris M. Goal-directed secondary motor tasks: their effects on gait in subjects with Parkinson disease. *Arch Phys Med Rehabil* 2000; 81(1):110-116.
- (54) O'Shea S, Morris ME, Iansek R. The effect of secondary cognitive and motor tasks on gait in Parkinson's disease. *Mov Disord* 2000; 15(2):403.
- (55) Rochester L, Hetherington V, Jones D, Nieuwboer A, Willems AM, Kwakkel G et al. Attending to the task: Interference effects of functional tasks on walking in Parkinson's disease and the roles of cognition, depression, fatigue, and balance. *Archives of Physical Medicine and Rehabilitation* 2004; 85(10):1578-1585.
- (56) Koerts J, Van Beilen M, Leenders KL, Brouwer WH. Controlled behavior in Parkinson's disease: initiative, planning and multi-task performance. In: Koerts J, editor. *Parkinson's Disease: Neuroimaging and clinical studies on cognition and depression*. [PhD thesis Groningen University]. Groningen: J Koerts; 2009.
- (57) Hausdorff JM, Balash J, Giladi N. Effects of cognitive challenge on gait variability in patients with Parkinson's disease. *J Geriatr Psychiatry Neurol* 2003; 16(1):53-58.
- (58) Marchese R, Bove M, Abbruzzese G. Effect of cognitive and motor tasks on postural stability in Parkinson's disease: A posturographic study. *Mov Disord* 2003; 18(6):652-658.
- (59) Bloem BR, Van Rossum MAC, Speelman AD, Munneke M. PD-related gait, posture, and balance disorders. In: Wolters E Ch, Van Laar T and Berends HW, editors. *Parkinsonism and Related Disorders*. VU University Press: Amsterdam; 2007. 57-77.
- (60) Bloem BR, Grimbergen YA, Cramer M, Willemsen M, Zwinderman AH. Prospective assessment of falls in Parkinson's disease. *J Neurol* 2001; 248(11):950-958.
- (61) Pickering RM, Grimbergen YA, Rigney U, Ashburn A, Mazibrada G, Wood B et al. A meta-analysis of six prospective studies of falling in Parkinson's disease. *Mov Disord* 2007; 22(13):1892-1900.
- (62) Mak MK, Pang MY. Fear of falling is independently associated with recurrent falls in patients with Parkinson's disease: a 1-year prospective study. *J Neurol* 2009; 256:1689-1695.
- (63) Wood BH, Bilclough JA, Bowron A, Walker RW. Incidence and prediction of falls in Parkinson's disease: a prospective multidisciplinary study. *J Neurol Neurosurg Psychiatry* 2002; 72(6):721-725.
- (64) Gray P, Hildebrand K. Fall risk factors in Parkinson's disease. *J Neurosci Nurs* 2000; 32(4):222-228.

Reference List

- (65) Stack E, Ashburn A. Fall events described by people with Parkinson's disease: implications for clinical interviewing and the research agenda. *Physiother Res Int* 1999; 4(3):190-200.
- (66) Ashburn A, Stack E, Pickering RM, Ward CD. Predicting fallers in a community-based sample of people with Parkinson's disease. *Gerontology* 2001; 47(5):277-281.
- (67) Wenning GK, Ebersbach G, Verny M, Chaudhuri KR, Jellinger K, McKee A et al. Progression of falls in postmortem-confirmed parkinsonian disorders. *Mov Disord* 1999; 14(6):947-950.
- (68) Allcock LM, Rowan EN, Steen IN, Wesnes K, Kenny RA, Burn DJ. Impaired attention predicts falling in Parkinson's disease. *Parkinsonism Relat Disord* 2009; 15(2):110-115.
- (69) Ashburn A, Stack E, Pickering RM, Ward CD. A community-dwelling sample of people with Parkinson's disease: characteristics of fallers and non-fallers. *Age Ageing* 2001; 30(1):47-52.
- (70) Carpenter MG, Allum JH, Honegger F, Adkin AL, Bloem BR. Postural abnormalities to multidirectional stance perturbations in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2004; 75(9):1245-1254.
- (71) Balash Y, Peretz C, Leibovich G, Herman T, Hausdorff JM, Giladi N. Falls in outpatients with Parkinson's disease - Frequency, impact and identifying factors. *Journal of Neurology* 2005; 252(11):1310-1315.
- (72) Sato Y, Manabe S, Kuno H, Oizumi K. Amelioration of osteopenia and hypovitaminosis D by 1alpha-hydroxyvitamin D3 in elderly patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1999; 66(1):64-68.
- (73) Johnell O, Melton LJ, III, Atkinson EJ, O'Fallon WM, Kurland LT. Fracture risk in patients with parkinsonism: a population-based study in Olmsted County, Minnesota. *Age Ageing* 1992; 21(1):32-38.
- (74) Schrag A, Hovris A, Morley D, Quinn N, Jahanshahi M. Caregiver-burden in Parkinson's disease is closely associated with psychiatric symptoms, falls, and disability. *Parkinsonism & Related Disorders* 2006; 12(1):35-41.
- (75) Adkin AL, Frank JS, Jog MS. Fear of falling and postural control in Parkinson's disease. *Mov Disord* 2003; 18(5):496-502.
- (76) Franchignoni F, Martignoni E, Ferriero G, Pasetti C. Balance and fear of falling in Parkinson's disease. *Parkinsonism Relat Disord* 2005; 11(7):427-433.
- (77) Fertl E, Doppelbauer A, Auff E. Physical activity and sports in patients suffering from Parkinson's disease in comparison with healthy seniors. *J Neural Transm Park Dis Dement Sect* 1993; 5(2):157-161.
- (78) Schrag A, Jahanshahi M, Quinn N. What contributes to quality of life in patients with Parkinson's disease? *Journal of Neurology Neurosurgery and Psychiatry* 2000; 69(3):308-312.
- (79) Visser M, van Rooden SM, Verbaan D, Marinus J, Stiggelbout AM, van Hilten JJ. A comprehensive model of health-related quality of life in Parkinson's disease. *J Neurol* 2008; 255(10):1580-1587.
- (80) Rahman S, Griffin HJ, Quinn NP, Jahanshahi M. Quality of life in Parkinson's disease: the relative importance of the symptoms. *Mov Disord* 2008; 23(10):1428-1434.
- (81) Fox SH, Brotchie JM, Lang AE. Non-dopaminergic treatments in development for Parkinson's disease. *Lancet Neurol* 2008; 7(10):927-938.
- (82) KNGF. [Professional profile of the physical therapist]. Royal Dutch Society for Physical Therapy. www.fysionet.nl [2008].
- (83) Plant R, Ashburn A, Lovgreen B, Maehle V, Handford F, Kinnear E. Physiotherapy for people with Parkinson's disease: UK best practice. Short Report. Newcastle upon Tyne: Institute of Rehabilitation; 2000.
- (84) Muller V, Mohr B, Rosin R, Pulvermuller F, Muller F, Birbaumer N. Short-term effects of behavioral treatment on movement initiation and postural control in Parkinson's disease: a controlled clinical study. *Mov Disord* 1997; 12(3):306-314.
- (85) Nieuwboer A, De Weertd W, Dom R, Truyen M, Janssens L, Kamsma Y. The effect of a home physiotherapy program for persons with Parkinson's disease. *J Rehabil Med* 2001; 33(6):266-272.
- (86) Thaut MH, McIntosh GC, Rice RR, Miller RA, Rathbun J, Brault JM. Rhythmic auditory stimulation in gait training for Parkinson's disease patients. *Mov Disord* 1996; 11(2):193-200.
- (87) Cubo E, Moore CG, Goetz CG. Wheeled and stand-aid walkers in Parkinson's disease patients with gait freezing. *Annals of Neurology* 2001; 50(3):S50.

- (88) Plant R, Walton G, Ashburn A, Lovgreen B, Handford F, Kinnear E. Guidelines for physiotherapy practice in Parkinson's disease. Newcastle, U.K.: University of Northumbria, Institute of Rehabilitation; 2001.
- (89) Bloem BR, Beckley DJ, van Dijk JG, Zwiderman AH, Remler MP, Roos RA. Influence of dopaminergic medication on automatic postural responses and balance impairment in Parkinson's disease. *Mov Disord* 1996; 11(5):509-521.
- (90) Bloem BR, Beckley DJ, van Dijk JG. Are automatic postural responses in patients with Parkinson's disease abnormal due to their stooped posture? *Exp Brain Res* 1999; 124(4):481-488.
- (91) Auff E, Fertl E, Schnider P. [Parkinson disease and neurologic rehabilitation]. *Wien Med Wochenschr* 1995; 145(13):302-305.
- (92) Bohannon RW. Physical rehabilitation in neurologic diseases. *Curr Opin Neurol* 1993; 6(5):765-772.
- (93) Handford F. Towards a rational basis for physiotherapy in Parkinson's disease. *Baillieres Clin Neurol* 1993; 2(1):141-158.
- (94) Szekeley BC, Kosanovich NN, Sheppard W. Adjunctive treatment in Parkinson's disease: physical therapy and comprehensive group therapy. *Rehabil Lit* 1982; 43(3-4):72-76.
- (95) Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ* 1996; 312(7023):71-72.
- (96) Centre for Evidence Based Medicine. What is EBM? www.cebm.net [2009; cited 2009 Apr. 29].
- (97) Schunemann HJ, Oxman AD, Brozek J, Glasziou P, Jaeschke R, Vist GE et al. Grading quality of evidence and strength of recommendations for diagnostic tests and strategies. *BMJ* 2008; 336(7653):1106-1110.
- (98) Ng DC. Parkinson's disease. Diagnosis and treatment. *West J Med* 1996; 165(4):234-240.
- (99) Martin JP. The basal ganglia and posture. London: Pitman Medical; 1967.
- (100) Morris ME, Iansek R, Matyas TA, Summers JJ. The pathogenesis of gait hypokinesia in Parkinson's disease. *Brain* 1994; 117(5):1169-1181.
- (101) Wolters EC, Francot CM. Mental dysfunction in Parkinson's disease. *Parkinsonism Relat Disord* 1998; 4(3):107-112.
- (102) Patti F, Reggio A, Nicoletti F, Sellaroli T, Deinite G, Nicoletti F. Effects of rehabilitation therapy on Parkinsons' disability and functional independence. *J Neurol Rehabil* 1996; 14(4):223-231.
- (103) Schenkman M, Cutson TM, Kuchibhatla M, Chandler J, Pieper CF, Ray L et al. Exercise to improve spinal flexibility and function for people with Parkinson's disease: a randomized, controlled trial. *J Am Geriatr Soc* 1998; 46(10):1207-1216.
- (104) Mohr B, Muller V, Mattes R, Rosin R, Federmann B, Strehl U et al. Behavioral treatment of Parkinson's disease leads to improvement of motor skills and tremor reduction. *Behav Ther* 1996; 27:235-255.
- (105) Henneberg A. Additional therapies in Parkinson's disease patients: useful tools for the improvement of the quality of life or senseless loss of resources? *J Neurol* 1998; 245(Suppl 1):S23-S27.
- (106) Ulm G. The current significance of physiotherapeutic measures in the treatment of Parkinson's disease. *J Neural Transm Suppl* 1995; 46:455-460.
- (107) Van Emmerik REA, Wagenaar RC, de Goede CJT. Oefentherapie bij Parkinsonpatiënten. In: Bohn Stafleu Van Lochem, editor. *Jaarboek voor fysiotherapie / kinesithérapie*. Houten: 1995. 148-175.
- (108) Weiner WJ, Singer C. Parkinson's disease and nonpharmacologic treatment programs. *J Am Geriatr Soc* 1989; 37(4):359-363.
- (109) Homberg V. Motor training in the therapy of Parkinson's disease. *Neurology* 1993; 43(12 Suppl 6):S45-S46.
- (110) Dam M, Tonin P, Casson S, Bracco F, Piron L, Pizzolato G et al. Effects of conventional and sensory-enhanced physiotherapy on disability of Parkinson's disease patients. *Adv Neurol* 1996; 69:551-555.
- (111) Kwakkel G, Wagenaar RC, Koelman TW, Lankhorst GJ, Koetsier JC. Effects of intensity of rehabilitation after stroke. A research synthesis. *Stroke* 1997; 28(8):1550-1556.

Reference List

- (112) Cambach W, Wagenaar RC, Koelman TW, van Keimpema AR, Kemper HC. The long-term effects of pulmonary rehabilitation in patients with asthma and chronic obstructive pulmonary disease: a research synthesis. *Arch Phys Med Rehabil* 1999; 80(1):103-111.
- (113) Cooper H, Hedges LV. The handbook of research synthesis. New York: Russell Sage Foundation; 1994.
- (114) Crossley SM. Intensive physiotherapy for Parkinson's disease in a holiday environment. *Physiother* 1986; 72(8):383-384.
- (115) Flewitt B, Capildeo R, Rose FC. Physiotherapy and assessments in Parkinson's disease using the polarised light goniometer. In: Capildeo R, Rose FC, editors. *Research in progress in Parkinson's disease*. London: Pitman Medical; 1981. 404-415.
- (116) Franklin S, Kohout LJ, Stern GM, Dunning M. Physiotherapy in Parkinson's disease. In: Capildeo R, Rose FC, editors. *Research in progress in Parkinson's disease*. London: Pitman Medical; 1981. 397-400.
- (117) Gibberd FB, Page NG, Spencer KM, Kinnear E, Hawksworth JB. Controlled trial of physiotherapy and occupational therapy for Parkinson's disease. *Br Med J (Clin Res Ed)* 1981; 282(6271):1196.
- (118) Mitchell PH, Mertz MA, Catanzaro ML. Group exercise: a nursing therapy in Parkinson's disease. *Rehabil Nurs* 1987; 12(5):242-245.
- (119) Pedersen SW, Oberg B, Insulander A, Vretman M. Group training in parkinsonism: quantitative measurements of treatment. *Scand J Rehabil Med* 1990; 22(4):207-211.
- (120) Comella CL, Stebbins GT, Brown-Toms N, Goetz CG. Physical therapy and Parkinson's disease: a controlled clinical trial. *Neurology* 1994; 44(3):376-378.
- (121) Gauthier L, Dalziel S, Gauthier S. The benefits of group occupational therapy for patients with Parkinson's disease. *Am J Occup Ther* 1987; 41(6):360-365.
- (122) Palmer SS, Mortimer JA, Webster DD, Bistevins R, Dickinson GL. Exercise therapy for Parkinson's disease. *Arch Phys Med Rehabil* 1986; 67(10):741-745.
- (123) Formisano R, Pratesi L, Modarelli FT, Bonifati V, Meco G. Rehabilitation and Parkinson's disease. *Scand J Rehabil Med* 1992; 24(3):157-160.
- (124) Kwakkel G, Wagenaar RC, Twisk JW, Lankhorst GJ, Koetsier JC. Intensity of leg and arm training after primary middle-cerebral-artery stroke: a randomised trial. *Lancet* 1999; 354(9174):191-196.
- (125) Bonnet AM, Loria Y, Saint-Hilaire MH, Lhermitte F, Agid Y. Does long-term aggravation of Parkinson's disease result from nondopaminergic lesions? *Neurology* 1987; 37(9):1539-1542.
- (126) Giladi N, Treves TA, Simon ES, Shabtai H, Orlov Y, Kandinov B et al. Freezing of gait in patients with advanced Parkinson's disease. *J Neural Transm* 2001; 108(1):53-61.
- (127) Bloem BR, van Vugt JP, Beckley DJ. Postural instability and falls in Parkinson's disease. *Adv Neurol* 2001; 87:209-223.
- (128) Schrag A, Selai C, Jahanshahi M, Quinn NP. The EQ-5D—a generic quality of life measure—is a useful instrument to measure quality of life in patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2000; 69(1):67-73.
- (129) Alkhani A, Lozano AM. Pallidotomy for parkinson disease: a review of contemporary literature. *J Neurosurg* 2001; 94(1):43-49.
- (130) Okun MS, Stover NP, Subramanian T, Gearing M, Wainer BH, Holder CA et al. Complications of gamma knife surgery for Parkinson disease. *Arch Neurol* 2001; 58(12):1995-2002.
- (131) Boer AGd, Sprangers MA, Speelman HD, de Haes HC. Predictors of health care use in patients with Parkinson's disease: a longitudinal study. *Mov Disord* 1999; 14(5):772-779.
- (132) Chartered Society of Physiotherapy. *Neurology: Parkinson's disease, multiple sclerosis and severe traumatic brain injury. Physiotherapy Effectiveness Bulletin* 2001; 3:1-11.
- (133) Marinus J, Visser M, Stiggelbout AM, Rabey JM, Bonuccelli U, Kraus PH et al. Activity-based diary for Parkinson's disease. *Clin Neuropharmacol* 2002; 25(1):43-50.
- (134) Hobson JP, Edwards NI, Meara RJ. The Parkinson's Disease Activities of Daily Living Scale: a new simple and brief subjective measure of disability in Parkinson's disease. *Clin Rehabil* 2001; 15(3):241-246.

- (135) Global Parkinson's Disease Survey Steering Committee. Factors impacting on quality of life in Parkinson's disease: results from an international survey. *Mov Disord* 2002; 17(1):60-67.
- (136) Clarke CE, Zobkiw RM, Gullaksen E. Quality of life and care in Parkinson's disease. *Br J Clin Pract* 1995; 49(6):288-293.
- (137) Cutson TM, Laub KC, Schenkman M. Pharmacological and nonpharmacological interventions in the treatment of Parkinson's disease. *Phys Ther* 1995; 75(5):363-373.
- (138) de Goede CJ, Keus SH, Kwakkel G, Wagenaar RC. The effects of physical therapy in Parkinson's disease: a research synthesis. *Arch Phys Med Rehabil* 2001; 82(4):509-515.
- (139) Deane KH, Ellis-Hill C, Jones D, Whurr R, Ben Shlomo Y, Playford ED et al. Systematic review of paramedical therapies for Parkinson's disease. *Mov Disord* 2002; 17(5):984-991.
- (140) Thompson AJ, Playford ED. Rehabilitation for patients with Parkinson's disease. *Lancet* 2001; 357(9254):410-411.
- (141) Yarrow S. Survey of members of the Parkinson's Disease Society. London: Parkinson's Disease Society and Policy Studies Institute; 1999.
- (142) Bloem BR, Keus SHJ, Munneke M. PROMISE-NL: Preparatory studies. 1st Scientific conference of the Association of Physiotherapists in Parkinson's Disease Europe (APPDE) 2003; Southampton.
- (143) Central Bureau of Statistics. [Use of medical services]. *statline.cbs.nl* [2003; cited 2003 Dec. 5].
- (144) Dodel RC, Singer M, Kohne-Volland R, Selzer R, Scholz W, Rathay B et al. [Cost of illness in Parkinson disease. A retrospective 3-month analysis of direct costs]. *Nervenarzt* 1997; 68(12):978-984.
- (145) Parkinson Disease Society. Parkinson's Disease. Meeting a need? Discussion Document 1994; 11.
- (146) National Institute for Public Health and the Environment (RIVM). [Costs of diseases]. *www.rivm.nl/kostenvanziekten* [2002; cited 2003 Sept. 15].
- (147) Deane KH, Jones D, Playford ED, Ben Shlomo Y, Clarke CE. Physiotherapy versus placebo or no intervention in Parkinson's disease. *The Cochrane Library* [2002] (Issue 3) Available from: URL:PM:11687029.
- (148) Willemsen MD, Grimbergen YA, Slabbekoorn M, Bloem BR. [Falling in Parkinson disease: more often due to postural instability than to environmental factors]. *Ned Tijdschr Geneesk* 2000; 144(48):2309-2314.
- (149) Bloem BR. Preventie van vallen bij de ziekte van Parkinson. In: Roos RAC, Mulder WJ, editors. *Voorwaarts bij CVA en ziekte van Parkinson*. Leiden: Boerhaave Commissie; 2000. 71-92.
- (150) Rubinstein TC, Giladi N, Hausdorff JM. The power of cueing to circumvent dopamine deficits: A review of physical therapy treatment of gait disturbances in Parkinson's disease. *Mov Disord* 2002; 17(6):1148-1160.
- (151) Bloem BR, Munneke M, Plant R. A randomised controlled trial of physiotherapy in PD - the PROMISE trial. Multidisciplinary care in Parkinson's disease and parkinsonism, from science to practice 2001;8.
- (152) Kwakkel G, de Goede C. [Research proposal RES-CUE valued 'excellent': Huge grant for VUmc]. *Fysiother* 2002; 2:34-37.
- (153) Bloem BR. Postural instability in Parkinson's disease. *Clin Neurol Neurosurg* 1992; 94(Suppl):S41-S45.
- (154) Management of Parkinson's disease: an evidence-based review. *Mov Disord* 2002; 17(Suppl 4):S1-166.
- (155) Keus SHJ, Bloem BR, Verbaan D, de Jonge PA, Hofman M, van Hilten BJ et al. Physiotherapy in Parkinson's disease: utilisation and patient satisfaction. *J Neurol* 2004; 251(6):680-687.
- (156) Deane KH, Jones D, Playford ED, Ben-Shlomo Y, Clarke CE. Physiotherapy for patients with Parkinson's Disease: a comparison of techniques. *Cochrane Database Syst Rev* 2001;(3):CD002817.
- (157) AGREE Collaboration. Appraisal of Guidelines for Research & Evaluation (AGREE) Instrument. *www.agreecollaboration.org* [2001 [cited 2005 Jan. 13].
- (158) Hendriks HJM, Bekkering GE, van Erteekoven H, Brandsma JW, van der Wees PhJ, de Bie RA. Development and implementation of national practice guidelines: A prospect for continuous quality improvement in physiotherapy. Introduction to the method of guideline development. *Physiother* 2000; 86:535-547.
- (159) Nieuwboer A, De Weertd W, Nuyens G, Hantson L, Feys H. [A review of the efficacy of physiotherapy in Parkinson's disease]. *Ned Tijdschr Fysiother* 1994; 5:122-128.

Reference List

- (160) Bergen JL, Toole T, Elliott III RG, Wallace B, Robinson K, Maitland CG. Aerobic exercise intervention improves aerobic capacity and movement initiation in Parkinson's disease patients. *NeuroRehabilitation* 2002; 17(2):161-168.
- (161) Bridgewater KJ, Sharpe M. Trunk muscle training and early parkinson's disease. *Physiother Th Pract* 1997; 13(2):139-153.
- (162) de Goede CJT, Kwakkel G. [The efficacy of physiotherapy group treatment in Parkinson's disease: a cross-over trial]. *Ned Tijdschr Fysiother* 2004; 114(3):78-82.
- (163) Hirsch MA, Toole T, Maitland CG, Rider RA. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch Phys Med Rehabil* 2003; 84(8):1109-1117.
- (164) Hurwitz A. The benefit of a home exercise regimen for ambulatory Parkinson's disease patients. *J Neurosci Nurs* 1989; 21(3):180-184.
- (165) Marchese R, Diverio M, Zucchi F, Lentino C, Abbruzzese G. The role of sensory cues in the rehabilitation of parkinsonian patients: a comparison of two physical therapy protocols. *Mov Disord* 2000; 15(5):879-883.
- (166) Miyai I, Fujimoto Y, Ueda Y, Yamamoto H, Nozaki S, Saito T et al. Treadmill training with body weight support: its effect on Parkinson's disease. *Arch Phys Med Rehabil* 2000; 81(7):849-852.
- (167) Pacchetti C, Mancini F, Aglieri R, Fundaro C, Martignoni E, Nappi G. Active music therapy in Parkinson's disease: an integrative method for motor and emotional rehabilitation. *Psychosom Med* 2000; 62(3):386-393.
- (168) Stallibrass C, Sissons P, Chalmers C. Randomized controlled trial of the Alexander technique for idiopathic Parkinson's disease. *Clin Rehabil* 2002; 16(7):695-708.
- (169) Toole T, Hirsch MA, Forkink A, Lehman DA, Maitland CG. The effects of a balance and strength training program on equilibrium in Parkinsonism: A preliminary study. *NeuroRehabilitation* 2000; 14(3):165-174.
- (170) Yekutieli MP, Pinhasof A, Shahar G, Sroka H. A clinical trial of the re-education of movement in patients with Parkinson's disease. *Clin Rehabil* 1991; 5:207-214.
- (171) Cerri C., Arosio A, Biella AM, Premoselli S, Piccini L. Physical exercise therapy of Parkinson's. *Mov Disord* 1994; 9(Suppl 1):68.
- (172) Chandler C, Plant R. A targeted physiotherapy service for people with Parkinson's disease from diagnosis to end stage: a pilot study. In: Percival R, Hobson P, editors. *Parkinson's disease: study in physiological and social care*. Leicester: BPS Books; 1999. 256-269.
- (173) Forkink A, Toole T, Hirsch MA, Lehman DA, Maitland CG. The effects of a balance and strengthening program on equilibrium in Parkinsonism. Working paper series (PI-96-33). Tallahassee: Florida Sate University; 1996.
- (174) Homann CN, Crevenna R, Kojnig H, et al. Can physiotherapy improve axial symptoms in parkinsonian patients? A pilot study with the computerized movement analysis battery Zebris. *Mov Disord* 1998; 14(Suppl 2):234.
- (175) Koseoglu F, Inan L, Ozel S, Deviren SD, Karabiyikoglu G, Yorgancioglu R et al. The effects of a pulmonary rehabilitation program on pulmonary function tests and exercise tolerance in patients with Parkinson's disease. *Funct Neurol* 1997; 12(6):319-325.
- (176) Shiba Y, Obuchi S, Toshima H, Yamakita H. Comparison between visual and auditory stimulation in gait training of patients with idiopathic Parkinson's disease. World Conference of Physical Therapy Conference 1999.
- (177) Wade DT. Outcome measures for clinical rehabilitation trials: impairment, function, quality of life, or value? *Am J Phys Med Rehabil* 2003; 82(10):S26-S31.
- (178) Beurskens AJ, de Vet HC, Koke AJ. Responsiveness of functional status in low back pain: a comparison of different instruments. *Pain* 1996; 65(1):71-76.
- (179) Wright JG, Young NL. The patient-specific index: asking patients what they want. *J Bone Joint Surg Am* 1997; 79(7):974-983.
- (180) McIntosh GC, Brown SH, Rice RR, Thaut MH. Rhythmic auditory-motor facilitation of gait patterns in patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1997; 62(1):22-26.
- (181) Thaut MH, McIntosh GC. Music therapy in mobility training with the elderly: a review of current research. *Care Manag J* 1999; 1(1):71-74.
- (182) Thaut MH, McIntosh KW, McIntosh GC, Hoemberg V. Auditory rhythmicity enhances movement and speech motor control in patients with Parkinson's disease. *Funct Neurol* 2001; 16(2):163-172.

- (183) Dietz MA, Goetz CG, Stebbins GT. Evaluation of a modified inverted walking stick as a treatment for parkinsonian freezing episodes. *Mov Disord* 1990; 5(3):243-247.
- (184) Dunne JW, Hankey GJ, Edis RH. Parkinsonism: upturned walking stick as an aid to locomotion. *Arch Phys Med Rehabil* 1987; 68(6):380-381.
- (185) Lewis GN, Byblow WD, Walt SE. Stride length regulation in Parkinson's disease: the use of extrinsic, visual cues. *Brain* 2000; 123 (Pt 10):2077-2090.
- (186) Morris ME, Ianssek R, Matyas TA, Summers JJ. Stride length regulation in Parkinson's disease. Normalization strategies and underlying mechanisms. *Brain* 1996; 119 (Pt 2):551-568.
- (187) Keus SHJ, Hendriks HJM, Bloem BR, Bredero-Cohen AB, de Goede CJT, van Haaren M et al. KNGF Guidelines for physical therapy in patients with Parkinson's disease. *Ned Tijdschr Fysiother* 2004; 114(3 (Suppl):www.kngf.nl-www.cebp.nl.
- (188) Keus SH, Bloem BR, van Hilten JJ, Ashburn A, Munneke M. Effectiveness of physiotherapy in Parkinson's disease: the feasibility of a randomised controlled trial. *Parkinsonism Relat Disord* 2007; 13(2):115-121.
- (189) Brichetto G, Pelosin E, Marchese R, Abbruzzese G. Evaluation of physical therapy in parkinsonian patients with freezing of gait: a pilot study. *Clinical Rehabilitation* 2006; 20(1):31-35.
- (190) Caglar AT, Gurses HN, Mutluay FK, Kiziltan G. Effects of home exercises on motor performance in patients with Parkinson's disease. *Clin Rehabil* 2005; 19(8):870-877.
- (191) Dibble LE, Nicholson DE, Shultz B, MacWilliams BA, Marcus RL, Moncur C. Sensory cueing effects on maximal speed gait initiation in persons with Parkinson's disease and healthy elders. *Gait Posture* 2004; 19(3):215-225.
- (192) Ellis T, de Goede CJ, Feldman RG, Wolters EC, Kwakkel G, Wagenaar RC. Efficacy of a physical therapy program in patients with Parkinson's disease: A randomized controlled trial. *Arch Phys Med Rehabil* 2005; 86(4):626-632.
- (193) Farley BG, Koshland GF. Training BIG to move faster: the application of the speed-amplitude relation as a rehabilitation strategy for people with Parkinson's disease. *Experimental Brain Research* 2005; 167(3):462-467.
- (194) Frenkel-Toledo S, Giladi N, Peretz C, Herman T, Gruendlinger L, Hausdorff JM. Treadmill walking as an external pacemaker to improve gait rhythm and stability in Parkinson's disease. *Mov Disord* 2005; 20(9):1109-1114.
- (195) Inzelberg R, Peleg N, Nisipeanu P, Magadle R, Carasso RL, Weiner P. Inspiratory muscle training and the perception of dyspnea in Parkinson's disease. *Can J Neurol Sci* 2005; 32(2):213-217.
- (196) Jobges M, Heuschkel G, Pretzel C, Illhardt C, Renner C, Hummelsheim H. Repetitive training of compensatory steps: a therapeutic approach for postural instability in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2004; 75(12):1682-1687.
- (197) Lehman DA, Toole T, Lofald D, Hirsch MA. Training with verbal instructional cues results in near-term improvement of gait in people with Parkinson disease. *J Neurol Phys Ther* 2005; 29(1):2-8.
- (198) Lun V, Pullan N, Labelle N, Adams C, Suchowersky O. Comparison of the effects of a self-supervised home exercise program with a physiotherapist-supervised exercise program on the motor symptoms of Parkinson's disease. *Mov Disord* 2005; 20(8):971-975.
- (199) Paterson J. Chinese exercise modalities in Parkinson's disease. www.clinicaltrials.gov/show/NCT00029809 [2003].
- (200) Pellecchia MT, Grasso A, Biancardi LG, Squillante M, Bonavita V, Barone P. Physical therapy in Parkinson's disease: an open long-term rehabilitation trial. *J Neurol* 2004; 251(5):595-598.
- (201) Pohl M, Rockstroh G, Ruckriem S, Mrass G, Mehrholz J. Immediate effects of speed-dependent treadmill training on gait parameters in early Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 2003; 84(12):1760-1766.
- (202) Protas EJ, Mitchell K, Williams A, Qureshy H, Caroline K, Lai EC. Gait and step training to reduce falls in Parkinson's disease. *Neurorehabilitation* 2005; 20(3):183-190.
- (203) Schallow G, Paasuke M, Ereline J, Gapeyeva H. Improvement in Parkinson's disease patients achieved by coordination dynamics therapy. *Electromyogr Clin Neurophysiol* 2004; 44(2):67-73.

Reference List

- (204) Sidaway B, Anderson J, Danielson G, Martin L, Smith G. Effects of long-term gait training using visual cues in an individual with Parkinson disease. *Phys Ther* 2006; 86(2):186-194.
- (205) Suteerawattananon M, Morris GS, Etnyre BR, Jankovic J, Protas EJ. Effects of visual and auditory cues on gait in individuals with Parkinson's disease. *J Neurol Sci* 2004; 219(1-2):63-69.
- (206) Svircev A, Craig LH, Juncos JL. A pilot study examining the effects of neuromuscular therapy on patients with Parkinson's disease. *J Am Osteopath Assoc* 2005; 105(1):26.
- (207) Tuite P, Anderson N, Konczak J. Constraint-induced movement therapy in Parkinson's disease. *Mov Disord* 2005; 20(7):910-911.
- (208) Smidt N, de Vet HC, Bouter LM, Dekker J, Arendzen JH, de Bie RA et al. Effectiveness of exercise therapy: a best-evidence summary of systematic reviews. *Aust J Physiother* 2005; 51(2):71-85.
- (209) Keus SHJ, Bloem BR, Nijkrake M, Lim I, Roos RAC, Kwakkel G et al. Improving the efficiency of physiotherapeutic care in Parkinson's disease: the PARKNET trial. *Gait Posture* 2005; 21(Suppl 1):S95.
- (210) Olanow CW, Watts RL, Koller WC. An algorithm (decision tree) for the management of Parkinson's disease (2001): treatment guidelines. *Neurology* 2001; 56(11):S1-S88.
- (211) Keus SH, Bloem BR, Hendriks EJ, Bredero-Cohen AB, Munneke M. Evidence-based analysis of physical therapy in Parkinson's disease with recommendations for practice and research. *Mov Disord* 2007; 22(4):451-460.
- (212) Bekkering GE, Engers AJ, Wensing M, Hendriks HJM, van Tulder MW, Oostendorp RAB et al. Development of an implementation strategy for physiotherapy guidelines on low back pain. *Australian Journal of Physiotherapy* 2003; 49(3):208-214.
- (213) Ross S, Grant A, Counsell C, Gillespie W, Russell I, Prescott R. Barriers to participation in randomised controlled trials: a systematic review. *J Clin Epidemiol* 1999; 52(12):1143-1156.
- (214) Beurskens AJ, de Vet HC, Koke AJ, Lindeman E, van der Heijden GJ, Regtop W et al. A patient-specific approach for measuring functional status in low back pain. *J Manipulative Physiol Ther* 1999; 22(3):144-148.
- (215) Nieuwboer A, De Weerd W, Dom R, Bogaerts K, Nuyens G. Development of an activity scale for individuals with advanced Parkinson disease: reliability and "on-off" variability. *Phys Ther* 2000; 80(11):1087-1096.
- (216) Peto V, Jenkinson C, Fitzpatrick R. PDQ-39: a review of the development, validation and application of a Parkinson's disease quality of life questionnaire and its associated measures. *J Neurol* 1998; 245(Suppl 1):S10-S14.
- (217) Verhoeven AC, Boers M, van der LS. Validity of the MACTAR questionnaire as a functional index in a rheumatoid arthritis clinical trial. *The McMaster Toronto Arthritis J Rheumatol* 2000; 27(12):2801-2809.
- (218) Wright JG, Young NL, Waddell JP. The reliability and validity of the self-reported patient-specific index for total hip arthroplasty. *J Bone Joint Surg Am* 2000; 82(6):829-837.
- (219) Hedges LV, Olkin I. Statistical methods for meta-analysis. Orlando, FL: Academic Press; 1985.
- (220) Hollis S, Campbell F. What is meant by intention to treat analysis? Survey of published randomised controlled trials. *BMJ* 1999; 319(7211):670-674.
- (221) Peto V, Jenkinson C, Fitzpatrick R. Determining minimally important differences for the PDQ-39 Parkinson's disease questionnaire. *Age Ageing* 2001; 30(4):299-302.
- (222) Ashburn A, Jones D, Plant R, Lovgreen B, Kinnear E, Handford F et al. Physiotherapy for people with Parkinson's disease in the UK: an exploration of practice. *Int J Ther Rehabil* 2004; 11(4):160-167.
- (223) Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; 12(3):189-198.
- (224) Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Phys Ther* 2000; 80(9):896-903.
- (225) Giladi N, Hausdorff JM. The role of mental function in the pathogenesis of freezing of gait in Parkinson's disease. *J Neurol Sci* 2006; 248(1-2):173-176.
- (226) Nieuwboer A, Herman T, Rochester L, Ehab Emil G, Giladi N. The new revised freezing of gait questionnaire, a reliable and valid instrument to measure freezing in Parkinson's disease. *Parkinsonism Relat Disord* 2008; 14 (Suppl 1):S68.

- (227) Nieuwboer A, Kwakkel G, Rochester L, Jones D, Van WE, Willems AM et al. Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *J Neurol Neurosurg Psychiatry* 2007; 78(2):134-140.
- (228) Van Wegen E, Lim I, de Goede C, Nieuwboer A, Willems A, Jones D et al. The effects of visual rhythms and optic flow on stride patterns of patients with Parkinson's disease. *Parkinsonism & Related Disorders* 2006; 12(1):21-27.
- (229) Ashburn A, Fazakarley L, Ballinger C, Pickering R, McLellan LD, Fitton C. A randomised controlled trial of a home-based exercise programme to reduce the risk of falling among people with Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2007; 78(7):678-684.
- (230) Kwakkel G, de Goede CJ, Van Wegen EE. Impact of physical therapy for Parkinson's disease: A critical review of the literature. *Parkinsonism Relat Disord* 2007; 13(Suppl 3):S478-S487.
- (231) Siderowf A, McDermott M, Kieburtz K, Blindauer K, Plumb S, Shoulson I. Test-retest reliability of the unified Parkinson's disease rating scale in patients with early Parkinson's disease: results from a multicenter clinical trial. *Mov Disord* 2002; 17(4):758-763.
- (232) Qutubuddin AA, Pegg PO, Cifu DX, Brown R, McNamee S, Carne W. Validating the Berg Balance Scale for patients with Parkinson's disease: a key to rehabilitation evaluation. *Arch Phys Med Rehabil* 2005; 86(4):789-792.
- (233) Kegelmeier DA, Kloos AD, Thomas KM, Kostyk SK. Reliability and validity of the Tinetti Mobility Test for individuals with Parkinson disease. *Phys Ther* 2007; 87(10):1369-1378.
- (234) Okuma Y, Yanagisawa N. The clinical spectrum of freezing of gait in Parkinson's disease. *Mov Disord* 2008; 23(Suppl 2):S426-S430.
- (235) Bekkering GE, Hendriks HJM, Koes BW, Oostendorp RAB, Ostelo RWJG, Thomassen JMC et al. [National practice guidelines for physical therapy in patients with low back pain]. *Ned Tijdschr Fysiother* 2005; 115(Suppl).
- (236) Hoehn MM, Yahr MD. Parkinsonism: onset, progression and mortality. *Neurology* 1967; 17(5):427-442.
- (237) Nijkrake MJ, Keus SH, Oostendorp RA, Overeem S, Mulleners W, Bloem BR et al. Allied health care in Parkinson's disease: Referral, consultation, and professional expertise. *Mov Disord* 2009; 24(2):282-286.
- (238) Jolles BM, Buchbinder R, Beaton DE. A study compared nine patient-specific indices for musculoskeletal disorders. *J Clin Epidemiol* 2005; 58(8):791-801.
- (239) Feinstein AR, Cicchetti DV. High agreement but low kappa: I. The problems of two paradoxes. *J Clin Epidemiol* 1990; 43(6):543-549.
- (240) Hanna-Pladdy B. Dysexecutive syndromes in neurologic disease. *J Neurol Phys Ther* 2007; 31(3):119-127.
- (241) Shulman LM, Pretzer-Aboff I, Anderson KE, Stevenson R, Vaughan CG, Gruber-Baldini AL et al. Subjective report versus objective measurement of activities of daily living in Parkinson's disease. *Mov Disord* 2006; 21(6):794-799.
- (242) Tully M, Cantrill J. The test-retest reliability of the modified Patient Generated Index
2. *J Health Serv Res Policy* 2002; 7(2):81-89.
- (243) Munneke M, Keus S, Nijkrake M, Kwakkel G, Berendse H, Roos R et al. Efficiency of evidence-based physiotherapy for Parkinson's disease: The ParkinsonNet trial. *Mov Disord* 2008; 23(Suppl 1):S220.
- (244) Goodwin VA, Richards SH, Taylor RS, Taylor AH, Campbell JL. The effectiveness of exercise interventions for people with Parkinson's disease: A systematic review and meta-analysis. *Mov Disord* 2008; 23(5):631-640.
- (245) Nijkrake MJ, Keus SH, Ewalds H, Overeem S, Braspenning JC, Oostendorp RA et al. Quality indicators for physiotherapy in Parkinson's disease. *Eur J Phys Rehabil Med* 2009; 45(2):239-245.
- (246) Nijkrake MJ, Keus SHJ, Overeem S, Oostendorp RAB, Vliet VLieland TPM, Mulleners W et al. The ParkinsonNet concept: development, implementation and initial experience. *Mov Disord* 2010; (in press).
- (247) Campbell MK, Elbourne DR, Altman DG. CONSORT statement: extension to cluster randomised trials. *BMJ* 2004; 328(7441):702-708.
- (248) Beattie P, Turner C, Dowda M, Michener L, Nelson R. The MedRisk instrument for measuring patient satisfaction with physical therapy care: a psychometric analysis. *J Orthop Sports Phys Ther* 2005; 35(1):24-32.

Reference List

- (249) Nijkrake MJ, Keus SHJ, Quist-Anholts GWL, Overeem S, De Roode MH, Lindeboom R et al. Evaluation of a Patient-Specific Index as an outcome measure for physiotherapy in Parkinson's disease. *Eur J Phys Rehabil Med* 2009; 45(4):507-512.
- (250) Keus SH, Nieuwboer A, Bloem BR, Borm GF, Munneke M. Clinimetric analyses of the Modified Parkinson Activity Scale. *Parkinsonism Relat Disord* 2008; 15(4):263-269.
- (251) Drummond M, Weatherly H, Ferguson B. Economic evaluation of health interventions. *BMJ* 2008; 337:a1204.
- (252) Esselink RA, de Bie RM, de Haan RJ, Lenders MW, Nijssen PC, Staal MJ et al. Unilateral pallidotomy versus bilateral subthalamic nucleus stimulation in PD: a randomized trial. *Neurology* 2004; 62(2):201-207.
- (253) Donders A, van der Heijden G, Stijnen T, Moons K. A gentle introduction to imputation of missing values. *Journal of Clinical Epidemiology* 2009; 59:1087-1091.
- (254) Oostenbrink JB, Bouwmans CAM, Koopmanschap MA, Rutten FFH. [Manual for costing research]. Amstelveen, the Netherlands: Health Care Insurance Board; 2004.
- (255) Brouwer WB, Koopmanschap MA. The friction-cost method : replacement for nothing and leisure for free? *Pharmacoeconomics* 2005; 23(2):105-111.
- (256) Gold MR, Siegel JE, Russel LB, Weinstein MC. *Cost-Effectiveness in Health and Medicine*. New York: Oxford University Press; 1996.
- (257) Keus SH, Munneke M, Nijkrake MJ, Kwakkel G, Bloem BR. Physical therapy in Parkinson's disease: Evolution and future challenges. *Mov Disord* 2009; 24(1):1-14.
- (258) Kwakkel G, de Goede CJ, Van Wegen EE. Impact of physical therapy for Parkinson's disease: A critical review of the literature. *Parkinsonism Relat Disord* 2008; 13(Suppl 3):S478-S487.
- (259) Ashburn A, Ruth MP, Louise F, Claire B, Lindsay MD, Carolyn F. Recruitment to a clinical trial from the databases of specialists in Parkinson's disease. *Parkinsonism Relat Disord* 2007; 13(1):35-39.
- (260) Smith SM, Allwright S, O'Dowd T. Effectiveness of shared care across the interface between primary and specialty care in chronic disease management. *Cochrane Database Syst Rev* 2007;(3):CD004910-[accessed March 5, 2009].
- (261) Noyes K, Liu HS, Li Y, Holloway R, Dick AW. Economic burden associated with Parkinson's disease on elderly medicare beneficiaries. *Mov Disord* 2006; 21(3):362-372.
- (262) Cubo E, Alvarez E, Morant C, De Pedro CJ, Martinez MP, Genova R et al. Burden of disease related to Parkinson's disease in Spain in the year 2000. *Mov Disord* 2005; 11:1481-1487.
- (263) Dodel RC, Berger K, Oertel WH. Health-related quality of life and healthcare utilisation in patients with Parkinson's disease: impact of motor fluctuations and dyskinesias. *Pharmacoeconomics* 2001; 19(10):1013-1038.
- (264) Giladi N, Balash Y. The clinical approach to gait disturbances in Parkinson's disease; maintaining independent mobility. *J Neural Transm Suppl* 2006;(70):327-332.
- (265) Rascol O, Goetz C, Koller W, Poewe W, Sampaio C. Treatment interventions for Parkinson's disease: an evidence based assessment. *Lancet* 2002; 359(9317):1589-1598.
- (266) Nijkrake MJ, Bloem BR, Keus SH, Mulleners W. Quality of allied health care in Parkinson's disease. *Mov Disord* 2006; 21:S131.
- (267) Mutch WJ, Strudwick A, Roy SK, Downie AW. Parkinsons-Disease - Disability, Review, and Management. *Br Med J* 1986; 293(6548):675-677.
- (268) Debaere F, Wenderoth N, Sunaert S, van HP, Swinnen SP. Internal vs external generation of movements: differential neural pathways involved in bimanual coordination performed in the presence or absence of augmented visual feedback. *Neuroimage* 2003; 19(3):764-776.
- (269) Cochrane Eyes and Vision Group. Quick guide for handsearchers. www.cochraneeyes.org [2008; cited 2008 Jan. 7].
- (270) U.S. National Institutes of Health Registry. <http://clinicaltrials.gov> [2007; cited 8 A.D. Feb. 21].
- (271) metaRegister of Controlled Trials (mRCT). www.controlled-trials.com/mrct [2008; cited 2008 Feb. 21].
- (272) Australian New Zealand Clinical Trials Registry. www.anzctr.org.au [2008; cited 2008 Feb. 21].
- (273) University Hospital Medical Information Network. www.umin.ac.jp [2008; cited 2008 Feb. 21].
- (274) Netherlands Trial Register. www.trialregister.nl [2008; cited 2008 Feb. 21].

- (275) United Kingdom's National Health Service (NHS). The National Research Register (NRR). www.nrr.nhs.uk, Issue 3, July 2007 [2007; cited 2008 Feb. 21].
- (276) Burini D, Farabollini B, Iacucci S, Rimatori C, Riccardi G, Capecci M et al. A randomised controlled cross-over trial of aerobic training versus Qigong in advanced Parkinson's disease. *Eura Medicophys* 2006; 42(3):231-238.
- (277) Cakit BD, Saracoglu M, Genc H, Erdem HR, Inan L. The effects of incremental speed-dependent treadmill training on postural instability and fear of falling in Parkinson's disease. *Clin Rehabil* 2007; 21(8):698-705.
- (278) Craig LH, Svircev A, Haber M, Juncos JL. Controlled pilot study of the effects of neuromuscular therapy in patients with Parkinson's disease. *Mov Disord* 2006; 21(12):2127-2133.
- (279) Dibble LE, Hale TF, Marcus RL, Droge J, Gerber JP, LaStayo PC. High-intensity resistance training amplifies muscle hypertrophy and functional gains in persons with Parkinson's disease. *Mov Disord* 2006; 21(9):1444-1452.
- (280) Mak MK, Hui-Chan CW. Cued task-specific training is better than exercise in improving sit-to-stand in patients with Parkinson's disease: A randomized controlled trial. *Mov Disord* 2008; 23(4):501-509.
- (281) Qutubuddin AA, Cifu DX, Armistead-Jehle P, Carne W, McGuirk TE, Baron MS. A comparison of computerized dynamic posturography therapy to standard balance physical therapy in individuals with Parkinson's disease: a pilot study. *Neurorehabilitation* 2007; 22(4):261-265.
- (282) Tamir R, Dickstein R, Huberman M. Integration of motor imagery and physical practice in group treatment applied to subjects with Parkinson's disease. *Neurorehabil Neural Repair* 2007; 21(1):68-75.
- (283) Toole T, Maitland CG, Warren E, Hubmann MF, Panton L. The effects of loading and unloading treadmill walking on balance, gait, fall risk, and daily function in Parkinsonism. *Neurorehabilitation* 2005; 20(4):307-322.
- (284) CONSORT Group. Consolidated Standards of Reporting Trials (CONSORT) Statement. <http://www.consort-statement.org> [2007; cited 2008 Mar. 10].
- (285) Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomised trials. *Lancet* 2001; 357(9263):1191-1194.
- (286) Crizzle AM, Newhouse IJ. Is physical exercise beneficial for persons with Parkinson's disease? *Clin J Sport Med* 2006; 16(5):422-425.
- (287) Deane KH, Jones D, Ellis-Hill C, Clarke CE, Playford ED, Ben Shlomo Y. A comparison of physiotherapy techniques for patients with Parkinson's disease. The Cochrane Library [2002 (Issue 3):[CD002815] Available from: URL:PM:11279765.
- (288) Gage H, Storey L. Rehabilitation for Parkinson's disease: a systematic review of available evidence. *Clin Rehabil* 2004; 18(5):463-482.
- (289) Lim I, Van Wegen E, de Goede C, Deutekom M, Nieuwboer A, Willems A et al. Effects of external rhythmic cueing on gait in patients with Parkinson's disease: a systematic review. *Clin Rehabil* 2005; 19(7):695-713.
- (290) Suchowersky O, Gronseth G, Perlmutter J, Reich S, Zesiewicz T, Weiner WJ. Practice Parameter: Neuroprotective strategies and alternative therapies for Parkinson disease (an evidence-based review) Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006; 66(7):976-982.
- (291) Taylor NF, Dodd KJ, Shields N, Bruder A. Therapeutic exercise in physiotherapy practice is beneficial: a summary of systematic reviews 2002-2005. *Australian Journal of Physiotherapy* 2007; 53(1):7-16.
- (292) Treatment of Parkinson's disease. *Neurol Sci* 2003; 24 Suppl 3:S165-S213.
- (293) Horstink M, Tolosa E, Bonuccelli U, Deuschl G, Friedman A, Kanovsky P et al. Review of the therapeutic management of Parkinson's disease. Report of a joint task force of the European Federation of Neurological Societies and the Movement Disorder Society- European Section. Part I: Early (uncomplicated) Parkinson's disease. *Eur J Neurol* 2006; 13(11):1170-1185.
- (294) Horstink M, Tolosa E, Bonuccelli U, Deuschl G, Friedman A, Kanovsky P et al. Review of the therapeutic management of Parkinson's disease. Report of a joint task force of the European Federation of Neurological Societies (EFNS) and the Movement Disorder Society-European Section (MDS-ES). Part II: late (complicated) Parkinson's disease. *Eur J Neurol* 2006; 13(11):1186-1202.
- (295) Rochester L, Hetherington V, Jones D, Nieuwboer A, Willems AM, Kwakkel G et al. The effect of external rhythmic cues (auditory and visual) on walking during a functional task in homes of people with Parkinson's disease. *Arch Phys Med Rehabil* 2005; 86(5):999-1006.

Reference List

- (296) Rochester L, Nieuwboer A, Baker K, Hetherington V, Willems AM, Chavret F et al. The attentional cost of external rhythmical cues and their impact on gait in Parkinson's disease: effect of cue modality and task complexity. *J Neural Transm* 2007; 114(10):1243-1248.
- (297) Kwakkel G. Impact of intensity of practice after stroke: issues for consideration. *Disabil Rehabil* 2006; 28(13-14):823-830.
- (298) Fried BJ, Boers M, Baker PRA. A Method for Achieving Consensus on Rheumatoid-Arthritis Outcome Measures - the Omeract Conference Process. *Journal of Rheumatology* 1993; 20(3):548-551.
- (299) U.S Food and Drug Administration. Draft guidance for industry on patient-reported outcome measures: use in medicinal product development to support labelling claims. www.fda.gov/cber/gdlns/probl.pdf [2006; cited 2008 Mar. 10].
- (300) Hobart JC, Cano SJ, Zajick JP, Thompson AJ. Rating scales as outcome measures for clinical trials in neurology: problems, solutions, and recommendations. *Lancet Neurol* 2007; 6(12):1094-1105.
- (301) Tugwell P, Bombardier C, Buchanan WW, Goldsmith CH, Grace E, Hanna B. The MACTAR Patient Preference Disability Questionnaire--an individualized functional priority approach for assessing improvement in physical disability in clinical trials in rheumatoid arthritis. *J Rheumatol* 1987; 14(3):446-451.
- (302) Stratford P, Gill C, Westaway M, Binkley J. Assessing disability and change on individual patients: a report of a patient specific measure. *Physiother Canada* 1995; 47:258-263.
- (303) Law M, Polatajko H, Pollock N, McColl MA, Carswell A, Baptiste S. Pilot testing of the Canadian Occupational Performance Measure: clinical and measurement issues. *Can J Occup Ther* 1994; 61(4):191-197.
- (304) Engers AJ, Wensing M, van Tulder MW, Timmermans A, Oostendorp RAB, Koes BW et al. Implementation of the Dutch low back pain guideline for general practitioners - A cluster randomized controlled trial. *Spine* 2005; 30(6):595-600.
- (305) Grimshaw J, Eccles M, Thomas R, MacLennan G, Ramsay C, Fraser C et al. Toward evidence-based quality improvement - Evidence (and its limitations) of the effectiveness of guideline dissemination and implementation strategies 1966-1998. *J Gen Intern Med* 2006; 21:S14-S20.
- (306) Munneke M, Keus SHJ, Nijkrake M, Kwakkel G, Berendse HW, Roos RAC et al. ParkNet: a multifaceted strategy to implement practice guidelines for physiotherapy in Parkinson's disease. *Parkinsonism Rel Disord* 2008; 14(Suppl 1):S64.
- (307) Wantland DJ, Portillo CJ, Holzemer WL, Slaughter R, McGhee EM. The effectiveness of Web-based vs. non-Web-based interventions: a meta-analysis of behavioral change outcomes. *J Med Internet Res* 2004; 6(4):e40.
- (308) Bloem BR, van Laar T, Keus SHJ, de Beer H, Poot E, Buskens E et al. [Multidisciplinary guidelines for Parkinson's disease]. Alphen a/d Rijn: Van Zuiden Communications, 2010.
- (309) Chowdhury MM, Dagash H, Pierro A. A systematic review of the impact of volume of surgery and specialization on patient outcome. *Br J Surg* 2007; 94(2):145-161.
- (310) Davoli M, Amato L, Minozzi S, Bargagli AM, Vecchi S, Perucci CA. [Volume and health outcomes: an overview of systematic reviews]. *Epidemiol Prev* 2005; 29(3-4 Suppl):3-63.