



Universiteit
Leiden
The Netherlands

Radio emission from merging galaxy clusters : characterizing shocks, magnetic fields and particle acceleration

Weeren, R.J. van

Citation

Weeren, R. J. van. (2011, December 20). *Radio emission from merging galaxy clusters : characterizing shocks, magnetic fields and particle acceleration.* Retrieved from <https://hdl.handle.net/1887/18259>

Version: Corrected Publisher's Version

[Licence agreement concerning inclusion of doctoral
thesis in the Institutional Repository of the University
of Leiden](#)

License: <https://hdl.handle.net/1887/18259>

Note: To cite this publication please use the final published version (if applicable).

Radio emission from merging galaxy clusters

characterizing shocks, magnetic fields and particle acceleration

Radio emission from merging galaxy clusters

characterizing shocks, magnetic fields and particle acceleration

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. P. F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op dinsdag 20 december 2011
klokke 15:00 uur

door

Reinout Johannes van Weeren

geboren te De Bilt
in 1980

Promotiecommissie

Promotores: Prof. dr. H. J. A. Röttgering
Prof. dr. G. K. Miley

Overige leden: Prof. dr. M. Brüggen (Jacobs University Bremen, Germany)
Prof. dr. A. G. de Bruyn (Stichting ASTRON, Dwingeloo;
Rijksuniversiteit Groningen)
Prof. dr. H. Falcke (Radboud Universiteit Nijmegen)
Prof. dr. M. A. Garrett (Stichting ASTRON, Dwingeloo;
Universiteit Leiden)
Prof. dr. W. Jaffe
Prof. dr. K. H. Kuijken

The cover shows a GMRT 150 MHz radio image of the field around the galaxy cluster CIZA J2242.8+5301. LOFAR antennas are visible at the bottom.

GMRT image by Huib Intema.

Cover design by Brigitta van Weeren.

ISBN: 978-94-6191-099-8

Contents

1	Introduction	1
1.1	Cosmology & large-scale structure formation	1
1.2	Galaxy clusters	2
1.3	Radio emission from galaxy clusters	2
1.3.1	Radio halos	3
1.3.1.1	Origin of radio halos	3
1.3.2	Radio relics	4
1.4	This thesis	5
1.5	Future prospects	8
2	A search for steep spectrum radio relics and halos with the GMRT	11
2.1	Introduction	12
2.2	Sample selection	14
2.3	Observations & data reduction	14
2.4	Results	16
2.4.1	Individual sources	16
2.4.1.1	VLSS J1133.7+2324	16
2.4.1.2	VLSS J1431.8+1331	18
2.4.1.3	VLSS J2217.5+5943, 24P73	19
2.4.1.4	VLSS J0004.9–3457	21
2.4.1.5	VLSS J0717.5+3745, MACS J0717.5+3745	21
2.4.1.6	VLSS J0915.7+2511	23
2.4.1.7	VLSS J1515.1+0424, Abell 2048	24
2.5	Spectral index modeling	24
2.6	Discussion	27
2.7	Conclusions	32
2.8	Appendix: Other sources in the sample	35

3	Diffuse steep-spectrum sources from the 74 MHz VLSS survey	41
3.1	Introduction	42
3.2	Observations & data reduction	43
3.2.1	GMRT 325 MHz observations	43
3.2.2	VLA 1.4 GHz observations	44
3.2.3	WSRT 1.3 – 1.7 GHz observations of 24P73	46
3.2.4	Optical WHT & INT imaging	47
3.3	Results	47
3.3.1	VLSS J1133.7+2324, 7C 1131+2341	48
3.3.2	VLSS J1431.8+1331, MaxBCG J217.95869+13.53470	50
3.3.3	VLSS J2217.5+5943, 24P73	52
3.3.4	VLSS J0004.9–3457	53
3.3.5	VLSS J0915.7+2511, MaxBCG J138.91895+25.19876	55
3.3.6	VLSS J1515.1+0424, Abell 2048	55
3.4	Optical imaging around five compact steep-spectrum sources	59
3.4.1	VLSS J2043.9–1118	59
3.4.2	VLSS J1117.1+7003	61
3.4.3	VLSS J2209.5+1546	61
3.4.4	VLSS J0516.2+0103	61
3.4.5	VLSS J2241.3–1626	62
3.5	Discussion	62
3.6	Conclusions	64
4	Radio continuum observations of new radio halos and relics from the NVSS and WENSS surveys: Relic orientations, cluster X-ray luminosity and redshift distributions	67
4.1	Introduction	68
4.2	Observations & data reduction	69
4.2.1	Radio observations	69
4.2.2	Optical WHT & INT images	71
4.3	Results	71
4.3.1	Abell 1612	71
4.3.2	Abell 746	76
4.3.3	Abell 523	76
4.3.4	Abell 697	79
4.3.5	Abell 2061	80
4.3.6	Abell 3365	84
4.3.7	CIZA J0649.3+1801	86
4.3.8	CIZA J0107.7+5408	89
4.3.9	Abell 2034	89
4.3.10	RXC J1053.7+5452	90
4.4	Discussion	90
4.4.1	Comparison with the REFLEX and NORAS X-ray clusters	94
4.4.2	X-ray peak and galaxy distribution separation	96
4.5	Conclusions	99

5 Diffuse radio emission in the merging cluster MACS J0717.5+3745: the discovery of the most powerful radio halo	101
5.1 Introduction	102
5.2 Observations & data reduction	103
5.3 Results: images, equipartition magnetic field & spectral index maps	104
5.3.1 Equipartition magnetic field strength	106
5.3.2 Spectral index	107
5.4 Discussion	109
5.4.1 Alternative explanations for the elongated radio structures	109
5.4.2 Origin of the radio relic	110
5.5 Conclusions	111
6 Radio observations of ZwCl 2341.1+0000: a double radio relic cluster	113
6.1 Introduction	114
6.2 Observations & data reduction	116
6.3 Results	121
6.4 Radio, X-ray, and galaxy distribution comparison	123
6.5 Spectral index & equipartition magnetic field strength	125
6.5.1 Spectral index	125
6.5.2 Equipartition magnetic field	126
6.6 Discussion	127
6.6.1 Alternative explanations	127
6.6.2 Comparison of spectral indices and magnetic field strengths with other double relics	128
6.6.3 Origin of the double relic	129
6.7 Conclusions	130
6.8 Appendix: Compact sources at 610 MHz and optical counterparts	132
7 A double radio relic in the merging galaxy cluster ZwCl 0008.8+5215	135
7.1 Introduction	136
7.2 Observations & data reduction	137
7.2.1 GMRT observations	137
7.2.2 WSRT 1.3–1.7 GHz observations	138
7.2.3 Optical images & spectroscopy	139
7.3 Results	140
7.3.1 Redshift of ZwCl 0008.8+5215	140
7.3.2 Thermal ICM and galaxy distribution	140
7.3.3 Radio continuum maps	141
7.3.4 Spectral index and polarization maps	144
7.3.5 Radio galaxies in the cluster	145
7.4 Discussion	147
7.4.1 Origin of the double radio relic	147
7.4.2 Radio luminosity profile for the eastern relic	151
7.4.3 Simulated radio luminosity and spectral index profiles	152
7.4.4 Equipartition magnetic field strength	153
7.5 Conclusions	154

8 Particle acceleration on megaparsec scales in a merging galaxy cluster	157
8.1 Appendix: Data reduction	165
9 The “toothbrush-relic”: evidence for a coherent linear 2-Mpc scale shock wave in a massive merging galaxy cluster?	169
9.1 Introduction	170
9.2 Observations & data reduction	173
9.2.1 GMRT observations	173
9.2.2 WSRT observations	174
9.2.3 WHT spectroscopy & imaging	174
9.3 Results: redshift, X-rays, and radio continuum maps	174
9.3.1 Redshift of 1RXS J0603.3+4214	174
9.3.2 X-ray emission from the ICM	175
9.3.3 Radio continuum maps	175
9.4 Radio spectra	178
9.4.1 Spectral index maps	178
9.4.2 Integrated radio spectra	182
9.5 Radio color-color diagrams	182
9.5.1 Spectral models	185
9.5.2 Effect of resolution on surface brightness	186
9.5.3 Color-color diagrams	187
9.5.3.1 Color-color diagrams for B1 and B2+B3	187
9.5.3.2 The effect of resolution and mixing	188
9.5.4 Global spectrum	188
9.5.4.1 Shift diagrams	188
9.6 RM-synthesis & polarization maps	189
9.6.1 Polarization maps	189
9.6.2 RM-synthesis	189
9.6.2.1 Application to the L-band WSRT data	195
9.6.3 The depolarization properties of the radio relic	195
9.7 Discussion	197
9.7.1 Relic spectra	198
9.7.2 Alternative models to explain the relativistic electrons from the radio relic	199
9.7.3 Magnetic field	200
9.8 Conclusions	200
10 Using double radio relics to constrain galaxy cluster mergers: A model of double radio relics in CIZA J2242.8+5301	203
10.1 Introduction	204
10.2 Overview: Simulations of galaxy cluster mergers	205
10.3 Numerical method	207
10.3.1 Radio emission from shocks	207
10.3.2 Initial conditions	209
10.4 Results	210
10.4.1 Radio maps	211
10.4.1.1 Mass ratio	211

10.4.1.2	Impact parameter	216
10.4.1.3	Viewing angle	216
10.4.1.4	β -model	219
10.4.1.5	Cool core	219
10.4.2	Spectral index	219
10.5	Substructure & clumping	221
10.6	Discussion	223
10.6.1	Effect of dark matter dynamics	223
10.6.2	Relic width and brightness profiles	223
10.6.3	Origin of single radio relics	224
10.6.4	A quantitative metric for the goodness of fit	224
10.7	Summary	225
11	The discovery of diffuse steep spectrum sources in Abell 2256	227
11.1	Introduction	228
11.2	Observations & data reduction	228
11.3	Results	230
11.3.1	Spectral indices	231
11.4	Discussion	232
11.5	Conclusions	233
12	LOFAR Abell 2256 observations between 18 and 67 MHz	235
12.1	Introduction	236
12.2	Observations & data reduction	237
12.2.1	Primary beam correction, absolute flux-scale and self-calibration	240
12.2.2	Radio images	242
12.2.3	Spectral index map and integrated fluxes	243
12.3	Discussion	245
12.3.1	Source F	245
12.3.2	Source AG+AH	246
12.3.3	Radio relic and halo	247
12.4	Conclusions	249
Bibliography		251
Nederlandse samenvatting		265
Curriculum vitae		271
Nawoord / Acknowledgements		273

