



Universiteit  
Leiden

The Netherlands

## **Magnetic resonance imaging in neonatal hypoxic-ischemic brain injury**

Liauw, L.

### **Citation**

Liauw, L. (2009, March 19). *Magnetic resonance imaging in neonatal hypoxic-ischemic brain injury*. Retrieved from <https://hdl.handle.net/1887/13690>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13690>

**Note:** To cite this publication please use the final published version (if applicable).

## *Chapter 6*

### **Do Apparent Diffusion Coefficient measurements predict outcome in children with neonatal hypoxic-ischemic encephalopathy?**

Lishya Liauw  
Gerda van Wezel-Meijler  
Sylvia Veen  
Mark van Buchem  
Jeroen van der Grond

American Journal of Neuroradiology, in press

## **Background and purpose**

Diffusion-weighted imaging (DWI) permits early detection and quantification of hypoxic-ischemic brain lesions. Our aim was to assess the predictive value of DWI and apparent diffusion coefficient (ADC) measurements for outcome in children with perinatal asphyxia.

## **Materials and methods**

Term neonates underwent MR imaging within 10 days after birth because of asphyxia. MR imaging examinations were retrospectively evaluated for hypoxic-ischemic brain damage. ADC was measured in 30 standardized brain regions and in visibly abnormal areas on DWI. In survivors, developmental outcome until early school age was graded into the following categories: 1) normal, 2) mildly abnormal, and 3) definitely abnormal. For analysis, category 3 and death were labeled 'adverse', 1 and 2 were 'favorable', and 2-3 and death were 'abnormal' outcome. Differences in outcome between infants with and without DWI abnormalities were analyzed using Chi-Squared tests. The nonparametric Mann-Whitney U test analyzed whether ADC values in visible DWI abnormalities correlated with age at imaging. Logistic regression analysis tested the predictive value for outcome of the ADC in each standardized brain region. Receiver Operating Characteristic analysis was used to find optimal ADC cutoff values for each region for the various outcome scores.

## **Results**

Twenty-four infants (13 male) were included. Mean age at MR imaging was 4.3 days (range 1-9 days). Seven infants died. There was no difference in outcome between infants with and without visible DWI abnormalities. Only ADC of the posterior limb of the internal capsule correlated with age. ADC in visibly abnormal DWI regions did not have a predictive value for outcome. Of all measurements performed, only the ADC in the normal-appearing basal ganglia and brain stem correlated significantly with outcome; low ADC values were associated with abnormal/adverse outcome, and higher ADC values, with normal/favorable outcome (basal ganglia:  $p = 0.03$  for abnormal;  $p = 0.01$  for adverse outcome; brain stem:  $p = 0.006$  for abnormal;  $p = 0.03$  for adverse outcome).

## **Conclusions**

ADC values in normal-appearing basal ganglia and brain stem correlated with outcome, independently of all MR imaging findings including those of DWI. ADC values in visibly abnormal brain tissue on DWI did not show a predictive value for outcome.

## **Acknowledgement**

We like to thank S. Servaas, MSc, for his help with data collection.

## Introduction

Neonatal hypoxic-ischemic brain injury is one of the most important causes of neurologic disabilities during childhood, accounting for more than half of the cases of cerebral palsy (1,2). MR imaging is an established tool to detect neonatal brain injury, and it plays an important role in the prediction of neurologic outcome in neonates with hypoxic-ischemic brain injury (2-6). MR imaging findings of the brains of neonates with hypoxic-ischemic encephalopathy have been described by many authors (1,3,4,6,7). Conventional T1- and T2-weighted images are used most often but findings may be normal despite significant brain injury, especially when imaging is performed early, during the first few days after the event (1,3,6,7).

Diffusion-weighted imaging (DWI) often reveals hypoxic-ischemic brain injury at an earlier stage than conventional MR imaging (8-11); in addition, it provides quantitative apparent diffusion coefficient (ADC) values in brain tissue (12-16). It has been proposed that these quantitative measures may have additional diagnostic value, especially in cases with diffuse lesions that may remain below the threshold of visual detection. Some articles have published normal values for ADC in different regions of the normal brain in young infants (12,13,16,17). Others have reported abnormal ADC values in severely abnormal (as shown by MR imaging) white matter and deep grey matter, during the first week after the hypoxic-ischemic event (11-13,18-22). However, in none of these studies were ADC values related to longer follow-up (11-13,18-22). In addition, in none of these studies outcomes were related to ADC in normal-appearing brain regions. To our knowledge, only one relatively small study in 11 neonates described the predictive value of DWI and ADC measurements for outcome at 24 months of age in children born asphyxiated (23). Predictive values of DWI or ADC were not calculated. In that study, DWI was measured only in the basal ganglia and periventricular white matter. No discrimination between regions with visible DWI abnormalities and normal-appearing regions was made.

In the present study, we investigated the value of DWI for outcome prediction until early school age in neonates with hypoxic-ischemic encephalopathy. Our aim was to assess the following: 1) the relation between visible DWI abnormalities and outcome, and 2) the relation to outcome of ADC values in regions with and without visible DWI abnormalities.

## Materials and methods

### Subjects

The institutional review board approved this retrospective study and waived informed consent. Brain MR imaging studies performed between January 2001 and December 2003 in infants who were born at a gestational age of > 37 weeks and who underwent MR imaging to assess brain injury due to perinatal asphyxia

were selected as previously described (24). All MR imaging examinations were performed within 10 days after birth. All 24 infants included in our study were born at or transferred to our institution and were admitted to our tertiary neonatal intensive care unit. They were included if they fulfilled all the following criteria: 1) signs of fetal distress before delivery (abnormal cardiotocographic recording such as decreased variability, late deceleration, baseline bradycardia); 2) Apgar score of < 7 after 5 minutes; 3) umbilical cord blood pH level of < 7.2; 4) clinical signs of hypoxic-ischemic encephalopathy (25); and 5) radiologic evidence of hypoxic-ischemic brain injury on MR imaging as reported by the attending neuroradiologist.

Evaluation of the clinical history, retrospectively based on data from individual patient documentation, was performed by one author (S.V., having 23 years of experience in neonatology and neonatal follow-up). This author was blinded to the results of the brain MR imaging examinations and ADC measurements. Neonatal clinical classification of hypoxic-ischemic encephalopathy (HIE) was retrospectively graded according to Sarnat and Sarnat (25). HIE stage 1 indicates a mild encephalopathy, lasting < 24 hours with normal findings on electroencephalography (EEG). HIE stage 2 indicates a moderate encephalopathy, with the infant being lethargic and hypotonic and having seizures. In HIE stage 3, there is a severe encephalopathy with the infant being comatose and severely hypotonic with decreased or absent reflex activity and a severely depressed EEG finding (25).

Surviving infants visited the outpatient clinic of our institution or the referring hospital at regular intervals, at the ages of three months, one year, two years, and five years. If necessary as judged by the clinical condition, additional visits were scheduled. All surviving infants underwent developmental and neurologic examinations performed by experienced and especially trained neonatologists, pediatricians, or child neurologists. In surviving children, developmental outcome at clinical follow-up was graded as follows (21): 1) normal – that is, normal neurologic examination findings, normal cognitive developmental history, and normal Van Wiechen examination findings (a Dutch child health care developmental assessment tool based on the developmental scales of Gesell and adapted by Touwen and Hempel (26,27)); 2) mildly abnormal – that is, mildly abnormal neurologic findings (mild hypertonia, hypotonia and/or asymmetry) and/or mildly abnormal cognitive developmental history (such as mild speech and language delay, mild behavioral disorders and/or the need for remedial teaching at school) and/or mildly abnormal Van Wiechen examination findings; and 3) definitely abnormal – that is, children with major neurologic problems, such as cerebral palsy and/or severe cognitive function disorders and/or epilepsy and/or severely abnormal Van Wiechen examination findings. Outcome was classified in category 4 in case of neonatal death. For analysis, outcome categories 1 and 2 were labeled 'favorable'; categories 3 and 4, 'adverse'; and categories 2-4, 'abnormal'. Outcome was retrospectively graded by one author (S.V.), again blinded to the MR imaging findings and also to the neonatal clinical classification. Outcome classification was based on a compilation of all available information of all follow-up visits.

## MR Imaging

If necessary, the infants were sedated. In ventilated sedated infants or infants on antiepileptic medication, this sedation was performed by increasing the dosage of the sedative or antiepileptic medication just before the MR imaging procedure. Those who were not on sedatives or antiepileptic medication received chloral hydrate, 50-70 mg/kg, administered orally 30 minutes before imaging. Infants were positioned supine and snugly swaddled-up in blankets during the imaging procedure. Ear protection was used and consisted of commercially available neonatal ear muffs (MiniMuffs; Natus Medical, San Carlos, Calif) placed over the ears. The infant's head was immobilized by molded foam put around the head during the imaging procedure. The temperature was maintained and heart rate and oxygen saturation were monitored throughout the procedure. A physician experienced in neonatal resuscitation was always present during imaging and transportation to and from the MR imaging unit.

Images were obtained with superconducting magnets (Gyrosan ACS-NT 15; Philips Medical Systems, Best, the Netherlands) operating at a field-strength of 1.5 T. T1-weighted spin-echo sequences (TR/TE, 550-560/14-20 ms), T2-weighted spin-echo sequences (TR/TE, 5406-6883/100-120 ms), fluid-attenuated inversion recovery sequences (TR/TE, 6000-8000/110-120 ms; TI, 2000 ms), T1-weighted gadolinium-enhanced spin-echo sequences (TR/TE, 550-560/14-20 ms), and DWI were performed. DWI was performed by using a single-shot spin-echo echo-planar sequence (TR/TE, 5132-4200/74 ms) with a b-value of 1000 s/mm<sup>2</sup>. All sequences were performed in transverse planes. Section thickness was 4 - 5 mm with an intersection gap of 0.4 - 0.5 mm. For DWI, section thickness was 6 mm. Because this was a retrospective study, MR spectroscopy was not routinely performed in this patient group.

## MR Imaging Data Collection

All MR images were retrospectively evaluated by two authors, a neonatologist and neuroradiologist, both experienced in neonatal neuroimaging (G.v.W.-M. and L.L.; G.v.W.-M. having sixteen and L.L. having nine years of experience). Both reviewers were blinded to the infant's identities and thus to the clinical history of the infants. DWI abnormalities were independently assessed on a 5-point scale. The confidence of the investigator for the detection of the abnormality was scored with a diagnostic confidence score, assigned by using a 5-point scale; definitely abnormal, probably abnormal, equivocal, probably normal, or definitely normal. In a consecutive consensus reading, final DWI abnormalities were assessed. In this consensus reading, conventional T1-and T2-weighted images were also assessed, and it was indicated whether brain injury was present and which type of injury pattern was present in each infant. Consensus was reached in all cases (24).

## ADC measurements

ADC measurements were independently performed in 30 standardized regions of the brain by one author (L.L.) who was blinded to the patient's identity, by using Philips proprietary software. In addition, ADC measurements were performed in brain regions that appeared visibly abnormal on DWI. All measurements were

retrospectively performed on the MR imaging operator console. All MR imaging examinations were reloaded from the archives. Trace maps were used to calculate ADC values in all regions of interest. Because of image deformation in the DWI acquisition, regions of interest were identified on the original T2-weighted images and visually matched and positioned on the b=0 image of the DWI acquisition, which can be considered as fast T2-weighted echo-planar MR imaging. From there, the region of interest was copied onto the corresponding ADC map. Oval regions of interest were used for all regions. Regions of interest were drawn to include as much of the structure as possible. Care was taken to avoid inclusion of cerebrospinal fluid in any region of interest. In both hemispheres, regions of interest were drawn in the white matter of the semioval center anteriorly and posteriorly, close to the central sulcus, head of the caudate nucleus, putamen and globus pallidus, posterior limb of the internal capsule (PLIC), frontal and posterior white matter at the level of the basal ganglia, medial and lateral part of the thalamus, middle cerebellar peduncle, temporal and occipital white matter, white matter in the cerebellar hemispheres, and the cerebral peduncles in the brain stem at the level of the red nucleus. Care was taken not to place the region of interest in the dentate nucleus or the cerebellar folia when measuring the cerebellar white matter. The area of each region of interest was 0.3 –1 cm<sup>2</sup>, except in the PLIC and central sulcus, where the region of interest areas were 0.15-0.2 cm<sup>2</sup>.

### **Statistical analysis**

All statistical analyses were performed by one author (J.v.d.G.) by using the Statistical Package for the Social Sciences software (SPSS for Windows, Version 14.0.2; SPSS, Chicago, Ill). To study whether outcome (adverse - favorable, abnormal - normal) differed between infants with and without visible DWI abnormalities, we used Chi-Squared tests. The nonparametric Mann-Whitney U test was used to analyze whether ADC values in visible DWI abnormalities and in normal-appearing brain regions displaying no visible abnormalities on conventional MR imaging, DWI, and ADC images correlated with the postnatal age at imaging. To investigate the predictive value on outcome of ADC values in each brain region, we used logistic regression analysis in which age at imaging was entered as a covariate. Receiver Operating Characteristic analysis was used to find optimal ADC cutoff values for each region for the various developmental outcome scores. A p < 0.05 was considered statistically significant.

## **Results**

### **Outcome**

Twenty-four infants (13 male) were included in this study. Mean age at MR imaging was 4.3 days (range, 1-9 days), mean postconceptional age was 41 weeks (range, 37<sup>+4</sup> -43<sup>+3</sup> weeks). Of the infants who were included in this study, ten were sedated; the other 15 infants did not require (additional) medication. MR imaging showed no prevailing hypoxic-ischemic injury pattern. Table 1 shows outcome in

relation to the neonatal clinical classification. Outcome of the surviving infants was scored at the minimum age of two years (range, 2.08 - 5.75 years; mean, 3.75 years; median, 3.46 years; standard deviation, 1.28). In all seven children (four males) who died (mean, 8.4 days after birth; median, 7 days; range, 4-13 days), the cause of death was respiratory failure and/or multi-organ failure due to perinatal asphyxia.

**Table 1** Outcome in relation to neonatal clinical classification.

Neonatal classification according to Sarnat and Sarnat <sup>25</sup>	No	Outcome			
		N n=6	MA n=1	DA n=10	D n=7
<b>HIE grade 1</b>	3	2	0	1	0
<b>HIE grade 2</b>	18	4	1	9	4
<b>HIE grade 3</b>	3	0	0	0	3

N= normal

MA= mildly abnormal


DA= definitely abnormal

D= died

### DWI findings

Despite considerable artifacts in the posterior fossa on DWI images, the cerebellum, and brain stem could be assessed well for their signal intensity (SI). In 21 of the 24 neonates, the following 36 abnormalities were seen on DWI (Figure 1): 5 SI changes in the basal ganglia, 1 SI change in the thalamus, and 6 SI changes in the PLIC. Five abnormalities were recognized as arterial infarction; 6, as borderzone infarction; and 3 lesions, as punctate white matter lesions. In ten infants, the white matter of the cerebral hemispheres had diffusely abnormal signal intensity on DWI.

There was no difference in outcome between infants with and without visible DWI abnormalities. A significant correlation between ADC values in visible DWI abnormalities (in  $10^{-6}/\text{mm}^2/\text{s}$ ) and age at imaging was found only for the PLIC, showing a strong correlation between ADC values and postnatal age at imaging ( $R=0.823$ ,  $p=0.003$ ). ADC values in visibly abnormal regions on DWI did not have any predictive value for outcome (corrected for age at imaging).



**Figure 1**

(**a**) T1-weighted image: TR/TE, 550/14 ms; signals acquired, two; matrix, 205/256; section thickness, 5 mm; section gap, 0.5 mm; FOV, 16 cm. (**b**) T2-weighted image: TR/TE, 5405/120 ms; signals acquired, two; matrix, 205/256; section thickness, 5 mm; section gap, 0.5 mm; FOV, 16 cm. (**c**) B=1000 DWI image: TR/TE, 4299/74 ms; section thickness, 6 mm; b-value of 1000 s/mm<sup>2</sup>. (**d**) ADC image. These images are of an infant of outcome category 3 (definitely abnormal outcome at four years nine months of age). The T1-weighted image (**a**) shows irregular high signal intensity in and around the peri-Rolandic cortex (arrow). The T2-weighted image (**b**) shows abnormal high signal intensity in the subcortical white matter around the peri-Rolandic area (arrow). The b=1000 DWI (**c**) shows high signal intensity in the peri-Rolandic area and the central white matter (arrows). The ADC image (**d**) shows low signal intensity in the peri-Rolandic area and the central white matter (arrows).

## Normal-appearing brain tissue findings

Normal-appearing brain tissue displayed no visible abnormalities on conventional MR imaging, DWI, and ADC images. In four normal-appearing brain regions, ADC was significantly related to age at imaging. Higher ADC values were found with increasing age in the occipital white matter ( $R = 0.622$ ,  $p = 0.011$ ), temporal white matter ( $R = 0.578$ ,  $p = 0.024$ ), PLIC ( $R = 0.501$ ,  $p = 0.025$ ), and medial thalamus ( $R = 0.451$ ,  $p = 0.027$ ). In none of the other normal-appearing regions was a significant correlation found.

Table 2 shows the ADC values in the normal-appearing regions (on conventional MR imaging and DWI) for neonates of the favorable (categories 1 - 2) and adverse (category 3 - death) outcome groups. In the normal-appearing basal ganglia and brain stem, we found significantly lower ADC values in the subgroup of neonates with abnormal/adverse outcome than in infants with normal/favorable outcome. The ADC in both regions correlated significantly with outcome (corrected for age at imaging: basal ganglia,  $p = 0.03$  for abnormal,  $p = 0.01$  for adverse outcome; and brain stem,  $p = 0.006$  for abnormal,  $p = 0.03$  for adverse outcome). For none of the other normal-appearing brain regions were significant correlations found with outcome.

Tables 3-6 show the results of the Receiver Operating Characteristic analysis analyses (sensitivity, specificity, positive- and negative predictive values (PPVs, NPVs), and multiple cutoff values of ADC of the basal ganglia and brain stem) for abnormal (outcome categories, 2-3 and death) and adverse (outcome category 3 and death) outcomes, respectively. For abnormal outcome, the ADC cutoff level of the basal ganglia for a 100% sensitivity (0% specificity) is  $1357 \times 10^{-6}/\text{mm}^2/\text{s}$  and for a 100% specificity (67% sensitivity)  $1018 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.035$ ). For adverse outcome, the ADC cutoff level for a 100% sensitivity (0% specificity) is  $1357 \times 10^{-6}/\text{mm}^2/\text{s}$  and for a 100% specificity (71% sensitivity)  $1018 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.014$ ). For abnormal outcome, the ADC cutoff level of the brain stem for a 100% sensitivity (33% specificity) is  $1182 \times 10^{-6}/\text{mm}^2/\text{s}$  and for a 100% specificity (67% sensitivity)  $988 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.006$ ). For adverse outcome, the ADC cutoff level for a 100% sensitivity (29% specificity) is  $1182 \times 10^{-6}/\text{mm}^2/\text{s}$  and for a 100% specificity (47% sensitivity)  $897 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.03$ ).

For abnormal outcome, the ADC cutoff level of the basal ganglia for a 100% PPV (55% NPV) is  $1018 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.035$ ). For adverse outcome, the ADC cutoff level for a 100% PPV (64% NPV) is  $1018 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.014$ ). For abnormal outcome, the ADC cutoff level of the brain stem for a 100% PPV (50% NPV) is  $988 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.006$ ). For adverse outcome, the ADC cutoff level for a 100% PPV (44% NPV) is  $897 \times 10^{-6}/\text{mm}^2/\text{s}$  ( $p = 0.03$ ).

**Table 2** ADC data of the normal-appearing brain tissue for the *favorable* and *adverse* outcome categories. Data are shown calculated from the ADC value (in  $10^{-6}/\text{mm}^2/\text{s}$ ).

	Mean		Std. deviation		Range	
	Favorable	Adverse	Favorable	Adverse	Favorable	Adverse
<b>Semioval center anterior</b>	1394.6	1337	62.3	228	1331-1479	979-1644
<b>Semioval center posterior</b>	1295.3	1282.5	41.8	258.8	1219-1346	943-1775
<b>Central sulcus</b>	983.9	977.9	84.6	147	806-1122	724-1248
<b>Basal ganglia</b>	1115.3*	994*	62.5	181.2	1033-1206	718-1388
<b>Caudate head</b>	1160.5	1120.4	77.8	173.9	1017-1298	838-1443
<b>Frontal wm</b>	1561.9	1499.1	158.6	273.8	1311-1768	1057-1942
<b>Posterior wm</b>	1483.1	1550.5	207.4	176.9	1133-1794	1247-1791
<b>Medial thalamus</b>	953.5	996.1	81.9	183.1	826-1103	756-1401
<b>Lateral thalamus</b>	930.6	939.7	83	205.6	810-1033	664-1412
<b>PLIC</b>	965.9	904.7	77.2	176.4	866-1099	626-1160
<b>Brain stem</b>	1075*	952.7*	108.6	137.2	920-1242	724-1182
<b>Cerebellum</b>	1106.6	1217.2	158.9	110.3	820-1384	975-1447
<b>Occipital wm</b>	1475	1332.4	156.4	198.8	1270-1706	984-1628
<b>Temporal wm</b>	1360	1318.6	62.2	144.5	1289-1474	1038-1611
<b>Middle cerebellar peduncle</b>	878.9	878.7	119.9	109.7	727-1115	655-1151

wm = white matter

\*: p <0.05 between Favorable and Adverse outcome

**Table 3** ROC analysis showing sensitivity, specificity, positive and negative predictive values for *abnormal* outcome of multiple cutoff levels of ADC (in  $10^{-6}/\text{mm}^2/\text{s}$ ) of the basal ganglia\*.

Cutoff level	Sensitivity	95% CI†	Specificity	95% CI†	PPV	NPV
<b>&lt;574</b>	6.67	1.1 - 32.0	100.00	54.1 - 100.0	100.0	30.0
<b>&lt;1018.5</b>	66.67	38.4 - 88.1	100.00	54.1 - 100.0	100.0	54.5
<b>&lt;1036.5</b>	66.67	38.4 - 88.1	83.33	36.1 - 97.2	90.9	50.0
<b>&lt;1109.5</b>	80.00	51.9 - 95.4	50.00	12.4 - 87.6	80.0	50.0
<b>&lt;1174</b>	86.67	59.5 - 98.0	16.67	2.8 - 63.9	72.2	33.3
<b>&lt;1239</b>	93.33	68.0 - 98.9	0.00	0.0 - 45.9	70.0	0.0

Note:-ROC indicates receiver Operating Characteristic analysis; CI, confidence interval.

\* Area under the ROC curve=0.767. Significance level P (area=0.5)=0.0347.

† 95% CI=0.533-0.920.

**Table 4** ROC analysis showing sensitivity, specificity, positive and negative predictive values for *adverse* outcome of multiple cutoff levels of ADC (in  $10^{-6}$  mm<sup>2</sup>/s) of the basal ganglia\*.

Cutoff level	Sensitivity	95% CI†	Specificity	95% CI†	PPV	NPV
<574	7.14	1.2 – 33.9	100.00	58.9 – 100.0	100.0	35.0
<1018.5	71.43	41.9 – 91.4	100.00	58.9 – 100.0	100.0	63.6
<1061.5	78.57	49.2 – 95.1	71.43	29.3 – 95.5	84.6	62.5
<1104.5	78.57	49.2 – 95.1	57.14	18.8 – 89.6	78.6	57.1
<1148	85.71	57.2 – 97.8	28.57	4.5– 70.7	70.6	50.0
<1239	92.86	66.1 – 98.8	0.00	0.0 – 41.1	65.0	0.0

\* Area under the ROC curve=0.786. Significance level P (area=0.5)=0.0135.

† 95% CI=0.554-0.931.

**Table 5** ROC analysis showing sensitivity, specificity, positive and negative predictive values for *abnormal* outcome of multiple cutoff levels of ADC (in  $10^{-6}$  mm<sup>2</sup>/s) of the brain stem\*.

Cutoff level	Sensitivity	95% CI†	Specificity	95% CI†	PPV	NPV
<736.5	5.56	0.9 – 27.4	100.00	54.1 – 100.0	100.0	26.1
<988	66.67	41.0 – 86.6	100.00	54.1 – 100.0	100.0	50.0
<995	66.67	41.0 – 86.6	83.33	36.1 – 97.2	92.3	45.5
<1093	83.33	58.6 – 96.2	50.00	12.4 – 87.6	83.3	50.0
<1151.5	94.44	72.6 – 99.1	33.33	5.3 – 77.3	81.0	66.7
<1182	100.00	81.3 – 100.0	33.33	5.3 – 77.3	81.8	100.0

\* Area under the ROC curve=0.815. Significance level P (area=0.5)=0.0061.

† 95% CI=0.605-0.941.

**Table 6** ROC analysis showing sensitivity, specificity, positive and negative predictive values for *adverse* outcome of multiple cutoff levels of ADC (in  $10^{-6}$  mm<sup>2</sup>/s) of the brain stem\*.

Cutoff level	Sensitivity	95% CI†	Specificity	95% CI†	PPV	NPV
<736.5	5.88	1.0 – 28.8	100.00	58.9 – 100.0	100.0	30.4
<897	47.06	23.0 – 72.1	100.00	58.9 – 100.0	100.0	43.7
<942.5	47.06	23.0 – 72.1	85.71	42.2 – 97.6	88.9	40.0
<1151.5	94.12	71.2 – 99.0	28.57	4.5 – 70.7	76.2	66.7
<1182	100.00	80.3 – 100.0	28.57	4.5 – 70.7	77.3	100.0

\* Area under the ROC curve=0.756. Significance level P (area=0.5)=0.0305.

† 95% CI=0.540-0.906.

## Discussion

The aim of this study was to investigate the value of DWI and ADC for outcome prediction until early school age in neonates with hypoxic-ischemic encephalopathy. While ADC values in visibly abnormal brain tissue on DWI did not correlate with outcome, ADC of the normal-appearing basal ganglia and brain stem correlated with outcome, independently of all MR imaging findings, including DWI.

Apart from case reports, only a few published studies have so far addressed the predictive value of ADC measurements in neonates with hypoxic-ischemic brain damage (10,12,13,21,23). Wolf et al. (12) studied 13 term neonates with suggested hypoxic-ischemic encephalopathy within 12 days after birth. Clinical follow-up was short, but not specified. ADC values were measured in predefined brain regions; no distinction was made between regions with visible DWI abnormalities and normal-appearing regions. Wolf et al. (12) found decreased ADC values in the PLIC, corona radiata, posterior frontal white matter, and parietal white matter in patients with hypoxic-ischemic encephalopathy compared with control subjects, but they did not assess the predictive value of ADC values for outcome.

Rutherford et al. (13) studied 63 term neonates who presented with early seizures, of whom 49 with hypoxic-ischemic encephalopathy had undergone repetitive MR imaging. Reduced ADC in the centrum semiovale, anterior and posterior white matter at the level of the basal ganglia, central sulcus, lentiform nucleus, medial and lateral thalamus, PLIC, brainstem, and cerebellum allowed early detection of infarction. However, this study did not investigate the relation between ADC and long-term clinical follow-up.

Boichot et al. (21) studied the predictive value of conventional MR imaging and DWI for neurologic outcome in 30 term neonates with severe hypoxic-ischemic encephalopathy (21). Imaging was performed in the first twelve days after birth; outcome was evaluated after at least twelve months. A mixed pattern of cortical and basal ganglia signal intensity abnormalities on MR imaging was associated with unfavorable outcome. Significantly reduced ADC values were noted in all measured brain regions, especially the basal ganglia, in the unfavorable outcome group but only if MR imaging was performed during the first six days after birth.

To our knowledge, the study by L'Abée et al. (23) is the only study relating DWI to longer follow-up until 24 months of age in a group of 11 asphyxiated neonates. ADC values were measured in the basal ganglia and the periventricular white matter of the parietooccipital region, and neurodevelopment was assessed in survivors at 24 months. Nine of the 11 neonates had a poor outcome. ADC values in the basal ganglia were abnormally low in five and in the parietal white matter in four of the seven patients who died. Although low ADC values had predictive value, normal ADC values were also seen in children with poor outcome. In none of these studies ADC was related to follow-up longer than infancy.

The most important result of our study is that ADC values in the basal ganglia and brain stem were predictive of outcome. The ADC values in the *normal-appearing* brain stem and basal ganglia in infants with perinatal asphyxia correlated with abnormal or adverse outcome, *independently* of all MR imaging findings including DWI; low ADC values were associated with abnormal/adverse outcome and higher ADC values, with normal/favorable outcome. Normal-appearing brain tissue displayed no visible abnormalities on conventional MR imaging, DWI, and ADC images. Our finding is in accordance with several *conventional* MR imaging studies showing motor outcome to be related to the severity of basal ganglia and internal capsule involvement in infants with hypoxic-ischemic encephalopathy (4,5,28-30).

Our finding is also in agreement with the results of several other studies on ADC measurements in neonatal hypoxic-ischemic encephalopathy. Boichot et al. (21) found reduced ADC in the basal ganglia in infants with abnormal outcomes after neonatal hypoxic-ischemic encephalopathy. However, they did not discriminate between regions with visible DWI abnormalities and normal-appearing regions. L'Abée et al. (23) found abnormally low ADC values in the basal ganglia in five of seven patients who died. Again, no discrimination between regions with visible DWI abnormalities and normal-appearing regions was made. In a group of neonates with hypoxic-ischemic encephalopathy, Vermeulen et al. (20) showed lower ADC in the brain stem compared with that of controls. They did not relate ADC values to outcome. Our study also seems to be in agreement with Barkovich et al. (31), who showed that on follow-up imaging, abnormalities may be seen in regions of formerly normal-appearing white and grey matter. In other words, brain tissue may be abnormal despite normal appearance on (early) MR imaging. We, therefore, hypothesize that quantitative ADC measurements are more reliable in detecting abnormal brain tissue than visual analysis of conventional and DWI images.

In our clinical setting, the other important finding was that ADC values in visibly abnormal regions on DWI did not have any significant predictive value for outcome (corrected for age at imaging). There are two probable explanations for this remarkable finding. First, we studied a group of infants with hypoxic-ischemic encephalopathy having various hypoxic-ischemic brain lesions. Possibly, if we had selected a large group of infants with a specific type of brain injury, we would have found a predictive value of ADC for outcome in abnormal-appearing regions as did Hunt et al. (19). They showed that high ADC values in the PLIC were associated with survival in term infants with hypoxic-ischemic encephalopathy. The lowest ADC values in the PLIC were found in survivors with severe disability and in survivors with global brain injury. Second, we imaged infants between one and ten days after birth. Although 13 of 24 infants underwent imaging between one and four days after birth and only two infants were older than seven but younger than ten days, timing of imaging was still variable. It is well known that ADC values vary with time after hypoxia-ischemia as a result of the ongoing pathologic processes and that areas of abnormal diffusivity may 'pseudonormalize' within the first two weeks of life (31).

We found a strong association only between the ADC value of the PLIC, either visibly abnormal or normal, and postnatal age at imaging. Other authors have reported abnormal ADC values in injured brain tissue during the first week after the hypoxic-ischemic event; ADC values later became higher before returning to normal values (11-13,18-22). Our finding may be explained by the fact that our group was relatively small and heterogeneous. It is possible that if a larger number of infants had been studied, we would have found associations between ADC values and postnatal age at imaging in more structures.

As mentioned above, an imperfection of our study may be the timing of the MR imaging examinations, taking place up to ten days after birth, whereas on DWI, signal intensity abnormalities gradually diminish and are most obvious between one and four days after the incident (11,13,32). Although postnatal age at imaging was used as a covariate in our analyses, it would have been more appropriate to consider time since the inciting event. However, in most cases of perinatal asphyxia, the exact time of inciting event is difficult to determine (1). Another limitation is that we did not perform follow-up imaging while in the first two weeks after birth; normal ADC values do not exclude the presence of abnormal tissue in neonates with hypoxic-ischemic brain damage (13,31). Finally, because of the heterogeneity of the group we studied, showing various injury patterns, the number of infants with specific injury patterns was limited. This may have influenced our results, showing no predictive value of ADC in brain tissue appearing abnormal on DWI.

## **Conclusions**

We conclude that in neonates with hypoxic-ischemic encephalopathy, low ADC values of the normal-appearing basal ganglia and brain stem correlate with abnormal or adverse outcome, independently of all imaging findings. In this group of neonates with heterogeneous injury patterns, ADC values in visibly abnormal brain tissue on DWI did not have predictive value for outcome.

# References

1. Cowan F, Rutherford M, Groenendaal F, et al. Origin and timing of brain lesions in term infants with neonatal encephalopathy. *Lancet* 2003;361:736-742.
2. Cowan F. Outcome after intrapartum asphyxia in term infants. *Semin Neonatol* 2000;5:127-140.
3. Rutherford MA. The asphyxiated term infant. In: Rutherford MA, ed. *MRI of the neonatal brain*. London: W.B. Saunders; 2002;99-128.
4. Rutherford MA, Pennock JM, Counsell SJ, et al. Abnormal magnetic resonance signal in the internal capsule predicts poor neurodevelopmental outcome in infants with hypoxic-ischemic encephalopathy. *Pediatrics* 1998;102:323-328.
5. Cowan FM, de Vries LS. The internal capsule in neonatal imaging. *Semin Fetal Neonatal Med* 2005;10:461-474.
6. Barkovich AJ. Brain and spine injuries in infancy and childhood. In: Barkovich AJ, ed. *Pediatric Neuroimaging*. Philadelphia, PA: Lippincott, Williams & Wilkins; 2005;190-290.
7. Rutherford M, Pennock J, Schwieso J, Cowan F, Dubowitz L. Hypoxic-ischaemic encephalopathy: early and late magnetic resonance imaging findings in relation to outcome. *Arch Dis Child Fetal Neonatal Ed* 1996;75:F145-F151.
8. Cowan FM, Pennock JM, Hanrahan JD, Manji KP, Edwards AD. Early detection of cerebral infarction and hypoxic ischemic encephalopathy in neonates using diffusion-weighted magnetic resonance imaging. *Neuropediatrics* 1994;25:172-175.
9. Inder T, Huppi PS, Zientara GP, et al. Early detection of periventricular leukomalacia by diffusion-weighted magnetic resonance imaging techniques. *J Pediatr* 1999;134:631-634.
10. Liu AY, Zimmerman RA, Haselgrove JC, Bilaniuk LT, Hunter JV. Diffusion-weighted imaging in the evaluation of watershed hypoxic-ischemic brain injury in pediatric patients. *Neuroradiology* 2001;43:918-926.
11. Forbes KP, Pipe JG, Bird R. Neonatal hypoxic-ischemic encephalopathy: detection with diffusion-weighted MR imaging. *AJNR* 2000;21:1490-1496.
12. Wolf RL, Zimmerman RA, Clancy R, Haselgrove JH. Quantitative apparent diffusion coefficient measurements in term neonates for early detection of hypoxic-ischemic brain injury: initial experience. *Radiology* 2001;218:825-833.
13. Rutherford M, Counsell S, Allsop J, et al. Diffusion-weighted magnetic resonance imaging in term perinatal brain injury: a comparison with site of lesion and time from birth. *Pediatrics* 2004;114:1004-1014.
14. Neil JJ, Shiran SI, McKinstry RC, et al. Normal brain in human newborns: apparent diffusion coefficient and diffusion anisotropy measured by using diffusion tensor MR imaging. *Radiology* 1998;209:57-66.
15. Lovblad KO, Schneider J, Ruoss K, Steinlin M, Fusch C, Schroth G. Isotropic apparent diffusion coefficient mapping of postnatal cerebral development. *Neuroradiology* 2003;45:400-403.
16. Tanner SF, Ramenghi LA, Ridgway JP, et al. Quantitative comparison of intrabrain diffusion in adults and preterm and term neonates and infants. *AJR* 2000;174:1643-1649.
17. Toft PB, Leth H, Peitersen B, Lou HC, Thomsen C. The apparent diffusion coefficient of water in gray and white matter of the infant brain. *J Comput Assist Tomogr* 1996;20:1006-1011.
18. Barkovich AJ, Westmark KD, Bedi HS, Partridge JC, Ferriero DM, Vigneron DB. Proton spectroscopy and diffusion imaging on the first day of life after perinatal asphyxia: preliminary report. *AJNR* 2001;22:1786-1794.
19. Hunt RW, Neil JJ, Coleman LT, Kean MJ, Inder TE. Apparent diffusion coefficient in the posterior limb of the internal capsule predicts outcome after perinatal asphyxia. *Pediatrics* 2004;114:999-1003.
20. Vermeulen RJ, Fetter WP, Hendriks L, van Schie PE, van der Knaap MS, Barkhof F. Diffusion-weighted MRI in severe neonatal hypoxic ischaemia: the white cerebrum. *Neuropediatrics* 2003;34:72-76.
21. Boichot C, Walker PM, Durand C, et al. Term neonate prognoses after perinatal asphyxia: contributions of MR imaging, MR spectroscopy, relaxation times, and apparent diffusion coefficients. *Radiology* 2006;239:839-848.

22. Zarifi MK, Astrakas LG, Poussaint TY, Plessis AA, Zurakowski D, Tzika AA. Prediction of adverse outcome with cerebral lactate level and apparent diffusion coefficient in infants with perinatal asphyxia. *Radiology* 2002;225:859-870.
23. L'Abée C, de Vries LS, van der Grond J, Groenendaal F. Early diffusion-weighted MRI and 1H-Magnetic Resonance Spectroscopy in asphyxiated full-term neonates. *Biol Neonate* 2005;88:306-312.
24. Liauw L, van der Grond J, van den Berg-Huysmans AA, Palm-Meinders IH, van Buchem MA, van Wezel-Meijler G. Hypoxic-ischemic encephalopathy: diagnostic value of conventional MR imaging pulse sequences in term-born neonates. *Radiology* 2008;247:204-212.
25. Sarnat HB, Sarnat MS. Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. *Arch Neurol* 1976;33:696-705.
26. Gesell A, Amatruda CS. Developmental diagnosis: normal and abnormal child development. In: Knobloch H, Pasamanick B, Hagerstown MD, eds. *Developmental diagnosis; the evaluation and management of normal and abnormal neuropsychologic development in infancy and early childhood*. 3rd ed. New York: Harper and Row; 1974.
27. Brouwers-de Jong EA, Burgmeijer RJF, Laurent de Angulo MS. *Ontwikkelingsonderzoek op het consultatiebureau. Handboek bij het vernieuwde Van Wiechenonderzoek*. Assen, the Netherlands: van Gorcum; 1996: 1-343 (in Dutch).
28. Krageloh-Mann I, Helber A, Mader I, et al. Bilateral lesions of thalamus and basal ganglia: origin and outcome. *Dev Med Child Neurol* 2002;44:477-484.
29. Mercuri E, Barnett AL. Neonatal brain MRI and motor outcome at school age in children with neonatal encephalopathy: a review of personal experience. *Neural Plast* 2003;10:51-57.
30. Rutherford M, Srinivasan L, Dyet L, et al. Magnetic resonance imaging in perinatal brain injury: clinical presentation, lesions and outcome. *Pediatr Radiol* 2006;36:582-592.
31. Barkovich AJ, Miller SP, Bartha A, et al. MR imaging, MR spectroscopy, and diffusion tensor imaging of sequential studies in neonates with encephalopathy. *AJNR* 2006;27:533-547.
32. Mader I, Schoning M, Klose U, Kuker W. Neonatal cerebral infarction diagnosed by diffusion-weighted MRI: pseudonormalization occurs early. *Stroke* 2002;33:1142-1145.