



Universiteit
Leiden

The Netherlands

Magnetic resonance imaging in neonatal hypoxic-ischemic brain injury

Liauw, L.

Citation

Liauw, L. (2009, March 19). *Magnetic resonance imaging in neonatal hypoxic-ischemic brain injury*. Retrieved from <https://hdl.handle.net/1887/13690>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/13690>

Note: To cite this publication please use the final published version (if applicable).

Curriculum vitae

The author of this thesis was born August 10, 1964 in Bandung, Indonesia. As a young infant she moved with her parents to the Netherlands, growing up in Voorburg, a suburb of The Hague. After finishing High School (Gymnasium beta) at the St. Maartenscollege in Voorburg (1976-1982), she started Medical School at the Leiden University Medical Centre (LUMC). The last part of her medical studies were spent in research at the Otobiological Research Laboratories, Ohio State University, Columbus, Ohio, USA (Dr. D.J. Lim) in association with the department of Otorhinolaryngology of the LUMC (Prof. Dr. J.J. Grote). In 1990 she graduated from Medical School with honors. After working as an intern in general surgery (Groene Hart Ziekenhuis Gouda) and neuroradiology (LUMC), she started her Radiology residency at the LUMC in 1992 (Prof. Dr. A.E. van Voorthuisen and Prof. Dr. J.L. Bloem). Following her fellowship in Neuroradiology and Head & Neck Radiology (1997-1998), she worked as a junior radiologist at the department of Radiology at the LUMC, in the areas of Neuroradiology and Head & Neck radiology, ultrasonography, mammography, and pediatric radiology. From December 2005, she worked as a radiologist at the department of Radiology, Radboud University Nijmegen Medical Centre (Prof. Dr. G.J. Blickman). She was secretary of the Head & Neck section of the Radiological Society of the Netherlands (2002-2008). Since October 2008 she holds position of senior consultant radiologist in Singapore. Work on this thesis (prof. Dr. M.A. van Buchem) commenced in 2000, after a false start in 1996, with the inspiring collaboration of Dr. G. van Wezel-Meijler (department of Neonatology, LUMC) and Dr. J. van der Grond (department of Radiology, LUMC).

List of publications

Liauw L, Lim DJ, Bakaletz LO, van Blitterswijk C. Effect of Vitamin A on the growth and differentiation of rat external auditory canal epithelium in organ culture. *Am J Otolaryngol* 1991; 12 (2): 67- 75.

Pieters MSM, Jenekens-Schinkel A, Stijnen Th, Edelbroek PM, Brouwer OF, **Liauw L**, Heyer A, Lanser JBK, Peters ACB. Carbamazepine (CBZ) controlled release compared with conventional CBZ: a controlled study of attention and vigilance in children with epilepsy. *Epilepsia* 1992; 33: 1137-1144.

Wintzen AR, Badrising UA, Roos RAC, Vielvoye J, **Liauw L**, Pauwels EKJ. Dysphagia in ambulant patients with Parkinson's disease: common, not dangerous. *Can J Neurol Sci* 1994; 21: 53-56.

Wintzen AR, Badrising UA, Roos RAC, Vielvoye J, **Liauw L**. Influence of bolus volume on hyoid movements in normal individuals and patients with Parkinson's disease. *Can J Neurol Sci* 1994; 21; 57-59.

Liauw L, Vielvoye GJ, de Keizer RJW, van Duinen SG. Optic nerve glioma mimicking an optic nerve meningioma. *Clin Neur and Neurosurgery* 1996; 98: 258-261.

Kluin-Nelemans JC, **Liauw L**, van Amsterdam J, van Eck-Smit BLF, Pauwels EKJ. Nuevas modalidades para el estadiaje y seguimiento de los pacientes con enfermedad de Hodgkin (New modalities for staging and follow-up of patients with Hodgkin's disease). *Revisiones en Cancer* 1996; 10: 45-51.

Liauw L, van Buchem MA, Feuth JDM, van Vugt AB, Bode PJ. Cerebral fat embolism. *Eur Radiol* 1998; 8: 1059.

Liauw L, van Buchem MA, Spilt A, de Bruine FT, van den Berg R, Hermans J, Wasser MNJM. MR angiography of the intracranial venous system. *Radiology* 2000; 214: 678-682.

Jeurissen SER, **Liauw L**, Willems LNA, Huizinga TWJ. Jaw claudication in Takayasu's disease. *Ann Rheum Dis* 2003; 62 (10): 922.

D.H.van der Dussen, B.R.Bloem, **L.Liauw**, M.D.Ferrari. Ophthalmoplegic migraine: migrainous or inflammatory? *Cephalalgia* 2004; 24 (4); 312-5.

Leijser LM, Klein RH, Veen S, **Liauw L**, van Wezel-Meijler G. Hyperechogenicity of the thalamus and basal ganglia in very preterm infants: radiological findings and short-term neurological outcome. *Neuropediatrics* 2004; 35(5): 283-9.

Van den Berg R, Schepers A, de Bruine FT, **Liauw L**, Mertens BJA, van der Mey AGL, van Buchem MA. The value of MR angiography techniques in the detection of head and neck paragangliomas. *EJR* 2004; 52(3): 240-245.

Scherjon SA, **Liauw L**, Kant SG. MRI supported diagnosis and counselling in a family with a probably autosomal recessive form of pachygyria. *Prenatal Diagnosis* 2005; 25:1097-1101.

Liauw L, Meinders-Palm IH, van der Grond J, Leijser LM, le Cessie S, Laan AEM, Heeres BC, van Buchem MA, van Wezel-Meijler G. Differentiating normal myelination from hypoxic-ischemic encephalopathy on T1-weighted MR images; a new approach. *AJNR* 2007;28 (4):660-665.

Leijser LM, Vein AA, **Liauw L**, Strauss T, Veen S, van Wezel-Meijler G. Prediction of short-term neurological outcome in full-term neonates with hypoxic-ischaemic encephalopathy based on combined use of electroencephalogram and neuroimaging. *Neuropediatrics* 2007; 38 (5): 219-227.

Liauw L, van der Grond J, Slooff V, Wiggers-de Bruine FT, Laan LAEM, le Cessie S, van Buchem MA, van Wezel-Meijler G. Differentiation between peritrigonal terminal zones and hypoxic-ischemic white matter injury on MRI. *European Journal of Radiology* 2008; 65 (3):395-401.

Liauw L, van der Grond J, van den Berg-Huysmans AA, Palm-Meinders IH, van Buchem MA, van Wezel-Meijler G. Hypoxic-ischemic encephalopathy: diagnostic value of conventional MR imaging pulse sequences in term-born neonates. *Radiology* 2008; 247 (1): 204-212.

Leijser LM, **Liauw L**, Veen S, de Boer IP, Walther FJ, van Wezel-Meijler G. Comparing brain white matter on sequential cranial ultrasound and early MRI in very preterm infants. *Neuroradiology* 2008; 50 (9): 799-811.

Liauw L, van der Grond J, van den Berg-Huysmans AA, Laan LAEM, van Buchem MA, van Wezel-Meijler G. Is there a way to predict outcome in (near) term neonates with hypoxic-ischemic encephalopathy based on MR imaging? *AJNR* 2008; 29 (9): 1789-1794.

Liauw L, van Wezel-Meijler G, Veen S, van Buchem MA, van der Grond J. Do ADC measurements predict outcome in children with hypoxic-ischemic encephalopathy? *AJNR*, in press.

Pennings RJE, **Liauw L**, Cremers CWRJ. A spontaneous otogenic extradural pneumocephalus. *Otology & Neurotology*, in press.

