



Universiteit  
Leiden  
The Netherlands

## Child and parental adaptation to pediatric oncology

Vrijmoet-Wiersma, J.

### Citation

Vrijmoet-Wiersma, J. (2010, January 14). *Child and parental adaptation to pediatric oncology*. Retrieved from <https://hdl.handle.net/1887/14561>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14561>

**Note:** To cite this publication please use the final published version (if applicable).



# Psychometric qualities of the Dutch version of the Pediatric Inventory for Parents (PIP): a multi-center study

C.M.J. Vrijmoet-Wiersma<sup>1</sup>, J.E.H.M. Hoekstra-Weebers<sup>2</sup>, W.M.G. de Peinder<sup>3</sup>, H.M. Koopman<sup>4</sup>, W.J.E. Tissing<sup>5</sup>, Ph.D.A. Treffers<sup>6</sup>, M.B. Bierings<sup>7</sup>, N.C.A. Jansen<sup>7</sup>, M.A. Grootenhuis<sup>8</sup>, R. M. Egeler<sup>1</sup>

<sup>1</sup>Leiden University Medical Center, The Netherlands, <sup>2</sup>Wenckebach Institute, University Medical Center Groningen, University of Groningen

The Netherlands, <sup>3</sup>Health psychology, Leiden University, The Netherlands

<sup>4</sup>Medical Psychology, Leiden University Medical Center, The Netherlands

<sup>5</sup>University Medical Center Groningen, University of Groningen,

The Netherlands, <sup>6</sup>Academic Center for Child and Adolescent Psychiatry

Curium; Leiden University Medical Center, The Netherlands

<sup>7</sup>Wilhelmina Children's Hospital, University Medical Center Utrecht,

The Netherlands, <sup>8</sup>Academic Medical Center, Amsterdam, The Netherlands

**Psychooncology 2009, Apr. (8) (Epub)**

## Abstract

*Goals of work.* Diagnosis and treatment of childhood cancer are continuous stressors in the lives of the entire family involved. Disease-related tools for the assessment of parental stress and adaptation are scarce. For that reason, the Pediatric Inventory for Parents (PIP), a disease-related measure, was translated into Dutch and its psychometric qualities were determined to prove its value. *Methods.* The PIP and three other measures (State-Trait Anxiety Inventory, General Health Questionnaire and Parenting Stress Index, Short Form) were administered to 174 parents of 107 children diagnosed with cancer in three university medical centers in the Netherlands. *Results.* Internal consistency (Cronbach's  $\alpha = .94$  and  $.95$ ) and test-retest reliability (Pearson's  $r$  between  $.67$  and  $.87$ ) of the Dutch PIP Total Scales are satisfactory. Validity was illustrated by a high correlation between PIP scores and anxiety and general stress. Confirmatory factor analysis showed acceptable fit to the data for the original four-factor and the one-factor models; the four-factor model showed slightly better fit. *Conclusion.* The PIP can be used in clinical practice to assess disease-related parental stress. Further psychometric testing is highly recommended.

## Introduction

When parents are confronted with a cancer diagnosis in their child, they often report to feel as if their world has fallen apart. Although most parents show remarkable resilience over time, for a subgroup of parents, levels of psychological distress remain high throughout the entire treatment period and thereafter [14-16,34,41]. Heightened levels of depression [1,44], anxiety [5,6,30,33], stress [21,30], a decreased quality of life [12], marital distress [10,13] and post-traumatic stress symptoms (e.g. [18,25]) have been reported in parents of pediatric cancer patients. This effect has been found to persist in a substantial proportion of the parents, five to even ten years or longer [21,42]. Distress levels are highest around and shortly after diagnosis [41]. Parents experiencing most emotional problems at diagnosis and during treatment continue to report high levels of distress, even after treatment ends [21,36]. Mothers tend to report more stress than fathers [42,44] and younger parents and parents of younger children report more stress than parents of older children [35].

To obtain a better understanding of parental stress related to pediatric cancer, multidimensional assessment specific to the circumstances of parents of these children is needed [22,36]. In most studies, generic measures of psychological maladjustment have been used to assess parental distress. Significant differences with the reference group were found in some [27], but not in all studies [23]. Some authors argue that these traditional instruments are not sensitive enough to assess emotional and behavioral changes related to medical conditions [3]. However, when no differences are found, one cannot simply conclude that the measures are insufficient; it could also mean that there are in fact no differences between the groups.

Using disease-related measures in combination with generic measures could provide additional information that can be used to study the impact of psychosocial interventions and that will help to guide psychosocial interventions [22]. Several questionnaires have been developed specifically for parents of children with cancer or children undergoing stem cell transplantation [7,11,19,28,32]. The Pediatric Inventory for Parents (PIP) [35] was designed to examine areas of stress and concern in parents of children with a medical illness. It has been proven a reliable instrument for examining parent's report of stress related to caring for a child with a serious illness, such as cancer [35,36], diabetes [37] and sickle cell disease [24]. One of the assets of the PIP is that parents are asked to rate both the frequency of stressful illness-related events and the difficulty they experience with these events. PIP total scores correlated significantly with a generic measure of state anxiety and parenting stress within a childhood oncology population [35].

The availability of assessment instruments in more than one language is low. This is unfortunate, because it would facilitate international multi-center studies and allow for cross-cultural comparisons. However, the translation in another language and culture is a lengthy and laborious process and is not always carried out adequately and/or documented properly in research articles.

The present study evaluates the psychometric qualities of the Dutch PIP, specifically item distribution, test-retest reliability, construct validity (by calculating correlations between the PIP and three other measures) and discriminative validity (i.e. the ability of the PIP to distinguish between known groups). Using confirmatory factor analyses, we evaluated the original four-factor model of the PIP by examining the goodness-of-fit to the data. As the total scales are regularly used and previous studies [31] found substantial correlations (ranging from 0.45 – 0.83) between the four subscales, we also evaluated the comparative fit of the four-factor model versus the one-factor model.

## Method

### Participants

All parents of children -aged 0 through 18 years- diagnosed with a malignancy 1-18 months ago between January 2005 - February 2007 in three medical centers (Leiden University Medical Centre, Wilhelmina Children's Hospital Utrecht and University Medical Centre Groningen) were asked by letter to participate in the study. We chose to include parents of children diagnosed between 1 and 18 months ago in order to obtain a sample that would be more or less homogeneous with regards to 'time since diagnosis'. Parents of deceased children were excluded from the study. In total 268 parents were approached (78 in Leiden, 60 in Utrecht and 130 in Groningen).

### Procedure

The study was approved by the Medical Ethics Committee of all three medical centers. The PIP was translated by a team of five persons. An English 'native speaker' was asked to make a back-translation, according to the procedure described by Van Widenfelt et al. [39]. The author of the PIP provided feedback on the back-translated questionnaire. The translated version was piloted with three couples and adaptations were made if necessary.

Eligible parents received information about the study and an informed consent form. In Leiden, part of the parents (40%) received the questionnaires by mail and part (60%) filled in the questionnaires in the clinic. No differences between the methods were found. Parents were instructed to fill in the questionnaires separately and not to consult each other. In Utrecht and Groningen, questionnaires were mailed to the parents' homes

along with the study information and consent forms.

One week after filling in the PIP and other questionnaires, a random half of the parents (every even numbered returned booklet) received the PIP again, to obtain test-retest reliability data. Parents who did not want to participate in the study were asked to supply demographic and illness-related data.

### Measures

*Parental disease-related stress* was measured using the *Pediatric Inventory for Parents (PIP)*. The PIP is a 42-item self-report questionnaire that measures parental stress related to the serious illness of the child with respect to (a) communication with the child and the medical team (9 items), (b) emotional distress (15 items), (c) medical care (8 items) and (d) role function (10 items). Each of the 42 items is rated on two 5-point Likert-type scales. Parents need to respond to the items twice: the first time to assess the frequency of each stressor; the second time to assess how difficult the issue has been for the parent. Parents are asked to consider last week when responding to each item. Examples of PIP-items: 'Learning upsetting news' or 'Speaking with the doctor'. Higher scores refer to more stress. Adequate internal consistency ( $\alpha = .80-.96$ ) and construct validity of the original version of the PIP (scale scores range 42-210) have been reported [35].

*Parental anxiety* was measured using the *State Trait Anxiety Index, state and trait version (STAI)*, a 40-item questionnaire that measures the respondent's transitory emotional condition of stress and the general inclination towards anxiety. Dutch reference data and information on reliability (Cronbach's  $\alpha$ . 95 and .94) and validity are available [2]. In our study, the alphas were .95 (state anxiety) and .94 (trait anxiety).

*Parental psychological distress* was assessed using the *General Health Questionnaire (GHQ)*, a 12-item version self-report measure of non psychotic psychiatric disorders that can be used as a general measure for psychological distress. The psychometric properties of the Dutch version of the scale are reported to be good [20] and the questionnaire has been used frequently in research and patient care [40,43]. In our study, the alpha was .87.

*Parental stress* associated with raising children (i.e. parenting stress) was assessed using the *Parental Stress Index, short form (PSI-SF), Dutch version* [8]. The PSI-SF is derived from the full 123-item PSI. The PSI-SF is a reliable and valid measure and contains 25 items that are scored on a five-point continuum from strongly agree to strongly disagree. The PSI-SF differentiates well between clinical and non-clinical groups and has been used in various studies [38]. In our study, the alpha was .93.

*Demographic and clinical information:* Gender, age, marital status, educational level of the parent, gender and age of the target child, the child's medical diagnosis, current treatment status and the number of weeks since diagnosis were recorded.

### Statistical analyses

To assess differences between responders and non-responders, we used independent T-tests and chi square tests for categorical variables. We evaluated the normal distribution of the PIP with the test of normality and we calculated skewness and kurtosis. Confirmatory factor analysis (CFA) with weighted least squares means and variance adjusted estimation (WLSMV), applied to the polychoric correlation matrix, was used to evaluate the fit to the data of the original and the modified four- and one-factor models. WLSMV has been shown to perform well with ordinal variables and rather small samples [4]. The Mplus program version 2.02 [29] was used for factor analyses.

The fit of the models was assessed using practical fit indices, the values of which were evaluated according to the guidelines formulated by Marsh, Hau, and Wen [26]. The indices included the normed comparative fit index (CFI), the Tucker-Lewis index (TLI), the root mean square error of approximation (RMSEA) and the standardized root mean square residual (SRMR). RMSEA and SRMR values  $>0.10$  are regarded as indication of bad fit, as were CFI and TLI values  $< 0.85$ . We used the Cheung and Rensvold [9] CFI-criteria to test the difference between models: changes in CFI ( $\Delta$ CFI) of  $-.01$  or less indicate that the hypothesis of equal fit should not be rejected, when  $\Delta$ CFI lie between  $-.01$  and  $-.02$ , differences may exist and definite differences between models exist when  $\Delta$ CFI is greater than  $-.02$ .

To compute test-retest reliability we used Pearson's  $r$ . To assess internal consistency of the PIP total and the four domain scales Cronbach's coefficient  $\alpha$  was calculated. Alpha values of  $.7$  and above were considered adequate. Construct validity was examined by conducting correlation analyses between the PIP and the other psychological measures. The intercorrelations of the PIP subscales were calculated by use of Pearson correlations. To test the discriminative validity, independent T-tests were used to examine the effect of demographic and illness-related variables on stress reported on the PIP.

## Results

### Participants

The overall response rate was 66% (72% in Leiden, 65% in Utrecht and 61% in Groningen). Six parents were excluded because of missing data. In total, 174 parents of 107 children participated. Of 15 single-parent families and 25 families, only one parent participated.

Reasons for refusal were the experience of too many stressful events (26%), lack of time (18%), the illness and treatment of the child was considered to have been completed too long ago (16%), too busy with work (15%), language problems (13%) or it

was too confronting to the parents (10%). Non-participating parents did not differ from participating parents with regard to age, marital status, educational level, sex and age of the child. However, we did find significant differences with respect to parent gender, parent ethnicity and treatment status. In the non-participating group, the percentage of fathers, non-Dutch parents and parents with a child off treatment was higher. See Table 1.

Results on the outcome measures and demographic characteristics did not significantly differ between the three medical centers, hence we analyzed all data together.

### Item Distribution and interscale correlation

First, the test of normality was performed on the different subscales and total scales of the PIP. Both total scales showed normality, the subscales significantly deviated from normality, except for two scales. Kurtosis was found for Communication Frequency (3.25,  $p < 0.001$ ) and Emotional distress Frequency (2.76,  $p < 0.01$ ). Skewness was found for the scales Communication Frequency, Emotional distress Difficulty and Role function Frequency. Interscale correlations of the PIP-sub scales varied from  $.50$  (Medical care Difficulty with Communication Frequency) to  $.82$  (Communication Frequency with Emotional distress Difficulty).

### Reliability

Cronbach's alphas for the Total scales (PIP-Frequency =  $.94$ , PIP-Difficulty =  $.95$ ) and for the subscales Medical Care, Emotional Distress and Role Function ( $\geq .80$ ) were adequate. The alpha-value for the Communication scale was low for the PIP-Frequency and PIP-Difficulty scales (see table 2a). After deletion of item number 2 *Arguing with family member(s)*, the alpha was acceptable ( $.65$ ).

When analyzing mothers and fathers separately, we found similar reliability scores (see table 2b). Mean scores between mothers and fathers differed significantly (mothers scoring higher,  $p < .05$ ) on the Frequency scales Communication, Medical care and Emotional distress and the Total Frequency score. The Difficulty scale Emotional distress and Total Difficulty scores were also significantly higher for mothers than for fathers.

Test-retest reliability after 14 days was adequate ( $.68 \leq r \leq .87$ ), based on returned questionnaires from 78 parents (33 fathers, 45 mothers) of the 111 parents approached to fill in the PIP twice (70% response rate).

**Table 1.** Demographic characteristics of parents and children

	Responders (N=107)			Non-responders (N=39)		
	No. of children: (N=174)			No. of children: (N=94)		
Parent characteristics	M	Range	SD	M	Range	SD
Age	41.0	22-65	7.0	40.0	28-60	6.8
	N	%		N	%	
Sex						
Male	74	42.5		52	55.3*	
Female	100	57.5		42	44.7	
Educational level						
Lower	41	23.6		33	35.1	
middle	76	43.7		30	31.9	
higher	57	32.8		31	33.0	
Ethnicity						
Dutch	161	92.5		73	77.7*	
Non-Dutch	13	7.5		21	22.3	
Marital status						
Married/living with partner	161	92.5		87	92.6	
Divorced/widowed/single	13	7.5		7	7.4	
Hospital						
Leiden	56	32.2		22	23.4	
Utrecht	39	22.4		21	22.3	
Groningen	79	45.4		51	54.3	
<b>Child characteristics</b>	<b>M</b>	<b>Range</b>	<b>SD</b>	<b>M</b>	<b>Range</b>	<b>SD</b>
		(N=174)			(N=94)	
Age at assessment (months)	115.0	8-218	61.8	106.2	17-230	59.3
Time since diagnosis (weeks)	40.5	5-110	25.3	42.4	8-110	22.0
	N	%		N	%	
Sex						
Male	57	53.3		20	51.3	
Female	50	46.7		19	48.7	
Diagnosis						
ALL/ JMML	41	38.3		12	30.8	
AML	8	7.5		2	5.1	
(Non)Hodgkin's lymphoma	19	17.7		6	15.4	
Bone tumors <sup>a</sup>	14	13.1		3	7.7	
Brain tumor	11	10.3		4	10.3	
Neuroblastoma	5	4.7		1	2.6	
Wilm's tumor	4	3.7		3	7.7	
Other	5	4.7		8	20.5	
Treatment status						
On treatment	86	80.4		17	43.6*	
Off treatment	21	19.6		22	56.4	

<sup>a</sup>Ewing sarcoma, osteosarcoma and synovia sarcoma

ALL = acute lymphatic leukaemia, JMML = juvenile myelomonocytic leukaemia, AML = acute myeloid leukaemia

\* Significant difference between responders and non-responders ( $p < 0.05$ )

**Table 2a.** Means, Standard Deviations, and Internal Consistency of the Pediatric Inventory for Parents (N = 174)

	PIP-F			PIP-D		
	M	(SD)	$\alpha$	M	(SD)	$\alpha$
I. Communication	23.2	(4.5)	.60	18.6	(5.4)	.73
II. Medical care	24.0	(7.2)	.85	16.9	(6.2)	.84
III. Em. distress	43.1	(10.1)	.88	44.6	(12.6)	.91
IV. Role function	25.1	(7.1)	.80	21.6	(7.4)	.82
Total	115.4	(26.0)	.94	101.7	(28.5)	.95

PIP F = PIP Frequency, PIP D = PIP Difficulty, Em. = emotional

**Table 2b.** Means, Standard Deviations, and Internal Consistency of the Pediatric Inventory for Parents, mothers (N = 100) and fathers (N=74)

	PIP-F mothers			PIP-D mothers			PIP-F fathers			PIP-D fathers		
	M	(SD)	$\alpha$	M	(SD)	$\alpha$	M	(SD)	$\alpha$	M	(SD)	$\alpha$
I. Communication	24.3*	(4.9)	.52	19.1	(5.1)	.73	22.0	(4.6)	.69	17.9	(5.7)	.82
II. Medical care	25.3*	(7.1)	.83	17.5	(5.9)	.81	22.2	(6.9)	.84	16.2	(6.6)	.83
III. Em. distress	44.2*	(11.3)	.75	46.7*	(12.1)	.90	40.8	(9.5)	.82	41.7	(12.7)	.85
IV. Role function	25.8	(6.9)	.76	22.1	(7.1)	.78	24.1	(7.3)	.88	20.9	(7.7)	.92
Total	120.6*	(26.3)	.95	105.4*	(27.4)	.94	109.3	(25.7)	.95	96.8	(29.3)	.99

PIP F = PIP Frequency, PIP D = PIP Difficulty, Em. = emotional

\* Significant difference between mothers and fathers ( $p < .05$ )

### Confirmatory Factor Analyses

Two identical factor models for the PIP-frequency and the PIP-difficulty items were (comparatively) evaluated: a four- and a one-factor model. The one-factor model, constituted by all 42 items of the PIP, reflects the possibility that one single latent dimension underlies the items. The factors of the four-factor model represent the original four scales; the factors were allowed to correlate. The fit index values are presented in Table III. Results indicated adequate fit of all models on the TLI but a bad fit on the remaining indices for all models except the four-factor model for the Difficulty items.

We evaluated areas of strain in the factor models using the modification indices of ML estimation. Error-correlations  $> 0.100$  between item pairs, indicating not modeled minor factors, were found for 5 item pairs in all models: between item pairs 14-16 (indicating a minor factor 'distress over child suffering'), and between the item-pairs 24-29, 24-39, 26-29, 29-36 (indicating a minor factor 'worrying about the child's future'). After adding the 5 error-correlations to the models, good fit on TLI and (nearly) acceptable fit on the remaining indices was found for all models (see Table 3).

The differences in fit to the data between the adjusted four- and one-factor models were negligible, as indicated by  $\Delta CFI < 0.10$ . Therefore, the one-factor models, representing the Total scales, may be preferred.

**Table 3.** Goodness of Fit Indices for PIP Four-Factor and One-Factor Models\*

Models	$\chi^2$	df	CFI	TLI	RMSEA	SRMR
<i>Frequency</i>						
Four-Factor Model	270.04	88	0.869	0.933	0.109	0.091
One-Factor Model	292.91	88	0.852	0.924	0.105	0.091
Four-Factor Model with $\theta_{14,16}$ , $\theta_{24,29}$ , $\theta_{24,36}$ , $\theta_{26,29}$ , $\theta_{29,36}$ free	239.35	88	0.891	0.944	0.099	0.084
One-Factor Model with $\theta_{14,16}$ , $\theta_{24,29}$ , $\theta_{24,36}$ , $\theta_{26,29}$ , $\theta_{29,36}$ free	249.95	88	0.883	0.940	0.103	0.086
<i>Difficulty</i>						
Four-Factor Model	235.73	88	0.895	0.956	0.098	0.081
One-Factor Model	253.06	87	0.882	0.950	0.105	0.085
Four-Factor Model with $\theta_{14,16}$ , $\theta_{24,29}$ , $\theta_{24,36}$ , $\theta_{26,29}$ , $\theta_{29,36}$ free	204.44	87	0.909	0.962	0.091	0.079
One-Factor Model with $\theta_{14,16}$ , $\theta_{24,29}$ , $\theta_{24,36}$ , $\theta_{26,29}$ , $\theta_{29,36}$ free	227.85	87	0.900	0.957	0.096	0.082

\*: weighted least squares means and variance adjusted estimation was applied to the polychoric correlation matrix; CFI = Comparative Fit Index, TLI = Tucker-Lewis index, RMSEA = Root Mean Square Error of Approximation, SRMR = Standardized Root Mean Square Residual.

### Construct Validity

Correlations between the PIP total scales and the other instruments were calculated. For all parents, the PIP-Frequency scale was strongly associated with STAI state and trait ( $r = .52$  and  $r = .55$ ,  $p < .01$ ) and with the GHQ ( $r = .54$ ,  $p < .01$ ) and weakly with the PSI-SF ( $r = .19$ ,  $p < .05$ ). The PIP-Difficulty scale was strongly related to STAI state and trait and GHQ ( $r = .59$ ,  $r = .66$  and  $r = .51$  respectively, all  $p < .01$ ) and weakly with the PSI-SF ( $r = .24$ ,  $p < .01$ ). When analyzing fathers and mothers separately, the correlations were comparable.

### Discriminative validity

Mothers reported higher scores than fathers (PIP-Frequency,  $p < .01$ ,  $t = 2.84$ ). Older fathers (i.e. fathers above the mean age of 41 years at assessment) reported significantly more distress than younger fathers (PIP-Difficulty,  $p < .05$ ,  $t = 2.19$ ). Interestingly, older fathers also reported significantly higher state anxiety levels. For mothers, no age effect was found. Parents of younger children (under versus over 115 months) reported higher stress scores than parents of older children (PIP-Difficulty,  $p < .05$ ,  $t = 2.11$ ).

Parents of children on treatment had significantly higher PIP scores than parents whose children had completed treatment (PIP-Frequency,  $p < .05$ ,  $t = 2.92$ ) and parents of children diagnosed more recently –less than 40 weeks ago– reported more stress (PIP-Frequency,  $p < .05$ ,  $t = 2.49$ ) than parents of children who were diagnosed longer ago.

### Item-subscale and item-total correlations

Item-subscale correlations varied from  $-.07$  to  $.67$  in the Frequency Scale, with a mean of  $.46$ . For the Difficulty items, item-subscale correlations varied from  $.23$  to  $.79$  with a mean of  $.53$ . Lowest item-subscale correlations were found for the scale Communication. Item-total correlations varied from  $.02$  to  $.70$  for the Frequency items (mean  $.47$ ) and from  $.32$  to  $.77$  for the Difficulty items (mean  $.55$ ).

## Discussion

Gaining insight into parents' stress following pediatric cancer is increasingly important in order to deliver adequate psychosocial care to the entire family. Disease-related measures can add important information about parental adaptation to stressful illness-related situations. Results regarding the Dutch version of the Pediatric Inventory for Parents (PIP), a disease-related measure of parental stress, are satisfactory. We found adequate (test-retest) reliability scores for the PIP Total scales and three of the four subscales (Medical care, Emotional distress and Role function). The fourth subscale, Communication, needs improvement. This last finding is not in line with the results from the original study [35]. Cultural differences with regards to communicating with hospital staff and family could perhaps explain part of this difference in results.

PIP scores correlated strongly with a generic measure of anxiety and general psychological functioning. This means that disease-related distress, although it measures a different construct, can have considerable overlap with general well-being and anxiety. The added value of the PIP however is that it asks parents about their stress concerning disease-related situations. Scores on the PIP could be transformed into an individual 'stress profile', which could be used to tailor psychosocial support.

The low correlation of PIP scores with parenting stress scores suggests that stress resulting from difficulties disciplining and setting limits to one's child (parenting stress) differs from stress associated with having a child with a serious illness (parental stress). In various studies, e.g. [17], the PSI is used as a measure of parental stress instead of stress associated with parenting. This strategy might well result in drawing the wrong conclusions about the stress reactions parents can have as a result of their child's illness.

As expected and in line with other research [33,44], mothers showed higher PIP-scores than fathers, parents closer to diagnosis and parents of younger children reported more stress. Older fathers reported higher stress levels. For mothers, no age effect was found. This result is contrary to the findings of the original PIP-study, in which younger parents (both mothers and fathers) reported more stress [35].

Confirmatory factor analyses showed that not only the four factor model, representing the four subscales of the PIP, but also the one-factor model, representing the Total scales, showed acceptable fit to the data after three items were dropped from the models and the error-correlation of one item-pair was added. The Total scales may be regarded as sufficient for practical purposes as the difference in fit between the two models were minor, and very strong correlations were found between the four latent factors.

One of the advantages of the PIP is subdivision of parental stress levels into Frequency and Difficulty scores (even though the scales correlate highly and thus outcomes might be partly overlapping), which enables the psychosocial team to target interventions more precisely to the needs of parents in different phases of their child's treatment. In our study, PIP-Frequency scores discriminated between parents of children on treatment and parents whose children have ended treatment. However, PIP-Difficulty scores for the two groups were equal. This finding may imply that although the frequency of stressful disease-related events is lower in parents of children off treatment, the perceived difficulty of these events remains similar.

### Limitations and practical implications

Despite the results of the present study, there are some limitations that need mentioning. Firstly, the age range of the children in our study group varied widely, making comparison of parental stress levels difficult. Being the parent of an ill baby or toddler versus a teenager will render different sources of stress. Secondly, a substantial proportion of parents refused participation. Reasons for non-participation ranged from being too stressed to considering the treatment of the child to have been completed too long ago or being too busy with other things like work. It is unclear if this caused an under- or overestimation of parental stress levels.

Lastly, the procedure of administering the questionnaires was different in the three hospitals. Approaching parents face to face in the clinic or the outpatient's ward yielded a higher response rate than mailing the questionnaires. However, this did not seem to influence reported levels of stress.

One of the assets of the study is the relatively large, multi center study group. Furthermore, we managed to include a large percentage of fathers in our study. The PIP could be used in regular patient care to assess all parents of newly diagnosed children at

critical time points in therapy: shortly after diagnosis, then again after 4-6 months (usually seen as the 'stabilization phase') and by the end of treatment. These time points seem to cover the process of parental stress through the phases of childhood cancer well.

In summary, the Dutch PIP is a reliable and valid assessment tool to gain insight in stress experienced by parents during the course of their child's cancer treatment. Continuous psychometric testing is recommended in different populations and at different time points.

### Acknowledgements

We would like to thank the parents for taking the time to complete the questionnaires. Secondly, we express our thanks to the members of the translation team (Natasja van Lang, Bas Berger and Brigit van Widenfelt) and to Arnold Goedhart for his help with the analyses. Lastly, we thank Randi Streisand for her consent to translate the PIP in Dutch and for her comments on the manuscript.



## References

1. Allen R, Newman SP, Souhami RL (1997) Anxiety and depression in adolescent cancer: findings in patients and parents at the time of diagnosis. *Eur J Cancer* 33(8):1250-1255.
2. Bakker FC, van Wieringen PCW, van der Ploeg HM, Spielberger CD. Manual for the Dutch version of the state-trait anxiety inventory for children (STAIC). Lisse: Swets & Zeitlinger; 1989.
3. Barrera M, D'Agostino NM, Gibson J, Gilbert T, Weksberg R, Malkin D (2004) Predictors and mediators of psychological adjustment in mothers of children newly diagnosed with cancer. *Psychooncology* 13(9):630-641.
4. Beauducel A, Herzberg PI (2006) On the performance of maximum likelihood versus means and variance adjusted weighted least squares estimation in CFA. *Structural Equation Modeling* 13(2):186-203.
5. Best M, Streisand R, Catania L, Kazak AE (2001) Parental distress during pediatric leukemia and posttraumatic stress symptoms (PTSS) after treatment ends. *J Pediatr Psychol* 26(5):299-307.
6. Boman K, Lindahl A, Bjork O (2003) Disease-related distress in parents of children with cancer at various stages after the time of diagnosis. *Acta Oncol* 42(2):137-146.
7. Bonner MJ, Hardy KK, Guill AB, McLaughlin C, Schweitzer H, Carter K (2006) Development and validation of the parent experience of child illness. *J Pediatr Psychol* 31(3):310-321.
8. Brock de AJLL, Vermulst AA, Gerris JRM, Abidin RR. Handleiding Nijmeegse ouderlijke stress index. Lisse: Swets & Zeitlinger; 1992.
9. Cheung GW, Rensvold RB (2002) Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling* 9:233-255.
10. Dahlquist LM, Czyzewski DI, Copeland KG, Jones CL, Taub E, Vaughan JK (1993) Parents of children newly diagnosed with cancer: anxiety, coping, and marital distress. *J Pediatr Psychol* 18(3):365-376.

11. Derogatis LR (1986) The psychosocial adjustment to illness scale (PAIS). *J Psychosom Res* 30(1):77-91.
12. Eiser C, Eiser JR, Stride C (2005) Quality of life in children newly diagnosed with cancer and their mothers. *Health and Quality of Life Outcomes* 3(1):29.
13. Fife B, Norton J, Groom G (1987) The Family's adaptation to childhood leukemia. *Soc Sci Med* 24(2):159-168.
14. Grootenhuis MA, Last BF (1997) Adjustment and coping by parents of children with cancer: a review of the literature. *Support Care Cancer* 5(6):466-484.
15. Hoekstra-Weebers JE, Jaspers JP, Kamps WA, Klip EC (1999) Risk factors for psychological maladjustment of parents of children with cancer. *J Am Acad Child Adolesc Psychiatry* 38(12):1526-1535.
16. Hoekstra-Weebers JE, Jaspers JP, Kamps WA, Klip EC (2001) Psychological adaptation and social support of parents of pediatric cancer patients: a prospective longitudinal study. *J Pediatr Psychol* 26(4):225-235.
17. Hung JW, Wu YH, Yeh CH (2004) Comparing stress levels of parents of children with cancer and parents of children with physical disabilities. *Psychooncology* 13(12):898-903.
18. Kazak AE, Alderfer M, Rourke MT, Simms S, Streisand R, Grossman JR (2004) Posttraumatic stress disorder (PTSD) and posttraumatic stress symptoms (PTSS) in families of adolescent childhood cancer survivors. *J Pediatr Psychol* 29(3):211-219.
19. Kazak AE, Prusak A, McSherry M, Simms S, Beele D, Rourke MT et al. (2001) The psychosocial assessment tool (PAT): pilot data on a brief screening instrument for identifying high-risk families in pediatric oncology. *Families, Systems and Health* 19(3):303-317.
20. Koeter MWJ, Ormel J. General Health Questionnaire, Dutch manual. Lisse: Swets & Zeitlinger; 1991.

21. Kupst MJ, Natta MB, Richardson CC, Schulman JL, Lavigne JV, Das L (1995) Family coping with pediatric leukemia: ten years after treatment. *J Pediatr Psychol* 20(5):601-617.
22. La Greca AM, Lemanek KL (1996) Assessment as a process in pediatric psychology. *J Pediatr Psychol* 21(2):137-151.
23. Lahteenmaki PM, Sjoblom J, Korhonen T, Salmi TT (2004) The life situation of parents over the first year after their child's cancer diagnosis. *Acta Paediatr* 93(12):1654-1660.
24. Logan DE, Radcliffe J, Smith-Whitley K (2002) Parent factors and adolescent sickle cell disease: associations with patterns of health service use. *J Pediatr Psychol* 27(5):475-484.
25. Manne S, DuHamel K, Redd WH (2000) Association of psychological vulnerability factors to post-traumatic stress symptomatology in mothers of pediatric cancer survivors. *Psychooncology* 9(5):372-384.
26. Marsh HW, Hau KT, Wen Z (2004) In search of golden rules: Comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's (1999) findings. *Structural Equation Modeling* 11:320-341.
27. Maurice-Stam H, Oort FJ, Last BF, Grootenhuis MA (2008) Emotional functioning of parents of children with cancer: the first five years of continuous remission after the end of treatment. *Psychooncology* 17:448-459.
28. McCubbin MA, Svavarsdottir EK. *The Care of My Child with Cancer Questionnaire*. Reykjavik, Iceland: University of Iceland, Faculty of Nursing; 1999.
29. Muthén LK, Muthén BO. *Mplus User's Guide*. Los Angeles, CA: Muthén & Muthén; 2005.
30. Norberg AL, Lindblad F, Boman KK (2005) Parental traumatic stress during and after paediatric cancer treatment. *Acta Oncol* 44(4):382-388.

31. Pai ALH, Patino-Fernandez AM, McSherry M, Steele D, Alderfer MA, Reilly AT et al. (2008) The psychosocial assessment tool (PAT2.0): Psychometric properties of a screener for psychosocial distress in families of children newly diagnosed with cancer. *J Pediatr Psychol* 33(1):50-62.
32. Phipps S, Dunavant M, Lensing S, Rai SN (2005) Psychosocial predictors of distress in parents of children undergoing stem cell or bone marrow transplantation. *J Pediatr Psychol* 30(2):139-153.
33. Santacrose S (2002) Uncertainty, anxiety, and symptoms of posttraumatic stress in parents of children recently diagnosed with cancer. *J Pediatr Oncol Nurs* 19(3):104-111.
34. Steele RG, Long A, Reddy KA, Luhr M, Phipps S (2003) Changes in maternal distress and child-rearing strategies across treatment for pediatric cancer. *J Pediatr Psychol* 28(7):447-452.
35. Streisand R, Braniecki S, Tercyak KP, Kazak AE (2001) Childhood illness-related parenting stress: the pediatric inventory for parents. *J Pediatr Psychol* 26(3):155-162.
36. Streisand R, Kazak AE, Tercyak KP (2003) Pediatric-specific parenting stress and family functioning in parents of children treated for cancer. *Children's Health Care* 32:245-256.
37. Streisand R, Swift E, Wickmark T, Chen R, Holmes CS (2005) Pediatric parenting stress among parents of children with type 1 diabetes: the role of self-efficacy, responsibility, and fear. *J Pediatr Psychol* 30(6):513-521.
38. Thomas KA, Renaud MT, Depaul D (2004) Use of the parenting stress index in mothers of preterm infants. *Adv Neonatal Care* 4(1):33-41.
39. van Widenfelt BM, Treffers PhDA, de Beurs E, Siebelink BM, Koudijs E (2005) Translation and cross-cultural adaptation of assessment instruments used in psychological research with children and families. *Clinical Child and Family Psychology Review* 8(2):135-147.
40. Vance Y, Morse R, Jenney ME, Eiser C (2001) Issues in measuring quality of life in childhood cancer: measures, proxies, and parental mental health. *J Child Psychol Psychiatry* 42(5):661-667.

41. Vrijmoet-Wiersma CMJ, van Klink JMM, Kolk AM, Koopman HM, Ball LM, Maarten Egeler R (2008) Assessment of parental psychological stress in pediatric cancer: A review. *J Pediatr Psychol* 33(7):694-706.
42. Wijnberg-Williams BJ, Kamps WA, Klip EC, Hoekstra-Weebers JE (2006) Psychological adjustment of parents of pediatric cancer patients revisited: five years later. *Psychooncology* 15(1):1-8.
43. Wijnberg-Williams BJ, Kamps WA, Klip EC, Hoekstra-Weebers JE (2006) Psychological distress and the impact of social support on fathers and mothers of pediatric cancer patients: long-term prospective results. *J Pediatr Psychol* 31(8):785-792.
44. Yeh CH (2002) Gender differences of parental distress in children with cancer. *J Adv Nurs* 38(6):598-606.