



Universiteit
Leiden
The Netherlands

The ubiquitin proteasome system in Huntington disease : impairment of the proteolytic machinery aggravates huntingtin aggregation and toxicity

Pril, R. de

Citation

Pril, R. de. (2011, February 23). *The ubiquitin proteasome system in Huntington disease : impairment of the proteolytic machinery aggravates huntingtin aggregation and toxicity*. Retrieved from <https://hdl.handle.net/1887/16530>

Version: Corrected Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/16530>

Note: To cite this publication please use the final published version (if applicable).

Abbreviations

AD	- Alzheimer disease
ALS	- amyotrophic lateral sclerosis
AMPA	- α -amino-3-hydroxy-5 methyl-4-isoxazolepropionicacid
BDNF	- brain derived neurotrophic factor
CBP	- CREB binding protein
CHIP	- C-terminus of Hsc70 interacting protein
CREB	- cyclic AMP response element binding protein
DRPLA	- dentatorubropallidolysian atrophy
DS	- Down syndrome
DUB	- deubiquitinating enzyme
ERAD	- endoplasmatic reticulum associated degradation
GFAP	- glial fibrillaric acid protein
Hap1	- huntingtin associated protein 1
HD	- Huntington disease
HECT	- homologous to E6-AP, E3-domain type
Hip1	- huntingtin interacting protein 1
Hip1R	- Hip1-related
Hippi	- Hip1 protein interactor
Hsp	- heat shock protein
Htt	- huntingtin protein
LTP	- long-term potentiation
MCBP	- multiubiquitin chain binding proteins
NEDD8	- neural precursor cell-expressed developmentally down-regulated gene
NGFR	- nerve growth factor receptor
NII	- neuronal intranuclear inclusions
PA 28/200	- proteasome activator 28 / 200
Pak-1/2	- p21-activated kinase-1 / 2
PD	- Parkinson disease
PGPH	- peptidyl-glutamyl peptide hydrolizing
PHD	- plant homeo domain, E3-domain type
PQBP-1	- polyglutamine binding protein-1
RING	- really interesting new gene, E3-domain type
SBMA	- spinobulbar muscular atrophy
SCA	- spinocerebellar ataxia
Sp1	- specificity protein-1
SUMO	- small ubiquitin-like modifier
TBP	- TATA-binding protein
UBA	- ubiquitin-associated domain
UBB ⁺¹	- ubiquitin-B ⁺¹
Ubl	- ubiquitin like domain
UCH-L1	- ubiquitin C-terminal hydrolase L1
UHFR-2	- ubiquitin like with PHD and Ring finger domain-2
UPR	- unfolded protein response
UPS	- ubiquitin proteasome system

References

- Adachi, H., Katsuno, M., Minamiyama, M., Sang, C., Pagoulatos, G., Angelidis, C., Kusakabe, M., Yoshiki, A., Kobayashi, Y., Doyu, M. and Sobue, G. (2003) Heat shock protein 70 chaperone overexpression ameliorates phenotypes of the spinal and bulbar muscular atrophy transgenic mouse model by reducing nuclear-localized mutant androgen receptor protein. *J Neurosci* **23**: 2203-2211.
- Alves-Rodrigues, A., Gregori, L. and Figueiredo-Pereira, M.E. (1998) Ubiquitin, cellular inclusions and their role in neurodegeneration. *Trends Neurosci* **21**: 516-520.
- Arrasate, M., Mitra, S., Schweitzer, E.S., Segal, M.R. and Finkbeiner, S. (2004) Inclusion body formation reduces levels of mutant huntingtin and the risk of neuronal death. *Nature* **431**: 805-810.
- Askanas, V. and Engel, W.K. (2003) Proposed pathogenetic cascade of inclusion-body myositis: Importance of amyloid- β , misfolded proteins, predisposing genes, and aging. *Curr Opin Rheumatol* **15**: 737-744.
- Aziz, N.A., Jurgens, C.K., Landwehrmeyer, G.B., EHDN Registry Study Group, van Roos-Mom, W.M.C., van Ommen, G.J.B., Stijnen, T. and Roos, R.A.C. (2009) Normal and mutant HTT interact to affect clinical severity and progression in Huntington disease. *Neurology* **73**: 1280-1285.
- Bailey, C.K., Andriola, I.F.M., Kampinga, H.H. and Merry, D.E. (2002) Molecular chaperones enhance the degradation of expanded polyglutamine repeat androgen receptor in a cellular model of spinal and bulbar muscular atrophy. *Hum Mol Genet* **11**: 515-523.
- Banerjee, M., Datta, M., Majumder, P., Mukhopadhyay, D. and Bhattacharyya, N.P. (2010) Transcription regulation of caspase-1 by R393 of HIPPI and its molecular partner HIP-1. *Nucl Acids Res* **38**: 878-892.
- Bardag-Gorce, F., Riley, N., Nguyen, V., Montgomery, R.O., French, B.A., Li, J., van Leeuwen, F.W., Lungo, W., McPhaul, L.W. and French, S.W. (2003) The mechanism of cytoskeleton aggresome formation: The role of mutant ubiquitin (UBB+1). *Exp Mol Pathol* **74**: 160-167.
- Bence, N.F., Sampat, R.M. and Kopito, R.R. (2001) Impairment of the ubiquitin-proteasome system by protein aggregation. *Science* **292**: 1552-1555.
- Benn, C.L., Sun, T., Sadri-Vakili, G., McFarland, K.N., DiRocco, D.P., Yohrling, G.J., Clark, T.W., Bouzou, B. and Cha, J.H.J. (2008) Huntingtin modulates transcription, occupies gene promoters in vivo, and binds directly to DNA in a polyglutamine-dependent manner. *J Neurosci* **28**: 10720-10733.
- Bennett, E.J., Shaler, T.A., Woodman, B., Ryu, K.Y., Zaitseva, T.S., Becker, C.H., Bates, G.P., Schulman, H. and Kopito, R.R. (2007) Global changes to the ubiquitin system in Huntington's disease. *Nature* **448**: 704-708.
- Bett, J.S., Cook, C., Petrucelli, L. and Bates, G.P. (2009) The ubiquitin-proteasome reporter GFPu does not accumulate in neurons of the R6/2 transgenic mouse model of Huntington's disease. *PLoS ONE* **4**: e5128.

References

- Bett, J.S., Goellner, G.M., Woodman, B., Pratt, G., Rechsteiner, M. and Bates, G.P. (2006) Proteasome impairment does not contribute to pathogenesis in R6/2 Huntington's disease mice: Exclusion of proteasome activator REGy as a therapeutic target. *Hum Mol Genet* **15**: 33-44.
- Bjorkoy, G., Lamark, T., Brech, A., Outzen, H., Perander, M., Overvatn, A., Stenmark, H. and Johansen, T. (2005) P62/SQSTM1 forms protein aggregates degraded by autophagy and has a protective effect on huntingtin-induced cell death. *J Cell Biol* **171**: 603-614.
- Brummelkamp, T.R., Nijman, S.M.B., Dirac, A.M.G. and Bernards, R. (2003) Loss of the cylindromatosis tumour suppressor inhibits apoptosis by activating NF- κ B. *Nature* **424**: 797-801.
- Carrard, G., Bulteau, A.L., Petropoulos, I. and Friguet, B. (2002) Impairment of proteasome structure and function in aging. *Int J Biochem Cell Biol* **34**: 1461-1474.
- Carrell, R.W. and Lomas, D.A. (1997) Conformational disease. *The Lancet* **350**: 134-138.
- Chai, Y., Koppenhafer, S.L., Shoesmith, S.J., Perez, M.K. and Paulson, H.L. (1999) Evidence for proteasome involvement in polyglutamine disease: Localization to nuclear inclusions in SCA3/MJD and suppression of polyglutamine aggregation in vitro. *Hum Mol Genet* **8**: 673-682.
- Chan, E.Y.W., Luthi-Carter, R., Strand, A., Solano, S.M., Hanson, S.A., DeJohn, M.M., Kooperberg, C., Chase, K.O., DiFiglia, M., Young, A.B., Leavitt, B.R., Cha, J.H.J., Aronin, N., Hayden, M.R. and Olson, J.M. (2002) Increased huntingtin protein length reduces the number of polyglutamine-induced gene expression changes in mouse models of Huntington's disease. *Hum Mol Genet* **11**: 1939-1951.
- Chan, H.Y.E., Warrick, J.M., Andriola, I., Merry, D. and Bonini, N.M. (2002) Genetic modulation of polyglutamine toxicity by protein conjugation pathways in Drosophila. *Hum Mol Genet* **11**: 2895-2904.
- Chevalier-Larsen, E.S., O'Brien, C.J., Wang, H., Jenkins, S.C., Holder, L., Lieberman, A.P. and Merry, D.E. (2004) Castration restores function and neurofilament alterations of aged symptomatic males in a transgenic mouse model of spinal and bulbar muscular atrophy. *J Neurosci* **24**: 4778-4786.
- Ciechanover, A. and Brundin, P. (2003) The ubiquitin proteasome system in neurodegenerative diseases. Sometimes the chicken, sometimes the egg. *Neuron* **40**: 427-446.
- Colin, E., Zala, D., Liot, G., Rangone, H., Borrell-Pages, M., Li, X.J., Saudou, F. and Humbert, S. (2008) Huntingtin phosphorylation acts as a molecular switch for anterograde/retrograde transport in neurons. *EMBO J* **27**: 2124-2134.
- Cookson, M.R. (2004) Roles of the proteasome in neurodegenerative disease: Refining the hypothesis. *Ann Neurol* **56**: 315-316.
- Cuervo, A.M. (2004) Autophagy: In sickness and in health. *Trends Cell Biol* **14**: 70-77.
- Cui, L., Jeong, H., Borovecki, F., Parkhurst, C.N., Tanese, N. and Krainc, D. (2006) Transcriptional repression of PGC-1 α by mutant huntingtin leads to mitochondrial dysfunction and neurodegeneration. *Cell* **127**: 59-69.
- Cummings, C.J., Mancini, M.A., Antalfy, B., DeFranco, D.B., Orr, H.T. and Zoghbi, H.Y. (1998) Chaperone suppression of aggregation and altered subcellular proteasome localization imply protein misfolding in SCA1. *Nat Genet* **19**: 148-154.

- Cummings, C.J., Reinstein, E., Sun, Y., Antalffy, B., Jiang, Y., Ciechanover, A., Orr, H.T., Beaudet, A.L. and Zoghbi, H.Y. (1999) Mutation of the E6-AP ubiquitin ligase reduces nuclear inclusion frequency while accelerating polyglutamine-induced pathology in SCA1 mice. *Neuron* **24**: 879-892.
- Cummings, C.J., Sun, Y., Opal, P., Antalffy, B., Mestril, R., Orr, H.T., Dillmann, W.H. and Zoghbi, H.Y. (2001) Over-expression of inducible HSP70 chaperone suppresses neuropathology and improves motor function in SCA1 mice. *Hum Mol Genet* **10**: 1511-1518.
- De Cristofaro, T., Affaitati, A., Cariello, L., Avvedimento, E.V. and Varrone, S. (1999) The length of polyglutamine tract, its level of expression, the rate of degradation, and the transglutaminase activity influence the formation of intracellular aggregates. *Biochem Biophys Res Commun* **260**: 150-158.
- De Pril, R., Fischer, D.F., Maat-Schieman, M.L.C., Hobo, B., de Vos, R.A.I., Brunt, E.R., Hol, E.M., Roos, R.A.C. and van Leeuwen, F.W. (2004) Accumulation of aberrant ubiquitin induces aggregate formation and cell death in polyglutamine diseases. *Hum Mol Genet* **13**: 1803-1813.
- De Vrij, F.M., Sluijs, J.A., Gregori, L., Fischer, D.F., Hermens, W.T., Goldgaber, D., Verhaagen, J., Van Leeuwen, F.W. and Hol, E.M. (2001) Mutant ubiquitin expressed in Alzheimer's disease causes neuronal death. *FASEB J* **15**: 2680-2688.
- De Vrij, F.M.S., Fischer, D.F., van Leeuwen, F.W. and Hol, E.M. (2004) Protein quality control in Alzheimer's disease by the ubiquitin proteasome system. *Prog Neurobiol* **74**: 249-270.
- Desai, U.A., Pallos, J., Ma, A.A.K., Stockwell, B.R., Thompson, L.M., Marsh, J.L. and Diamond, M.I. (2006) Biologically active molecules that reduce polyglutamine aggregation and toxicity. *Hum Mol Genet* **15**: 2114-2124.
- Diaz-Hernandez, M., Hernandez, F., Martin-Aparicio, E., Gomez-Ramos, P., Moran, M.A., Castano, J.G., Ferrer, I., Avila, J. and Lucas, J.J. (2003) Neuronal induction of the immunoproteasome in Huntington's disease. *J Neurosci* **23**: 11653-11661.
- DiFiglia, M., Sapp, E., Chase, K.O., Davies, S.W., Bates, G.P., Vonsattel, J.P. and Aronin, N. (1997) Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science* **277**: 1990-1993.
- Diguet, E., Petit, F., Escartin, C., Cambon, K., Bizat, N., Dufour, N., Hantraye, P., Deglon, N. and Brouillet, E. (2009) Normal aging modulates the neurotoxicity of mutant huntingtin. *PLoS ONE* **4**: e4637.
- Ding, Q., Lewis, J.J., Strum, K.M., Dimayuga, E., Bruce-Keller, A.J., Dunn, J.C. and Keller, J.N. (2002) Polyglutamine expansion, protein aggregation, proteasome activity, and neural survival. *J Biol Chem* **277**: 13935-13942.
- Dompierre, J.P., Godin, J.D., Charrin, B.C., Cordelieres, F.P., King, S.J., Humbert, S. and Saudou, F. (2007) Histone deacetylase 6 inhibition compensates for the transport deficit in Huntington's disease by increasing tubulin acetylation. *J Neurosci* **27**: 3571-3583.
- Donaldson, K.M., Li, W., Ching, K.A., Batalov, S., Tsai, C.C. and Joazeiro, C.A.P. (2003) Ubiquitin-mediated sequestration of normal cellular proteins into polyglutamine aggregates. *Proc Natl Acad Sci U S A* **100**: 8892-8897.
- Donoso, G., Herzog, V. and Schmitz, A. (2005) Misfolded BiP is degraded by a proteasome-independent endoplasmic-reticulum-associated degradation pathway. *Biochem J* **387**: 897-903.

References

- Doumanis, J., Wada, K., Kino, Y., Moore, A.W. and Nukina, N. (2009) RNAi screening in *Drosophila* cells identifies new modifiers of mutant huntingtin aggregation. *PLoS ONE* **4**: e7275.
- Dragatsis, I., Levine, M.S. and Zeitlin, S. (2000) Inactivation of Hdh in the brain and testis results in progressive neurodegeneration and sterility in mice. *Nat Genet* **26**: 300-306.
- Dunah, A.W., Jeong, H., Griffin, A., Kim, Y.M., Standaert, D.G., Hersch, S.M., Mouradian, M.M., Young, A.B., Tanese, N. and Krainc, D. (2002) Sp1 and TAFII130 transcriptional activity disrupted in early Huntington's disease. *Science* **296**: 2238-2243.
- Duyao, M.P., Auerbach, A.B., Ryan, A., Persichetti, F., Barnes, G.T., McNeil, S.M., Ge, P., Vonsattel, J.P., Gusella, J.F., Joyner, A.L. and et al. (1995) Inactivation of the mouse Huntington's disease gene homolog Hdh. *Science* **269**: 407-410.
- Ehlers, M.D. (2003) Ubiquitin and synaptic dysfunction: Ataxic mice highlight new common themes in neurological disease. *Trends Neurosci* **26**: 4-7.
- Elsasser, S., Chandler-Militello, D., Muller, B., Hanna, J. and Finley, D. (2004) Rad23 and Rpn10 serve as alternative ubiquitin receptors for the proteasome. *J Biol Chem* **279**: 26817-26822.
- Encinas, M., Iglesias, M., Liu, Y., Wang, H., Muñaisen, A., Cena, V., Gallego, C. and Comella, J.X. (2000) Sequential treatment of SH-SY5Y cells with retinoic acid and brain-derived neurotrophic factor gives rise to fully differentiated, neurotrophic factor-dependent, human neuron-like cells. *J Neurochem* **75**: 991-1003.
- Evans, D.A., van der Kleij, A.A., Sonnemans, M.A., Burbach, J.P. and van Leeuwen, F.W. (1994) Frameshift mutations at two hotspots in vasopressin transcripts in post-mitotic neurons. *Proc Natl Acad Sci U S A* **91**: 6059-6063.
- Evans, D.A.P., Burbach, J.P.H., Swaab, D.F. and Van Leeuwen, F.W. (1996) Mutant vasopressin precursors in the human hypothalamus: Evidence for neuronal somatic mutations in man. *Neuroscience* **71**: 1025-1030.
- Everett, C.M. and Wood, N.W. (2004) Trinucleotide repeats and neurodegenerative disease. *Brain* **127**: 2385-2405.
- Fan, J., Cowan, C.M., Zhang, L.Y.J., Hayden, M.R. and Raymond, L.A. (2009) Interaction of postsynaptic density protein-95 with NMDA receptors influences excitotoxicity in the yeast artificial chromosome mouse model of Huntington's disease. *J Neurosci* **29**: 10928-10938.
- Feecke, W., Gianfriddo, M., Gaviraghi, G., Terstappen, G.C. and Heitz, F. (2009) Small molecule drug discovery for Huntington's disease. *Drug Discovery Today* **14**: 453-464.
- Field, J., Nikawa, J., Broek, D., MacDonald, B., Rodgers, L., Wilson, I.A., Lerner, R.A. and Wigler, M. (1988) Purification of a RAS-responsive adenylyl cyclase complex from *Saccharomyces cerevisiae* by use of an epitope addition method. *Mol Cell Biol* **8**: 2159-2165.
- Fischer, D.F., de Pril, R., van Steenhoven, D.M.P.C., Janssen, R.A.J., Kwak, S., Howland, D.S. and Signer, E. (2008) Molecular targets and compounds, and methods to identify the same, useful in the treatment of neurodegenerative diseases. *US patent*: WO2009098196 and WO2009098197.
- Fischer, D.F., de Vos, R.A.I., van Dijk, R., de Vrij, F.M.S., Proper, E.A., Sonnemans, M.A.F., Verhage, M.C., Sluijs, J.A., Hobo, B., Zouambia, M., Jansen Steur, E.N.H., Kamphorst, W., Hol, E.M. and van Leeuwen, F.W. (2003) Disease-specific accumulation of mutant ubiquitin as a marker for proteasomal dysfunction in the brain. *FASEB J* **17**: 2014-2024.

- Fischer, D.F., van Dijk, R., van Tijn, P., Hobo, B., Verhage, M.C., van der Schors, R.C., Li, K.W., van Minnen, J., Hol, E.M. and van Leeuwen, F.W. (2009) Long-term proteasome dysfunction in the mouse brain by expression of aberrant ubiquitin. *Neurobiol Aging* **30**: 847-863.
- Fratta, P., Engel, W.K., van Leeuwen, F.W., Hol, E.M., Vattemi, G. and Askanas, V. (2004) Mutant ubiquitin UBB⁺¹ is accumulated in sporadic inclusion-body myositis muscle fibers. *Neurology* **63**: 1114-1117.
- French, B.A., van Leeuwen, F., Riley, N.E., Yuan, Q.X., Bardag-Gorce, F., Gaal, K., Lue, Y.H., Marceau, N. and French, S.W. (2001) Aggresome formation in liver cells in response to different toxic mechanisms: Role of the ubiquitin-proteasome pathway and the frameshift mutant of ubiquitin. *Exp Mol Pathol* **71**: 241-246.
- Fruh, K., Gossen, M., Wang, K., Bujard, H., Peterson, P.A. and Yang, Y. (1994) Displacement of housekeeping proteasome subunits by MHC-encoded LMPs: A newly discovered mechanism for modulating the multicatalytic proteinase complex. *EMBO J* **13**: 3236-3244.
- Furukawa, Y., Vigouroux, S., Wong, H., Guttman, M., Rajput, A.H., Ang, L., Briand, M., Kish, S.J. and Briand, Y. (2002) Brain proteasomal function in sporadic Parkinson's disease and related disorders. *Ann Neurol* **51**: 779-782.
- Gauthier, L.R., Charrin, B.C., Borrell-Pages, M., Dompierre, J.P., Rangone, H., Cordelieres, F.P., De Mey, J., MacDonald, M.E., Lessmann, V., Humbert, S. and Saudou, F. (2004) Huntingtin controls neurotrophic support and survival of neurons by enhancing BDNF vesicular transport along microtubules. *Cell* **118**: 127-138.
- Gerez, L., de Haan, A., Hol, E.M., Fischer, D.F., van Leeuwen, F.W., van Steeg, H. and Benne, R. (2005) Molecular misreading: The frequency of dinucleotide deletions in neuronal mRNAs for β -amyloid precursor protein and ubiquitin-B. *Neurobiol Aging* **26**: 145-155.
- Gervais, F.G., Singaraja, R., Xanthoudakis, S., Gutekunst, C.A., Leavitt, B.R., Metzler, M., Hackam, A.S., Tam, J., Vaillancourt, J.P., Houtzager, V., Rasper, D.M., Roy, S., Hayden, M.R. and Nicholson, D.W. (2002) Recruitment and activation of caspase-8 by the huntingtin-interacting protein Hip-1 and a novel partner Hippi. *Nat Cell Biol* **4**: 95-105.
- Giasson, B.I. and Lee, V.M.Y. (2003) Are ubiquitination pathways central to Parkinson's disease? *Cell* **114**: 1-8.
- Glickman, M.H. and Ciechanover, A. (2002) The ubiquitin-proteasome proteolytic pathway: Destruction for the sake of construction. *Physiol Rev* **82**: 373-428.
- Goehler, H., Lalowski, M., Stelzl, U., Waelter, S., Stroedicke, M., Worm, U., Droege, A., Lindenberg, K.S., Knoblich, M., Haenig, C., Herbst, M., Suopanki, J., Scherzinger, E., Abraham, C., Bauer, B., Hasenbank, R., Fritzsche, A., Ludewig, A.H., Buessow, K., Coleman, S.H., Gutekunst, C.A., Landwehrmeyer, B.G., Lehrach, H. and Wanker, E.E. (2004) A protein interaction network links GIT1, an enhancer of huntingtin aggregation, to Huntington's disease. *Mol Cell* **15**: 853-865.
- Graham, R.K., Deng, Y., Slow, E.J., Haigh, B., Bissada, N., Lu, G., Pearson, J., Shehadeh, J., Bertram, L., Murphy, Z., Warby, S.C., Doty, C.N., Roy, S., Wellington, C.L., Leavitt, B.R., Raymond, L.A., Nicholson, D.W. and Hayden, M.R. (2006) Cleavage at the caspase-6 site is required for neuronal dysfunction and degeneration due to mutant huntingtin. *Cell* **125**: 1179-1191.

References

- Gray, M., Shirasaki, D.I., Cepeda, C., Andre, V.M., Wilburn, B., Lu, X.H., Tao, J., Yamazaki, I., Li, S.H., Sun, Y.E., Li, X.J., Levine, M.S. and Yang, X.W. (2008) Full-length human mutant huntingtin with a stable polyglutamine repeat can elicit progressive and selective neuropathogenesis in BACHD mice. *J Neurosci* **28**: 6182-6195.
- Gunawardena, S., Her, L.S., Brusch, R.G., Laymon, R.A., Niesman, I.R., Gordesky-Gold, B., Sintasath, L., Bonini, N.M. and Goldstein, L.S. (2003) Disruption of axonal transport by loss of huntingtin or expression of pathogenic polyQ proteins in Drosophila. *Neuron* **40**: 25-40.
- Guterman, A. and Glickman, M.H. (2004) Deubiquitinating enzymes are IN/(trinsic to proteasome function). *Curr Protein Pept Sci* **5**: 201-211.
- Hackam, A.S., Hodgson, J.G., Singaraja, R., Zhang, T., Gan, L., Gutekunst, C.A., Hersch, S.M. and Hayden, M.R. (1999) Evidence for both the nucleus and cytoplasm as subcellular sites of pathogenesis in Huntington's disease in cell culture and in transgenic mice expressing mutant huntingtin. *Philos Trans R Soc Lond B Biol Sci* **354**: 1047-1055.
- Haldeman, M.T., Xia, G., Kasperek, E.M. and Pickart, C.M. (1997) Structure and function of ubiquitin conjugating enzyme E2-25K: The tail is a core-dependent activity element. *Biochemistry (Mosc)* **36**: 10526-10537.
- Halliday, G.M., McRitchie, D.A., Macdonald, V., Double, K.L., Trent, R.J. and McCusker, E. (1998) Regional specificity of brain atrophy in Huntington's disease. *Exp Neurol* **154**: 663-672.
- Hay, D.G., Sathasivam, K., Tobaben, S., Stahl, B., Marber, M., Mestrel, R., Mahal, A., Smith, D.L., Woodman, B. and Bates, G.P. (2004) Progressive decrease in chaperone protein levels in a mouse model of Huntington's disease and induction of stress proteins as a therapeutic approach. *Hum Mol Genet* **13**: 1389-1405.
- Hegde, A.N. (2004) Ubiquitin-proteasome-mediated local protein degradation and synaptic plasticity. *Prog Neurobiol* **73**: 311-357.
- Heng, M.Y., Detloff, P.J. and Albin, R.L. (2008) Rodent genetic models of Huntington disease. *Neurobiol Dis* **32**: 1-9.
- Hodges, A., Strand, A.D., Aragaki, A.K., Kuhn, A., Sengstag, T., Hughes, G., Elliston, L.A., Hartog, C., Goldstein, D.R., Thu, D., Hollingsworth, Z.R., Collin, F., Synek, B., Holmans, P.A., Young, A.B., Wexler, N.S., Delorenzi, M., Kooperberg, C., Augood, S.J., Faull, R.L.M., Olson, J.M., Jones, L. and Luthi-Carter, R. (2006) Regional and cellular gene expression changes in human Huntington's disease brain. *Hum Mol Genet* **15**: 965-977.
- Hol, E.M., Roelofs, R.F., Moraal, E., Sonnemans, M.A., Sluijs, J.A., Proper, E.A., de Graan, P.N., Fischer, D.F. and van Leeuwen, F.W. (2003) Neuronal expression of GFAP in patients with Alzheimer pathology and identification of novel GFAP splice forms. *Mol Psychiatry* **8**: 786-796.
- Holmberg, C.I., Staniszewski, K.E., Mensah, K.N., Matouschek, A. and Morimoto, R.I. (2004) Inefficient degradation of truncated polyglutamine proteins by the proteasome. *EMBO J* **23**: 4307-4318.
- Hope, A.D., De Silva, R., Fischer, D.F., Hol, E.M., Van Leeuwen, F.W. and Lees, A.J. (2003) Alzheimer's associated variant ubiquitin causes inhibition of the 26S proteasome and chaperone expression. *J Neurochem* **86**: 394-404.

- Hu, J., Matsui, M., Gagnon, K.T., Schwartz, J.C., Gabillet, S., Arar, K., Wu, J., Bezprozvanny, I. and Corey, D.R. (2009) Allele-specific silencing of mutant huntingtin and ataxin-3 genes by targeting expanded CAG repeats in mRNAs. *Nat Biotechnol* **27**: 478-484.
- Iijima-Ando, K., Wu, P., Drier, E.A., Iijima, K. and Yin, J.C.P. (2005) cAMP-response element-binding protein and heat-shock protein 70 additively suppress polyglutamine-mediated toxicity in *Drosophila*. *Proc Natl Acad Sci U S A* **102**: 10261-10266.
- Ishikawa, K., Fujigasaki, H., Saegusa, H., Ohwada, K., Fujita, T., Iwamoto, H., Komatsuzaki, Y., Toru, S., Toriyama, H., Watanabe, M., Ohkoshi, N., Shoji, S., Kanazawa, I., Tanabe, T. and Mizusawa, H. (1999) Abundant expression and cytoplasmic aggregations of α 1A voltage-dependent calcium channel protein associated with neurodegeneration in spinocerebellar ataxia type 6. *Hum Mol Genet* **8**: 1185-1193.
- Iwata, A., Nagashima, Y., Matsumoto, L., Suzuki, T., Yamanaka, T., Date, H., Deoka, K., Nukina, N. and Tsuji, S. (2009) Intranuclear degradation of polyglutamine aggregates by the ubiquitin-proteasome system. *J Biol Chem* **284**: 9796-9803.
- Iwata, A., Riley, B.E., Johnston, J.A. and Kopito, R.R. (2005) HDAC6 and microtubules are required for autophagic degradation of aggregated huntingtin. *J Biol Chem* **280**: 40282-40292.
- Jana, N.R., Dikshit, P., Goswami, A., Kotliarov, S., Murata, S., Tanaka, K. and Nukina, N. (2005) Co-chaperone CHIP associates with expanded polyglutamine protein and promotes their degradation by proteasomes. *J Biol Chem* **280**: 11635-11640.
- Jana, N.R., Zemskov, E.A., Wang, G.H. and Nukina, N. (2001) Altered proteasomal function due to the expression of polyglutamine-expanded truncated N-terminal huntingtin induces apoptosis by caspase activation through mitochondrial cytochrome c release. *Hum Mol Genet* **10**: 1049-1059.
- Jeong, H., Then, F., Melia, T.J., Mazzulli, J.R., Cui, L., Savas, J.N., Voisine, C., Paganetti, P., Tanese, N., Hart, A.C., Yamamoto, A. and Krainc, D. (2009) Acetylation targets mutant huntingtin to autophagosomes for degradation. *Cell* **137**: 60-72.
- Kalchman, M.A., Graham, R.K., Xia, G., Koide, H.B., Hodgson, J.G., Graham, K.C., Goldberg, Y.P., Gietz, R.D., Pickart, C.M. and Hayden, M.R. (1996) Huntingtin is ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. *J Biol Chem* **271**: 19385-19394.
- Kalchman, M.A., Koide, H.B., McCutcheon, K., Graham, R.K., Nichol, K., Nishiyama, K., Kazemi-Esfarjani, P., Lynn, F.C., Wellington, C., Metzler, M., Goldberg, Y.P., Kanazawa, I., Gietz, R.D. and Hayden, M.R. (1997) HIP1, a human homologue of *S. Cerevisiae* Sla2p, interacts with membrane-associated huntingtin in the brain. *Nat Genet* **16**: 44-53.
- Kaltenbach, L.S., Romero, E., Becklin, R.R., Chettier, R., Bell, R., Phansalkar, A., Strand, A., Torcassi, C., Savage, J., Hurlburt, A., Cha, G.H., Ukani, L., Chepanoske, C.L., Zhen, Y., Sahasrabudhe, S., Olson, J., Kurschner, C., Ellerby, L.M., Peltier, J.M., Botas, J. and Hughes, R.E. (2007) Huntingtin interacting proteins are genetic modifiers of neurodegeneration. *PLoS Genet* **3**: e82.
- Katayama, T., Imaizumi, K., Manabe, T., Hitomi, J., Kudo, T. and Tohyama, M. (2004) Induction of neuronal death by ER stress in Alzheimer's disease. *J Chem Neuroanat* **28**: 67-78.
- Keck, S., Nitsch, R., Grune, T. and Ullrich, O. (2003) Proteasome inhibition by paired helical filament-tau in brains of patients with Alzheimer's disease. *J Neurochem* **85**: 115-122.

References

- Keller, J.N., Dimayuga, E., Chen, Q., Thorpe, J., Gee, J. and Ding, Q. (2004) Autophagy, proteasomes, lipofuscin, and oxidative stress in the aging brain. *Int J Biochem Cell B* **36**: 2376-2391.
- Keller, J.N., Gee, J. and Ding, Q. (2002) The proteasome in brain aging. *Ageing Res Rev* **1**: 279-293.
- Keller, J.N., Hanni, K.B. and Markesberry, W.R. (2000) Impaired proteasome function in Alzheimer's disease. *J Neurochem* **75**: 436-439.
- Kennedy, W.R., Alter, M. and Sung, J.H. (1968) Progressive proximal spinal and bulbar muscular atrophy of late onset: A sex-linked recessive trait. *Neurology* **18**: 671-680.
- Kerstens, H.M., Poddighe, P.J. and Hanselaar, A.G. (1995) A novel in situ hybridization signal amplification method based on the deposition of biotinylated tyramine. *J Histochem Cytochem* **43**: 347-352.
- Kim, H.C. and Huibregtse, J.M. (2009) Polyubiquitination by HECT E3s and the determinants of chain type specificity. *Mol Cell Biol* **29**: 3307-3318.
- Kim, I., Mi, K. and Rao, H. (2004) Multiple interactions of Rad23 suggest a mechanism for ubiquitylated substrate delivery important in proteolysis. *Mol Biol Cell* **15**: 3357-3365.
- Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S., Yokochi, M., Mizuno, Y. and Shimizu, N. (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* **392**: 605-608.
- Klement, I.A., Skinner, P.J., Kaytor, M.D., Yi, H., Hersch, S.M., Clark, H.B., Zoghbi, H.Y. and Orr, H.T. (1998) Ataxin-1 nuclear localization and aggregation: Role in polyglutamine-induced disease in SCA1 transgenic mice. *Cell* **95**: 41-53.
- Ko, S., Kang, G.B., Song, S.M., Lee, J.G., Shin, D.Y., Yun, J.H., Sheng, Y., Cheong, C., Jeon, Y.H., Jung, Y.K., Arrowsmith, C.H., Avvakumov, G.V., Dhe-Paganon, S., Yoo, Y.J., Eom, S.H. and Lee, W. (2010) Structural basis of E2-25K/UBB+1 interaction leading to proteasome inhibition and neurotoxicity. *J Biol Chem* **285**: 36070-36080.
- Koide, R., Ikeuchi, T., Onodera, O., Tanaka, H., Igarashi, S., Endo, K., Takahashi, H., Kondo, R., Ishikawa, A., Hayashi, T. and et al. (1994) Unstable expansion of CAG repeat in hereditary dentatorubral-pallidoluysian atrophy (DRPLA). *Nat Genet* **6**: 9-13.
- Konishi, Y., Beach, T., Sue, L.I., Hampel, H., Lindholm, K. and Shen, Y. (2003) The temporal localization of frame-shift ubiquitin-B and amyloid precursor protein, and complement proteins in the brain of non-demented control patients with increasing Alzheimer's disease pathology. *Neurosci Lett* **348**: 46-50.
- Kopito, R.R. and Sitia, R. (2000) Aggresomes and Russell bodies. Symptoms of cellular indigestion? *EMBO Rep* **1**: 225-231.
- Korolchuk, V.I., Mansilla, A., Menzies, F.M. and Rubinsztein, D.C. (2009) Autophagy inhibition compromises degradation of ubiquitin-proteasome pathway substrates. *Mol Cell* **33**: 517-527.
- Kostova, Z. and Wolf, D.H. (2003) For whom the bell tolls: Protein quality control of the endoplasmic reticulum and the ubiquitin-proteasome connection. *EMBO J* **22**: 2309-2317.
- Kovacs, J.J., Murphy, P.J.M., Gaillard, S., Zhao, X., Wu, J.T., Nicchitta, C.V., Yoshida, M., Toft, D.O., Pratt, W.B. and Yao, T.P. (2005) HDAC6 regulates Hsp90 acetylation and chaperone-dependent activation of glucocorticoid receptor. *Mol Cell* **18**: 601-607.

- Lam, Y.A., Lawson, T.G., Velayutham, M., Zweier, J.L. and Pickart, C.M. (2002) A proteasomal ATPase subunit recognizes the polyubiquitin degradation signal. *Nature* **416**: 763-767.
- Lam, Y.A., Pickart, C.M., Alban, A., Landon, M., Jamieson, C., Ramage, R., Mayer, R.J. and Layfield, R. (2000) Inhibition of the ubiquitin-proteasome system in Alzheimer's disease. *Proc Natl Acad Sci U S A* **97**: 9902-9906.
- Landles, C. and Bates, G.P. (2004) Huntingtin and the molecular pathogenesis of Huntington's disease. *EMBO Rep* **5**: 958-963.
- Lappe-Siefke, C., Loebrich, S., Hevers, W., Waidmann, O.B., Schweizer, M., Fehr, S., Fritschy, J.M., Dikic, I., Eilers, J., Wilson, S.M. and Kneussel, M. (2009) The ataxia *ax^l* mutation causes abnormal GABA_A receptor turnover in mice. *PLoS Genet* **5**: e1000631.
- Lee, B.H., Lee, M.J., Park, S., Oh, D.C., Elsasser, S., Chen, P.C., Gartner, C., Dimova, N., Hanna, J., Gygi, S.P., Wilson, S.M., King, R.W. and Finley, D. (2010) Enhancement of proteasome activity by a small-molecule inhibitor of USP14. *Nature* **467**: 179-184.
- Levine, A.J. (1997) P53, the cellular gatekeeper for growth and division. *Cell* **88**: 323-331.
- Li, J.L., Hayden, M.R., Almqvist, E.W., Brinkman, R.R., Durr, A., Dode, C., Morrison, P.J., Suchowersky, O., Ross, C.A., Margolis, R.L., Rosenblatt, A., Gomez-Tortosa, E., Cabrero, D.M., Novelletto, A., Frontali, M., Nance, M., Trent, R.J., McCusker, E., Jones, R., Paulsen, J.S., Harrison, M., Zanko, A., Abramson, R.K., Russ, A.L., Knowlton, B., Djousse, L., Mysore, J.S., Tariot, S., Gusella, M.F., Wheeler, V.C., Atwood, L.D., Cupples, L.A., Saint-Hilaire, M., Cha, J.H., Hersch, S.M., Koroshetz, W.J., Gusella, J.F., MacDonald, M.E. and Myers, R.H. (2003) A genome scan for modifiers of age at onset in Huntington disease: The HD MAPS study. *Am J Hum Genet* **73**: 682-687.
- Li, S.H., Cheng, A.L., Zhou, H., Lam, S., Rao, M., Li, H. and Li, X.J. (2002) Interaction of Huntington disease protein with transcriptional activator Sp1. *Mol Cell Biol* **22**: 1277-1287.
- Li, S.H., Lam, S., Cheng, A.L. and Li, X.J. (2000) Intranuclear huntingtin increases the expression of caspase-1 and induces apoptosis. *Hum Mol Genet* **9**: 2859-2867.
- Li, X., Wang, C.E., Huang, S., Xu, X., Li, X.J., Li, H. and Li, S. (2010) Inhibiting the ubiquitin-proteasome system leads to preferential accumulation of toxic N-terminal mutant huntingtin fragments. *Hum Mol Genet* **19**: 2445-2455.
- Lindsten, K. and Dantuma, N.P. (2003) Monitoring the ubiquitin/proteasome system in conformational diseases. *Ageing Res Rev* **2**: 433-449.
- Lindsten, K., de Vrij, F.M., Verhoef, L.G., Fischer, D.F., van Leeuwen, F.W., Hol, E.M., Masucci, M.G. and Dantuma, N.P. (2002) Mutant ubiquitin found in neurodegenerative disorders is a ubiquitin fusion degradation substrate that blocks proteasomal degradation. *J Cell Biol* **157**: 417-427.
- Lomas, D.A. and Carrell, R.W. (2002) Serpinopathies and the conformational dementias. *Nat Rev Genet* **3**: 759-768.
- Lu, C., Cantin, M., Seidah, N. and Chretien, M. (1982) Immunohistochemical localization of human pituitary glycoprotein (HPGP)-like immunoreactivity in the hypothalamus and pituitary of normal and homozygous diabetes insipidus (Brattleboro) rats. *J Histochem Cytochem* **30**: 999-1003.
- Lunkes, A. and Mandel, J. (1998) A cellular model that recapitulates major pathogenic steps of Huntington's disease. *Hum Mol Genet* **7**: 1355-1361.

References

- Luo, S., Mizuta, H. and Rubinsztein, D.C. (2008) P21-activated kinase 1 promotes soluble mutant huntingtin self-interaction and enhances toxicity. *Hum Mol Genet* **17**: 895-905.
- Luo, S. and Rubinsztein, D.C. (2009) Huntingtin promotes cell survival by preventing Pak2 cleavage. *J Cell Sci* **122**: 875-885.
- Luthi-Carter, R., Hanson, S.A., Strand, A.D., Bergstrom, D.A., Chun, W., Peters, N.L., Woods, A.M., Chan, E.Y., Kooperberg, C., Krainc, D., Young, A.B., Tapscott, S.J. and Olson, J.M. (2002) Dysregulation of gene expression in the R6/2 model of polyglutamine disease: Parallel changes in muscle and brain. *Hum Mol Genet* **11**: 1911-1926.
- Luthi-Carter, R., Strand, A., Peters, N.L., Solano, S.M., Hollingsworth, Z.R., Menon, A.S., Frey, A.S., Spektor, B.S., Penney, E.B., Schilling, G., Ross, C.A., Borchelt, D.R., Tapscott, S.J., Young, A.B., Cha, J.H. and Olson, J.M. (2000) Decreased expression of striatal signaling genes in a mouse model of Huntington's disease. *Hum Mol Genet* **9**: 1259-1271.
- Luthi-Carter, R., Strand, A.D., Hanson, S.A., Kooperberg, C., Schilling, G., La Spada, A.R., Merry, D.E., Young, A.B., Ross, C.A., Borchelt, D.R. and Olson, J.M. (2002) Polyglutamine and transcription: Gene expression changes shared by DRPLA and Huntington's disease mouse models reveal context-independent effects. *Hum Mol Genet* **11**: 1927-1937.
- Luthi-Carter, R., Taylor, D.M., Pallos, J., Lambert, E., Amore, A., Parker, A., Moffitt, H., Smith, D.L., Runne, H., Gokce, O., Kuhn, A., Xiang, Z., Maxwell, M.M., Reeves, S.A., Bates, G.P., Neri, C., Thompson, L.M., Marsh, J.L. and Kazantsev, A.G. (2010) SIRT2 inhibition achieves neuroprotection by decreasing sterol biosynthesis. *Proc Natl Acad Sci U S A* **107**: 7927-7932.
- Maat-Schieman, M.L., Dorsman, J.C., Smoor, M.A., Siesling, S., van Duinen, S.G., Verschueren, J.J., den Dunnen, J.T., van Ommen, G.J. and Roos, R.A.C. (1999) Distribution of inclusions in neuronal nuclei and dystrophic neurites in Huntington disease brain. *J Neuropathol Exp Neurol* **58**: 129-137.
- Mangiarini, L., Sathasivam, K., Seller, M., Cozens, B., Harper, A., Hetherington, C., Lawton, M., Trottier, Y., Lehrach, H., Davies, S.W. and Bates, G.P. (1996) Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice. *Cell* **87**: 493-506.
- Mantamadiotis, T., Lemberger, T., Bleckmann, S.C., Kern, H., Kretz, O., Martin Villalba, A., Tronche, F., Kellendonk, C., Gau, D., Kapfhammer, J., Otto, C., Schmid, W. and Schutz, G. (2002) Disruption of CREB function in brain leads to neurodegeneration. *Nat Genet* **31**: 47-54.
- Matsumoto, M., Yada, M., Hatakeyama, S., Ishimoto, H., Tanimura, T., Tsuji, S., Kakizuka, A., Kitagawa, M. and Nakayama, K.I. (2004) Molecular clearance of ataxin-3 is regulated by a mammalian E4. *EMBO J* **23**: 659-669.
- Maynard, C.J., Bottcher, C., Ortega, Z., Smith, R., Florea, B.I., Diaz-Hernandez, M., Brundin, P., Overkleef, H.S., Li, J.Y., Lucas, J.J. and Dantuma, N.P. (2009) Accumulation of ubiquitin conjugates in a polyglutamine disease model occurs without global ubiquitin/proteasome system impairment. *Proc Natl Acad Sci U S A* **106**: 13986-13991.
- McDonough, H. and Patterson, C. (2003) CHIP: A link between the chaperone and proteasome systems. *Cell Stress Chaperones* **8**: 303-308.
- McNaught, K.S.P. (2004) Proteolytic dysfunction in neurodegenerative disorders. *Int Rev Neurobiol*: 95-119.

- McNeil, S., Novelletto, A., Srinidhi, J., Barnes, G., Kornbluth, I., Altherr, M., Wasmuth, J., Gusella, J., MacDonald, M. and Myers, R. (1997) Reduced penetrance of the Huntington's disease mutation. *Hum Mol Genet* **6**: 775-779.
- McPhaul, L.W., Wang, J., Hol, E.M., Sonnemann, M.A.F., Riley, N., Nguyen, V., Yuan, Q.X., Lue, Y.H., van Leeuwen, F.W. and French, S.W. (2002) Molecular misreading of the ubiquitin B gene and hepatic mallory body formation. *Gastroenterology* **122**: 1878-1885.
- Menalled, L.B., Sison, J.D., Wu, Y., Olivieri, M., Li, X.J., Li, H., Zeitlin, S. and Chesselet, M.F. (2002) Early motor dysfunction and striosomal distribution of huntingtin microaggregates in Huntington's disease knock-in mice. *J Neurosci* **22**: 8266-8276.
- Menzies, F.M., Huebener, J., Renna, M., Bonin, M., Riess, O. and Rubinsztein, D.C. (2010) Autophagy induction reduces mutant ataxin-3 levels and toxicity in a mouse model of spinocerebellar atrophy type 3. *Brain* **133**: 93-104.
- Metzger, S., Rong, J., Nguyen, H.P., Cape, A., Tomiuk, J., Soehn, A.S., Propping, P., Freudenberg-Hua, Y., Freudenberg, J., Tong, L., Li, S.H., Li, X.J. and Riess, O. (2008) Huntington-associated protein-1 is a modifier of the age-at-onset of Huntington's disease. *Hum Mol Genet* **17**: 1137-1146.
- Michalik, A. and Van Broeckhoven, C. (2003) Pathogenesis of polyglutamine disorders: Aggregation revisited. *Hum Mol Genet* **12**: 173R-186R.
- Michalik, A. and Van Broeckhoven, C. (2004) Proteasome degrades soluble expanded polyglutamine completely and efficiently. *Neurobiol Dis* **16**: 202-211.
- Miller, R.J. and Wilson, S.M. (2003) Neurological disease: UPS stops delivering! *Trends Pharmacol Sci* **24**: 18-23.
- Miller, V.M., Nelson, R.F., Gouvion, C.M., Williams, A., Rodriguez-Lebron, E., Harper, S.Q., Davidson, B.L., Rebagliati, M.R. and Paulson, H.L. (2005) CHIP suppresses polyglutamine aggregation and toxicity in vitro and in vivo. *J Neurosci* **25**: 9152-9161.
- Miller, V.M., Xia, H., Marrs, G.L., Gouvion, C.M., Lee, G., Davidson, B.L. and Paulson, H.L. (2003) Allele-specific silencing of dominant disease genes. *Proc Natl Acad Sci U S A* **100**: 7195-7200.
- Mitra, S., Tsvetkov, A.S. and Finkbeiner, S. (2009) Single neuron ubiquitin-proteasome dynamics accompanying inclusion body formation in Huntington disease. *J Biol Chem* **284**: 4398-4403.
- Mitsui, K., Nakayama, H., Akagi, T., Nekooki, M., Ohtawa, K., Takio, K., Hashikawa, T. and Nukina, N. (2002) Purification of polyglutamine aggregates and identification of elongation factor-1 α and heat shock protein 84 as aggregate-interacting proteins. *J Neurosci* **22**: 9267-9277.
- Muchowski, P.J., Schaffar, G., Sittler, A., Wanker, E.E., Hayer-Hartl, M.K. and Hartl, F.U. (2000) Hsp70 and hsp40 chaperones can inhibit self-assembly of polyglutamine proteins into amyloid-like fibrils. *Proc Natl Acad Sci U S A* **97**: 7841-7846.
- Muchowski, P.J. and Wacker, J.L. (2005) Modulation of neurodegeneration by molecular chaperones. *Nat Rev Neurosci* **6**: 11-22.
- Munoz, E., Rey, M.J., Mila, M., Cardozo, A., Ribalta, T., Tolosa, E. and Ferrer, I. (2002) Intranuclear inclusions, neuronal loss and CAG mosaicism in two patients with Machado-Joseph disease. *J Neurol Sci* **200**: 19-25.

References

- Nakamura, K., Jeong, S.Y., Uchihara, T., Anno, M., Nagashima, K., Nagashima, T., Ikeda, S., Tsuji, S. and Kanazawa, I. (2001) SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. *Hum Mol Genet* **10**: 1441-1448.
- Naldini, L., Blomer, U., Gallay, P., Ory, D., Mulligan, R., Gage, F.H., Verma, I.M. and Trono, D. (1996) In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science* **272**: 263-267.
- Nasir, J., Floresco, S.B., O'Kusky, J.R., Diewert, V.M., Richman, J.M., Zeisler, J., Borowski, A., Marth, J.D., Phillips, A.G. and Hayden, M.R. (1995) Targeted disruption of the Huntington's disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes. *Cell* **81**: 811-823.
- Neumann, M. and Gabel, D. (2002) Simple method for reduction of autofluorescence in fluorescence microscopy. *J Histochem Cytochem* **50**: 437-439.
- Nixon, R.A. (2005) Endosome function and dysfunction in Alzheimer's disease and other neurodegenerative diseases. *Neurobiol Aging* **26**: 373-382.
- Nollen, E.A.A., Garcia, S.M., van Haaften, G., Kim, S., Chavez, A., Morimoto, R.I. and Plasterk, R.H.A. (2004) Genome-wide RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. *Proc Natl Acad Sci U S A* **101**: 6403-6408.
- Novina, C.D. and Sharp, P.A. (2004) The RNAi revolution. *Nature* **430**: 161-164.
- Nucifora, F.C., Jr., Ellerby, L.M., Wellington, C.L., Wood, J.D., Herring, W.J., Sawa, A., Hayden, M.R., Dawson, V.L., Dawson, T.M. and Ross, C.A. (2003) Nuclear localization of a non-caspase truncation product of atrophin-1, with an expanded polyglutamine repeat, increases cellular toxicity. *J Biol Chem* **278**: 13047-13055.
- Oddo, S., Billings, L., Kesslak, J.P., Cribbs, D.H. and LaFerla, F.M. (2004) A β immunotherapy leads to clearance of early, but not late, hyperphosphorylated tau aggregates via the proteasome. *Neuron* **43**: 321-332.
- Okamoto, S., Pouladi, M.A., Talantova, M., Yao, D., Xia, P., Ehrnhoefer, D.E., Zaidi, R., Clemente, A., Kaul, M., Graham, R.K., Zhang, D., Vincent Chen, H.S., Tong, G., Hayden, M.R. and Lipton, S.A. (2009) Balance between synaptic versus extrasynaptic NMDA receptor activity influences inclusions and neurotoxicity of mutant huntingtin. *Nat Med* **15**: 1407-1413.
- Okazawa, H., Rich, T., Chang, A., Lin, X., Waragai, M., Kajikawa, M., Enokido, Y., Komuro, A., Kato, S., Shibata, M., Hatanaka, H., Mouradian, M.M., Sudol, M. and Kanazawa, I. (2002) Interaction between mutant ataxin-1 and PQBP-1 affects transcription and cell death. *Neuron* **34**: 701-713.
- Orlowski, R.Z. and Kuhn, D.J. (2008) Proteasome inhibitors in cancer therapy: Lessons from the first decade. *Clin Cancer Res* **14**: 1649-1657.
- Ortega, Z., Diaz-Hernandez, M., Maynard, C.J., Hernandez, F., Dantuma, N.P. and Lucas, J.J. (2010) Acute polyglutamine expression in inducible mouse model unravels ubiquitin/proteasome system impairment and permanent recovery attributable to aggregate formation. *J Neurosci* **30**: 3675-3688.
- Pandey, U.B., Nie, Z., Batlevi, Y., McCray, B.A., Ritson, G.P., Nedelsky, N.B., Schwartz, S.L., DiProspero, N.A., Knight, M.A., Schuldiner, O., Padmanabhan, R., Hild, M., Berry, D.L., Garza, D., Hubbert, C.C., Yao, T.P., Baehrecke, E.H. and Taylor, J.P. (2007) HDAC6 rescues neurodegeneration and provides an essential link between autophagy and the UPS. *Nature* **447**: 859-863.

- Pankiv, S., Clausen, T.H., Lamark, T., Brech, A., Bruun, J.A., Outzen, H., Overvatn, A., Bjorkoy, G. and Johansen, T. (2007) P62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* **282**: 24131-24145.
- Pankiv, S., Lamark, T., Bruun, J.A., Overvatn, A., Bjorkoy, G. and Johansen, T. (2010) Nucleocytoplasmic shuttling of p62/SQSTM1 and its role in recruitment of nuclear polyubiquitinated proteins to promyelocytic leukemia bodies. *J Biol Chem* **285**: 5941-5953.
- Paulson, H.L., Das, S.S., Crino, P.B., Perez, M.K., Patel, S.C., Gotsdiner, D., Fischbeck, K.H. and Pittman, R.N. (1997) Machado-Joseph disease gene product is a cytoplasmic protein widely expressed in brain. *Ann Neurol* **41**: 453-462.
- Paulson, H.L., Perez, M.K., Trottier, Y., Trojanowski, J.Q., Subramony, S.H., Das, S.S., Vig, P., Mandel, J.L., Fischbeck, K.H. and Pittman, R.N. (1997) Intranuclear inclusions of expanded polyglutamine protein in spinocerebellar ataxia type-3. *Neuron* **19**: 333-344.
- Paxinos, G. and Franklin, K. (2001) The mouse brain in stereotaxic coordinates. Academic Press, p. 264.
- Perutz, M.F., Johnson, T., Suzuki, M. and Finch, J.T. (1994) Glutamine repeats as polar zippers: Their possible role in inherited neurodegenerative diseases. *Proc Natl Acad Sci U S A* **91**: 5355-5358.
- Peters, M.F., Nucifora, F.C., Jr., Kushi, J., Seaman, H.C., Cooper, J.K., Herring, W.J., Dawson, V.L., Dawson, T.M. and Ross, C.A. (1999) Nuclear targeting of mutant huntingtin increases toxicity. *Mol Cell Neurosci* **14**: 121-128.
- Peth, A., Besche, H.C. and Goldberg, A.L. (2009) Ubiquitinated proteins activate the proteasome by binding to Usp14/Ubp6, which causes 20S gate opening. *Mol Cell* **36**: 794-804.
- Petrucelli, L., Dickson, D., Kehoe, K., Taylor, J., Snyder, H., Grover, A., De Lucia, M., McGowan, E., Lewis, J., Prihar, G., Kim, J., Dillmann, W.H., Browne, S.E., Hall, A., Voellmy, R., Tsuboi, Y., Dawson, T.M., Wolozin, B., Hardy, J. and Hutton, M. (2004) CHIP and Hsp70 regulate tau ubiquitination, degradation and aggregation. *Hum Mol Genet* **13**: 703-714.
- Pfister, E.L., Kennington, L., Straubhaar, J., Wagh, S., Liu, W., DiFiglia, M., Landwehrmeyer, B., Vonsattel, J.P., Zamore, P.D. and Aronin, N. (2009) Five siRNAs targeting three SNPs may provide therapy for three-quarters of Huntington's disease patients. *Curr Biol* **19**: 774-778.
- Pichler, A., Knipscheer, P., Oberhofer, E., van Dijk, W.J., Korner, R., Olsen, J.V., Jentsch, S., Melchior, F. and Sixma, T.K. (2005) SUMO modification of the ubiquitin-conjugating enzyme E2-25K. *Nat Struct Mol Biol* **12**: 264-269.
- Pickart, C.M. (2001) Mechanisms underlying ubiquitination. *Annu Rev Biochem* **70**: 503-533.
- Pickart, C.M. and Cohen, R.E. (2004) Proteasomes and their kin: Proteases in the machine age. *Nat Rev Mol Cell Biol* **5**: 177-187.
- Pickart, C.M. and Fushman, D. (2004) Polyubiquitin chains: Polymeric protein signals. *Curr Opin Chem Biol* **8**: 610-616.
- Qin, Z.H., Wang, Y., Kegel, K.B., Kazantsev, A., Apostol, B.L., Thompson, L.M., Yoder, J., Aronin, N. and DiFiglia, M. (2003) Autophagy regulates the processing of amino terminal huntingtin fragments. *Hum Mol Genet* **12**: 3231-3244.

References

- Ratovitski, T., Gucek, M., Jiang, H., Chighladze, E., Waldron, E., D'Ambola, J., Hou, Z., Liang, Y., Poirier, M.A., Hirschhorn, R.R., Graham, R., Hayden, M.R., Cole, R.N. and Ross, C.A. (2009) Mutant huntingtin N-terminal fragments of specific size mediate aggregation and toxicity in neuronal cells. *J Biol Chem* **284**: 10855-10867.
- Ravikumar, B., Vacher, C., Berger, Z., Davies, J.E., Luo, S., Oroz, L.G., Scaravilli, F., Easton, D.F., Duden, R., O'Kane, C.J. and Rubinsztein, D.C. (2004) Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease. *Nat Genet* **36**: 585-595.
- Rechsteiner, M. and Hill, C.P. (2005) Mobilizing the proteolytic machine: Cell biological roles of proteasome activators and inhibitors. *Trends Cell Biol* **15**: 27-33.
- Rideout, H.J., Lang-Rollin, I. and Stefanis, L. (2004) Involvement of macroautophagy in the dissolution of neuronal inclusions. *Int J Biochem Cell B* **36**: 2551-2562.
- Ross, C.A. and Pickart, C.M. (2004) The ubiquitin-proteasome pathway in Parkinson's disease and other neurodegenerative diseases. *Trends Cell Biol* **14**: 703-711.
- Ross, C.A. and Poirier, M.A. (2004) Protein aggregation and neurodegenerative disease. *Nat Med* **10**: S10-17.
- Rubinsztein, D.C., Leggo, J., Coles, R., Almqvist, E., Biancalana, V., Cassiman, J.J., Chotai, K., Connarty, M., Crauford, D., Curtis, A., Curtis, D., Davidson, M.J., Differ, A.M., Dode, C., Dodge, A., Frontali, M., Ranen, N.G., Stine, O.C., Sherr, M., Abbott, M.H., Franz, M.L., Graham, C.A., Harper, P.S., Hedreen, J.C., Hayden, M.R. and et al. (1996) Phenotypic characterization of individuals with 30-40 CAG repeats in the Huntington disease (HD) gene reveals HD cases with 36 repeats and apparently normal elderly individuals with 36-39 repeats. *Am J Hum Genet* **59**: 16-22.
- Salehi, A. and Swaab, D.F. (1999) Diminished neuronal metabolic activity in Alzheimer's disease. *J Neural Transm* **106**: 955-986.
- Saudou, F., Finkbeiner, S., Devys, D. and Greenberg, M.E. (1998) Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions. *Cell* **95**: 55-66.
- Scherzinger, E., Lurz, R., Turmaine, M., Mangiarini, L., Hollenbach, B., Hasenbank, R., Bates, G.P., Davies, S.W., Lehrach, H. and Wanker, E.E. (1997) Huntingtin-encoded polyglutamine expansions form amyloid-like protein aggregates in vitro and in vivo. *Cell* **90**: 549-558.
- Schilling, G., Becher, M.W., Sharp, A.H., Jinnah, H.A., Duan, K., Kotzuk, J.A., Slunt, H.H., Ratovitski, T., Cooper, J.K., Jenkins, N.A., Copeland, N.G., Price, D.L., Ross, C.A. and Borchelt, D.R. (1999) Intranuclear inclusions and neuritic aggregates in transgenic mice expressing a mutant N-terminal fragment of huntingtin. *Hum Mol Genet* **8**: 397-407.
- Schilling, G., Savonenko, A.V., Klevytska, A., Morton, J.L., Tucker, S.M., Poirier, M., Gale, A., Chan, N., Gonzales, V., Slunt, H.H., Coonfield, M.L., Jenkins, N.A., Copeland, N.G., Ross, C.A. and Borchelt, D.R. (2004) Nuclear-targeting of mutant huntingtin fragments produces Huntington's disease-like phenotypes in transgenic mice. *Hum Mol Genet* **13**: 1599-1610.
- Schmidt, T., Lindenberg, K.S., Krebs, A., Schols, L., Laccone, F., Herms, J., Rechsteiner, M., Riess, O. and Landwehrmeyer, G.B. (2002) Protein surveillance machinery in brains with spinocerebellar ataxia type 3: Redistribution and differential recruitment of 26S proteasome subunits and chaperones to neuronal intranuclear inclusions. *Ann Neurol* **51**: 302-310.

- Schwartz, A.L. and Ciechanover, A. (2009) Targeting proteins for destruction by the ubiquitin system: Implications for human pathobiology. *Annu Rev Pharmacol Toxicol* **49**: 73-96.
- Schwartz, D.C. and Hochstrasser, M. (2003) A superfamily of protein tags: Ubiquitin, SUMO and related modifiers. *Trends Biochem Sci* **28**: 321-328.
- Seo, H., Sonntag, K.C. and Isacson, O. (2004) Generalized brain and skin proteasome inhibition in Huntington's disease. *Ann Neurol* **56**: 319-328.
- Sherman, M.Y. and Goldberg, A.L. (2001) Cellular defenses against unfolded proteins: A cell biologist thinks about neurodegenerative diseases. *Neuron* **29**: 15-32.
- Shimohata, T., Nakajima, T., Yamada, M., Uchida, C., Onodera, O., Naruse, S., Kimura, T., Koide, R., Nozaki, K., Sano, Y., Ishiguro, H., Sakoe, K., Ooshima, T., Sato, A., Ikeuchi, T., Oyake, M., Sato, T., Aoyagi, Y., Hozumi, I., Nagatsu, T., Takiyama, Y., Nishizawa, M., Goto, J., Kanazawa, I., Davidson, I., Tanese, N., Takahashi, H. and Tsuji, S. (2000) Expanded polyglutamine stretches interact with TAFII130, interfering with CREB-dependent transcription. *Nat Genet* **26**: 29-36.
- Shintani, T. and Klionsky, D.J. (2004) Autophagy in health and disease: A double-edged sword. *Science* **306**: 990-995.
- Sieradzan, K.A., Mechan, A.O., Jones, L., Wanker, E.E., Nukina, N. and Mann, D.M. (1999) Huntington's disease intranuclear inclusions contain truncated, ubiquitinated huntingtin protein. *Exp Neurol* **156**: 92-99.
- Sinadinos, C., Burbidge-King, T., Soh, D., Thompson, L.M., Marsh, J.L., Wyttenbach, A. and Mudher, A.K. (2009) Live axonal transport disruption by mutant huntingtin fragments in Drosophila motor neuron axons. *Neurobiol Dis* **34**: 389-395.
- Sipione, S., Rigamonti, D., Valenza, M., Zuccato, C., Conti, L., Pritchard, J., Kooperberg, C., Olson, J.M. and Cattaneo, E. (2002) Early transcriptional profiles in huntingtin-inducible striatal cells by microarray analyses. *Hum Mol Genet* **11**: 1953-1965.
- Song, S. and Jung, Y.K. (2004) Alzheimer's disease meets the ubiquitin-proteasome system. *Trends Mol Med* **10**: 565-570.
- Song, S., Kim, S.Y., Hong, Y.M., Jo, D.G., Lee, J.Y., Shim, S.M., Chung, C.W., Seo, S.J., Yoo, Y.J., Koh, J.Y., Lee, M.C., Yates, A.J., Ichijo, H. and Jung, Y.K. (2003) Essential role of E2-25K/Hip-2 in mediating amyloid- β neurotoxicity. *Mol Cell* **12**: 553-563.
- Steffan, J.S., Agrawal, N., Pallos, J., Rockabrand, E., Trotman, L.C., Slepko, N., Illes, K., Lukacsovich, T., Zhu, Y.Z., Cattaneo, E., Pandolfi, P.P., Thompson, L.M. and Marsh, J.L. (2004) SUMO modification of huntingtin and Huntington's disease pathology. *Science* **304**: 100-104.
- Steffan, J.S., Bodai, L., Pallos, J., Poelman, M., McCampbell, A., Apostol, B.L., Kazantsev, A., Schmidt, E., Zhu, Y.Z., Greenwald, M., Kurokawa, R., Housman, D.E., Jackson, G.R., Marsh, J.L. and Thompson, L.M. (2001) Histone deacetylase inhibitors arrest polyglutamine-dependent neurodegeneration in Drosophila. *Nature* **413**: 739-743.
- Steffan, J.S., Kazantsev, A., Spasic-Boskovic, O., Greenwald, M., Zhu, Y.Z., Gohler, H., Wanker, E.E., Bates, G.P., Housman, D.E. and Thompson, L.M. (2000) The Huntington's disease protein interacts with p53 and CREB-binding protein and represses transcription. *Proc Natl Acad Sci U S A* **97**: 6763-6768.

References

- Stenoien, D.L., Cummings, C.J., Adams, H.P., Mancini, M.G., Patel, K., DeMartino, G.N., Marcelli, M., Weigel, N.L. and Mancini, M.A. (1999) Polyglutamine-expanded androgen receptors form aggregates that sequester heat shock proteins, proteasome components and SRC-1, and are suppressed by the HDJ-2 chaperone. *Hum Mol Genet* **8**: 731-741.
- Strand, A.D., Baquet, Z.C., Aragaki, A.K., Holmans, P., Yang, L., Cleren, C., Beal, M.F., Jones, L., Kooperberg, C., Olson, J.M. and Jones, K.R. (2007) Expression profiling of Huntington's disease models suggests that brain-derived neurotrophic factor depletion plays a major role in striatal degeneration. *J Neurosci* **27**: 11758-11768.
- Strehlow, A.N.T., Li, J.Z. and Myers, R.M. (2007) Wild-type huntingtin participates in protein trafficking between the Golgi and the extracellular space. *Hum Mol Genet* **16**: 391-409.
- Subramaniam, S., Sixt, K.M., Barrow, R. and Snyder, S.H. (2009) Rhes, a striatal specific protein, mediates mutant-huntingtin cytotoxicity. *Science* **324**: 1327-1330.
- Suhr, S.T., Senut, M.C., Whitelegge, J.P., Faull, K.F., Cuizon, D.B. and Gage, F.H. (2001) Identities of sequestered proteins in aggregates from cells with induced polyglutamine expression. *J Cell Biol* **153**: 283-294.
- Szebenyi, G., Morfini, G.A., Babcock, A., Gould, M., Selkoe, K., Stenoien, D.L., Young, M., Faber, P.W., MacDonald, M.E., McPhaul, M.J. and Brady, S.T. (2003) Neuropathogenic forms of huntingtin and androgen receptor inhibit fast axonal transport. *Neuron* **40**: 41-52.
- Takiyama, Y., Oyanagi, S., Kawashima, S., Sakamoto, H., Saito, K., Yoshida, M., Tsuji, S., Mizuno, Y. and Nishizawa, M. (1994) A clinical and pathologic study of a large Japanese family with Machado-Joseph disease tightly linked to the DNA markers on chromosome 14q. *Neurology* **44**: 1302-1308.
- Tank, E.M.H. and True, H.L. (2009) Disease-associated mutant ubiquitin causes proteasomal impairment and enhances the toxicity of protein aggregates. *PLoS Genet* **5**: e1000382.
- Taylor, J.P., Tanaka, F., Robitschek, J., Sandoval, C.M., Taye, A., Markovic-Plese, S. and Fischbeck, K.H. (2003) Aggresomes protect cells by enhancing the degradation of toxic polyglutamine-containing protein. *Hum Mol Genet* **12**: 749-757.
- Terashima, T., Kawai, H., Fujitani, M., Maeda, K. and Yasuda, H. (2002) SUMO-1 co-localized with mutant atrophin-1 with expanded polyglutamines accelerates intranuclear aggregation and cell death. *Neuroreport* **13**: 2359-2364.
- Terry, R.D. and Katzman, R. (2001) Life span and synapses: Will there be a primary senile dementia? *Neurobiol Aging* **22**: 347-348.
- Thomas, E.A., Coppola, G., Desplats, P.A., Tang, B., Soragni, E., Burnett, R., Gao, F., Fitzgerald, K.M., Borok, J.F., Herman, D., Geschwind, D.H. and Gottesfeld, J.M. (2008) The HDAC inhibitor 4b ameliorates the disease phenotype and transcriptional abnormalities in Huntington's disease transgenic mice. *Proc Natl Acad Sci U S A* **105**: 15564-15569.
- Thrower, J.S., Hoffman, L., Rechsteiner, M. and Pickart, C.M. (2000) Recognition of the polyubiquitin proteolytic signal. *EMBO J* **19**: 94-102.
- Trottier, Y., Lutz, Y., Stevanin, G., Imbert, G., Devys, D., Cancel, G., Saudou, F., Weber, C., David, G., Tora, L. and et al. (1995) Polyglutamine expansion as a pathological epitope in Huntington's disease and four dominant cerebellar ataxias. *Nature* **378**: 403-406.

- Tsai, Y.C., Fishman, P.S., Thakor, N.V. and Oyler, G.A. (2003) Parkin facilitates the elimination of expanded polyglutamine proteins and leads to preservation of proteasome function. *J Biol Chem* **278**: 22044-22055.
- Twelvetrees, A.E., Yuen, E.Y., Arancibia-Carcamo, I.L., MacAskill, A.F., Rostaing, P., Lumb, M.J., Humbert, S., Triller, A., Saudou, F., Yan, Z. and Kittler, J.T. (2010) Delivery of GABAARs to synapses is mediated by HAP1-KIF5 and disrupted by mutant huntingtin. *Neuron* **65**: 53-65.
- Valtin, H. (1982) The discovery of the Brattleboro rat, recommended nomenclature, and the question of proper control. In: Sokol H, Valtin H (eds.) *The Brattleboro rat*. New York Academy of Sciences, pp. 1-9.
- Van den Hurk, W.H., Willems, H.J.J., Bloemen, M. and Martens, G.J.M. (2001) Novel frameshift mutations near short simple repeats. *J Biol Chem* **276**: 11496-11498.
- Van Leeuwen, F., van der Beek, E., Seger, M., Burbach, P. and Ivell, R. (1989) Age-related development of a heterozygous phenotype in solitary neurons of the homozygous Brattleboro rat. *Proc Natl Acad Sci U S A* **86**: 6417-6420.
- Van Leeuwen, F.W., de Kleijn, D.P., van den Hurk, H.H., Neubauer, A., Sonnemans, M.A., Sluijs, J.A., Koycu, S., Ramdjielal, R.D., Salehi, A., Martens, G.J., Grosveld, F.G., Peter, J., Burbach, H. and Hol, E.M. (1998) Frameshift mutants of β -amyloid precursor protein and ubiquitin-B in Alzheimer's and Down patients. *Science* **279**: 242-247.
- Van Leeuwen, F.W., Hol, E.M., Hermanussen, R.W., Sonnemans, M.A., Moraal, E., Fischer, D.F., Evans, D.A., Chooi, K.F., Burbach, J.P. and Murphy, D. (2000) Molecular misreading in non-neuronal cells. *FASEB J* **14**: 1595-1602.
- Van Nocker, S., Sadis, S., Rubin, D., Glickman, M., Fu, H., Coux, O., Wefes, I., Finley, D. and Vierstra, R. (1996) The multiubiquitin-chain-binding protein Mcb1 is a component of the 26S proteasome in *Saccharomyces cerevisiae* and plays a nonessential, substrate-specific role in protein turnover. *Mol Cell Biol* **16**: 6020-6028.
- Van Raamsdonk, J.M., Murphy, Z., Slow, E.J., Leavitt, B.R. and Hayden, M.R. (2005) Selective degeneration and nuclear localization of mutant huntingtin in the YAC128 mouse model of Huntington disease. *Hum Mol Genet* **14**: 3823-3835.
- Van Tijn, P., de Vrij, F.M.S., Schuurman, K.G., Dantuma, N.P., Fischer, D.F., van Leeuwen, F.W. and Hol, E.M. (2007) Dose-dependent inhibition of proteasome activity by a mutant ubiquitin associated with neurodegenerative disease. *J Cell Sci* **120**: 1615-1623.
- Van Wijk, S.J.L., de Vries, S.J., Kemmeren, P., Huang, A., Boelens, R., Bonvin, A.M.J.J. and Timmers, H.T.M. (2009) A comprehensive framework of E2-RING E3 interactions of the human ubiquitin-proteasome system. *Mol Syst Biol* **5**: 1-16.
- Venkatraman, P., Wetzel, R., Tanaka, M., Nukina, N. and Goldberg, A.L. (2004) Eukaryotic proteasomes cannot digest polyglutamine sequences and release them during degradation of polyglutamine-containing proteins. *Mol Cell* **14**: 95-104.
- Verhoef, L.G., Lindsten, K., Masucci, M.G. and Dantuma, N.P. (2002) Aggregate formation inhibits proteasomal degradation of polyglutamine proteins. *Hum Mol Genet* **11**: 2689-2700.
- Verhoef, L.G.G.C., Heinen, C., Selivanova, A., Halff, E.F., Salomons, F.A. and Dantuma, N.P. (2009) Minimal length requirement for proteasomal degradation of ubiquitin-dependent substrates. *FASEB J* **23**: 123-133.

References

- Verma, R., Oania, R., Graumann, J. and Deshaies, R.J. (2004) Multiubiquitin chain receptors define a layer of substrate selectivity in the ubiquitin-proteasome system. *Cell* **118**: 99-110.
- Vogel, G. (1998) Possible new cause of Alzheimer's disease found. *Science* **279**: 174.
- von Horsten, S., Schmitt, I., Nguyen, H.P., Holzmann, C., Schmidt, T., Walther, T., Bader, M., Pabst, R., Kobbe, P., Krotova, J., Stiller, D., Kask, A., Vaermann, A., Rathke-Hartlieb, S., Schulz, J.B., Grasshoff, U., Bauer, I., Vieira-Saecker, A.M.M., Paul, M., Jones, L., Lindenberg, K.S., Landwehrmeyer, B., Bauer, A., Li, X.J. and Riess, O. (2003) Transgenic rat model of Huntington's disease. *Hum Mol Genet* **12**: 617-624.
- Vonsattel, J.P., Myers, R.H., Stevens, T.J., Ferrante, R.J., Bird, E.D. and Richardson, E.P., Jr. (1985) Neuropathological classification of Huntington's disease. *J Neuropathol Exp Neurol* **44**: 559-577.
- Wacker, J.L., Huang, S.Y., Steele, A.D., Aron, R., Lotz, G.P., Nguyen, Q., Giorgini, F., Roberson, E.D., Lindquist, S., Masliah, E. and Muchowski, P.J. (2009) Loss of Hsp70 exacerbates pathogenesis but not levels of fibrillar aggregates in a mouse model of Huntington's disease. *J Neurosci* **29**: 9104-9114.
- Wacker, J.L., Zareie, M.H., Fong, H., Sarikaya, M. and Muchowski, P.J. (2004) Hsp70 and Hsp40 attenuate formation of spherical and annular polyglutamine oligomers by partitioning monomer. *Nat Struct Mol Biol* **11**: 1215-1222.
- Warrick, J.M., Morabito, L.M., Bilen, J., Gordesky-Gold, B., Faust, L.Z., Paulson, H.L. and Bonini, N.M. (2005) Ataxin-3 suppresses polyglutamine neurodegeneration in Drosophila by a ubiquitin-associated mechanism. *Mol Cell* **18**: 37-48.
- Wellington, C.L., Ellerby, L.M., Gutekunst, C.A., Rogers, D., Warby, S., Graham, R.K., Loubser, O., van Raamsdonk, J., Singaraja, R., Yang, Y.Z., Gafni, J., Bredesen, D., Hersch, S.M., Leavitt, B.R., Roy, S., Nicholson, D.W. and Hayden, M.R. (2002) Caspase cleavage of mutant huntingtin precedes neurodegeneration in Huntington's disease. *J Neurosci* **22**: 7862-7872.
- Wexler, N.S., Lorimer, J., Porter, J., Gomez, F., Moskowitz, C., Shackell, E., Marder, K., Penchaszadeh, G., Roberts, S.A., Gayan, J., Brocklebank, D., Cherny, S.S., Cardon, L.R., Gray, J., Dlouhy, S.R., Wiktorowski, S., Hodes, M.E., Conneally, P.M., Penney, J.B., Gusella, J., Cha, J.H., Irizarry, M., Rosas, D., Hersch, S., Hollingsworth, Z., MacDonald, M., Young, A.B., Andresen, J.M., Housman, D.E., De Young, M.M., Bonilla, E., Stillings, T., Negrette, A., Snodgrass, S.R., Martinez-Jaurrieta, M.D., Ramos-Arroyo, M.A., Bickham, J., Ramos, J.S., Marshall, F., Shoulson, I., Rey, G.J., Feigin, A., Arnheim, N., Acevedo-Cruz, A., Acosta, L., Alvir, J., Fischbeck, K., Thompson, L.M., Young, A., Dure, L., O'Brien, C.J., Paulsen, J., Brickman, A., Krch, D., Peery, S., Hogarth, P., Higgins, D.S., Jr. and Landwehrmeyer, B. (2004) Venezuelan kindreds reveal that genetic and environmental factors modulate Huntington's disease age of onset. *Proc Natl Acad Sci U S A* **101**: 3498-3503.
- Weydt, P., Pineda, V.V., Torrence, A.E., Libby, R.T., Satterfield, T.F., Lazarowski, E.R., Gilbert, M.L., Morton, G.J., Bammmer, T.K., Strand, A.D., Cui, L., Beyer, R.P., Easley, C.N., Smith, A.C., Krainc, D., Luquet, S., Sweet, I.R., Schwartz, M.W. and La Spada, A.R. (2006) Thermoregulatory and metabolic defects in Huntington's disease transgenic mice implicate PGC-1 α in Huntington's disease neurodegeneration. *Cell Metab* **4**: 349-362.
- Wilbur, J.D., Chen, C.Y., Manalo, V., Hwang, P.K., Fletterick, R.J. and Brodsky, F.M. (2008) Actin binding by Hip1 (huntingtin-interacting protein 1) and Hip1R (Hip1-related protein) is regulated by clathrin light chain. *J Biol Chem* **283**: 32870-32879.

- Willingham, S., Outeiro, T.F., DeVit, M.J., Lindquist, S.L. and Muchowski, P.J. (2003) Yeast genes that enhance the toxicity of a mutant huntingtin fragment or α -synuclein. *Science* **302**: 1769-1772.
- Wilson, S.M., Bhattacharyya, B., Rachel, R.A., Coppola, V., Tessarollo, L., Householder, D.B., Fletcher, C.F., Miller, R.J., Copeland, N.G. and Jenkins, N.A. (2002) Synaptic defects in ataxia mice result from a mutation in Usp14, encoding a ubiquitin-specific protease. *Nat Genet* **32**: 420-425.
- Wu, S.S., de Chadarevian, J.P., McPhaul, L., Riley, N.E., van Leeuwen, F.W. and French, S.W. (2002) Coexpression and accumulation of ubiquitin +1 and ZZ proteins in livers of children with $\alpha 1$ -antitrypsin deficiency. *Pediatr Dev Pathol* **5**: 293-298.
- Wyttensbach, A., Swartz, J., Kita, H., Thykjaer, T., Carmichael, J., Bradley, J., Brown, R., Maxwell, M., Schapira, A., Orntoft, T.F., Kato, K. and Rubinsztein, D.C. (2001) Polyglutamine expansions cause decreased CRE-mediated transcription and early gene expression changes prior to cell death in an inducible cell model of Huntington's disease. *Hum Mol Genet* **10**: 1829-1845.
- Xu, P., Duong, D.M., Seyfried, N.T., Cheng, D., Xie, Y., Robert, J., Rush, J., Hochstrasser, M., Finley, D. and Peng, J. (2009) Quantitative proteomics reveals the function of unconventional ubiquitin chains in proteasomal degradation. *Cell* **137**: 133-145.
- Yang, H., Zhong, X., Ballar, P., Luo, S., Shen, Y., Rubinsztein, D.C., Monteiro, M.J. and Fang, S. (2007) Ubiquitin ligase Hrd1 enhances the degradation and suppresses the toxicity of polyglutamine-expanded huntingtin. *Exp Cell Res* **313**: 538-550.
- Yang, S.H., Cheng, P.H., Banta, H., Piotrowska-Nitsche, K., Yang, J.J., Cheng, E.C.H., Snyder, B., Larkin, K., Liu, J., Orkin, J., Fang, Z.H., Smith, Y., Bachevalier, J., Zola, S.M., Li, S.H., Li, X.J. and Chan, A.W.S. (2008) Towards a transgenic model of Huntington's disease in a non-human primate. *Nature* **453**: 921-924.
- Yoshizawa, T., Yoshida, H. and Shoji, S. (2001) Differential susceptibility of cultured cell lines to aggregate formation and cell death produced by the truncated Machado-Joseph disease gene product with an expanded polyglutamine stretch. *Brain Res Bull* **56**: 349-352.
- Yu, Z.X., Li, S.H., Nguyen, H.P. and Li, X.J. (2002) Huntingtin inclusions do not deplete polyglutamine-containing transcription factors in HD mice. *Hum Mol Genet* **11**: 905-914.
- Zala, D., Colin, E., Rangone, H.l.n., Liot, G.r., Humbert, S. and Saudou, F.d.r. (2008) Phosphorylation of mutant huntingtin at S421 restores anterograde and retrograde transport in neurons. *Hum Mol Genet* **17**: 3837-3846.
- Zeitlin, S., Liu, J.P., Chapman, D.L., Papaioannou, V.E. and Efstratiadis, A. (1995) Increased apoptosis and early embryonic lethality in mice nullizygous for the Huntington's disease gene homologue. *Nat Genet* **11**: 155-163.
- Zhang, S., Binari, R., Zhou, R. and Perrimon, N. (2010) A genomewide RNA interference screen for modifiers of aggregates formation by mutant huntingtin in Drosophila. *Genetics* **184**: 1165-1179.
- Zhang, S., Feany, M.B., Saraswati, S., Littleton, J.T. and Perrimon, N. (2009) Inactivation of Drosophila huntingtin affects long-term adult functioning and the pathogenesis of a Huntington's disease model. *Dis Model Mech* **2**: 247-266.
- Zhang, Y., Engelman, J. and Friedlander, R.M. (2009) Allele-specific silencing of mutant Huntington's disease gene. *J Neurochem* **108**: 82-90.

References

- Zhang, Y., Leavitt, B.R., van Raamsdonk, J.M., Dragatsis, I., Goldowitz, D., MacDonald, M.E., Hayden, M.R. and Friedlander, R.M. (2006) Huntington inhibits caspase-3 activation. *EMBO J* **25**: 5896-5906.
- Zhou, H., Cao, F., Wang, Z., Yu, Z.X., Nguyen, H.P., Evans, J., Li, S.H. and Li, X.J. (2003) Huntingtin forms toxic N-terminal fragment complexes that are promoted by the age-dependent decrease in proteasome activity. *J Cell Biol* **163**: 109-118.
- Zhuchenko, O., Bailey, J., Bonnen, P., Ashizawa, T., Stockton, D.W., Amos, C., Dobyns, W.B., Subramony, S.H., Zoghbi, H.Y. and Lee, C.C. (1997) Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the α 1A-voltage-dependent calcium channel. *Nat Genet* **15**: 62-69.
- Zoghbi, H.Y. and Orr, H.T. (2000) Glutamine repeats and neurodegeneration. *Annu Rev Neurosci* **23**: 217-247.
- Zouambia, M., Fischer, D.F., Hobo, B., De Vos, R.A.I., Hol, E.M., Varndell, I.M., Sheppard, P.W. and Van Leeuwen, F.W. (2008) Proteasome subunit proteins and neuropathology in tauopathies and synucleinopathies: Consequences for proteomic analyses. *Proteomics* **8**: 1221-1236.
- Zuccato, C., Tartari, M., Crotti, A., Goffredo, D., Valenza, M., Conti, L., Cataudella, T., Leavitt, B.R., Hayden, M.R., Timmus, T., Rigamonti, D. and Cattaneo, E. (2003) Huntington interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes. *Nat Genet* **35**: 76-83.