



Universiteit  
Leiden  
The Netherlands

## Cardiac resynchronization therapy : determinants of patient outcome and emerging indications

Bommel, R.J. van

### Citation

Bommel, R. J. van. (2012, May 31). *Cardiac resynchronization therapy : determinants of patient outcome and emerging indications*. Retrieved from <https://hdl.handle.net/1887/19036>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/19036>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/19036> holds various files of this Leiden University dissertation.

**Author:** Bommel, Rutger Jan van

**Title:** Cardiac resynchronization therapy : determinants of patient outcome and emerging indications

**Issue Date:** 2012-05-31

# Chapter 17

## Diverse patterns of longitudinal and radial dyssynchrony in patients with advanced systolic heart failure

Zhang Q, **van Bommel RJ**, Chan YS, Delgado V, Liang Y, Schalij MJ, Bax JJ, Fang F, Wai-Kwok Yip G, Yu CM

*Heart* 2011;97(7):574-8

## ABSTRACT

**Background:** Little was known about the impact of QRS duration and etiology of heart failure on the pattern of left ventricular long- and short-axis dyssynchrony. This was investigated in the current study by using tissue Doppler imaging (TDI) and two-dimensional (2D) speckle-tracking imaging.

**Methods:** 448 heart failure patients ( $65 \pm 12$  years, 75% males) with ejection fraction  $\leq 35\%$  from the 2 cardiac centers were examined for the occurrence of longitudinal dyssynchrony by TDI and radial dyssynchrony by 2D speckle-tracking imaging. Region(s) of the latest mechanical contraction were also determined.

**Results:** Longitudinal dyssynchrony was identified in 263 (59%) patients while radial dyssynchrony in 186 (42%). There were 125 (28%) patients who had both longitudinal and radial dyssynchrony, 138 (31%) had only longitudinal, 61 (13%) had only radial, and 124 (28%) had neither form of dyssynchrony. By using TDI, the single most delayed segment was the septal, lateral, anterior, inferior, anteroseptal and posterior wall in 12%, 27%, 12%, 19%, 7% and 13% of patients while multi-segmental delay occurred in 10% of patients. These figures were 10%, 8%, 5%, 10%, 12%, 14% and 41% by using 2D speckle-tracking. When compared between patients with wide and narrow QRS complexes, both longitudinal (63% vs. 53%) and radial (49% vs. 36%) dyssynchrony parameters were slightly more frequently positive in the wide QRS group defined by QRS duration  $\geq 120$  ms (both  $p < 0.05$ ). When compared between non-ischemic and ischemic patients, the prevalence of longitudinal dyssynchrony was comparable (61% vs. 57%,  $p = 0.467$ ), while radial dyssynchrony was marginally more common in the non-ischemic group (47% vs. 37%,  $p = 0.049$ ). However, the distributions of the most delayed segment between the sub-groups were highly similar when assessed by the same echocardiographic method.

**Conclusions:** In patients with advanced systolic heart failure, the patterns of longitudinal and radial dyssynchrony are heterogeneous, and mechanical dyssynchrony is slightly more prevalent in the wide QRS group or the non-ischemic group.

## INTRODUCTION

Systolic dyssynchrony has been recognized as a common mechanical phenomenon in patients with advanced heart failure. It not only occurs in heart failure with electrical conduction delay as evidenced by a wide QRS complex, but also exists in those with a normal QRS duration.<sup>1,2</sup> With the development of advanced and quantitative echocardiographic imaging techniques, mechanical dyssynchrony could further be investigated for different components according to the axis of left ventricular (LV) motion. Among various echocardiographic techniques, tissue Doppler imaging (TDI) has been validated for assessing the timing of regional myocardial motion along the LV longitudinal axis, while two-dimensional (2D) speckle-tracking imaging has been adopted for measuring radial axis dyssynchrony.<sup>3,4,5</sup>

Cardiac resynchronization therapy (CRT) is an established device therapy for patients with refractory heart failure who have electromechanical delay.<sup>6,7,8,9</sup> While the QRS duration on surface ECG failed to predict a favorable response, the assessment of dyssynchrony by echocardiographic techniques showed a close relationship with improvement after CRT.<sup>10,11,12,13</sup> Furthermore, it has recently been suggested that identifying the location of the latest mechanical activation might provide insight into the optimal location of the LV pacing lead, apart from deploying it at the postero-lateral wall.<sup>14,15</sup> Therefore, in the current study, we examined the occurrence of longitudinal and/or radial dyssynchrony and the pattern of the latest contraction in heart failure patients with LV ejection fraction (LVEF)  $\leq 35\%$  despite optimal medical therapy.

## METHODS

### Patients

From Jan, 2006 to Jan, 2009, patients who presented congestive heart failure with an LVEF  $\leq 35\%$  despite optimal medical therapy were recruited from 2 cardiac centers (CUHK and LUMC). Patients were excluded from the study if they had a previous pacemaker implantation and/or chronic atrial fibrillation. Ischemic etiology of heart failure was confirmed by a history of myocardial infarction and/or a positive result by coronary angiogram. The study protocol was approved by the ethic committees of the 2 institutions and informed consent was obtained from each participant.

Comprehensive echocardiography with two-dimensional (2D), Doppler and TDI was performed in each patient (Vivid 7, Vingmed-General Electric, Horten, Norway). At least 3 consecutive beats were stored and the images were analyzed offline with the aid of a customized software package (EchoPac PC, version 7.0.0; Vingmed-General Electric, Horten, Norway). LV

end-diastolic volume (LVEDV), LV end-systolic volume (LVESV) and LVEF were calculated from the apical 4- and 2-chamber views, using the biplane Simpson's method.<sup>16</sup>

Color TDI images were acquired in the apical 4-chamber, 2-chamber and 3-chamber views for assessing LV longitudinal motion. The myocardial velocity curve was reconstituted offline in each segment of the 6-basal, 6-mid LV model, where the time to peak systolic velocity during ejection phase (Ts) was measured.<sup>17</sup> Longitudinal systolic dyssynchrony was calculated as the standard deviation of Ts among the 12 LV segments (Ts-SD) and a cut-off value of  $\geq 33$ ms was adopted as previously described.<sup>18</sup> The segment(s) with the latest contraction was determined by the longest Ts.

Gray-scale 2D images at the mid-cavity level of LV short-axis view were analyzed for radial deformation by using 2D speckle-tracking technology. The myocardial radial strain curves were reconstituted in each of the 6 mid-LV segments from which the time to maximal radial strain was measured. Radial dyssynchrony was calculated as the time difference between the antero-septal and posterior segment (antero-septal-to-posterior delay) and a cut-off of  $\geq 130$  ms was adopted to identify significant dyssynchrony.<sup>18,19</sup> The most delayed segment(s) was that with the longest time to maximal radial strain.

## Statistical analysis

Statistical analysis was performed using SPSS 13.0 (SPSS, Inc., Chicago, Illinois, USA). Parametric variables are expressed as mean $\pm$ SD while categorical variables are expressed as numbers and percentages. The paired sample t-test or McNemar test was adopted when appropriate in comparisons between longitudinal and radial dyssynchrony. The distributions of most delayed segment between longitudinal and radial dyssynchrony were compared by Chi-square test. The Chi-square test or unpaired sample t-test was used when appropriate to compare between sub-groups, including wide vs. narrow QRS and non-ischemic vs. ischemic heart failure. A p-value  $< 0.05$  was considered statistically significant.

## RESULTS

### Demographics of the study population

A total of 448 patients were recruited in this observational study. All patients were put on optimal medical therapy for heart failure which included diuretics, angiotensin converting enzyme inhibitors or angiotensin receptor blockers,  $\beta$ -blockers and Spironolactone. Table 1 shows baseline characteristics of the study population.


**Table 1.** Baseline characteristics of the study population (n = 448)

|                                      |                               |
|--------------------------------------|-------------------------------|
| Age                                  | 65 ± 12years                  |
| Gender (male)                        | 75%                           |
| Heart rhythm (sinus)                 | 100%                          |
| NYHA class, II / III / IV            | 27 (6%) / 403 (90%) / 18 (4%) |
| Etiology of heart failure (ischemic) | 250 (56%)                     |
| QRS duration ≥120 ms                 | 241 (54%)                     |
| Left bundle branch block             | 174 (39%)                     |
| Right bundle branch block            | 30 (7%)                       |
| Intraventricular conduction delay    | 37 (8%)                       |
| LV end-diastolic volume (ml)         | 191 ± 71                      |
| LV end-systolic volume (ml)          | 141 ± 60                      |
| LV ejection fraction (%)             | 27 ± 7                        |

### Prevalence and pattern of systolic dyssynchrony in heart failure

By using TDI assessment, the mean Ts-SD of the study population was 37.7±16.9 ms and significant longitudinal dyssynchrony occurred in 263 (59%) patients. In contrast, by using 2D speckle-tracking imaging, the mean antero-septal-to-posterior delay was 136±115 ms and significant radial dyssynchrony occurred in 186 (42%) patients. Therefore, longitudinal dyssynchrony parameter was significantly more frequently positive than radial dyssynchrony parameter ( $\chi^2 = 29.4$ ,  $p < 0.001$ ). Furthermore, there were 125 (28%) patients who had both longitudinal and radial dyssynchrony, 138 (31%) patients had only longitudinal dyssynchrony, 61 (13%) patients had only radial dyssynchrony, and 124 (28%) patients had neither form of dyssynchrony. The occurrence of longitudinal and radial dyssynchrony is concordant in 249 (56%) of the study patients.

By using TDI, the single most delayed segment was found in the septum for 52 (12%) patients, in the lateral wall for 120 (27%) patients, in the anterior wall for 55 (12%) patients, in the inferior wall for 85 (19%) patients, in the antero-septum for 30 (7%), and in the posterior wall for 60 (13%) patients. Forty-six (10%) patients had multi-segmental delay. Conversely, by using 2D speckle-tracking imaging, these figures were 45 (10%), 36 (8%), 22 (5%), 45 (10%), 54 (12%), 63 (14%) and 183 (41%) patients, respectively (Figure 1). A concordance in the most delayed segment between TDI and 2D speckle-tracking methods was found in 134 (30%) patients.



**Figure 1.** Distributions of the most delayed segment detected by tissue Doppler imaging (—●—) and two-dimensional speckle-tracking imaging (---■---) in the whole study population. S = septum; AS = antero-septum; A = anterior segment; L = lateral segment; P = posterior segment; I = inferior segment, Multi = multi-segmental delay.

### Pattern of systolic dyssynchrony in heart failure with wide vs. narrow QRS complexes

When comparing between patients with wide and narrow QRS complexes, the wide QRS group had a larger LVEDV ( $198 \pm 79$  ml vs.  $182 \pm 60$  ml,  $p = 0.014$ ), LVESV ( $148 \pm 68$  ml vs.  $132 \pm 50$  ml,  $p = 0.005$ ) and a lower LVEF ( $26.4 \pm 7.6\%$  vs.  $28.1 \pm 7.1\%$ ,  $p = 0.019$ ). The wide QRS group also had a more prolonged dyssynchrony indices of TDI ( $T_s\text{-SD}$ :  $39.2 \pm 16.0$  ms vs.  $36.0 \pm 17.8$  ms,  $p = 0.044$ ) and 2D speckle-tracking (anteroseptal-to-posterior delay:  $151 \pm 122$  ms vs.  $118 \pm 104$  ms,  $p = 0.003$ ) methods. The difference in the occurrence of positive longitudinal dyssynchrony parameter in the wide QRS group, when compared to narrow QRS group, was marginally significant (63% vs. 53%,  $p = 0.043$ ). Similarly, radial dyssynchrony parameter was slightly more frequently positive in wide QRS group (49% vs. 36%,  $p = 0.039$ ).

In the wide QRS group, the most delayed segment using TDI was the septum in 30 (12%) patients, the lateral wall in 67 (28%), the anterior wall in 27 (11%), the inferior wall in 50 (21%), the antero-septum in 17 (7%), and the posterior wall in 34 (14%). Multi-segmental delay was found in 16 (7%) patients. These figures were similar in the narrow QRS group, which were 22 (11%), 53 (26%), 28 (14%), 35 (17%), 13 (6%), 26 (12%) and 30 (14%) patients, respectively ( $p = 0.173$ ). Similar distributions between the wide and narrow QRS groups were also found using 2D speckle-tracking ( $p = 0.511$ ) (Figure 2A&2B).

Figure 2A





Figure 2B



**Figure 2.** Distributions of the most delayed segment detected by tissue Doppler imaging (A) and two-dimensional speckle-tracking imaging (B) in the wide QRS (solid line) and narrow QRS (dashed line) sub-groups.  
S = septum; AS = anteroseptum; A = anterior segment; L = lateral segment; P = posterior segment; I = inferior segment, Multi = multi-segmental delay.

### Pattern of systolic dyssynchrony in ischemic vs. non-ischemic patients

When comparing between non-ischemic and ischemic patients, there was no significant difference in the LVEDV ( $194 \pm 74$  ml vs.  $182 \pm 67$  ml,  $p = 0.391$ ), LVESV ( $144 \pm 65$  ml vs.  $138 \pm 57$  ml,  $p = 0.272$ ) and LVEF ( $26.6 \pm 7.6\%$  vs.  $27.6 \pm 7.2\%$ ,  $p = 0.637$ ). A wide QRS complex was found in 111 (56%) patients in the non-ischemic group and in 130 (52%) patients in the ischemic group ( $\chi^2 = 0.733$ ,  $p = 0.392$ ). Although dyssynchrony index by TDI was not different between the 2 groups ( $38.3 \pm 17.0$  ms vs.  $37.2 \pm 17.0$  ms,  $p = 0.504$ ), the anteroseptal-to-posterior delay by 2D speckle-tracking imaging was more prolonged in non-ischemic group ( $153 \pm 125$  ms vs.  $122 \pm 104$  ms,  $p = 0.005$ ). Longitudinal dyssynchrony by TDI was identified in 120 (61%) patients in the non-ischemic group and 143 patients (57%,  $p = 0.467$ ) in the ischemic group. In contrast, radial dyssynchrony by 2D speckle-tracking imaging was identified in 93 (47%) patients in the non-ischemic group and 92 patients (37%,  $p = 0.049$ ) in the ischemic group.

In the non-ischemic group, TDI identified the most delayed segment as the septum in 27 (14%), the lateral wall in 54 (27%), the anterior wall in 19 (10%), the inferior wall in 43 (21%), the anteroseptum in 10 (5%), and the posterior wall in 29 (15%) patients. Multi-segmental delay was observed in 16 (8%) patients. The figures were 25 (10%), 66 (26%), 36 (14%), 42 (19%), 20 (8%), 31 (12%) and 30 (11%) patients respectively in the ischemic group ( $p = 0.220$ ). Similarly, 2D speckle-tracking demonstrated no difference in the prevalence of delay in the septum, lateral wall, anterior wall, inferior wall, anteroseptum, posterior wall and multi-segmental delay between the non-ischemic (13%, 11%, 6%, 11%, 10%, 15% and 34%) and ischemic (8%, 5%, 4%, 9%, 14%, 14% and 46%) groups ( $p = 0.128$ ) (Figure 3A&3B).

Figure 3A





Figure 3B



**Figure 3.** Distributions of the most delayed segment detected by tissue Doppler imaging (A) and two-dimensional speckle-tracking imaging (B) in the non-ischemic (solid line) and ischemic (dashed line) sub-groups. S = septum; AS = anteroseptum; A = anterior segment; L = lateral segment; P = posterior segment; I = inferior segment, Multi = multi-segmental delay.

## DISCUSSION

In this study we show that longitudinal and radial dyssynchrony, as defined by two previously published and most commonly used parameters, was present in 59% and 42% of patients with heart failure and LV ejection fraction of  $\leq 35\%$ . However, the concordance of dyssynchrony, as defined by these two parameters, was modest and occurred in only slightly more than half of the patients. Interestingly, when the two methods were compared in their ability to detect single most delayed segment, the concordance occurred in about one-third of the patients. Of note, dyssynchrony parameters were positive in patients with both narrow and wide QRS complex, although this was slightly more frequent in the presence of wide QRS complex, a finding which is consistent with our previous observation.<sup>20</sup> Finally, heart failure etiology (ischemic vs. non-ischemic) only marginally affected dyssynchrony parameters or the detection and location of the most delayed segment.

### Longitudinal vs. radial dyssynchrony

The current study chose 2 previously validated parameters, namely Ts-SD by TDI and anteroseptal-to-posterior delay by 2D speckle-tracking, to investigate the occurrence of longitudinal and radial dyssynchrony simultaneously in a large number of heart failure patients.<sup>17,19</sup> TDI has been widely used for assessment of LV longitudinal dyssynchrony, which has shown a close agreement with cardiac magnetic resonance imaging (MRI).<sup>21</sup> Furthermore, a number of TDI dyssynchrony parameters have been suggested helpful in predicting favorable response to CRT.<sup>18</sup> 2D speckle-tracking imaging is an angle-independent tool where the derived radial dyssynchrony has been found being able to identify CRT responders.<sup>19</sup> The diverse pattern of longitudinal and radial dyssynchrony displayed in the current study indicates the complex nature of mechanical dyssynchrony in heart failure as cardiac motion *per se* is complex and contains various components of action. The use of different echocardiographic technologies which can track longitudinal and radial motions is likely to provide comprehensive and incremental information on the extent of LV dyssynchrony. Interestingly, the wide QRS group had a higher prevalence of both longitudinal and radial dyssynchrony than the narrow QRS group, which probably reflected the additional impact of electrical propagation abnormalities within the LV of the former group.<sup>22</sup> On the other hand, the lack of difference in the prevalence of longitudinal dyssynchrony between etiologies of heart failure corroborates the findings of previous studies in patients undergoing CRT. These studies demonstrated a similar response rate in terms of LV reverse remodeling at 3-month follow up between ischemic and non-ischemic heart failure patients.<sup>23</sup> Of note, in heart failure patients with a wide QRS complex, the simultaneous assessment of longitudinal dyssynchrony by TDI and radial dyssynchrony by speckle-tracking resulted in a higher CRT responder rate of 93% than that

of only assessing either parameter in which the responder rate was only 57%.<sup>24</sup> This study and our findings in the present study suggest that developing a robust scheme of dyssynchrony assessment which including different cardiac axes would provide complementary information and potentially assist future decision making.

### Diversity of the most delayed segments

In a few small-scale CRT studies, identification of the segment with the latest contraction by 2D speckle-tracking imaging was suggested to be helpful in identifying an optimal position for the LV pacing lead.<sup>15,25</sup> However, it remains controversial whether this echocardiography-guided implantation would result in a better response to CRT, when compared with conventional placement of the LV lead at the LV postero-lateral wall. The present study demonstrated a diverse pattern of the most delayed segments in both longitudinal and radial dyssynchrony, as illustrated in Figure 1. There was also a trend of more patients with postero-lateral wall delay by TDI (about 40% in all the sub-groups), while by 2D speckle-tracking imaging the trend was in favor of more patients with multi-segmental delay. Intriguingly, within each sub-group of wide and narrow QRS complex or ischemic and non-ischemic etiology, the patterns of delay by TDI or 2D speckle-tracking imaging were distinctively concordant, as illustrated in Figures 2 and 3. These findings could have been consistent with previous single-center observations that CRT resulted in a similar early volumetric responder rate for patients with wide and narrow QRS complexes who had a similar extent of systolic dyssynchrony,<sup>26,27</sup> although it was not demonstrated in the multi-center RethinQ study.<sup>28</sup>

### Study limitations

The current study was the first one of this kind to describe the prevalence and pattern of longitudinal and radial dyssynchrony simultaneously within a large heart failure population. However, it was not envisaged to explore the underlying pathophysiological processes leading to the development of different components of mechanical dyssynchrony, nor to demonstrate the correlation between longitudinal or radial dyssynchrony and prognosis in a cross-sectional observational study where no follow up data available. Meanwhile, the inclusion of much fewer patients in NYHA class II or IV made it impossible to correlate the echocardiographic findings with clinical parameters. Hopefully, our understanding of the complex relations between the different forms of cardiac dyssynchrony would be improved with the assistance of other advanced imaging techniques such as contrast-enhanced MRI, and with the data from prospective clinical trials.

## REFERENCES

1. Auricchio A, Yu CM. Beyond the measurement of QRS complex toward mechanical dyssynchrony: cardiac resynchronisation therapy in heart failure patients with a normal QRS duration. *Heart* 2004;90:479-481.
2. Cho GY, Song JK, Park WJ et al. Mechanical dyssynchrony assessed by tissue Doppler imaging is a powerful predictor of mortality in congestive heart failure with normal QRS duration. *J Am Coll Cardiol* 2005;46:2237-2243.
3. Delgado V, Ypenburg C, van Bommel RJ et al. Assessment of left ventricular dyssynchrony by speckle-tracking strain imaging comparison between longitudinal, circumferential, and radial strain in cardiac resynchronization therapy. *J Am Coll Cardiol* 2008;51:1944-1952.
4. Helle-Valle T, Crosby J, Edvardsen T et al. New noninvasive method for assessment of left ventricular rotation: speckle-tracking echocardiography. *Circulation* 2005;112:3149-3156.
5. Marwick TH, Yu CM, Sun JP. Myocardial imaging-tissue Doppler and speckle-tracking. First ed. Wiley-Blackwell, 2007.
6. St John Sutton MG, Plappert T, Abraham WT et al. Effect of cardiac resynchronization therapy on left ventricular size and function in chronic heart failure. *Circulation* 2003;107:1985-1990.
7. Auricchio A, Abraham WT. Cardiac resynchronization therapy: current state of the art: cost vs. benefit. *Circulation* 2004;109:300-307.
8. Cleland JG, Daubert JC, Erdmann E et al. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med* 2005;352:1539-1549.
9. Bax JJ, Abraham T, Barold SS et al. Cardiac resynchronization therapy: Part 1--issues before device implantation. *J Am Coll Cardiol* 2005;46:2153-2167.
10. Abraham WT, Fisher WG, Smith AL et al. Cardiac resynchronization in chronic heart failure. *N Engl J Med* 2002;346:1845-1853.
11. Ansalone G, Giannantoni P, Ricci R et al. Doppler myocardial imaging in patients with heart failure receiving biventricular pacing treatment. *Am Heart J* 2001;142:881-896.
12. Alonso C, Leclercq C, Victor F et al. Electrocardiographic predictive factors of long-term clinical improvement with multisite biventricular pacing in advanced heart failure. *Am J Cardiol* 1999;84:1417-1421.
13. Ricci R, Pignalberi C, Ansalone G et al. Early and late QRS morphology and width in biventricular pacing: relationship to lead site and electrical remodeling. *J Interv Card Electrophysiol* 2002;6:279-285.
14. Murphy RT, Sigurdsson G, Mulamalla S et al. Tissue synchronization imaging and optimal left ventricular pacing site in cardiac resynchronization therapy. *Am J Cardiol* 2006;97:1615-1621.
15. Ypenburg C, van Bommel RJ, Delgado V et al. Optimal left ventricular lead position predicts reverse remodeling and survival after cardiac resynchronization therapy. *J Am Coll Cardiol* 2008;52:1402-1409.
16. Schiller NB, Shah PM, Crawford M et al. Recommendations for quantitation of the left ventricle by two-dimensional echocardiography. American Society of Echocardiography Committee on Standards, Subcommittee on Quantitation of Two-Dimensional Echocardiograms. *J Am Soc Echocardiogr* 1989;2:358-367.
17. Yu CM, Zhang Q, Yip GW et al. Diastolic and systolic asynchrony in patients with diastolic heart failure: a common but ignored condition. *J Am Coll Cardiol* 2007;49:97-105.
18. Gorcsan J, III, Abraham T, Agler DA et al. Echocardiography for cardiac resynchronization therapy: recommendations for performance and reporting--a report from the American Society of Echo-

- cardiography Dyssynchrony Writing Group endorsed by the Heart Rhythm Society. *J Am Soc Echocardiogr* 2008;21:191-213.
19. Suffoletto MS, Dohi K, Cannesson M et al. Novel speckle-tracking radial strain from routine black-and-white echocardiographic images to quantify dyssynchrony and predict response to cardiac resynchronization therapy. *Circulation* 2006;113:960-968.
20. Yu CM, Lin H, Zhang Q et al. High prevalence of left ventricular systolic and diastolic asynchrony in patients with congestive heart failure and normal QRS duration. *Heart* 2003;89:54-60.
21. Westenberg JJ, Lamb HJ, van der Geest RJ et al. Assessment of left ventricular dyssynchrony in patients with conduction delay and idiopathic dilated cardiomyopathy: head-to-head comparison between tissue Doppler imaging and velocity-encoded magnetic resonance imaging. *J Am Coll Cardiol* 2006;47:2042-2048.
22. Fung JW, Yu CM, Yip G et al. Variable left ventricular activation pattern in patients with heart failure and left bundle branch block. *Heart* 2004;90:17-19.
23. Zhang Q, Fung JW, Chan JY et al. Difference in long-term clinical outcome after cardiac resynchronisation therapy between ischaemic and non-ischaemic aetiologies of heart failure. *Heart* 2009;95:113-118.
24. Gorcsan J, III, Tanabe M, Bleeker GB et al. Combined longitudinal and radial dyssynchrony predicts ventricular response after resynchronization therapy. *J Am Coll Cardiol* 2007;50:1476-1483.
25. Becker M, Franke A, Breithardt OE et al. Impact of Left Ventricular Lead Position on the Efficacy of Cardiac Resynchronization Therapy. A Two-Dimensional Strain Echocardiography Study. *Heart* 2007;93:1197-1203.
26. Bleeker GB, Holman ER, Steendijk P et al. Cardiac resynchronization therapy in patients with a narrow QRS complex. *J Am Coll Cardiol* 2006;48:2243-2250.
27. Yu CM, Chan YS, Zhang Q et al. Benefits of cardiac resynchronization therapy for heart failure patients with narrow QRS complexes and coexisting systolic asynchrony by echocardiography. *J Am Coll Cardiol* 2006;48:2251-2257.
28. Beshai JF, Grimm RA, Nagueh SF et al. Cardiac-resynchronization therapy in heart failure with narrow QRS complexes. *N Engl J Med* 2007;357:2461-2471.