



Universiteit
Leiden
The Netherlands

Glycosaminoglycans and diabetic nephropathy

Pijl, J.W. van der

Citation

Pijl, J. W. van der. (1999, May 27). *Glycosaminoglycans and diabetic nephropathy*. Retrieved from <https://hdl.handle.net/1887/567>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/567>

Note: To cite this publication please use the final published version (if applicable).

Contents

Chapter 1	Introduction	1
Chapter 2	Extracellular matrix in human diabetic nephropathy: reduced expression of heparan sulphate in skin basement membrane <i>Diabetologia</i> 41: 791-798, 1998	43
Chapter 3	Improvement of heparan sulfate expression in the skin basement membrane after kidney and pancreas-kidney transplantation <i>Submitted</i>	59
Chapter 4	Danaparoid sodium lowers proteinuria in diabetic nephropathy <i>J Am Soc Nephrol</i> 8: 456-462, 1997	69
Chapter 5	Effect of danaparoid sodium on hard exudates in diabetic retinopathy <i>Lancet</i> 350: 1743-1745, 1997	85
Chapter 6	Effect of danaparoid sodium on proteinuria, von Willebrand factor, and hard exudates in patients with diabetes mellitus type 2 <i>J Am Soc Nephrol</i>, 1999 in press	93
Chapter 7	Summary and general discussion	107
Chapter 8	Samenvatting Curriculum vitae Acknowledgements	123
