

Cover Page



Universiteit Leiden




The handle <http://hdl.handle.net/1887/35155> holds various files of this Leiden University dissertation.

Author: Hussaarts, Leonie

Title: Immune modulation by schistosomes : mechanisms of T helper 2 polarization and implications for metabolic disorders

Issue Date: 2015-09-10



**SCHISTOSOME-DERIVED OMEGA-1
DRIVES TH2 POLARIZATION BY
SUPPRESSING PROTEIN SYNTHESIS
FOLLOWING INTERNALIZATION BY
THE MANNOSE RECEPTOR**

Bart Everts, Leonie Husaarts,* Nicole N. Driessen,*
Moniek H.J. Meevissen, Gabriele Schramm,
Alwin J. van der Ham, Barbara van der Hoeven,
Thomas Scholzen, Sven Burgdorf, Markus Mohrs,
Edward J. Pearce, Cornelis H. Hokke, Helmut Haas,
Hermelijn H. Smits and Maria Yazdanbakhsh

*both authors contributed equally

© 2012 Everts et al. *Journal of Experimental Medicine*.
209:10 1753-1767. doi: 10.1084/jem.20111381

2

ABSTRACT

Omega-1, a glycosylated T2 ribonuclease (RNase) secreted by *Schistosoma mansoni* eggs and abundantly present in soluble egg antigen (SEA), has recently been shown to condition dendritic cells (DCs) to prime Th2 responses. However, the molecular mechanisms underlying this effect remain unknown. We show here by site-directed mutagenesis of omega-1 that both the glycosylation and the RNase activity are essential to condition DCs for Th2 polarization. Mechanistically, we demonstrate that omega-1 is bound and internalized via its glycans by the mannose receptor (MR) and subsequently impairs protein synthesis by degrading both ribosomal and messenger RNA. These studies reveal an unrecognized pathway, involving MR and interference with protein synthesis that conditions DCs for Th2 priming.

INTRODUCTION

Dendritic cells (DCs) play a central role in the development and maintenance of immune responses during infection, as they govern both the activation and polarization of adaptive T helper (Th) cells. Classically, upon recognition of invading pathogens, resting DCs undergo a process of activation, so-called maturation, that involves stable presentation of peptides in the context of major histocompatibility complex (MHC)-I and -II, up-regulation of co-stimulatory molecules, and production of polarizing cytokines, that collectively enable DCs to potently activate and direct CD4⁺ T cell responses (1).

This paradigm is largely based on observations of responses towards pathogens, like bacteria, viruses and fungi. These pathogens harbor pathogen-associated molecular patterns that lead to classic DC activation by engaging several classes of innate pattern recognition receptors, including the Toll-like receptors (TLRs). Binding of pathogen-associated molecular patterns to these receptors initiates signaling cascades that generally result in the conditioning of DCs for priming of Th1- or Th17-biased responses which are instrumental in combating prokaryotic and single cell eukaryotic pathogens (2). In contrast to this classical view of DC activation, components derived from parasitic helminths, when co cultured with DCs, fail to induce the traditional signs of DC maturation. However, although overt maturation is not observed, unlike immature DCs, helminth antigen-treated DCs are altered such that they prime Th2-polarized immune responses (3).

Despite this consistent picture, the pathways through which helminth antigens manipulate DC function and drive Th2 responses are still poorly understood (4). The majority of the studies have been conducted with a complex mixture of soluble egg antigens (SEA) from the trematode *Schistosoma mansoni*. SEA is regarded as one of the most potent helminth-derived antigenic extracts that instruct DCs to drive Th2 polarization (3;5). So far these studies have mainly suggested that carbohydrate structures play a role in DC modulation by SEA, given that chemical modification of glycans on proteins present in SEA is known to abolish their capacity to induce Th2 polarization (6). In this respect, another class of pattern recognition receptors expressed by DCs, the carbohydrate-binding C-type lectin receptors (CLRs), have been suggested to play a role in modulation of DC function by SEA (7). For instance, SEA contains carbohydrate structures, such as Lewis-x (Le^x), that can be recognized by DC SIGN (DC-specific intercellular adhesion molecule-3-grabbing non-integrin; 8-10). Engagement of this receptor by components from pathogens such as *Helicobacter pylori* has been shown to suppress IL-12 production and modulate TLR-induced DC activation and T cell polarization (8;11). In addition, more recently it has been shown that SEA can modulate cytokine responses through another CLR, dectin-2 (12). Finally, a number of studies have raised the possibility that TLRs are involved in SEA-mediated Th2 induction (13;14). However, direct evidence for involvement of specific receptors or downstream pathways in SEA-driven Th2 polarization has been missing.

The recent identification of omega-1, a glycosylated T2 RNase, as the major component in schistosome eggs that is responsible for conditioning DCs for Th2 polarization (15-17), has allowed us to dissect the molecular pathways involved in a precise manner. Through site-

directed mutagenesis we show that both the RNase activity and the glycosylation of omega-1 are essential for programming of DCs for Th2 induction. Furthermore, we provide evidence that MR is critical for omega-1-driven Th2 responses and that internalization via this receptor is needed for biological activity of omega-1, as it allows omega-1 to interfere with translation, by degrading rRNA and mRNA, and thereby to condition these cells to prime Th2 responses.

RESULTS

Omega-1 requires both its glycosylation and RNase activity to condition DCs for priming of Th2 responses

The RNase activity of omega-1 has been proposed to play a role in the conditioning of DCs to prime Th2 responses (16). However, this was based on a chemical inactivation of the RNase activity by diethylpyrocarbonate-treatment, which can result in off-target modification of histidines as well as other amino acids that could alter the function or structure of the protein (18). Therefore, we addressed the role of RNase activity in a more stringent and specific manner by creating a mutant of recombinant wildtype (WT) omega-1 lacking RNase activity by site-directed mutagenesis. Specifically, a histidine residue in its catalytic domain, known from other T2 RNases to be essential for the enzymatic activity (19), was replaced by phenylalanine (omega-1 H58F) (Fig. 1 A). Apart from RNase activity, glycosylation of omega-1 may also be important for its Th2-priming capacity, since chemical modification of glycans on proteins present in SEA is known to abolish the ability of SEA to induce Th2 polarization (6). Moreover, potentially Th2-polarizing Le^x glycan motifs have recently been described to be present in glycans on omega-1 (20). To address the role of glycosylation in Th2 priming by omega-1, a glycosylation mutant was generated by a single amino-acid replacement at each of the two N-linked glycosylation sites (omega-1-N71/176Q) (Fig. 1 A) (17;20). An RNase assay showed that the RNase mutant did not have any RNase activity, while the RNase activity of the glycosylation mutant was unaffected (Fig. 1 B). In addition, the banding patterns of recombinant WT omega-1 and the mutants on silver stained SDS-PAGE and anti-omega-1 Western blots were in line with the absence of carbohydrates on the glycosylation mutant as evident by a single band instead of the three glycoforms of the recombinant WT omega-1 and the RNase mutant (Fig. 1 C). With regard to the glycans present on recombinant WT omega-1 and the RNase mutant, mass spectrometric analysis of tryptic glycopeptides showed the presence of N-glycans on Asn₁₇₆ with the monosaccharide composition Hex₃HexNAc₆Fuc_{2/3} (Fig. 1 D). This composition is indicative of the presence of GalNAcβ1-4(Fucα1-3)GlcNAc (LDN-F) antennae, a glycan element previously found on a protein from HEK293 cells (21), the cell type in which recombinant omega-1 is expressed. LDN-F motifs frequently occur on helminth glycoproteins, and the characteristics of LDN-F with respect to binding to CLRs are similar to those of the Le^x element (9;10;20;22).

To assess the role of glycosylation and RNase activity in omega-1-driven Th2 polarization, a well-established *in vitro* culture system of human monocyte-derived DCs (moDCs) and naïve CD4⁺ T cells was used, which mimics *in vivo* DC-mediated Th cell polarization (1). Similar to




Figure 1. Generation and evaluation of glycosylation and RNase mutants of recombinant omega-1.

(A) The amino acid sequence of omega-1 (Acc.No. ABB73003.1) is shown in which the mutation sites are depicted. The two conserved amino acid sequence (CAS)-domains essential for catalytic activity are marked in grey and the two N-linked glycosylation sites are depicted in white boxes. (B) RNA from PBMCs was incubated for 1 h with the different omega-1 variants (500 ng/ml and 100ng/ml) and analyzed on a 2% agarose gel for breakdown. RNase A was used as a positive control. One of two experiments is shown. (C) The omega-1 mutants were run under non-reducing conditions by SDS-PAGE and silver stained. A Western Blot by staining with a specific anti-omega-1 monoclonal antibody was in line with the absence of glycosylation only on the omega-1 glycosylation mutant. (D) MALDI-TOF mass spectrum of glycopeptides from a tryptic digest of recombinant WT omega-1, covering the glycosylation site N176. Recombinant omega-1 was subjected to SDS-PAGE under reducing conditions and stained with Colloidal blue. Stained bands were excised, subjected to reduction and alkylation and digested with trypsin. The MALDI-TOF-MS spectrum derived from the upper band in the SDS-PAGE pattern is depicted. Signals ($[M+H]^+$) are labeled with monoisotopic masses. Composition of the glycan moieties are given in terms of hexose (H), N-acetylhexosamine (N) and fucose (F). Differences in fucose content are indicated by double-headed arrows. Signals that cannot be assigned to glycopeptides are marked with asterisks (*).

natural omega-1 (15), recombinant WT omega-1 consistently and significantly suppressed in a concentration dependent manner the lipopolysaccharide (LPS)-induced upregulation of the costimulatory molecule CD86 (Fig. 2 A), as well as the production of IL-12 p70 following CD40 ligation (Fig. 2 B), which is an important characteristic of Th2-priming DCs (3). However, both the glycosylation mutant and the RNase mutant failed to alter LPS-induced CD86 expression or IL-12 production of DCs at any of the concentrations tested. Importantly, in contrast to DCs primed with recombinant WT omega-1, those conditioned with either mutant did not prime a Th2 response (Fig. 2 C+D). Similar results were obtained with cultures in which DCs were conditioned by the omega-1 mutants in the absence of LPS (Fig. 2 E). These data show that the RNase activity and the glycosylation of omega-1 are both essential, but as single property not sufficient, for the induction of Th2 responses via DCs.

Omega-1 requires both its glycosylation and RNase activity to prime Th2 responses *in vivo*

To test whether the *in vivo* Th2-priming capacity of omega-1 is dependent on glycosylation and RNase activity, recombinant WT omega-1 or its mutants were administered to 4get/KN2 IL-4 dual-reporter mice (23). In these mice IL-4-competent cells are GFP⁺ and IL-4-producing cells additionally express huCD2, allowing the direct visualization of Th2 differentiation and IL-4 production. Following the s.c. injection of the antigens into the footpad, the draining popliteal lymph nodes (LNs) were harvested on day 7 and CD4⁺CD44^{high} effector T cells were analyzed for the expression of GFP and huCD2 directly ex vivo. Injection of SEA resulted in a significant increase of GFP⁺ and huCD2⁺ cells, reflecting the induction of Th2 differentiation and IL-4 production *in vivo* (Fig. 3). Importantly, while recombinant WT omega-1 induced a marked Th2 response and the production of IL-4, both mutants were significantly impaired to prime this response as evident from lower frequencies (Fig. 3 A and B) as well as total numbers of huCD2⁺ T cells (Fig. 3 C) in the draining LN. Taken together, these data show that the glycosylation and the RNase activity of omega-1 play a crucial role in Th2 polarization induced by omega-1 *in vivo*.

Omega-1 is internalized by DCs via the mannose receptor (MR)

To get a better understanding of how glycosylation is involved in omega-1-driven Th2 polarization, we tested whether recognition of omega-1 by human DCs was dependent on the glycans present on omega-1. While human DCs were capable of binding fluorescently-labeled recombinant WT omega-1 or the RNase mutant as determined by FACS analysis, DCs failed to bind the glycosylation mutant, demonstrating that glycans present on omega-1 are essential for recognition by DCs (Fig. 4 A). Given the importance of glycosylation of omega-1 for binding to DCs, we explored the involvement of carbohydrate-binding CLRs in the recognition and uptake of omega-1. While DCs readily bound fluorescently-labeled omega-1, binding of natural omega-1 was totally prevented when DCs were pre-incubated with the calcium-chelator EGTA, which abolishes CLR binding to carbohydrate ligands (Fig. 4 B). In contrast, treatment of DCs with EGTA after 1 h incubation with omega-1, could not reduce the fluorescent signal

of omega-1, indicating that by then all bound omega-1 had been internalized. This suggests that DCs recognize and rapidly internalize omega-1 in a CLR dependent manner. SEA has been reported to be recognized and endocytosed by human moDCs via the CLR DC-SIGN and MR (7) that have the capacity to bind fucose-residues such as those found in Le^x (9;10;24), a glycan motif present on natural omega-1 (20). To determine whether MR and DC-SIGN are involved in recognition and internalization of natural omega-1, DCs were pre-incubated with




Figure 2. The glycosylation and RNase activity of omega-1 are essential for conditioning human DCs to prime Th2 responses. (A) Human moDCs were pulsed for 48 h with increasing concentrations of the mutant variants of recombinant omega-1 in combination with LPS (100 ng/ml) as a maturation factor and surface expression of CD86 was determined by FACS analysis. The expression levels, based on geometric mean fluorescence, are shown relative to the DCs stimulated with LPS alone, which is set to 1. Data are based on two independent experiments and shown as mean \pm SD. (B) Conditioned DCs were co-cultured for 24 h with a CD40-L expressing cell line, to mimic the interaction with T cells. IL-12p70 cytokine expression levels are shown relative to the DCs stimulated with LPS alone, which is set to 1. Data are representative of triplicate wells from one of 2 independent experiments and shown as mean \pm SD. (C) DCs conditioned as described in (A) were cultured with allogeneic naive CD4⁺ T cells for 12 days in the presence of staphylococcal enterotoxin B and IL-2. Intracellular cytokine production was assayed by FACS 6 h after the stimulation of primed T cells with phorbol 12-myristate 13-acetate (PMA) and ionomycin. Based on intracellular cytokine staining, the ratio of T cells single-positive for either IL-4 or IFN- γ was calculated relative to the control condition. Data are based on 2 independent experiments and shown as mean \pm SD. (D) An example of T cell polarization assay as described in (C) induced by the different recombinant omega-1 variants. The frequencies of each population are indicated as percentages in the plot. One representative result from 5 independent experiments is shown. (E) T cell polarization assay as described in (C) but in the absence of LPS. Data are representative of 3 independent experiments. Bars represent mean \pm SD. # $P < 0.05$, for significant differences compared to control conditions (*) or between test conditions (#) based on paired analysis (two-sided paired t -test). ω -1, omega-1; WT, wild type; H58F, RNase mutant; N71/176Q, glycosylation mutant.

mannan (a natural ligand that competes for binding to DC-SIGN and MR), or DC-SIGN- and MR-specific blocking antibodies, followed by a 1 h incubation with fluorescently-labeled SEA or natural omega-1. As reported previously (7), uptake of SEA by human moDCs could be reduced by mannan and either DC-SIGN or MR blocking antibodies in an additive manner (Fig. 4 C). With regard to omega-1, pre-treatment with mannan could fully block binding and uptake of omega-1 by DCs. Interestingly, binding and uptake of natural omega-1 were significantly reduced by MR but not by DC-SIGN blocking antibodies (Fig. 4 C). Pre-incubation with the combination of both blocking antibodies did not have any additional effect on the uptake of omega-1 as compared to pre-incubation with anti-MR antibody alone. In addition, we found that recombinant omega-1 was recognized and internalized by DCs in a similar MR-dependent fashion as natural omega-1 (Fig. 4 D). To further investigate the observations of selective recognition and uptake of omega-1 by MR, we made use of the K562 and 3T3 cell




Figure 3. Glycosylation and RNase activity are essential for omega-1 to prime Th2 responses *in vivo*. 4get/KN2 IL-4 dual reporter mice were injected s.c. with SEA (20 µg) or 3 µg WT mutant recombinant omega-1 into the footpad. After 7 days the frequency of GFP⁺ and huCD2⁺ within the CD4⁺CD44^{high} effector T cell population was determined by flow cytometry in the draining popliteal lymph nodes. Depicted are (A) concatenated FACS plots, (B) frequencies of huCD2⁺ within the CD4⁺CD44^{high} population and (C) total huCD2⁺ T cell numbers in draining lymph nodes of combined data of 4 mice per group. (A) The frequencies of each population are indicated as percentages in the plots. One of 3 independent experiments is shown. Bars represent mean ± SD. *, # P < 0.05, ** P < 0.01, *** P < 0.001 for values significantly different from the PBS control (*) or between test conditions (#) based on two-sided t-test. WT, wild type; H58F, RNase mutant; N71/176Q, glycosylation mutant.

lines selectively expressing human DC-SIGN and MR, respectively. Fluorescently-labeled SEA was readily bound by both the DC-SIGN- and MR-expressing cells, which was not observed upon pre-incubation with EGTA or in parental control cell lines lacking CLR expression. In line with the DC-binding and uptake data, omega-1 binding could be observed in the cell line expressing MR (Fig. 4 E), but not in the cell line expressing DC-SIGN (Fig. 4 F). It should be noted that in these uptake experiments (Fig. 4 C+D), blocking with anti-MR antibody was not complete ($\pm 40\%$ reduction). However, given that blocking the binding of omega-1 to the cell





Figure 4. Mannose Receptor (MR) mediates recognition and internalization of omega-1 by human DCs. (A) Human moDCs were incubated for 1 h with PF-647-labeled recombinant WT omega-1, the glycosylation mutant or the RNase mutant and analyzed for uptake of antigens by FACS analysis. One representative experiment with duplicate samples out of 2 experiments is shown. Bars represent mean \pm SD. (B) Human moDCs were incubated for 1 h with PF-647-labeled omega-1 and, where indicated, either pre-incubated ('pre') with EGTA to prevent omega-1 binding to CLR*s a priori*, or treated afterwards with EGTA ('post') to remove any CLR-bound omega-1 from the cell surface. One of two independent experiments is shown and data represent mean \pm SD of duplicates. (C+D) A binding/internalization assay of natural (C) and recombinant omega-1 (D) by immature moDCs was performed analogous to (B) following pre-incubation with indicated reagents. Binding and internalization are shown relative to control pre-treatment. (C) Data are based on 5 experiments and are shown as mean \pm SD. (D) One of two independent experiments is shown and data represent mean \pm SD of duplicates. (E) 3T3 cell-line expressing MR and (F) K-SIGN expressing DC-SIGN or parental control cell lines (3T3 and K-562) were incubated with PF-647-labeled omega-1 and SEA in the presence or absence of EGTA to determine specificity. One representative experiment based on duplicate samples out of 2 is shown. Bars represent mean \pm SD. *,# $P < 0.05$, **,## $P < 0.01$, *** $P < 0.001$ for significant differences between control conditions (*) or between test conditions (#) based on two-sided *t*-test. WT, wild type; H58F, RNase mutant; N71/176Q, glycosylation mutant.

line selectively expressing MR with the anti-MR antibody was not complete either (data not shown), it is likely that a low affinity of the anti-MR antibody accounts for this finding rather than that other receptors are involved. Taken together, our data show that recognition and internalization of omega-1 by human DCs is dependent on its glycosylation and that MR is a major CLR involved in this process.

Omega-1 suppresses DC function by interfering with protein synthesis

Next we examined the molecular mechanism through which the RNase activity of omega-1 exerts its modulatory effects on human DCs. We noted that omega-1-stimulated DCs in response to CD40 ligation were not only impaired in their capacity to produce IL-12 p70, as reported previously (15), but also to secrete other cytokines and chemokines (Fig. 5 A). This suggested that the suppression may not be gene specific, but could be the result of general inhibition of protein synthesis. Indeed, following exposure of DCs to omega-1 or SEA, a dose-dependent reduction of protein synthesis could be observed, similar to what is found in DCs exposed to ricin, a well-known protein synthesis inhibitor (25) (Fig. 5 B). In addition, this inhibition by omega-1 was time-dependent and observed in both the presence and absence of LPS (Fig. 5 C). The capacity to inhibit protein synthesis was dependent on its RNase activity and uptake via its glycans, since the RNase as well as the glycosylation mutant failed to interfere with protein synthesis (Fig 5 D). As several fungal ribonucleolytic proteins, so-called ribotoxins, have been described to inhibit protein synthesis through cleavage of ribosomal RNA (rRNA) following translocation into the cytosol (26), we first evaluated the localization of omega-1 in human DCs following uptake. We found that omega-1 was efficiently internalized

Figure 5. Omega-1 suppresses protein synthesis through breakdown of rRNA and mRNA. (A) After human moDCs had been pulsed for 40 h with omega-1 (125, 250 and 500 ng/ml) in combination with LPS (100 ng/ml), the cells were co-cultured for 24 h with the J558 cell-line, expressing CD40-L, to mimic the interaction with T cells. Bars represent mean \pm SD of triplicate wells of one of two independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ for values significantly different from the LPS control. (B) Following 16 h incubation of human DCs with a concentration range of indicated reagents in the presence of LPS (100 ng/ml), protein synthesis was assessed after a 2 h pulse with radioactively-labeled methionine. Ricin, as potent inhibitor of protein synthesis, was taken along as positive control (25). One of 2 experiments is shown. (C) As described in (B), protein synthesis by human DCs was followed over time after exposure to omega-1 (500 ng/ml) either in the presence or absence of LPS (100 ng/ml). Data are shown relative to unstimulated or LPS-stimulated controls as depicted by the dotted line. Data are representative of two independent experiments and are depicted as mean \pm SD. (D) Protein synthesis by human moDCs following exposure to increasing concentrations of the recombinant omega-1 variants was assessed as described in (B). Data are shown relative to LPS-stimulated DCs. Data are representative of two independent experiments and are depicted as mean \pm SD. (E) DCs were stimulated with FITC-labeled recombinant omega-1 for 1 h and uptake was visualized by confocal laser scanning microscopy. Nuclei were stained with Hoechst. One of three experiments is shown. Scale bar represents 10 μ m. See Fig. S1 for a video of Z-stacked images. (F) Cytoplasmic fractions of human DCs stimulated for 3 h with omega-1 were run under non-reducing conditions by SDS-PAGE and analyzed by Western Blot for presence of omega-1 or silver-stained to control for input. DCs incubated at 4°C were taken along as controls as these cells have surface-bound, but not internalized omega-1. One of two experiments is shown. (G) human DCs were stimulated with FITC-labeled omega-1 (1 μ g/ml) ►



- and after 2 h fixed and stained for rRNA. Subcellular localization of ω -1 was determined by confocal microscopy. One representative cell from three independent experiments is shown. Scale bar represents 10 μ m. (H) After rabbit reticulocyte lysate containing functional ribosomes was incubated for 1 h with ω -1 (1 and 5 μ g/ml), IPSE (1 and 5 μ g/ml) as negative control, or ES (schistosome egg excretory/secretory products) (25 μ g/ml), containing ω -1, isolated rRNA was analyzed for breakdown on a 2% agarose gel. The RNase α -sarcin was taken along as positive control as it should give a single rRNA cleavage product when incubated with functional ribosomes (white arrowhead) (28). One of three independent experiments is shown. (I) rRNA isolated from 24 h ω -1-stimulated human DCs and was visualized by running a lab-on-a-chip picogel. One of three experiments is shown. (J-K) rRNA or mRNA expression of indicated genes in DCs was assessed by real time-qPCR at different time points after stimulation with ω -1 (500 ng/ml and 2 μ g/ml) in the presence or absence of LPS (100 ng/ml). Data are shown relative to unstimulated or LPS-stimulated controls, which were set to 1. RNA expression was normalized based on a genomic real time-qPCR for *ccr5*. Data represent the mean of 3 independent experiments. ω -1, ω -1; WT, wild type; H58F, RNase mutant; N71/176Q, glycosylation mutant.

by DCs and present throughout the cell after 1 h (Fig. 5 E and Fig. S1). In addition, Western blots revealed the presence of omega-1 in the cytosolic fraction of omega-1-stimulated DCs after 3 h (Fig. 5 F). In line with this, co-localization experiments using immunofluorescence confocal microscopy by staining for rRNA, showed that 2 h after stimulation of DCs with omega-1, omega-1 partially co-localized with rRNA (Fig. 5 G). We next tested whether omega-1 could cleave rRNA in the context of functional ribosomes in a cell free assay. Omega-1 was able to break down rRNA, while IPSE, another *S. mansoni* egg-derived protein that lacks RNase activity but has identical glycans as omega-1 (27), did not induce any rRNA degradation (Fig. 5 H), indicating that omega-1 is able to interfere with ribosomal function by cleavage of rRNA. Analysis of the integrity of rRNA isolated from omega-1-exposed DCs by gel electrophoresis showed a preferential breakdown of 28S rRNA (Fig. 5 I). This was confirmed by real time-PCR, which additionally revealed breakdown of 18S at later time points (Fig. 5 J). Since the known T2 RNases have no sequence-specific RNase activity (19), we evaluated whether omega-1, which belongs to the T2 RNase family, may additionally impair protein synthesis by degrading not only rRNA but also mRNA transcripts in a generic manner. When DCs were stimulated with omega-1, both in the presence or absence of LPS, a concentration and time dependent loss in mRNA transcripts of housekeeping genes *TAF1* and *GAPDH* as well as *IL12B* could be observed (Fig. 5 K), suggesting that omega-1 targets mRNA transcripts in a general manner in DCs as well. Taken together, these data support the notion that the RNase activity enables omega-1 to modulate human DC function by interfering with protein synthesis through cleavage of rRNA and mRNA following translocation into the cytosol.

MR mediates omega-1-induced protein synthesis inhibition, DC modulation and Th2 polarization

To address the role of omega-1 binding by MR in mediating RNase-dependent DC modulation and Th2 priming by omega-1, we used blocking antibodies directed against MR or DC-SIGN. Blocking of MR during the stimulation of human DCs with omega-1 significantly prevented the inhibition of protein synthesis (Fig. 6 A), while blocking of DC-SIGN had no effect, showing that the interference with protein synthesis by omega-1 is dependent on MR. In line with these observations, blocking of MR significantly reduced the capacity of omega-1 to suppress LPS-induced CD86 expression (Fig. 6 B) and IL-12 production following CD40 ligation (Fig. 6 C) or to condition DCs to induce a Th2 response (Fig. 6 D). The importance of MR was further substantiated by the observations that in contrast to their WT counterparts, MR^{-/-} murine DCs, when conditioned with omega-1, failed to prime a Th2-skewed allogeneic T cell response *in vitro* (Fig. 6 E). These data establish that MR is essential for the omega-1-driven Th2 polarization via DCs *in vitro*.

Omega-1 requires MR to prime Th2 responses *in vivo*

Finally, to investigate the role of MR in Th2-priming by omega-1 *in vivo*, natural omega-1 or PBS were injected subcutaneously into the footpad of WT and MR^{-/-} mice. After 7 days

the draining popliteal LNs were harvested and restimulated *in vitro* with PBS, omega-1 or a polyclonal stimulus PHA and analyzed for cytokine production. Antigen-specific restimulation of omega-1-primed LNs from WT mice resulted in a Th2-polarized response as evidenced by elevated levels of Th2-associated cytokine IL-5 but not of Th1-associated cytokine IFN- γ , which was absent in LN cells derived from MR^{-/-} (Fig. 7 A). Furthermore, intracellular staining for IFN- γ and IL-4 following antigen-specific restimulation of CD4⁺ T cells from omega-1-primed LNs, showed a significant increase in IL-4-producing T cells from WT but not MR^{-/-} mice (Fig. 7 B). The failure of MR^{-/-} mice to prime a Th2-polarized response in response to omega-1 was not due to a general failure of MR^{-/-} cells to produce these cytokines as the responses to PHA were comparable in WT and MR^{-/-} mice (Fig. 7 A and B). Taken together, these data show that MR is essential for priming of Th2 responses by omega-1 *in vivo*.

DISCUSSION


Using omega-1, a single glycosylated T2 RNase secreted by *Schistosoma mansoni* eggs, we studied the molecular mechanisms involved in conditioning dendritic cells to induce Th2 responses. By generating mutants of omega-1 we could show that both the glycosylation and the RNase activity of omega-1 are essential for its potent Th2-inducing activity both *in vitro* and *in vivo*. The glycan structures on omega-1 suggested that CLRs might play a role in its interaction with DCs. Although both MR and DC-SIGN have been shown to mediate binding and uptake of fucosylated antigens by DCs (29) and omega-1 harbors fucose-containing Le^x-glycan moieties, we observed that omega-1 significantly bound only to a MR-, but not to a DC-SIGN-expressing cell-line and that internalization by DCs was mainly MR-dependent and did not involve DC-SIGN. Lack of strong binding and uptake of omega-1 by DC-SIGN might be explained by the fact that in most DC-SIGN binding studies polyvalent Le^x-containing beads or conjugates have been used, which may be bound by DC-SIGN with a higher affinity than soluble glycoproteins, such as omega-1, that would present Le^x at a low valency (22). In line with this observation, DC-SIGN blocking experiments suggest that interaction with DC-SIGN does not play a major role in omega-1-driven Th2 polarization via DCs. On the other hand, the importance of MR in recognition and uptake of omega-1 was substantiated by the finding that conditioning of both human and murine DCs for Th2 polarization by omega-1 were significantly impaired when MR was blocked or when the DCs were deficient for MR, respectively. Furthermore, we confirmed and extended the importance of MR in Th2 polarization by omega-1 *in vivo* by showing that an antigen-specific Th2 response induced in MR^{-/-} mice following footpad injection of omega-1 was strongly reduced compared to the response elicited in WT mice. In this respect it is important to note that human and murine MR have a similar carbohydrate binding specificity (30). Thus, this establishes that omega-1 relies on MR to drive Th2 polarization. Apart from schistosome egg-derived antigens, it was recently shown that MR can also recognize glycosylated antigens derived from schistosome larvae in the skin (31), and that MR^{-/-} mice display a Th1-biased antigen-specific T cell response in the skin-draining LNs following infection with cercaria. This study along with our data, indicate

that MR may play a role in shaping of Th2-polarized immune responses during different stages of schistosome infection.

In vitro studies with DCs have shown that MR-crosslinking with antibodies (32) or by mannosylated antigens (33;34) can drive an anti-inflammatory cytokine program in DCs away from a Th1-promoting profile (32) and that allergen-driven Th2 polarization by DCs is in part dependent on MR (35;36). These studies suggest that engagement of MR may be sufficient to promote Th2 polarization, potentially via signaling events. However, our data demonstrate that MR binding alone is not sufficient for Th2 induction by omega-1, since glycans present on omega-1, in absence of RNase activity, fail to program DCs to induce Th2 responses. This is in line with the observation that IPSE/alpha-1, another major glycoprotein secreted by *S. mansoni* eggs with identical glycosylation as omega-1 (27), which can bind the cell line expressing MR (unpublished data) but lacks RNase activity, is unable to prime Th2 responses (15).

Apart from its glycosylation, omega-1 requires its RNase activity to induce a Th2 response via modulation of human DCs. It was observed that omega-1 in an RNase-dependent manner impaired protein synthesis and that DCs exposed to omega-1 displayed a progressive reduction in mRNA content of several unrelated genes as well as in rRNA levels. The drop in mRNA transcripts from both housekeeping genes (*TAF1* and *GAPDH*), inducible genes (*IL12B*), as well as rRNA (28S and 18S), suggests that omega-1 does not degrade specific transcripts, but targets the global RNA pool in DCs. While it currently remains to be determined what the relative contribution of each of these processes and their relative timing is to the impairment of protein synthesis, it is most likely that the observed inhibition in protein synthesis is a combined effect of degradation of mRNA transcripts and interference with ribosomal integrity due to rRNA cleavage. These data support the view that as a consequence of RNA breakdown, reduced protein synthesis is the mode of action through which the RNase activity enables omega-1 to condition DCs for priming of Th2 responses. Some RNases have been linked to Th2 polarization before. The major birch pollen allergen, Bet v 1 (37), was identified as an RNase. Furthermore, some fungal RNases that appear to selectively cleave rRNA, such as mitogillin and Asp f 1, are known to be allergens (38). Interestingly, for Asp f-1 it was found that its allergenicity was lost when its capacity to interfere with ribosomal function was abolished (39). In addition, a report has linked an endogenous RNase, the eosinophil-derived neurotoxin, to DC-mediated Th2 polarization (40). Although these studies have not specifically addressed the role of RNase activity in direct priming of Th2 responses, they do highlight the possibility

Figure 6. MR mediates omega-1-induced DC modulation and Th2 polarization *in vitro*. Following 1 h pre-incubation with blocking antibodies against MR, DC-SIGN or an isotype control (20 µg/ml), human moDCs were pulsed for 16 h (A) or 48 h (B-D) with natural omega-1 (500 ng/ml) in combination with LPS (100 ng/ml). (A) Protein synthesis was assessed as described in Fig. 5 B. One representative experiment based on duplicate samples out of 3 experiments is shown. Data are shown as mean ± SD. (B) The expression levels of CD86 on human DCs assessed by FACS are based on geometric mean fluorescence, relative to the DCs stimulated with LPS alone, which is set to 1 (dashed line). Data are based on 3 independent experiments and shown as mean ± SD. (C) Conditioned human DCs were co-cultured for 24 h with a CD40-L expressing cell line, to mimic the interaction with T cells. IL-12p70 cytokine



► expression levels are shown relative to the DCs stimulated with LPS alone, which is set to 1 (dashed line). Data are based on 3 independent experiments and shown as mean ± SD. (D) Conditioned human moDCs were cultured with allogeneic naive CD4⁺ T cells for 12 days in the presence of staphylococcal enterotoxin B and IL-2, and T cell polarization was analyzed as described in figure 1. FACS plots of one representative experiment out of 6 is shown. Bar graphs are based on 6 independent experiments and represent mean ± SD. (E) Murine splenic WT or MR^{-/-} DCs from a C57BL/6 background were co-cultured with naive Balb/c CD4⁺ T cells in the presence 2 μg/ml omega-1. After an expansion with rIL-2 at d 3, T cells were restimulated on d 6 with PMA and ionomycin and analyzed for intracellular cytokines. 1 representative experiment out of 3 is shown. *,# P < 0.05, **,## P < 0.01, ***P < 0.001 for significant differences with the LPS control (*) or between test conditions (#) based on paired analysis (two-sided paired t-test). ω-1, omega-1.

that Th2 priming through interference with ribosomal function may not be a unique feature of *S. mansoni*-derived omega-1, but may be shared by other RNases as well. In this respect, it is interesting to note that RNase T2 homologs can be identified in the genomes of *S. japonicum* and *S. haematobium*, as well as of the nematodes *Brugia malayi*, *Loa loa* and *Ascaris suum* (41-43). However, it is currently unknown whether these parasites actually express these T2 RNases and if they do, whether they play any role in Th2 polarization in their host. Taken together, our data suggest that for an RNase to harbor a Th2-priming capacity, it needs to be recognized by DCs and routed to reach the cytosol where in turn its enzymatic activity would result in suppression of protein synthesis, yet without shutting down DC function altogether or inducing cell death before T cell priming has occurred.

It remains to be established how omega-1 would be able to reach the ribosomes present in the cytosol. Some ribosome-inactivating proteins have been shown to translocate from the ER into the cytosol after retrograde transport or by direct escape from endosomes into the cytosol (44). In this respect, since omega-1 is internalized via MR, it is interesting to note that cross-presentation of OVA by DCs, a process that requires translocation of the antigen from endosomes into the cytosol, has been shown to be dependent on MR (45;46). Mechanistically, it was demonstrated that binding of the MR to OVA leads to poly-ubiquitination of MR, resulting in the recruitment of the ATPase p97, a member of the ER-associated degradation machinery, towards the endosomal membrane. p97 in turn was found to provide the energy to pull out the MR ligand into the cytoplasm (47). This suggests that the MR itself can regulate the transport of its ligand, into the cytoplasm and provides a mechanism through which omega-1 could be translocated into the cytosol of DCs.

The suppression of protein synthesis in DCs by omega-1, would be in line with the documented inhibitory effects of omega-1 as well as SEA on DC activation and TLR-induced expression of co-stimulatory molecules and cytokines (15;16). In addition, this mode of action would also provide an explanation for the finding that omega-1 alters DC morphology as a result of cytoskeletal changes (16), since halting of translation and concomitant stress responses can affect actin rearrangements and thereby cell morphology (48). Importantly, during interactions with naïve T cells, omega-1-conditioned DCs will, in contrast to unconditioned DCs, be largely refractory to respond to CD40 ligation by T cells, as their protein synthesis machinery is impaired. As a consequence, T cells are primed in the absence of IL-12 and in the context of low antigen presentation, a situation that is known to favor the induction of Th2 responses (49;50). This mechanism would be different from a 'default hypothesis' for Th2 induction (51-53) as it represents a dominant and active suppression of signals during DC-T cell interactions. Such a model of active suppression of DC signals for Th2 polarization would be in line with recent data showing that SEA-pulsed DCs, although still capable of processing antigen to present it on MHC-II, are impaired in their upregulation of surface MHC-II and CD86 or expression of IL-12 in response to CD40 ligation (15;54), as well as with the observation that omega-1-primed DCs have a reduced capacity to form T cell-DC conjugates (16).

Taken together, based on our data we propose a model in which the glycans present on omega-1 do not play a dominant role in functional modulation of DC function for induction




Figure 7. MR is essential for omega-1-driven Th2 polarization *in vivo*. MR^{-/-} and WT BL/6 mice were injected s.c. with omega-1 (2 μg in 30 μl PBS) or PBS into the footpad. (A) After 7 days the cells from the draining lymph node (LNs) were restimulated *in vitro* for 4 days with PBS, omega-1 (2 μg/ml) or PHA (10 μg/ml), as polyclonal stimulus, after which cytokine production was determined by ELISA. (B) Intracellular cytokine production of the CD3⁺/CD4⁺ T cells from these LNs was assayed by FACS after an additional 6 h restimulation with PMA and ionomycin. FACS plots show concatenated data from 4 mice are shown. The bar graphs represent the percentage of T cells single-positive for either IL-4 or IFN-γ. One of 2 independent experiments is shown. Data are means ± s.e.m. of 4 mice per group based on pooled triplicate wells for each mouse. * P < 0.05, ** P < 0.01 for significant differences based on paired analysis (two-sided paired *t*-test). ω-1, omega-1.

of Th2 responses, but instead are essential for efficient recognition and internalization by DCs via the MR. Subsequently, following translocation into the cytosol, omega-1 programs DCs to drive Th2 polarization in an RNase-dependent manner by interfering with ribosomal function and protein synthesis. These studies have uncovered a novel mechanism through which DCs can be programmed to drive Th2 responses. It will be of great interest to study whether targeting of MR and the protein synthesis machinery to condition DCs for priming Th2 responses is unique to schistosome-driven Th2 polarization, or a mechanism that is also involved in the initiation of other Th2-polarized immune responses, found during other helminth infections or allergies. In addition, the insight may help the design of Th2-polarizing molecules, that could be used in the development of vaccines against parasitic worm infections or approaches to counterbalance unwanted Th1 responses in hyper-inflammatory diseases (55;56).

MATERIALS AND METHODS

Preparation and purification of *S. mansoni* egg-derived antigens

SEA, omega-1 and IPSE/alpha-1 were prepared and isolated as described previously (15;57). The purity of the preparations was controlled by SDS-PAGE and silverstaining. Protein concentrations were tested using the Bradford or BCA procedure. (58;59)

Generation and production of WT, glycosylation mutant and RNase mutant forms of recombinant omega-1

Site-directed mutagenesis was used to generate a glycosylation and RNase mutant by mutating the two putative N-linked glycosylation sites (N71/176Q) or by targeting a conserved amino-acid residue (H58F) that is known to be critical for enzymatic activity in homologous T2 RNases (17;19), respectively. H58F and N71/176Q mutants were created by polymerase chain reaction (PCR) using mutagenic primers on a DH5 α /pProExHtb plasmid (Invitrogen) containing the WT omega-1 sequence (Acc.No. ABB73003.1). Successful mutation was confirmed by DNA sequencing. Subsequently, using restriction enzymes Sfi I and Apal the templates for WT and mutant omega-1 were subcloned into a pSecTag2 plasmid (Invitrogen) for stable transfection into HEK cells (15). Secreted recombinant omega-1 forms were sequentially purified from the HEK cell culture medium by immobilized metal affinity chromatography and size exclusion chromatography as described previously (15).

Human DC culture, stimulation and analysis

Monocytes were isolated from venous blood from healthy volunteers according to protocols approved by the Institutional Review Board of Leiden University Medical Center by density centrifugation on ficoll followed by CD14⁺ MACS isolation (Miltenyi) or a Percoll gradient as described (60) and were cultured in RPMI medium supplemented with 10% FCS, human rGM-CSF (50 ng/ml, Invitrogen) and human rIL-4 (25 units/ml) (R&D Systems). On day 3, culture medium including the supplements was replaced and on day 6 immature DCs were stimulated with the indicated reagents in the presence of ultrapure LPS (100 ng/ml) (*E. coli* 0111 B4 strain, InvivoGen). For CLR blocking indicated cells were pre-incubated with 20 μ g/ml anti-DC-SIGN (clone AZN-D1, Beckman Coulter) or 20 μ g/ml anti-MR (clone 15.2, Biolegend) for 60 min at 37 °C. As a Th1 control DCs were also pulsed with IFN- γ (1000 U/ml). After 48 h, DCs were harvested for co-culture with naïve T cells. In addition, 1x10⁴ matured DCs were co-cultured with 1x10⁴ CD40L-expressing J558 cells for 24 h to determine cytokine production by the DCs following activation by CD40L. IL-12p70 concentrations were determined by ELISA using mouse anti-human IL-12, clone 20C2 as capture antibody and biotinylated mouse-anti-human IL-12, clone C8.6 as detection antibody (both BD) Concentrations of IL-10, TNF- α , MIP-1 β and RANTES were determined by a multiplex LUMINEX assay according to the manufacturer's instruction (InvivoGen). The expression of CD86 on pulsed DCs was determined by FACS (FACSCanto) through staining with CD86-FITC (BD).

Murine T cell polarization assay

Splenic CD11c⁺MHCII⁺ DCs and CD62L⁺ CD4⁺ T cells were isolated by sorting from naive splenocytes derived from C57BL/6 and Balb/c mice, respectively. 2.5×10^5 CD4⁺ T cells were co-cultured with 1.25×10^5 splenic DCs and stimulated with 2 µg/ml omega-1. At d 3, T cells were expanded with 30 u/ml rIL-2 (R&D systems) and at d 6 restimulated with 50 ng/ml phorbol 12-myristate 13-acetate plus 2 µg/ml ionomycin for 5 h. 10 µg/ml brefeldin A was added during the last 2 h (all Sigma). The cells were stained with a combination of IL-4-PE and IFN-γ-APC antibodies (BD).

Human T cell culture and determination of T cell polarization

To determine T cell polarization, 5×10^3 48 h-pulsed DCs were co-cultured with 2×10^4 naïve T cells that were purified using a human CD4⁺/CD45RO⁻ column kit (R&D) in the presence of staphylococcal enterotoxin B (10 pg/ml; Sigma) in 96-well flat-bottom plates (Corning). On day 5, rhIL-2 (10 U/ml, R&D) was added and the cultures were expanded for another 7 days. For intracellular cytokine production, the primed CD4⁺ T cells were restimulated with 50 ng/ml phorbol 12-myristate 13-acetate plus 2 µg/ml ionomycin for 6 h. 10 µg/ml brefeldin A was added during the last 2 h (all Sigma). The cells were stained with a combination of IL-4-PE and IFN-γ-FITC antibodies (BD).

DC-SIGN- and MR-expressing cell line

K562 cell line stably expressing DC-SIGN (a gift from K. Figdor, Radboud University Nijmegen Medical Centre, Nijmegen, Netherlands; (61)) or 3T3 cell line stably expressing human MR (a gift from J.L. Miller, University of Oxford, Oxford, England, UK and G. Brown, University of Aberdeen, Aberdeen, Scotland, UK; (62)) and their respective parental control cell lines were seeded overnight in a 96-well plate at 10.000 cells/well. Where indicated, cells were pre-incubated with 10mM EGTA for 30 min at 37 °C. Subsequently, cells were incubated with 2 µg/ml PF-647-labeled SEA or 500 ng/ml PF-647-labeled omega-1 at 37 °C for 1 h and washed in ice cold PBS before analysis using flowcytometry.

Protein synthesis inhibition

Immature DCs were seeded overnight in 96-well flat bottom plates before stimulation with indicated reagents in the presence of LPS. At indicated time points after stimulation protein synthesis was determined by a 2 h pulse at 37 °C with 3µCi /0,05 ml [³⁵S]-methionine (EasyTag Express Protein labeling mix, Perkin Elmer) in serum-, cysteine and L-methionine free RPMI-1640. After a double washing step in PBS, cells were lysed for 5 min in AV-lysis buffer (20mM Tris HCl, pH7.6, 150 mM NaCl, 0.5% DOC, 1.0% NP40, 0.1% SDS) in the presence of protease inhibitors Leupeptin and Aprotinin 200ug/ml. Lysates were transferred on a filter (Perkin Elmer) and dried. After radioactive labeled proteins were precipitated on the filter with trichloroacetic acid, filters were washed with 96% ethanol and dried. Using a liquid scintillation cocktail for aqueous solution the radioactivity present on the filters was measured in a β-counter.

RNase activity assay

RNA was extracted from PBMC using the RNeasy kit (Qiagen). RNA was incubated for 1 h at 37 °C with indicated antigens in 0.01M Tris 0.02% Cu. Subsequently, RNA breakdown was visualized by running the samples on a 2% agarose gel containing ethidium bromide.

Ribosomal RNA breakdown

Rabbit Reticulocyte Lysate (Promega) was incubated with antigens as described by others (28). Briefly, following 1 h incubation at 37 °C in Tris-HCl (15 mM NaCl, 50 mM KCl, 2,5 mM EDTA), the reaction was stopped with 10% SDS and RNA was extracted from the ribosomes with phenol/chloroform. Next, isolated ribosomal RNA was denatured at 95°C and visualized by running the samples on a 2% agarose gel containing ethidium bromide.

Analysis of ribosomal RNA integrity in human DCs

mRNA was isolated from DCs conditioned by omega-1 for indicated time points using RNeasy mini Kit (Qiagen) according to the manufacturers recommendations. Integrity of rRNA was visualized using Agilent RNA 6000 Pico Kit in a 2100 Bioanalyzer (Agilent) according to the manufacturers recommendations.

RNA and DNA isolation, DNase treatment and cDNA synthesis

Total DNA and RNA was simultaneously extracted using Qiagen DNeasy blood and tissue kit as per manufacturer's instruction, except for that no RNase reaction was performed. DNase treatment was done using RQ1 RNase-Free DNase (Promega) as per manufacturer's instruction. cDNA synthesis was performed following standard procedures.

Analysis of gene expression levels

Primers and Taqman probes were provided as a Taqman gene expression kit (Applied Biosystems) or designed using Primer Express (Applied Biosystems) and synthesized by Bioglegio. Primers for DNA PCR for CCR5 was a generous gift from E. Boon (Leiden University Medical Centre, Leiden, Netherlands (63)). Real time qPCR was performed using ABI PRISM 7700 Sequence Detection System (SDS, Applied Biosystems). mRNA expression levels were normalized based on DNA input as determined by CCR5 PCR. Data were visualized as heatmaps using Tableau software (<http://www.tableausoftware.com>)

Antigen uptake by human DCs

SEA and omega-1 were fluorescently labeled with PF-647 using the Promofluor labeling kit (Promokine) and according to the manufacturers recommendations. 10.000 immature DCs/well were seeded in a 96 well plate. Where indicated cells were pre-incubated with 10mM EGTA, 100 µg/ml Mannan (Sigma-Aldrich), 20µg/ml anti-DC-SIGN (clone AZN-D1, Beckman Coulter) or 20 µg/ml anti-MR (clone 15.2, Biolegend) for 60 min at 37 °C. Subsequently, cells were incubated with 2 µg/ml PF-647-labeled SEA or 500 ng/ml PF-647-labeled omega-1 at 37 °C for 1 h and washed in ice cold PBS before analysis using flowcytometry.

Confocal microscopy

For live cell imaging, purified HEK-omega-1 was fluorescently labeled with N-hydroxysuccinimide (NHS)-fluorescein (Thermo-Scientific) according to the manufacturer's instructions. After protein labeling, nonreacted NHS-fluorescein was removed using Zeba desalt spin columns (Thermo-Scientific). Live cell imaging was performed with a TCS Sp5 inverse confocal laser scanning microscope (Leica) and analyzed with LAS AF software. In detail, 2.5×10^4 DCs were added to a channel of an IV0.4 μ -slide (Ibidi) and incubated for 2 h at 37°C and 6% CO₂ with 10 μ l fluorescein-labeled HEK-omega-1 (0.3 mg/ml). Nuclei of the cells were then counterstained with Hoechst 33342 (1:10,000) for 30 min. For confocal microscopy of fixed DCs, cells were allowed to adhere to Poly-D-Lysine coated cover slips overnight at a concentration of 80-100.000 cells/2 ml in 10% FCS/RPMI. DCs were incubated with omega-1 for 2 h at 37 °C (1 μ g/ml). Incubated cells were washed three times in 1% BSA/RPMI, fixed for 15 minutes with 4% paraformaldehyde (Sigma) in PBS, and washed twice in PBS. Next, cells were permeabilized with 0.1% Triton-X in PBS for 1 minute, washed twice in PBS and blocked for 15 minutes with 1% BSA/PBS. Cells were subsequently incubated with antibodies against rRNA (Abcam), followed by a secondary incubation step with a G α M-AF546 antibody (Invitrogen) in 1% BSA/PBS. Cells were washed in PBS and cover slips were mounted on glass slides with Vectashield and analyzed by confocal microscopy. Leica AOBSP2 confocal laser scanning microscope (CLSM) system was used, containing a DM-IRE2 microscope with glycerol objective lens (PL APO 63x/NA1.30) and images were acquired using Leica confocal software (version 2.61).

Cytoplasmic omega-1 Western blot

Cytoplasmic extracts of omega-1-incubated DCs were prepared using Nuclear Extraction Kit (Active Motif) as per manufacturer's instructions. Cytoplasmic extracts were concentrated 10-fold and subjected to 12% SDS-PAGE followed by silver staining or blotting onto nitrocellulose membrane. For silver staining 30 μ g/cm were applied, for Western blotting 100 μ g/cm. Omega-1 was then detected by the monoclonal anti-omega-1 antibody 140-3E11 and an alkaline phosphatase-labeled goat anti-mouse IgG (1:10,000) detection antibody (Dianova). Visualization was done by the substrate/chromogen mixture of 0.033% (w/v) nitro blue tetrazolium and 0.017% (w/v) 5-bromo-4-chloro-indolyl phosphate (Serva) in 0.1 M Tris-buffered saline, pH 9.5.

In vivo experiments

4get/KN2 (64) mice were bred and housed in the animal facility of the Trudeau Institute and used at 8-12 weeks of age. MR^{-/-} mice on a C57BL/6 background were provided by Dr. M. C. Nussenzweig (Rockefeller University, New York, NY) and were bred and housed in the animal facility of the Institutes of Molecular Medicine and Experimental Immunology at the University Hospital, Bonn. All experimental procedures were approved by the Institutional Animal Care and Use Committee of the Trudeau Institute, and Molecular Medicine and Experimental Immunology at the University Hospital of Bonn and by the Animal Studies Committee of Washington University School of Medicine. Mice were immunized s.c. into one

hind footpad with SEA (20 µg), omega-1 (3 µg), in a volume of 50 µl and the draining popliteal lymph nodes were analyzed one week later.

***In vitro* restimulation of lymph node cells**

1.5×10^6 popliteal LN cells/ml from individual animals were restimulated with 10 µg/ml SEA or 2 µg/ml omega-1. IL-5, IL-4 and IFN- γ were measured by ELISA in day 4 supernatants according to the manufacturer's recommendations (R&D). Following removal of the supernatants, cells were restimulated with 50 ng/ml phorbol 12-myristate 13-acetate plus 2 µg/ml ionomycin for 6 h. 10 µg/ml brefeldin A was added during the last 2 h (all Sigma). The cells were stained with a combination of IL-4-PE and IFN- γ -FITC antibodies (BD).

Statistical analysis

Data were analyzed for statistical significance using a two-sided paired Student's *t*-test or where indicated a two-sided unpaired Student's *t*-test. All *p*-values < 0.05 were considered significant.

Online supplemental material

Online supplemental material can be found at: <http://jem.rupress.org/content/209/10/1753/suppl/DC1>. Fig S1 shows a movie of z-stacked images of live DCs internalizing omega-1.

ACKNOWLEDGEMENTS

The authors thank Krystelle Nganou-Mkamdop for performing DC-antigen uptake experiments, and Heike Rohweder and Daniela Barths for their technical assistance in the mutagenesis and cloning studies of omega-1. Furthermore, we thank Dr. Buschow and Dr. Miller for providing the DC-SIGN- and MR-expressing cell-lines, respectively. This work was supported by the Netherlands Foundation for the Advancement of Tropical Research (WOTRO), Grant No W93-385 20077, the Dutch Organization for Scientific Research (NWO), Grant No ZONMW 912-03-048, ZONMW-VENI 016.066.093 NWO-CW 700.55.013 and the National Institutes of Health grant AI53825.

REFERENCES

1. Kapsenberg ML. Dendritic-cell control of pathogen-driven T-cell polarization. *Nat Rev Immunol* 2003 Dec;3(12):984-93.
2. Akira S, et al. Pathogen Recognition and Innate Immunity. *Cell* 2006 Feb 24;124(4):783-801.
3. Carvalho L, et al. Review series on helminths, immune modulation and the hygiene hypothesis: mechanisms underlying helminth modulation of dendritic cell function. *Immunology* 2009 Jan;126(1):28-34.
4. MacDonald AS, et al. Alarming dendritic cells for Th2 induction. *J Exp Med* 2008 Jan 21;205(1):13-7.
5. Phytian-Adams AT, et al. CD11c depletion severely disrupts Th2 induction and development in vivo. *J Exp Med* 2010 Sep 27;207(10):2089-96.
6. Okano M, et al. Induction of Th2 responses and IgE is largely due to carbohydrates functioning as adjuvants on *Schistosoma mansoni* egg antigens. *J Immunol* 1999 Dec 15;163(12):6712-7.
7. van Liempt E, et al. *Schistosoma mansoni* soluble egg antigens are internalized by human dendritic cells through multiple C-type lectins and suppress TLR-induced dendritic cell activation. *Mol Immunol* 2007 Apr;44(10):2605-15.
8. Gringhuis SI, et al. Carbohydrate-specific signaling through the DC-SIGN signalosome tailors immunity to *Mycobacterium tuberculosis*, HIV-1 and *Helicobacter pylori*. *Nat Immunol* 2009 Oct;10(10):1081-8.
9. van Liempt E, et al. Specificity of DC-SIGN for mannose- and fucose-containing glycans. *FEBS Lett* 2006 Nov 13;580(26):6123-31.
10. Van Diel, et al. The dendritic cell-specific C-type lectin DC-SIGN is a receptor for *Schistosoma mansoni* egg antigens and recognizes the glycan antigen Lewis x. *Glycobiology* 2003 Jun;13(6):471-8.
11. Bergman MP, et al. *Helicobacter pylori* modulates the T helper cell 1/T helper cell 2 balance through phase-variable interaction between lipopolysaccharide and DC-SIGN. *J Exp Med* 2004 Oct 18;200(8):979-90.
12. Ritter M, et al. *Schistosoma mansoni* triggers Dectin-2, which activates the Nlrp3 inflammasome and alters adaptive immune responses. *Proc Natl Acad Sci U S A* 2010 Nov 23;107(47):20459-64.
13. Correale J, et al. Helminth antigens modulate immune responses in cells from multiple sclerosis patients through TLR2-dependent mechanisms. *J Immunol* 2009 Nov 1;183(9):5999-6012.
14. Thomas PG, et al. Maturation of dendritic cell 2 phenotype by a helminth glycan uses a Toll-like receptor 4-dependent mechanism. *J Immunol* 2003 Dec 1;171(11):5837-41.
15. Everts B, et al. Omega-1, a glycoprotein secreted by *Schistosoma mansoni* eggs, drives Th2 responses. *J Exp Med* 2009 Aug 3;206(8):1673-80.
16. Steinfelder S, et al. The major component in schistosome eggs responsible for conditioning dendritic cells for Th2 polarization is a T2 ribonuclease (omega-1). *J Exp Med* 2009 Aug 3;206(8):1681-90.
17. Fitzsimmons CM, et al. Molecular characterization of omega-1: a hepatotoxic ribonuclease from *Schistosoma mansoni* eggs. *Mol Biochem Parasitol* 2005 Nov;144(1):123-7.
18. Wolf B, et al. A mechanism of the irreversible inactivation of bovine pancreatic ribonuclease by diethylpyrocarbonate. A general reaction of diethylpyrocarbonate. A general reaction of diethylpyrocarbonate with proteins. *Eur J Biochem* 1970 Apr;13(3):519-25.
19. Irie M, et al. Ribonuclease T2. *Methods Enzymol* 2001;341:42-55.
20. Meevissen MH, et al. Structural Characterization of Glycans on Omega-1, a Major *Schistosoma mansoni* Egg Glycoprotein That Drives Th2 Responses. *J Proteome Res* 2010 Mar 18;9(5):2630-42.
21. Yan SB, et al. Novel Asn-linked oligosaccharides terminating in GalNAc beta (1-->4)[Fuc alpha (1-->3)]GlcNAc beta (1-->) are present in recombinant human protein C expressed in human kidney 293 cells. *Glycobiology* 1993 Dec;3(6):597-608.
22. Meevissen MH, et al. *Schistosoma mansoni* egg glycoproteins and C-type lectins of host immune cells: Molecular partners that shape immune responses. *Exp Parasitol* 2011 May 15;
23. Mohrs M, et al. Analysis of type 2 immunity in vivo with a bicistronic IL-4 reporter. *Immunity* 2001 Aug;15(2):303-11.
24. Taylor ME, et al. Contribution to ligand binding by multiple carbohydrate-recognition domains in the macrophage mannose

- receptor. *J Biol Chem* 1992 Jan 25;267(3):1719-26.
25. Montanaro L, et al. Inhibition by ricin of protein synthesis in vitro. Ribosomes as the target of the toxin. *Biochem J* 1973 Nov;136(3):677-83.
 26. Lacadena J, et al. Fungal ribotoxins: molecular dissection of a family of natural killers. *FEMS Microbiol Rev* 2007 Mar;31(2):212-37.
 27. Wuhler M, et al. IPSE/alpha-1, a major secretory glycoprotein antigen from schistosome eggs, expresses the Lewis X motif on core-difucosylated N-glycans. *FEBS J* 2006 May;273(10):2276-92.
 28. Kao R, et al. Mitogillin and related fungal ribotoxins. *Methods Enzymol* 2001;341:324-35.
 29. Frison N, et al. Oligolysine-based oligosaccharide clusters: selective recognition and endocytosis by the mannose receptor and dendritic cell-specific intercellular adhesion molecule 3 (ICAM-3)-grabbing nonintegrin. *J Biol Chem* 2003 Jun 27;278(26):23922-9.
 30. East L, et al. The mannose receptor family. *Biochim Biophys Acta* 2002 Sep 19;1572(2-3):364-86.
 31. Paveley RA, et al. The Mannose Receptor (CD206) is an important pattern recognition receptor (PRR) in the detection of the infective stage of the helminth *Schistosoma mansoni* and modulates IFN γ production. *Int J Parasitol* 2011 Nov;41(13-14):1335-45.
 32. Chieppa M, et al. Cross-linking of the mannose receptor on monocyte-derived dendritic cells activates an anti-inflammatory immunosuppressive program. *J Immunol* 2003 Nov 1;171(9):4552-60.
 33. Op den Brouw ML, et al. The mannose receptor acts as hepatitis B virus surface antigen receptor mediating interaction with intrahepatic dendritic cells. *Virology* 2009 Oct 10;393(1):84-90.
 34. Yamamoto Y, et al. Involvement of mannose receptor in cytokine interleukin-1 β (IL-1 β), IL-6, and granulocyte-macrophage colony-stimulating factor responses, but not in chemokine macrophage inflammatory protein 1 β (MIP-1 β), MIP-2, and KC responses, caused by attachment of *Candida albicans* to macrophages. *Infect Immun* 1997 Mar;65(3):1077-82.
 35. Royer PJ, et al. The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. *J Immunol* 2010 Aug 1;185(3):1522-31.
 36. Li J, et al. The dendritic cell mannose receptor mediates allergen internalization and maturation involving notch 1 signalling. *Clin Exp Immunol* 2010 Nov;162(2):251-61.
 37. Bufe A, et al. The major birch pollen allergen, Bet v 1, shows ribonuclease activity. *Planta* 1996;199(3):413-5.
 38. Kao R, et al. Mitogillin and related fungal ribotoxins. *Methods Enzymol* 2001;341:324-35.
 39. Garcia-Ortega L, et al. Production and characterization of a noncytotoxic deletion variant of the *Aspergillus fumigatus* allergen Asp f1 displaying reduced IgE binding. *FEBS J* 2005 May;272(10):2536-44.
 40. Yang D, et al. Eosinophil-derived neurotoxin acts as an alarmin to activate the TLR2-MyD88 signal pathway in dendritic cells and enhances Th2 immune responses. *J Exp Med* 2008 Jan 21;205(1):79-90.
 41. The *Schistosoma japonicum* genome reveals features of host-parasite interplay. *Nature* 2009 Jul 16;460(7253):345-51.
 42. Young ND, et al. Whole-genome sequence of *Schistosoma haematobium*. *Nat Genet* 2012 Feb;44(2):221-5.
 43. Hillwig MS, et al. Zebrafish RNase T2 genes and the evolution of secretory ribonucleases in animals. *BMC Evol Biol* 2009;9:170.
 44. Sandvig K, et al. Delivery into cells: lessons learned from plant and bacterial toxins. *Gene Ther* 2005 Jun;12(11):865-72.
 45. Burgdorf S, et al. Spatial and mechanistic separation of cross-presentation and endogenous antigen presentation. *Nat Immunol* 2008 May;9(5):558-66.
 46. Burgdorf S, et al. Distinct pathways of antigen uptake and intracellular routing in CD4 and CD8 T cell activation. *Science* 2007 Apr 27;316(5824):612-6.
 47. Zehner M, Chasan AI, Scheutte V, Embgenbroich M, Quast T, Kolanus W, et al. Mannose Receptor poly-ubiquitination regulates endosomal recruitment of p97 and cytosolic antigen translocation for cross-presentation. *Proc Natl Acad Sci U S A*. In press 2011.
 48. Pelling AE, et al. Mechanical dynamics of single cells during early apoptosis. *Cell Motil Cytoskeleton* 2009 Jul;66(7):409-22.

49. Oswald IP, et al. IL-12 inhibits Th2 cytokine responses induced by eggs of *Schistosoma mansoni*. *J Immunol* 1994 Aug 15;153(4):1707-13.
50. Constant S, et al. Extent of T cell receptor ligation can determine the functional differentiation of naive CD4+ T cells. *J Exp Med* 1995 Nov 1;182(5):1591-6.
51. Wang ZE, et al. CD4+ effector cells default to the Th2 pathway in interferon gamma-deficient mice infected with *Leishmania major*. *J Exp Med* 1994 Apr 1;179(4):1367-71.
52. Yates A, et al. Cytokine-modulated regulation of helper T cell populations. *J Theor Biol* 2000 Oct 21;206(4):539-60.
53. Jankovic D, et al. Mechanisms underlying helminth-induced Th2 polarization: default, negative or positive pathways? *Chem Immunol Allergy* 2006;90:65-81.:65-81.
54. Marshall FA, et al. Uncoupling of induced protein processing from maturation in dendritic cells exposed to a highly antigenic preparation from a helminth parasite. *J Immunol* 2008 Dec 1;181(11):7562-70.
55. Ricardo-Gonzalez RR, et al. IL-4/STAT6 immune axis regulates peripheral nutrient metabolism and insulin sensitivity. *Proc Natl Acad Sci USA* 2010 Dec 28;107(52):22617-22.
56. Jager A, et al. Effector and regulatory T-cell subsets in autoimmunity and tissue inflammation. *Scand J Immunol* 2010 Sep;72(3):173-84.
57. de Jong EC, et al. Microbial compounds selectively induce Th1 cell-promoting or Th2 cell-promoting dendritic cells in vitro with diverse th cell-polarizing signals. *J Immunol* 2002 Feb 15;168(4):1704-9.
58. Woodward MP, et al. Detection of monoclonal antibodies specific for carbohydrate epitopes using periodate oxidation. *J Immunol Methods* 1985 Apr 8;78(1):143-53.
59. Hamilton JV, et al. Periodate-sensitive immunological cross-reactivity between keyhole limpet haemocyanin (KLH) and serodiagnostic *Schistosoma mansoni* egg antigens. *Parasitology* 1999 Jan;118 (Pt 1):83-9.
60. Sallusto F, et al. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med* 1994 Apr 1;179(4):1109-18.
61. Geijtenbeek TB, et al. Identification of DC-SIGN, a novel dendritic cell-specific ICAM-3 receptor that supports primary immune responses. *Cell* 2000 Mar 3;100(5):575-85.
62. Miller JL, et al. The mannose receptor mediates dengue virus infection of macrophages. *PLoS Pathog* 2008 Feb 8;4(2):e17.
63. Boon EM, et al. Y chromosome detection by Real Time PCR and pyrophosphorolysis-activated polymerisation using free fetal DNA isolated from maternal plasma. *Prenat Diagn* 2007 Oct;27(10):932-7.
64. Mohrs K, et al. A two-step process for cytokine production revealed by IL-4 dual-reporter mice. *Immunity* 2005 Oct;23(4):419-29.

