

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/29450> holds various files of this Leiden University dissertation.

Author: Koens, Lianne

Title: Clinicopathologic and genetic aspects of primary cutaneous large B-cell lymphomas

Issue Date: 2014-10-29

MicroRNA profiling of primary cutaneous large B-cell lymphomas

*Public Library of Science ONE 2013;
8(12): e82471*

Lianne Koens¹

Yongjun Qin^{2,3}

Wai Y. Leung³

Willem E. Corver¹

Patty M. Jansen¹

Rein Willemze³

Maarten H. Vermeer³

Cornelis P. Tensen³

Departments of ¹Pathology; and ³Dermatology, Leiden University
Medical Center, Leiden, The Netherlands;

²Biotechnology Center, Shanxi Academy of Agricultural Sciences,
Taiyuan, China

Abstract

Aberrant expression of microRNAs is widely accepted to be pathogenetically involved in nodal diffuse large B-cell lymphomas (DLBCLs). However, the microRNAs profiles of primary cutaneous large B-cell lymphomas (PCLBCLs) are not yet described. Its two main subtypes, i.e., primary cutaneous diffuse large B-cell lymphoma, leg type (PCLBCL-LT) and primary cutaneous follicle center lymphoma (PCFCL) are characterized by an activated B-cell (ABC)-genotype and a germinal center B-cell (GCB)-genotype, respectively. We performed high-throughput sequencing analysis on frozen tumor biopsies from 19 cases of PCFCL and PCLBCL-LT to establish microRNA profiles. Cluster analysis of the complete microRNome could not distinguish between the two subtypes, but 16 single microRNAs were found to be differentially expressed. Single microRNA RT-qPCR was conducted on formalin-fixed paraffin-embedded tumor biopsies of 20 additional cases, confirming higher expression of miR-9-5p, miR-31-5p, miR-129-2-3p and miR-214-3p in PCFCL as compared to PCLBCL-LT. MicroRNAs previously described to be higher expressed in ABC-type as compared to GCB-type nodal DLBCL were not differentially expressed between PCFCL and PCLBCL-LT. In conclusion, PCFCL and PCLBCL-LT differ in their microRNA profiles. In contrast to their gene expression profile, they only show slight resemblance with the microRNA profiles found in GCB- and ABC-type nodal DLBCL.

3

Introduction

Primary cutaneous large B-cell lymphomas (PCLBCLs) are a group of malignant lymphoproliferative disorders presenting in the skin with no evidence of extracutaneous disease at the time of diagnosis.¹ In the latest World Health Organization (WHO) 2008 classification two main types of PCLBCL are distinguished: primary cutaneous follicle center lymphoma (PCFCL) and primary cutaneous diffuse large B-cell lymphoma, leg type (PCLBCL-LT).² PCFCL is considered an indolent type of lymphoma (5 year overall survival (OS) > 95%), whereas PCLBCL-LT has a more aggressive clinical course (5 year OS approximately 40%).³ The two subtypes also show marked differences at a molecular level by array comparative genomic hybridization⁴ and gene expression profiling. PCFCL and PCLBCL-LT have gene expression profiles corresponding to germinal center B-cell (GCB)-type diffuse large B-cell lymphoma (DLBCL) and activated B-cell (ABC)-type nodal DLBCL, respectively.⁵ Clinical distinction at the time of first diagnosis between these different subtypes of PCLBCL is important, as the first choice of treatment differs between the two entities. PCFCL can be adequately treated with radiotherapy, but the more aggressive behavior of PCLBCL-LT warrants for treatment with anthracyclin-based chemotherapy combined with rituximab.⁶

MicroRNAs are approximately 22 nucleotides long non-coding RNA molecules that can regulate translation of several specific target messenger RNAs (mRNAs). The two major responsible mechanisms are inhibition of translational initiation by blocking ribosomes and direct targeting of mRNA at the 3 prime untranslated region leading to degradation of the molecule.⁷ Alterations in the expression of microRNAs contribute to the pathogenesis of different types of malignancies, including malignant lymphomas, but are also involved in normal development of hematopoietic cells.⁸ Although a role for microRNAs in the pathogenesis of nodal DLBCLs is generally recognized, little is known about the presence and significance of microRNAs in PCLBCL. In one recently published study, a limited set of microRNAs was investigated in PCFCL.⁹

In nodal DLBCL, much effort has been given to differentiate between the two major molecular subtypes, i.e. GCB-type DLBCL and ABC-type or non-GCB-type DLBCL, by means of expression of specific microRNAs or microRNA profiles. However, conflicting results concerning this topic have been published. The most frequently encountered microRNA alterations are upregulation of miR-155, miR-21, miR-221 and miR-222 in ABC-type DLBCL as compared to GCB-type DLBCL as detected by microarray profiling and/or Real Time quantitative PCR (RT-qPCR).¹⁰⁻¹⁹ To date, the prognostic value of expression levels of specific microRNAs in nodal DLBCL remains unclear. Although higher expression of miR-222 correlating to a shorter progression-free survival (PFS) was reproduced in more than one study,^{12,15,20} a correlation was not consistently found for other microRNAs. In addition, reported microRNA profiles generated by complete microRNA profiling that would distinguish between the two subtypes of nodal DLBCL have no substantial similarities (Supplemental Table 1).^{12,14-18,21} To some extent, the observed variations might be explained by the different techniques used to quantify microRNAs, i.e. different microarrays or varying reference RNAs in RT-PCR experiments. Furthermore, the techniques used to distinguish the two molecular subtypes from each other are not uniform. The encountered differences might on the other hand also reflect the profound tumor heterogeneity within the group of nodal DLBCLs.

In search of differences in the pathogenesis of PCFCL and PCLBCL-LT we investigated the microRNA profiles in PCLBCL by performing microRNA high-throughput sequencing. Several single microRNAs that were differentially expressed between PCFCL and PCLBCL-LT in our high-throughput sequence data were validated by RT-qPCR. Furthermore, we were interested in comparing the microRNA profiles of PCFCL and PCLBCL-LT with the (known) microRNA profiles of GCB- and ABC-type nodal DLBCL, in search for common pathogenetical pathways and common targets for tailored therapy.

Materials and methods

Tumor samples

Frozen skin biopsy or excision material from primary tumors was collected from 6 patients with PCFCL and 13 patients with PCLBCL-LT from the archives of the Leiden University Medical Center (Leiden, The Netherlands). Formalin-fixed and paraffin-embedded (FFPE) skin biopsy material was available from 5 of these patients (2 PCFCL and 3 PCLBCL-LT cases) and an additional 8 patients with PCFCL and 7 patients with PCLBCL-LT. The diagnosis was based on the criteria of the WHO-EORTC and WHO 2008 classifications,^{1,22} and confirmed by a panel of hematopathologists and dermatologists, aided by several additional immunohistochemical stainings. The clinical data and expression of relevant immunohistochemical markers are summarized in Table 1. In all cases physical examination, computed tomography scan, bone marrow examination and peripheral blood counts did not show extracutaneous disease. Furthermore, FFPE lymph node excisions of 34 cases of nodal DLBCL were collected from the Leiden University Medical Center, the Diaconessenhuis (Leiden, The Netherlands) and the Reinier de Graaf Hospital (Delft, The Netherlands). Of those, 19 were ABC-type and 11 were GCB-type, according to both Hans and Choi immunohistochemical algorithms,^{23,24} the four samples with discrepancy between the two algorithms were discarded from further analysis. The tumor cell percentage was at least 75% in every tumor specimen, as shown by immunohistochemistry for CD20 and CD3. All tissue samples were handled in a coded fashion, according to the Dutch National Ethical guidelines (Code for Proper Secondary Use of Human Tissue, Dutch Federation of Medical Scientific Societies), waving the need for specific

approval of the study by the ethical committee and patient informed consent.

Human B-cells were purified from peripheral blood mononuclear cells isolated from buffy coats obtained from anonymous blood donors (Sanquin Bloodbank, Nijmegen, The Netherlands) and were *in vitro* activated subsequently (details are given in Supplemental Materials and Methods).

MicroRNA library preparation, high-throughput sequencing and data analysis

Total RNA was extracted from the frozen tumor samples and activated B-cells using TRIzol and according to the protocol described by the supplier (Invitrogen, Carlsbad, CA, USA). Library preparation was to a large extent based on the SOLiD Total RNA-Seq Kit protocol (SREK protocol, Applied Biosystems, Nieuwerkerk a/d IJssel, The Netherlands) as described previously (Supplemental Materials and Methods).²⁵ Subsequently, 0.5 nM per sample of the cDNA clusters generated was loaded onto the flow cells for massively parallel high-throughput sequencing using the Illumina Genome Analyzer II (Illumina) performing sequencing-by-synthesis technology. The raw sequencing data were extracted through standard Illumina pipelines. The primary sequencing data and are publicly available through the Gene Expression Omnibus (GEO) archive through accession GSE51359. Illumina sequence reads were aligned to the human genome (GRCh37/hg19) using the miRDeep2.0 algorithm,²⁶ which predicts precursor sequences and matches them to precursor microRNAs enlisted in the miRBase 18.0 database (<http://www.mirbase.org>). Only sequences with a minimum sequence length of 17 bases were analyzed, allowing for a maximum mismatch of one base. Samples with less than 100.000 microRNA reads were discarded from further analyses.²⁷ The different microRNAs were only included in the analyses if in at least half of the samples of at least one of the three analyzed groups (PCFCL, PCLBCL-LT or activated B-cells) the single microRNA read number exceeded 0,01% of the total number of microRNA reads (100 reads/million). The samples were then analyzed for differential expression between the groups by the Bioconductor package EdgeR,²⁸ considering only mature microRNAs. Differential expression was considered statistically significant when a *p*-value below 0.05 was reached after multiple testing correction according to the Benjamini-Hochberg method.²⁹ EdgeR transformed, normalized microRNA expression (in counts per million reads) were used for unsupervised hierarchical clustering, running under R 2.14.2 software (<http://www.r-project.org>). The NormFinder version 0.953 algorithm³⁰ was used on high-throughput sequencing data to identify the most stably expressed microRNAs within and between the analyzed subgroups suitable for RT-qPCR normalization.³¹

MicroRNA real-time qPCR

Total RNA was isolated from 10 μ m sections of FFPE tumor samples. Approximately 300 ng of total RNA from each sample (FFPE or frozen, same isolation as for the high-throughput sequencing) was reverse transcribed using the microRNA reverse transcription kit (Applied Biosystems). RT-qPCR was performed in duplicate using Taqman microRNA and control assays (Applied Biosystems). The output data were analyzed using CFX Manager software (Bio-Rad). The expression of the different microRNAs was analyzed using the $\Delta\Delta$ Ct method expressed relative to the most stably expressed endogenous controls according to GeNorm.³² For more details see Supplemental Materials and Methods. Statistical analyses were performed using the Mann Whitney U test in IBM SPSS version 20 (SPSS Inc., Chicago, IL, USA) and the data were visualized in GraphPad Prism (GraphPad Software Inc, CA, USA).

Table 1. Patient characteristics

Case	Experiment	Sex	Age at diagnosis	Primary tumor	Immunohistochemistry				BCR	Initial therapy	Recurrence	PFS (months)	Follow-up (months)	Current status
1 FC	HTS & RT-PCR	f	81	head	-	BCL2	BCL6	CD10	IRF4	-	yes	62	86	A ⁰
2 FC	HTS & RT-PCR	m	71	trunk	-	-	+	-	-	RT	no	89	90	A ⁰
3 FC	HTS	f	71	head	-	-	+	-	-	RT + o.p.	no	55	56	A ⁰
4 FC	HTS	m	72	head	+	-	+	-	-	RT	no	32	33	A ⁰
5 FC	HTS	m	67	head	-	-	+	-	-	RT	no	55	56	A ⁰
6 FC	HTS	m	59	arm	-	-	+	-	-	RT	yes	8	9	A ⁺
7 FC	RT-PCR	f	53	head	-	-	+	-	-	RT	yes	46	128	A ⁰
8 FC	RT-PCR	m	33	trunk	+	-	+	-	-	RT	yes	60	185	A ⁺
9 FC	RT-PCR	m	55	trunk	-	-	NA	-	-	RT	yes	5	105	A ⁰
10 FC	RT-PCR	f	50	trunk	-	-	-	-	-	RT	no	63	64	A ⁰
11 FC	RT-PCR	m	47	trunk	-	-	+	-	-	RT	yes	7	68	A ⁺
12 FC	RT-PCR	m	44	head	-	-	+	-	-	RT	no	52	53	A ⁰
13 FC	RT-PCR	m	35	trunk	-	-	+	-	-	RT + PCT	no	20	26	A ⁰
14 FC	RT-PCR	m	66	trunk	-	-	+	-	-	RT	yes	11	36	A ⁺
1 LT	HTS & RT-PCR	m	88	leg	+	+	+	-	+	igM, igD, κ	yes	2	12	D ⁺
2 LT	HTS & RT-PCR	m	81	leg	+	+	+	-	+	igM, igD, λ	no	0	2	D ⁺
3 LT	HTS & RT-PCR	m	68	leg	+	+	+	-	+	igM, igD, λ	yes	17	60	D ⁺
4 LT	HTS	f	89	leg	+	+	NA	-	+	NA	yes	21	32	A ⁺
5 LT	HTS	f	84	leg	+	+	+	-	+	igM, igD, κ	no	0	2	D ⁰
6 LT	HTS	f	83	leg	+	+	+	-	+	igM, igD, κ	yes	34	38	D ⁺
7 LT	HTS	m	83	leg	+	+	+	-	+	igM, igD, κ	no	49	50	A ⁰
8 LT	HTS	f	76	leg	+	+	-	-	+	igM, κ	no	0	13	D ⁺
9 LT	HTS	m	77	both legs	+	+	-	-	+	igM, igD	no	0	3	D ⁺
10 LT	HTS	f	79	leg	+	+	+	-	+	igM, κ	?	>20	32	A ⁺
11 LT	HTS	f	81	leg	+	+	-	-	+	igM	yes	20	39	D ⁺
12 LT	RT-PCR	f	73	leg	+	+	+	-	+	igM, igD, κ	no	79	85	D ⁰
13 LT	RT-PCR	f	71	leg	+	+	-	-	+	igM, λ	yes	34	73	A ⁰
14 LT	RT-PCR	m	47	leg	+	+	+	-	+	igM, igD, λ	no	43	49	A ⁰
15 LT	RT-PCR	f	75	leg	+	+	+	-	-	κ ^a	yes	2	29	D ⁺
16 LT	RT-PCR	f	77	leg	+	+	-	-	+	igM, λ	yes	10	19	D ⁺
17 LT	RT-PCR	f	75	both legs	+	+	-	-	-	igM, igD, κ	yes	1	12	D ⁺
18 LT	RT-PCR	f	84	leg	+	+	+	-	+	igM, igD, κ	no	0	3	D ⁺

Legend to Table 1.: FC: primary cutaneous follicle center lymphoma; PCLBCL-LT: primary cutaneous large B-cell lymphoma, leg type; BCR: B-cell receptor; PFS: progression-free survival; f: female; HTS: high-throughput sequencing; RT-PCR: Real Time polymerase chain reaction; m: male; NA: not assessed; RT: radiotherapy; o.p.: oral prednisone; PCT: polychemotherapy; A⁺: alive with disease; A⁰: alive without disease; D⁰: death with disease; D⁺: death without disease; ^a heavy chains not assessed.

Results

High-throughput sequencing

From 19 tumor samples (13 samples of PCLBCL-LT and 6 samples of PCFCL) and 4 samples of activated B-cells, microRNA libraries were generated and sequenced. Of these, two PCLBCL-LT samples yielded less than 100,000 total microRNA reads (7,302 and 55,383 reads) and were discarded from further analyses.²⁷ Mean microRNA read number in the other samples was 4,139,515 (standard deviation 9,348,501). After alignment of the sequence reads in miRDeep 2.0, 1921 different known precursor microRNAs were identified to be expressed in at least one sample. 328 precursor loci were identified as putative novel microRNAs. After discarding precursor microRNAs that showed low expression levels (see Materials and Methods), a total number of 238 different precursor microRNAs were included for further analysis.

MicroRNA profiles of PCFCL and PCLBCL-LT

The most abundantly expressed microRNAs of the analyzed subgroups are represented in Figure 1. In PCLBCL-LT, miR-19b-3p and miR-19a-3p show the highest expression levels, and comprise 12% and 11% of the total microRNome, respectively. In PCFCL, these microRNAs also constitute a substantial part of the microRNome (7% and 6%, respectively), but miR-150-5p is most abundantly expressed. The activated B-cells are to a large extent represented by miR-150-5p, comprising approximately one-fifth of the microRNome. Using NormFinder, miR-148b-3p, miR-25-3p, let-7c and let-7e-5p were the most stably expressed microRNAs in our complete study cohort (Supplemental Figure 1).

Unsupervised hierarchical clustering analysis of all samples using the 238 mature microRNAs of the precursor sequences identified did not result into distinct clustering of the two tumor types and/or the activated B-cells (Supplemental Figure 2).

Differential microRNA expression in PCLBCLs

Table 2 shows an overview of mature microRNAs that showed statistically significantly differential expression between PCFCL and PCLBCL-LT. Under the given conditions (allowing a mismatch of 1 base pair), MiRDeep 2.0 did not categorize miR-129-1-3p and miR-129-2-3p as separate microRNAs. Recounting of both variants in the aligned sequence data derived from miRDeep 2.0 showed a markedly higher expression of miR-129-2-3p in PCFCL as compared to PCLBCL-LT. Similarly, miRDeep 2.0 was not able to differentiate between miR-9-1, -2 and -3 variants, because of the relatively short aligned sequences (no sequences containing (part of) the precursor microRNA outside the mature microRNA region). For the 16 differentially expressed microRNAs, unsupervised hierarchical clustering analysis was applied (Figure 2). According to this analysis, two groups are separated from each other, one containing all cases of PCFCL and two cases of PCLBCL-LT, the other group comprising of the remaining cases of PCLBCL-LT. The two cases of PCLBCL-LT clustering with the PCFCL cases showed a clinical presentation, tumor cell morphology and immunohistochemical profile with classical features of PCLBCL-LT. One of the patients received polychemotherapy combined with radiotherapy and died of lymphoma after 38 months; the other patient was solely treated with radiotherapy and is still alive without disease recurrence after 50 months of follow-up.

Table 2. High-throughput sequencing: differential expression of microRNAs between PCFCL and PCLBCL-LT

MicroRNA		Reads per million, mean (SD)		Adjusted <i>p</i> -value ^a
		PCFCL	PCLBCL-LT	
miR-129-2-3p	^b	1.668 (3.260)	4 (8)	5.74E-15
miR-129-1-3p		184 (269)	18 (21)	2.10E-14
miR-1246		0 (0)	26 (45)	1.42E-5
miR-214-3p	^b	8.620 (14.534)	1.174 (1.232)	7.24E-4
miR-574-3p	^b	12.079 (14.180)	1.951 (1.392)	0.00266
miR-4485		6 (8)	77 (127)	0.00321
miR-222-5p		16 (14)	142 (126)	0.0108
miR-31-5p	^b	1.729 (995)	346 (385)	0.0129
miR-486-3p	^b	196 (194)	40 (43)	0.0133
miR-363-3p	^b	37 (27)	288 (535)	0.0153
miR-99a-5p	^b	9.843 (9.113)	2.168 (1.155)	0.0184
miR-574-5p		289 (184)	1.996 (5.654)	0.0442
miR-100-5p	^b	8.849 (6.850)	2.318 (1.400)	0.0442
miR-205-5p	^b	11.492 (12.787)	3.059 (2.818)	0.0442
miR-342-3p	^b	22.775 (24.636)	6.079 (3.992)	0.0442
miR-9-5p	^b	273 (309)	73 (74)	0.0442

^a *p*-values Benjamini-Hochberg corrected; ^b microRNA included in RT-qPCR validation; PCFCL: primary cutaneous follicle center lymphoma, PCLBCL-LT: primary cutaneous large B-cell lymphoma, leg type.

MicroRNA RT-qPCR validation

A selection of 11 differentially expressed microRNAs (marked in Table 3) was used for validation in a novel group of test cases. MiR-129-1-3p, miR-1246, miR-4485 and miR-222-5p were not validated because of the relatively low counts in both PCFCL and PCLBCL-LT samples. Of miR-574-5p, no Taqman PCR primer assay was available, because of dinucleotide repeats in the microRNA sequence that could interfere with the PCR. Assays for miR-148b-3p, miR-25-3p, let-7c and let-7e-5p were included as reference microRNAs, along with widely used reference RNAs U6 and RNU48. Of these six, miR-148b-3p, let-7e-5p and miR-25-3p were most stably expressed in the RT-qPCR experiments (according to GeNorm) and therefore used for normalization of the data. The expression of U6 and RNU48 was very variable in this sample set. Comparison of RT-qPCR performed on both RNA of frozen tumor material isolated for high-throughput sequence analysis and RNA isolated from FFPE sections of the same tumor sample (2 cases of PCFCL and 3 cases of PCLBCL-LT) showed high concordance (Supplemental Figure 3). With inclusion of all 20 FFPE PCLBCL samples in the analysis, including 10 PCFCL and 10 PCLBCL-LT cases, we confirmed higher expression of miR-129-2-3p, miR-214-3p, miR-31-5p and miR-9-5p in PCFCL as compared to PCLBCL-LT ($p < 0.05$) (Figure 3). For the other seven microRNAs, no statistically significantly differential expression was found between PCFCL and PCLBCL-LT. In addition to these differentially expressed microRNAs resulting from the deep sequence analysis, we tested whether four microRNAs frequently reported to show higher expression in ABC-type compared GCB-type nodal DLBCL (miR-21-5p, miR-155-5p, miR-221-3p and miR-222-3p) were higher expressed in PCLBCL-LT.

Figure 1. Most abundantly expressed microRNAs





Figure 2. Unsupervised hierarchical clustering analysis in PCLBCL samples of high-throughput sequencing, using 16 differentially expressed microRNAs



PCFCL and PCLBCL-LT tend to cluster together in two subgroups, with only 2 cases of PCLBCL-LT clustering together with the PCFCL cases.

However, no differential expression between PCLBCL-LT and PCFCL of these four microRNAs was observed (Figure 4). Above that, unsupervised hierarchical clustering analysis of the high-throughput sequence data of our PCLBCLs using microRNA signature profiles that should be able to distinguish GCB-type from ABC-type DLBCL^{12,14,17} did not set apart PCFCL from PCLBCL-LT.

MicroRNA RT-qPCR analysis of GCB- and ABC-subtypes of nodal DLBCLs

Because of the inconsistencies in literature concerning microRNA profiling of nodal DLBCL and the varying references used in these studies, a set of well-defined GCB- and ABC-subtype nodal DLBCLs was subjected to microRNA RT-qPCR analysis, using the same conditions as for our PCLBCL FFPE samples. As in PCLBCL, GeNorm analysis showed that miR-148b-3p, miR-25-3p, let-7e-5p were most stably expressed, and these three microRNAs were therefore used as reference microRNAs. We evaluated the expression of the four microRNAs that showed significantly higher expression in PCFCL as compared to PCLBCL-LT in our nodal DLBCL samples. The expression of miR-129-2-3p was indeed higher in GCB-type DLBCL as compared to the ABC-type ($p=0.00$), while the expression of miR-214-3p, miR-31-5p and miR-9-5p did not show statistically significant differences between both nodal subtypes. Four microRNAs frequently reported to show higher expression in ABC-type than in GCB-type nodal DLBCL (miR-21-5p, miR-155-5p, miR-221-3p and miR-222-3p) were included in the RT-qPCR analy-

sis. However, in our group of nodal DLBCLs, of these four microRNAs, we could only confirm higher expression levels of miR-155-5p ($p=0.01$) and miR-222-5p ($p=0.05$) in ABC-type as compared to GCB-type DLBCL (Figure 4).

Table 3. MicroRNAs in RT-qPCR validation: predicted targets

MicroRNA	Location ^a	Up ^b	Potential target(s) and function	
miR-129-2-3p	11p11.2	PCFCL	SOX4 CDK6	(B-cell) transcription factor regulator of cell cycle progression
miR-214-3p	1q24.3	PCFCL	PTEN EZH2	negative regulator of cell cycle progression polycomb group gene silencer
miR-574-3p	4p14	PCFCL	IL6	involved in NF- κ B signaling
miR-31-5p	9p21.3	PCFCL	PIK3C2A CXCL12	PI3K pathway component chemokine attractant
			RhoA	metastasis promoting gene
			NIK	NF- κ B signaling
miR-486-3p	8p11.21	PCFCL	PTEN E2F1	negative regulator of cell cycle progression regulator of cell cycle progression
miR-99a-5p	21q21.1	PCFCL	mTOR	PI3K family member
miR-100-5p	11q24.1	PCFCL	mTOR	PI3K family member
miR-205-5p	1q32.2	PCFCL	PTEN E2F1	negative regulator of cell cycle progression regulator of cell cycle progression
miR-342-3p	14q32.2	PCFCL	BCL2L10	BCL2 family member
miR-9-5p	1q22/5q14.3/15q26.1	PCFCL	PRDM1/BLIMP1 FOXP1	B-cell transcription factor (B-cell) transcription factor
			NF κ B1 = p50	NF- κ B signaling
miR-363-3p	Xq26.2	PCLBCL-LT	CDKN1A = p21	regulator of cell cycle progression


^a chromosomal location microRNA gene; ^b higher expression in subgroup of PCLBCL; PCFCL: primary cutaneous follicle center lymphoma; PCLBCL-LT: primary cutaneous diffuse large B-cell lymphoma, leg type.

Discussion

This is the first study to report complete genome-wide microRNA profiles of PCLBCLs, and comparing PCFCL and PCLBCL-LT subtypes. By use of next generation high-throughput microRNA sequence analysis, we provided microRNA profiles of these tumors on frozen biopsy samples, accurately exploring both quantitative and qualitative aspects of all microRNAs. The complete microRNomes of PCLBCL-LT and PCFCL did not allow their separation by unsupervised hierarchical clustering, and PCLBCL-LT did not cluster together with activated B-cells, its postulated normal counterpart (Supplemental Figure 1). The list of expressed and quantified microRNAs generated from this experiment was the basis for selecting specific microRNAs for RT-qPCR confirmation on FFPE tumor tissue of both lymphoma subtypes. The use of high-throughput

sequencing also enabled us to identify 328 putative novel microRNAs expressed in our samples, of which further validation was not performed and is warranted. Furthermore, these sequence data were also instrumental in selecting stably expressed reference microRNAs for subsequent confirmation experiments. We identified four microRNAs (miR-129-2-3p, miR-214-3p, miR-31-5p and miR-9-5p) of which statistically significantly higher expression in PCFCL as compared to PCLBCL-LT in high-throughput sequencing could be confirmed by RT-qPCR on a new FFPE study cohort.

Figure 3. RT-qPCR expression in FFPE samples (1)



The expression of the 4 microRNAs showing statistically significantly higher expression in PCFCL as compared to PCLBCL-LT are represented in dot plots. ΔΔCq was calculated with miR-148b-3p, let-7e-5p and miR-25-3p as stably expressed reference microRNAs. *P-value < 0.05 (Mann Whitney U test).

MicroRNomes of PCFCL and PCLBCL-LT

Considering all samples analyzed, miR-19a-3p and miR-19b-3p constituted a significant part of the microRNA profile, up to 12% for miR-19b-3p of the complete profile of PCLBCL-LT. Both microRNAs are members of the miR-17~92 cluster, the presence of which is critical for normal B-cell development.³³ It also acts as a potential oncogenic cluster in B-cell non-Hodgkin lymphomas,³⁴ with miR-19 components being essential in this oncogenic potential.³⁵ The other members of this cluster, miR-17-5p, miR-18a-5p, miR-20a-5p and miR-92a-3p, only

made up a very small proportion of the complete microRNome (with a maximum of 1%). The relatively low expression of miR-92a-3p was most remarkable, as this microRNA was found to be one of the most abundantly expressed microRNA in a recent study on nodal DLBCL and primary central nervous system DLBCL, an extranodal DLBCL with an ABC-like genotype.³⁶ Although in PCFCL, miR-19a-3p and miR-19b-3p also made up a substantial part of the microRNome, miR-150-5p was most abundantly expressed. MiR-150-5p is involved in normal B-cell development, targeting transcription factor and proto-oncogene c-MYB.³⁷

Differentially expressed microRNAs

By performing RT-qPCR in an independent cohort of FFPE tumor samples, the higher expression of miR-129-2-3p, miR-214-3p, miR-31-5p and miR-9-5p in PCFCL as compared to PCLBCL-LT in high-throughput sequence data could be confirmed.

MiR-129-2-3p targets transcription factor SOX4, which was validated in endometrial cancer.³⁸ SOX4 also has a critical function in normal B-cell development.³⁹ Because of the dichotomy in expression of miR-129-2-3p (11 cases without and 9 cases with expression of this microRNA in RT-qPCR), immunohistochemical staining for SOX4 was performed. However, no inverse correlation between RT-qPCR expression of miR-129-2-3p and immunohistochemical expression of SOX4 was observed (data not shown).


A previous array-based comparative genomic hybridization study performed by our group, showed low copy number gains of 1q23-q25 in PCFCL, but not in PCLBCL-LT cases.⁵ The miR-214 gene is located in this chromosomal region and correspondingly, higher levels of miR-214-3p were observed in PCFCL compared to PCLBCL-LT. MiR-214-3p has been reported as oncogenic microRNA in T-cells, negatively regulating tumor suppressor gene phosphatase and tensin homolog (PTEN),⁴⁰ and is involved in controlling cell cycle regulation. However, miR-214-3p has also been described to exert a tumor suppressor function in breast cancer cell lines, where reduced levels of this microRNA result in increased proliferation and invasion and in accumulation of EZH2,⁴¹ a polycomb group protein with the potential to promote cell proliferation.⁴²

The miR-31 gene is located on 9p21.3. This chromosomal region is often targeted by mono-allelic or biallelic deletions in PCLBCL-LT, but not in PCFCL.^{4,43} In accordance, the expression of miR-31-5p was lower in PCLBCL-LT in proportion to PCFCL. Correlation of previous multiplex ligation probe-dependent assay results of chromosomal location 9p21.3⁴³ to the current results showed that two of the PCLBCL-LT samples tested here had deletions in this region and low expression of miR-31-5p, while four samples did not have deletions and showed higher expression of this specific microRNA. Well characterized genes involved in this deletion are CDKN2A and CDKN2B, tumor suppressor genes involved in cell cycle control.⁴⁴ We previously reported that inactivation of CDKN2A in PCLBCL-LT is associated with inferior survival.^{4,43} It is however not clear whether the miR-31 gene is also part of the 9p21.3 deletion involved in this more aggressive behavior. Downregulation of miR-31-5p has been described in different types of T-cell non-Hodgkin lymphomas.⁴⁵ In these lymphomas, miR-31-5p was shown to be associated with activation of the nuclear factor (NF)-κB pathway, a signaling pathway often deregulated in several types of malignancies, by targeting NF-κB inducing kinase.⁴⁵

MiR-9-5p is involved in normal B-cell development, negatively regulating transcription factor PRDM1, an essential driver of plasma cell differentiation. In accordance, this microRNA

is more abundantly expressed in normal germinal center B-cells than plasma cells.²¹ As PCLBCL-LT is thought to be derived from B-cells in a stage just beyond the germinal center reaction, tending towards plasma cell differentiation, the lower expression of miR-9-5p might reflect a normal biological stage in B-cell maturation. In ovarian cancer, miR-9-5p has been demonstrated to act as a tumor suppressor microRNA by targeting the NF- κ B family member NF κ B1/p50.⁴⁶ Since the NF- κ B pathway has a well-known oncogenic potential in ABC-type nodal DLBCL,⁴⁷ it can be speculated that downregulation of miR-9-5p might have a pathogenic effect in PCLBCL-LT through reduced inhibition of NF- κ B signaling.

Figure 4. RT-qPCR expression in FFPE samples (2)



The expression of the 4 microRNAs that are repeatedly reported in previous studies to be more abundantly expressed in ABC-type than GCB-type nodal DLBCL are represented in dot plots. $\Delta\Delta Cq$ was calculated with miR-148b-3p, let-7e-5p and miR-25-3p as stably expressed reference microRNAs. *P-value < 0.05; ** P-value = 0.05 (Mann Whitney U test).

Correlation with nodal DLBCL subtypes

PCFCL and PCLBCL-LT have a very distinct gene expression profile, corresponding with gene expression profiles of GCB- and ABC-type nodal DLBCL.⁵ We therefore were interested in comparing both subtypes of PCLBCL with their molecular counterparts of nodal DLBCL. However, literature concerning microRNA expression in nodal DLBCL shows inconsistent results. Taken together, the most frequently reported microRNAs showing higher expression

in ABC-type as compared to GCB-type nodal DLBCL are miR-155-5p, miR-21-5p and miR-221-3p and miR-222-3p.^{11,12,15-19,48} In our PCLBCL samples, expression of these four microRNAs did not show significant differences between the two subtypes in both high-throughput sequencing and RT-qPCR. To accurately compare our PCLBCL samples with nodal DLBCL, and exclude external influences such as differing reference microRNAs, we performed RT-qPCR on well-defined subgroups of nodal DLBCL by using a set of three microRNAs (miR-148b-3p, let-7e-5p and miR-25-3p) that were stably expressed in our PCLBCL sequencing studies as a reference. These microRNAs were, in contrast to widely used references U6 and RNU48, stably expressed in our nodal DLBCLs. By using this approach of matching conditions, we confirmed higher expression of miR-155-5p and miR-222-3p in ABC-type as compared to GCB-type nodal DLBCL, but not of miR-21-5p and miR-221-3p.

The absence of higher expression of miR-155-5p in PCLBCL-LT as compared to PCFCL was most noteworthy. The role miR-155-5p has been studied extensively in nodal B-cell non-Hodgkin lymphomas and is linked to activation of the intracellular signaling pathway PI3K-Akt pathway¹⁰ and increased lymphoma cell motility,⁴⁹ and its expression is associated with NF- κ B activity in nodal DLBCL.⁴⁸ The finding that miR-155-5p was not elevated in PCLBCL-LT as compared to PCFCL suggests at least partially different pathogenetic mechanisms in PCLBCL-LT as compared to its nodal molecular counterpart.

In summary, we presented the first microRNA profiling results by next generation high-throughput sequencing and RT-qPCR comparing PCFCL and PCLBCL-LT. These two subtypes of PCLBCL did not show entirely distinct microRNA profiles, but could be distinguished by differential expression of microRNAs miR-129-2-3p, miR-214-3p, miR-31-5p and miR-9-5p. We also showed that the hitherto described microRNA signature of GCB- and ABC-subtype nodal DLBCLs could not be extrapolated to PCFCL and PCLBCL-LT. Therefore, we conclude that although PCFCL and PCLBCL-LT show strong resemblance at gene expression level with GCB- and ABC-type nodal DLBCL, respectively, their microRNA profiles are at least partially disparate from their nodal counterparts, suggesting distinct pathogenetic mechanisms.

References

1. Willemze R, Jaffe ES, Burg G, et al. (2005) WHO-EORTC classification for cutaneous lymphomas. *Blood* 105: 3768-85.
2. Swerdlow SH, Campo E, Harris NL, et al. (2008) WHO Classification of Tumours Haematopoietic and Lymphoid Tissues. Geneva, Switzerland: WHO PRESS.
3. Senff NJ, Hoefnagel JJ, Jansen PM, et al. (2007) Reclassification of 300 primary cutaneous B-Cell lymphomas according to the new WHO-EORTC classification for cutaneous lymphomas: comparison with previous classifications and identification of prognostic markers. *J Clin Oncol* 25: 1581-7.
4. Dijkman R, Tensen CP, Jordanova ES, et al. (2006) Array-based comparative genomic hybridization analysis reveals recurrent chromosomal alterations and prognostic parameters in primary cutaneous large B-cell lymphoma. *J Clin Oncol* 24: 296-305.
5. Hoefnagel JJ, Dijkman R, Basso K, et al. (2005) Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. *Blood* 105: 3671-8.
6. Senff NJ, Noordijk EM, Kim YH, et al. (2008) European Organization for Research and Treatment of Cancer and International Society for Cutaneous Lymphoma consensus recommendations for the management of cutaneous B-cell lymphomas. *Blood* 112: 1600-9.
7. Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116: 281-97.
8. Garzon R, Croce CM (2008) MicroRNAs in normal and malignant hematopoiesis. *Curr Opin Hematol* 15: 352-8.
9. Monsalvez V, Montes-Moreno S, Artiga MJ, et al. (2013) MicroRNAs as prognostic markers in indolent primary cutaneous B-cell lymphoma. *Mod Pathol* 26: 171-81.
10. Huang X, Shen Y, Liu M, et al. (2012) Quantitative proteomics reveals that miR-155 regulates the PI3K-AKT pathway in diffuse large B-cell lymphoma. *Am J Pathol* 181: 26-33.
11. Zhong H, Xu L, et al. (2012) Clinical and prognostic significance of miR-155 and miR-146a expression levels in formalin-fixed/paraffin-embedded tissue of patients with diffuse large B-cell lymphoma. *Exp Ther Med* 3: 763-70.
12. Montes-Moreno S, Martinez N, Sanchez-Espiridion B, et al. (2011) miRNA expression in diffuse large B-cell lymphoma treated with chemoimmunotherapy. *Blood* 118: 1034-40.
13. Jima DD, Zhang J, Jacobs C, et al. (2010) Deep sequencing of the small RNA transcriptome of normal and malignant human B cells identifies hundreds of novel microRNAs. *Blood* 116: e118-e127.
14. Culpin RE, Proctor SJ, Angus B, et al. (2010) A 9 series microRNA signature differentiates between germinal centre and activated B-cell-like diffuse large B-cell lymphoma cell lines. *Int J Oncol* 37: 367-76.
15. Malumbres R, Sarosiek KA, Cubedo E, et al. (2009) Differentiation stage-specific expression of microRNAs in B lymphocytes and diffuse large B-cell lymphomas. *Blood* 113: 3754-64.
16. Roehle A, Hoefig KP, Reipsilber D, et al. (2008) MicroRNA signatures characterize diffuse large B-cell lymphomas and follicular lymphomas. *Br J Haematol* 142: 732-44.
17. Lawrie CH, Chi J, Taylor S, et al. (2009) Expression of microRNAs in diffuse large B cell lymphoma is associated with immunophenotype, survival and transformation from follicular lymphoma. *J Cell Mol Med* 13: 1248-1260.
18. Lawrie CH, Soneji S, Marafioti T, et al. (2007) MicroRNA expression distinguishes between germinal center B cell-like and activated B cell-like subtypes of diffuse large B cell lymphoma. *Int J Cancer* 121: 1156-61.
19. Eis PS, Tam W, Sun L, et al. (2005) Accumulation of miR-155 and BIC RNA in human B cell lymphomas. *Proc Natl Acad Sci USA* 102: 3627-32.
20. Alencar AJ, Malumbres R, Kozloski GA, et al. (2011) MicroRNAs are independent predictors of outcome in diffuse

large B-cell lymphoma patients treated with R-CHOP. *Clin Cancer Res* 17: 4125-35.

21. Zhang J, Jima DD, Jacobs C, et al. (2009) Patterns of microRNA expression characterize stages of human B-cell differentiation. *Blood* 113: 4586-94.

22. Meijer CJLM, Vergier B, Duncan LM, Willemze R (2008) Primary cutaneous DLBCL, leg type. In: Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri S et al., editors. WHO classification of tumours of haematopoietic and lymphoid tissues. Lyon: IARC Press 242.

23. Hans CP, Weisenburger DD, Greiner TC, et al. (2004) Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood* 103: 275-82.

24. Choi WW, Weisenburger DD, Greiner TC, et al. (2009) A new immunostain algorithm classifies diffuse large B-cell lymphoma into molecular subtypes with high accuracy. *Clin Cancer Res* 15: 5494-502.

25. Qin Y, Buermans HP, van Kester MS, et al. (2012) Deep-sequencing analysis reveals that the miR-199a2/214 cluster within DNM3os represents the vast majority of aberrantly expressed microRNAs in Sezary syndrome. *J Invest Dermatol* 132: 1520-2.

26. Friedlander MR, Chen W, Adamidi C, et al. (2008) Discovering microRNAs from deep sequencing data using miRDeep. *Nat Biotechnol* 26: 407-15.

27. Volinia S, Galasso M, Sana ME, et al. (2012) Breast cancer signatures for invasiveness and prognosis defined by deep sequencing of microRNA. *Proc Natl Acad Sci USA* 109: 3024-9.

28. Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26: 139-40.

29. Benjamini YH, Y. (1995) Controlling the False Discovery Rate: A practical and powerful approach to multiple testing. *J R Statist Soc B* 57: 289-300.

30. Andersen CL, Jensen JL, Orntoft TF (2004) Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* 64: 5245-50.

31. Mestdagh P, Van Vlierberghe P, De Weer A, et al. (2009) A novel and universal method for microRNA RT-qPCR data normalization. *Genome Biol* 10: R64.

32. Vandesompele J, De Preter K, Pattyn F, et al. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034.

33. Ventura A, Young AG, Winslow MM, et al. (2008) Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family of miRNA clusters. *Cell* 132: 875-86.

34. He L, Thomson JM, et al. (2005) A microRNA polycistron as a potential human oncogene. *Nature* 435: 828-33.

35. Olive V, Bennett MJ, et al. (2009) miR19 is a key oncogenic component of mir17-92. *Genes Dev* 23: 2839-49.

36. Fischer L, Hummel M, Korfel A, et al. (2011) Differential micro-RNA expression in primary CNS and nodal diffuse large B-cell lymphomas. *Neuro Oncol* 13: 1090-8.

37. Xiao C, Calado DP, Galler G, et al. (2007) MiR-150 controls B cell differentiation by targeting the transcription factor c-Myb. *Cell* 131: 146-59.

38. Huang YW, Liu JC, Deatherage DE, et al. (2009) Epigenetic repression of microRNA-129-2 leads to overexpression of SOX4 oncogene in endometrial cancer. *Cancer Res* 69: 9038-46.

39. Smith E, Sigvardsson M (2004) The roles of transcription factors in B lymphocyte commitment, development, and transformation. *J Leukoc Biol* 75: 973-81.

40. Jindra PT, Bagley J, Godwin JG, Iacomini J (2010) Costimulation-dependent expression of microRNA-214 increases

the ability of T cells to proliferate by targeting Pten. *J Immunol* 185: 990-7.

41. Derfoul A, Juan AH, Difilippantonio MJ, et al. (2011) Decreased microRNA-214 levels in breast cancer cells coincides with increased cell proliferation, invasion and accumulation of the Polycomb Ezh2 methyltransferase. *Carcinogenesis* 32: 1607-14.

42. Muller H, Bracken AP, Vernell R, et al. (2001) E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis. *Genes Dev* 15: 267-85.

43. Senff NJ, Zoutman WH, Vermeer MH, et al. (2009) Fine-mapping chromosomal loss at 9p21: correlation with prognosis in primary cutaneous diffuse large B-cell lymphoma, leg type. *J Invest Dermatol* 129: 1149-55.

44. Kim WY, Sharpless NE (2006) The regulation of INK4/ARF in cancer and aging. *Cell* 127: 265-75.

45. Yamagishi M, Nakano K, Miyake A, et al. (2012) Polycomb-mediated loss of miR-31 activates NIK-dependent NF-kappaB pathway in adult T cell leukemia and other cancers. *Cancer Cell* 21: 121-35.

46. Guo LM, Pu Y, Han Z, et al. (2009) MicroRNA-9 inhibits ovarian cancer cell growth through regulation of NF-kappaB1. *FEBS J* 276: 5537-46.

47. Davis RE, Brown KD, Siebenlist U, Staudt LM (2001) Constitutive nuclear factor kappaB activity is required for survival of activated B cell-like diffuse large B cell lymphoma cells. *J Exp Med* 194: 1861-74.

48. Rai D, Karanti S, Jung I, Dahia PL, Aguiar RC (2008) Coordinated expression of microRNA-155 and predicted target genes in diffuse large B-cell lymphoma. *Cancer Genet Cytogenet* 181: 8-15.

49. Dagan LN, Jiang X, Bhatt S, et al. (2012) miR-155 regulates HGAL expression and increases lymphoma cell motility. *Blood* 119: 513-20.

Supplemental Materials and Methods

B-cell selection and activation

Human B-cells were purified from the separate lymphocyte fractions of peripheral blood (buffy coats) obtained from four healthy individuals by positive selection with magnetic CD19 MicroBeads (Miltenyi Biotec, Bergisch Gladbach, Germany). The B-cells were cultured in RPMI 1640 with 10% FCS and incubated with CpG (5 µg/ml) (ODN 2006, Invivogen, San Diego, USA) and monoclonal antibodies to CD40 (0.1 µg/ml) (BD Biosciences Pharmingen, Heidelberg, Germany) for 24 hours. After harvesting, a B-cell purity of more than 97% was confirmed by flow cytometric analysis with a mixture of monoclonal antibodies against B-cell (CD20, APC-H7), T-cell (CD3, PE-H7) and monocytes (CD14, APC). Simultaneously, the activation status was determined using anti-CD80 (PE) and anti-CD86 (FITC), respectively. All antibodies were obtained from BD Biosciences. Dead cells were excluded using 1 µM of DAPI. A BD Biosciences LSRII was used for acquisition. WinList 7 (Verity Software House, Topsham, ME, USA) was used for data analysis. Approximately 86% of the lymphocytes showed expression of these activation markers.


MicroRNA library preparation for high-throughput sequencing

Sequencing adaptors were ligated to total RNA on both ends of the RNA molecules and reverse-transcribed. The cDNA was pre-amplified with PCR primers containing specific sequence tags making the libraries compatible with the Illumina flow cells (Illumina, Son en Breugel, The Netherlands). Polyacrylamide *gel* electrophoresis size selection was performed, excising the 95-105 bp band containing the adaptor-ligated microRNA insert. Quantification and size verification was performed using a high sensitivity DNA chip (Agilent, Amstelveen, The Netherlands).


MicroRNA real-time qPCR

Total RNA was isolated from 10 µm sections of FFPE tumor samples (number of sections varying from 4 to 20 sections, according to the size of the specimen) using the RecoverAll Total Nucleic Acid Extraction Kit according to manufacturer's protocol (Ambion, Warrington, UK). Approximately 300 ng of total RNA from each sample (FFPE or frozen, same isolation as for the high-throughput sequencing) was reverse transcribed using the microRNA reverse transcription kit (Applied Biosystems) combined with the stem-loop Megaplex primer pool A v2.1 (Applied Biosystems), which allowed for simultaneous reverse transcription of 377 microRNAs and 4 endogenous controls. The reverse transcription was conducted on the LightCycler480 (Roche, Almere, The Netherlands), running 40 cycles with the following conditions: 2 minutes at 16°C, 1 minutes at 42°C and 1 second at 50°C, followed by 5 minutes holding at 85°C. RT-qPCR was performed using Taqman microRNA and control assays and 2x Universal PCR mastermix (Applied Biosystems). The reactions were run in duplicate on the CFX384 RT-qPCR Detection System, according to manufacturer's protocol (Bio-Rad Laboratories, Veenendaal, the Netherlands), with the following cycle parameters: 10 min at 95 °C, followed by 50 cycles denaturing for 15 s at 95 °C and annealing and extending for 60 s at 60 °C.


Supplemental Figure 1. Most stably expressed microRNAs in high-throughput sequencing according to GeNorm



Supplemental Figure 2: Unsupervised hierarchical clustering of the normalized expression of all analyzed microRNAs of high-throughput sequencing



Supplemental Figure 3. Internal validation



Supplemental Table 1. Complete microRNA profiling of nodal DLBCL: differences between ABC- and GCB-type nodal DLBCL

Author	Journal	Year	# pt	Technique	MicroRNA ↑ in ABC- vs GCB-DLBCL	MicroRNA ↑ in GCB- vs ABC-DLBCL	Subtyping DLBCLs
Lawrie	Int J Cancer	2007	5*	miRMAX microarray (252 microRNAs)	<u>miR-155</u> , <u>miR-21</u> , <u>miR-222</u>	none	established cell lines
Roehle	Br J Haematol	2008	58	qPCR assay (157 microRNAs)	<u>miR-155</u>	miR-27b, miR-28, miR-129, miR-133a, miR-133b, miR-138, miR-151	IHC: Hans
Lawrie†	J Cell Mol Med	2009	64	custom-made microarray (464 microRNAs)	miR-518a, miR-363, miR-21, miR-132, miR-340, miR-301, miR-30d, <u>miR-221</u> , miR-422b, miR-146b, <u>miR-155</u> , miR-190, miR-194, miR-660, miR-213, <u>miR-222</u> , miR-186	miR-620, miR-616, miR-199b, miR-421, miR-569, miR-653, miR-138, miR-520h, miR-129	IHC: Hans
Malumbrest†	Blood	2009	8*	LC Sciences microarray (711 microRNAs)	miR-146b-5p, miR-146a, miR-21, <u>miR-155</u> , miR-500, miR-222, miR-363, miR-574-5p, miR-574-3p	none	established cell lines
Zhang	Blood	2009	40	Exiqon microarray (789 microRNAs)	<i>(up- or downregulation not stated)</i> miR-142-3p, miR-16, miR-184, miR-191, miR-19a, miR-19b, miR-299-5p, miR-32, miR-30e*, miR-151-5p, miR-583, mghv-miR-M1-7-5p, miR-142-5p, miR-106b, miR-30e, miR-140-3p, miR-20a, miR-526b*, miR-28-5p, miR-30c		IHC: Hans
Culpin	Int J Oncol	2010	8*	Miltenyi Biotec microarray (1860 microRNAs)	miR-17, miR-19b, miR-20a, miR-29a, none	miR-28-5p, miR-30c	IHC: Hans
Montes-Moreno	Blood	2011	29	Agilent microarray (470 microRNAs)	miR-92a, miR-106a, miR-720, miR-1260, miR-1280	miR-17, miR-19b, miR-20a, miR-29a, none	IHC: Choi

* cell lines; † study generated a distinct microRNA subset separating ABC- from GCB-DLBCL by unsupervised hierarchical clustering analysis; # pt: number of patients; IHC: immunohistochemistry; ABC: activated B-cell; GCB: germinal center B-cell; DLBCL: diffuse large B-cell lymphoma, qPCR: quantitative polymerase chain reaction; NGS: next generation sequencing.

Supplemental Table 2. 238 microRNAs with sufficiently high enough read counts in next generation sequencing to be included in further analysis

let-7a-5p	miR-1275	miR-152	miR-195-5p	miR-223-5p	miR-31-5p	miR-378c	miR-574-3p
let-7b-5p	miR-128	miR-155-3p	miR-197-3p	miR-224-5p	miR-32-5p	miR-378d	miR-574-5p
let-7c	miR-1280	miR-155-5p	miR-199a-3p	miR-23a-3p	miR-320a	miR-378f	miR-577
let-7d-3p	miR-129-1-3p	miR-15a-3p	miR-199a-5p	miR-23b-3p	miR-320b	miR-378g	miR-582-5p
let-7d-5p	miR-129-2-3p	miR-15a-5p	miR-199b-3p	miR-23c	miR-324-5p	miR-378i	miR-590-3p
let-7e-5p	miR-1306-5p	miR-15b-3p	miR-199b-5p	miR-24-2-5p	miR-326	miR-423-3p	miR-590-5p
let-7f-5p	miR-1307-3p	miR-15b-5p	miR-19a-3p	miR-24-3p	miR-328	miR-423-5p	miR-598
let-7g-3p	miR-1307-5p	miR-16-2-3p	miR-19b-3p	miR-25-3p	miR-330-3p	miR-424-5p	miR-625-3p
let-7g-5p	miR-130a-3p	miR-16-5p	miR-200a-3p	miR-26a-5p	miR-331-3p	miR-425-3p	miR-652-3p
let-7i-3p	miR-130b-3p	miR-17-3p	miR-200b-3p	miR-26b-5p	miR-338-3p	miR-425-5p	miR-652-5p
let-7i-5p	miR-130b-5p	miR-17-5p	miR-200c-3p	miR-27a-3p	miR-339-3p	miR-4286	miR-660-5p
miR-100-5p	miR-138-5p	miR-181a-2-3p	miR-203	miR-27b-3p	miR-339-5p	miR-4485	miR-664-3p
miR-101-3p	miR-139-5p	miR-181a-3p	miR-205-5p	miR-28-3p	miR-33a-5p	miR-451a	miR-7-1-3p
miR-103a-3p	miR-140-3p	miR-181a-5p	miR-20a-3p	miR-28-5p	miR-342-3p	miR-4521	miR-7-5p
miR-106a-5p	miR-140-5p	miR-181b-5p	miR-20a-5p	miR-296-5p	miR-342-5p	miR-454-3p	miR-720
miR-106b-3p	miR-141-3p	miR-181d	miR-20b-5p	miR-29a-3p	miR-345-5p	miR-455-3p	miR-744-5p
miR-106b-5p	miR-142-3p	miR-182-5p	miR-21-3p	miR-29a-5p	miR-34a-5p	miR-484	miR-766-3p
miR-107	miR-142-5p	miR-183-5p	miR-21-5p	miR-29b-2-5p	miR-361-3p	miR-486-3p	miR-874
miR-10a-5p	miR-143-3p	miR-185-5p	miR-210	miR-29b-3p	miR-361-5p	miR-486-5p	miR-9-5p
miR-10b-5p	miR-144-3p	miR-186-5p	miR-214-3p	miR-29c-3p	miR-3613-3p	miR-497-5p	miR-92a-1-5p
miR-1246	miR-145-5p	miR-188-5p	miR-214-5p	miR-29c-5p	miR-3613-5p	miR-500a-3p	miR-92a-3p
miR-1247-5p	miR-146a-5p	miR-18a-5p	miR-215	miR-301a-3p	miR-362-3p	miR-501-3p	miR-92b-3p
miR-1249	miR-146b-5p	miR-18b-5p	miR-218-5p	miR-301b	miR-363-3p	miR-502-3p	miR-93-5p
miR-125a-5p	miR-148a-3p	miR-191-5p	miR-22-3p	miR-30a-3p	miR-365a-3p	miR-505-3p	miR-941
miR-125b-5p	miR-148b-3p	miR-192-5p	miR-22-5p	miR-30a-5p	miR-365b-3p	miR-505-5p	miR-96-5p
miR-126-3p	miR-150-3p	miR-193a-3p	miR-221-3p	miR-30b-5p	miR-374a-5p	miR-5100	miR-98
miR-126-5p	miR-150-5p	miR-193a-5p	miR-221-5p	miR-30c-5p	miR-374b-5p	miR-532-3p	miR-99a-5p
miR-1260a	miR-151a-3p	miR-193b-3p	miR-222-3p	miR-30d-5p	miR-376a-3p	miR-532-5p	miR-99b-5p
miR-1260b	miR-151a-5p	miR-193b-5p	miR-222-5p	miR-30e-3p	miR-376c	miR-548h-3p	
miR-1271-5p	miR-151b	miR-194-5p	miR-223-3p	miR-30e-5p	miR-378a-3p	miR-548z	

