



Universiteit
Leiden
The Netherlands

TRPM7, Calcium and the cytoskeleton

Langeslag, Michiel

Citation

Langeslag, M. (2006, October 11). *TRPM7, Calcium and the cytoskeleton*. Retrieved from <https://hdl.handle.net/1887/4863>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4863>

Note: To cite this publication please use the final published version (if applicable).

TRPM7, a Novel Regulator of Actomyosin Contractility and Cell Adhesion

Kristopher Clark, Michiel Langeslag, Bart van Leeuwen, Leonie Ran, Alexey G. Ryazanov, Carl G. Figdor, Wouter H. Moolenaar, Kees Jalink and Frank N. van Leeuwen

The EMBO Journal (2006) 25, 290–301

TRPM7, a Novel Regulator of Actomyosin Contractility and Cell Adhesion

Kristopher Clark¹, Michiel Langeslag², Bart van Leeuwen³, Leonie Ran³, Alexey G. Ryazanov⁴, Carl G. Figdor¹, Wouter H. Moolenaar³, Kees Jalink² and Frank N. van Leeuwen¹.

¹Department of Tumor Immunology, Nijmegen Centre for Molecular Life Sciences, Radboud University Nijmegen Medical Centre, PO Box 9101, 6500 HB Nijmegen, The Netherlands

²Division of Cell Biology, ³Division of Cellular Biochemistry and Center for Biomedical Genetics, the Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

⁴Department of Pharmacology, University of Medicine and Dentistry of New Jersey, Robert Wood Johnson Medical School, 675 Hoes lane, Piscataway, NJ, 08854, USA

Actomyosin contractility regulates various cell biological processes including cytokinesis, adhesion and migration. While in lower eukaryotes α -kinases control actomyosin relaxation, a similar role for mammalian α -kinases has yet to be established. Here, we examined whether TRPM7, a cation channel fused to an α -kinase, can affect actomyosin function. We demonstrate that activation of TRPM7 by bradykinin leads to a Ca^{2+} - and kinase-dependent interaction with the actomyosin cytoskeleton. Moreover, TRPM7 phosphorylates the myosin IIA heavy chain. Accordingly, low overexpression of TRPM7 increases intracellular Ca^{2+} levels accompanied by cell spreading, adhesion and the formation of focal adhesions. Activation of TRPM7 induces the transformation of these focal adhesions into podosomes by a kinase-dependent mechanism, an effect that can be mimicked by pharmacological inhibition of myosin II. Collectively, our results demonstrate that regulation of cell adhesion by TRPM7 is the combined effect of kinase-dependent and -independent pathways on actomyosin contractility.

Introduction

Actomyosin contractility in nonmuscle cells plays a fundamental role in regulating basic cellular functions such as cell shape, cytokinesis, adhesion and migration (Burridge and Wennerberg, 2004; De la Roche *et al*, 2002; Geiger and Bershadsky, 2002). Myosin II is the major motor protein driving contractility. Regulators of myosin II-based contractile responses include the Rho GTPase family and their

effectors, myosin light chain (MLC) kinases and phosphatases as well as myosin heavy chain (MHC) kinases (Burridge and Wennerberg, 2004; De la Roche *et al*, 2002). The *Dictyostelium* genome encodes several MHC kinases, which are essential for proper localization and assembly of myosin II during cell division and migration (Heid *et al*, 2004; Kolman *et al*, 1996; Rico and Egelhoff, 2003). Myosin II assembles into bipolar thick filaments that generate cortical tension by pulling together oppositely oriented actin filaments. MHC kinases inhibit myosin II function by phosphorylating the MHC tail on threonine residues leading to filament disassembly and consequently, a release in cortical tension (Egelhoff *et al*, 1993). The cell biological effects of overexpression or knockout of these MHC kinases are consistent with this paradigm. Defects in cytokinesis due to overexpression of MHC kinases are reversed by expressing a MHC mutant that cannot be phosphorylated and therefore forms stable myosin filaments (Rico and Egelhoff, 2003). Thus, MHC kinases play a key role in actomyosin remodeling in *Dictyostelium*.

Dictyostelium MHC kinases belong to a rare and novel class of kinases known as α -kinases that, with the exception of the ATP-binding site, show only limited sequence similarity with other members of the protein kinase superfamily (Ryazanov, 2002). Based on *in vivo* substrates identified to date, these kinases preferentially phosphorylate threonine residues presented within the context of an α -helix, hence the name α -kinases (Ryazanov, 2002). Whether mammalian α -kinases also regulate actomyosin contractility has yet to be established.

The mammalian genome encodes several α -kinases whose function, with the exception of elongation factor kinase, is unknown (Drennan and Ryazanov, 2004). Notably, TRPM6 and TRPM7 are two bifunctional proteins encoding a TRP cationic channel fused to a COOH-terminal α -kinase domain. Electrophysiological characterization of TRPM7 has suggested that it forms part of the channel responsible for the Magnesium Inhibited Current (MIC) and is implicated in cellular Mg^{2+} homeostasis (Kozak and Cahalan, 2003; Nadler *et al.*, 2001; Schmitz *et al.*, 2003). However, TRPM7 may also relay signals by phosphorylating downstream effector molecules. In line with a hypothesized role for TRPM7 as an active component in receptor mediated signal transduction, it was reported that TRPM7 channels are highly permeable to Ca^{2+} , that the TRPM7 COOH-terminus associates with phospholipase C (PLC) isoforms and that PLC activation regulates TRPM7 channel opening (Runnels *et al.*, 2001; Runnels *et al.*, 2002; Langeslag *et al.*, manuscript in preparation). We have previously found that bradykinin, a Gq-PLC coupled receptor agonist, induces Ca^{2+} -dependent phosphorylation of the MHC and disassembly of myosin IIA at the cell periphery, which correlates with cell spreading and adhesion in mammalian cells (van Leeuwen *et al.*, 1999). Thus, we hypothesized that the coupling of a cation channel to an α -kinase in a single polypeptide could explain the close relationship between Ca^{2+} signaling and Ca^{2+} -dependent actomyosin remodeling. Here, we investigated whether TRPM7 could link receptor-mediated signaling to cytoskeletal contractility and cell adhesion.

Results

TRPM7 Mediates Bradykinin-Induced Calcium Influx

To examine a role for TRPM7 in the regulation of actomyosin contractility, an HA-tagged TRPM7 cDNA encoding wild-type (TRPM7-WT) or a kinase-dead mutant (TRPM7-D1775A) were introduced into N1E-115 neuroblastoma cells by retroviral transduction (a technique that allows low, near physiological expression of recombinant proteins; supplementary Fig. S1). Both TRPM7-WT and TRPM7-D1775A were equally expressed at 2-3 times over endogenous TRPM7 levels (Fig. 1A, supplementary S2; Langeslag *et al.*, manuscript in

preparation). Both TRPM7-WT and TRPM7-D1775A run as doublets at 216 kDa due to differential glycosylation (supplementary Fig. S3). Moreover, proper localization of TRPM7-WT and TRPM7-D1775A to the plasma membrane was confirmed by fluorescence microscopy (supplementary Fig. S3; see also Fig. 4).




Figure 1 TRPM7 enhances receptor-operated calcium influx. (A) Expression of TRPM7-WT and TRPM7-D1775A in N1E-115 neuroblastoma cells was detected by IP-Western blot using antibodies against the HA-tag. Cells containing the empty vector were used as control in all experiments. (B) TRPM7 expression causes sustained Ca^{2+} influx following BK stimulation independently of kinase activity. Left panel, typical time course of BK-induced Ca^{2+} changes in control N1E-115 cells; other panels, BK-induced Ca^{2+} mobilizations in N1E-115/TRPM7 cells. The sustained influx depends on Ca^{2+} influx since it is acutely blocked by extracellular La^{3+} and Gd^{3+} (data not shown). (C) Quantification of Ca^{2+} data for control and TRPM7-overexpressing cells (mean \pm SEM from at least 20 experiments for each condition). Significant differences (*) are $P < 0.01$ from values obtained in control N1E-115 cells.




Figure 2 TRPM7 overexpression induces cell spreading and increases cell adhesion. (A) Phase contrast images of control, TRPM7-WT and TRPM7-D1775A expressing N1E-115 cells. Scale bar = 50 μm. (B) Cell surface area covered by N1E-115 cells expressing WT and kinase-dead TRPM7. Cells were stained for F-actin and cell surface area was calculated by quantifying the amount of pixels that exceeded a threshold ($n > 5$). (C) Quantification of cell adhesion of N1E-115 cells expressing WT and kinase-dead TRPM7 ($n = 3$). Significant differences (*) are $P < 0.01$ from values obtained in control N1E-115 cells.

Initially, we investigated by Ca^{2+} fluorometry whether TRPM7 is activated following stimulation with bradykinin (BK), a peptide agonist that promotes actomyosin relaxation in N1E-115 cells (van Leeuwen *et al.*, 1999). Low overexpression of TRPM7 in N1E-115 cells resulted in a moderate elevation of the resting Ca^{2+} levels (109 ± 9 nM vs. 85 ± 4 nM). The addition of BK triggered a rapid increase in cytosolic Ca^{2+} from internal stores in both parental and TRPM7 transduced cells. Strikingly in N1E-115/TRPM7 cells, this initial Ca^{2+} transient was followed by a sustained phase of elevated Ca^{2+} that lasted for several minutes before returning to basal levels and was not observed in the parental cells (Fig. 1B). Removal of extracellular Ca^{2+} by BAPTA immediately terminated the

plateau phase (data not shown). Patch-clamp studies and inhibitor profiling identified TRPM7 as the channel responsible for the Ca^{2+} influx (a detailed analysis of TRPM7 channel function will be published elsewhere; Langeslag *et al.*, manuscript in preparation). Finally, channel activation of kinase-dead TRPM7-D1775A by BK was not altered in comparison to TRPM7-WT (Fig. 1B,C), demonstrating that TRPM7 kinase activity is not required for BK-induced Ca^{2+} influx. From these results, we conclude that TRPM7 functions as a BK-regulated Ca^{2+} channel in N1E-115 cells.

TRPM7 Expression Promotes Cell Spreading and Increases Cell-Matrix Adhesion

Mouse N1E-115 neuroblastoma cells are an excellent model system to study the regulation of actomyosin contractility since signals that either activate contraction or promote relaxation are rapidly translated into cell rounding or cell spreading responses, respectively (Jalink *et al.*, 1994; van Leeuwen *et al.*, 1997; van Leeuwen *et al.*, 1999). Consistent with a role for TRPM7 in actomyosin regulation, N1E-115/TRPM7 cells showed enhanced spreading when compared to the parental cells (Fig. 2A,B). In addition to effects on cell spreading, low overexpression of TRPM7 increased cell-matrix adhesion of N1E-115 cells (Fig. 2A,C). Both effects of TRPM7 were independent of kinase activity, suggesting that Ca^{2+} -dependent mechanisms account for the morphological differences between parental and TRPM7-transduced cells.

TRPM7 Activation Leads to Podosome Formation

In light of the effects of TRPM7 on cell morphology and adhesion, we further investigated the effect of low TRPM7 overexpression and activation on the subcellular distribution of F-actin, myosin IIA and vinculin by confocal microscopy (Fig. 3A). Control N1E-115 cells displayed a predominantly cortical distribution of F-actin and myosin IIA, with little or no focal adhesions nor stress fibers. Following BK-induced cell spreading, a part of this peripheral actomyosin was lost while small focal adhesions could now be




Figure 3 Activation of TRPM7 induces actomyosin reorganization in conjunction with podosome formation. (A) TRPM7 activation induces actomyosin reorganization and formation of podosomes in N1E-115/TRPM7 cells. Cells were either serum starved (0.1% FCS) or stimulated with BK (10 nM; 30 min) and stained for actin (red), myosin IIA heavy chain (green) and vinculin (blue). Cells were analyzed by confocal microscopy. Scale bar = 20 μm . (B) Increase in cell surface area and production of podosomes occurs within minutes of BK stimulation in N1E-115/TRPM7 cells. N1E-115/TRPM7 cells expressing GFP- β -actin were followed by confocal microscopy before and after BK stimulation. Cell surface area was measured as described in *Materials and Methods*. Grayvalues are displayed using the Fire LookUpTable (LUT) of the public domain software Image J (National Institutes of Health, Bethesda, USA). Left panel, a resting cell; middle panel, the same cell 12 min post-BK stimulation; right panel, kinetics of BK-induced cell spreading. A movie is provided as supplementary data. Scale bar = 15 μm . (C) High magnification of myosin lattices found in N1E-115/TRPM7 cells following BK stimulation. N1E-115/TRPM7 cells were transfected with GFP-myosin, stimulated with BK and stained for actin. Scale bar = 2 μm .




Figure 4 TRPM7 is present at the cell surface and localizes to cell adhesion structures. N1E-115/TRPM7-WT cells were stimulated with BK (10 nM; 30 min). Subsequently, TRPM7 was detected using anti-HA (3F10) antibody followed by alexa 488-conjugated anti-rat IgG. The actin cytoskeleton was visualized using phalloidin-texas red. (A) TRPM7 is found in membrane ruffles. Scale bar = 10 μ m. (B) TRPM7 forms a ring surrounding the actin dense core of podosomes. Scale bar = 10 μ m. (C) High magnification of TRPM7 rings within podosomes. Scale bar = 5 μ m. A rat IgG control antibody did not show any specific staining (data not shown).

observed at the cell periphery by vinculin staining. In N1E-115/TRPM7 cells, a profound redistribution of actomyosin was observed. In contrast to the parental cells, focal adhesions were already present in unstimulated cells, whereas myosin IIA was dispersedly staining the cytoplasm. BK stimulation caused further cell spreading and increased the number and size of adhesion complexes. These responses were noticeable within 2 min and peaked at approximately 10 min after BK stimulation (Fig. 3B and supplementary movie 1). The architecture and size (~1 μ m) of these adhesion structures were remarkably different from the focal adhesions observed in unstimulated cells and resembled that of podosomes (Linder and Aepfelbacher, 2003). Similar to podosomes, the TRPM7-induced adhesions, found at the interface of the cell and matrix, had a diameter of approximately 1 μ m. Moreover, they consisted of an actin core surrounded by vinculin and conspicuous myosin lattices that were no longer associated with actin stress fibers (Fig. 3C). We conclude therefore, that

TRPM7 activation promotes the formation of podosomes.

TRPM7 is Present at the Cell Surface in Proximity to Cell Adhesion Structures

To examine whether TRPM7 may affect cell adhesion locally, we investigated the cellular distribution of TRPM7 by immunofluorescence. TRPM7 was clearly present within membrane ruffles indicating that the protein is localized at the cell surface (Fig. 4A). In addition, TRPM7 is enriched in cell adhesions where it is found in the podosome ring structure, together with myosin IIA (Fig. 4B,C). Our results suggest that TRPM7 regulates cell adhesion by directly affecting components within these structures.

Induction of Podosomes by TRPM7 Is Mediated by a Kinase-Dependent Mechanism

Although the global effects of TRPM7 on cell spreading and adhesion appear to be independent of kinase activity, we tested whether the TRPM7 kinase domain affects the subcellular organisation of the actomyosin cytoskeleton. By confocal microscopy, we compared N1E-115 cells expressing TRPM7-WT with cells expressing the kinase-dead TRPM7-D1775A mutant. No clear differences were noticed between unstimulated cells (data not shown). However to our surprise, cells expressing kinase-dead TRPM7 failed to induce podosomes in response to BK stimulation (Fig. 5). We conclude that TRPM7 activation




Figure 5 Regulation of podosome formation by TRPM7 is kinase-dependent. N1E-115/TRPM7-WT and N1E-115/TRPM7-D1775A cells were stimulated with BK (10 nM; 30 min) and stained for actin (red), myosin IIA heavy chain (green) and vinculin (blue). Cells were analyzed by confocal microscopy. Scale bar = 20 μ m.




Figure 6 Inhibition of myosin II function in N1E-115 cells expressing TRPM7-WT and TRPM7-D1775A leads to podosome formation. (A) N1E-115/TRPM7-WT and N1E-115/TRPM7-D1775A cells were incubated in the presence of vehicle control or 50 μ M blebbistatin for 30 min. Cells were stained for actin (red), myosin (green) and vinculin (blue) and visualized by confocal microscopy. Scale bar = 15 μ m. **(B)** High magnification view shows that the adhesion complexes induced by BK and blebbistatin in N1E-115/TRPM7-WT cells are architecturally similar since they consist of an actin dense core surrounded by a ring of vinculin and myosin IIA. Scale bar = 5 μ m.

results in a kinase-dependent remodeling of the actomyosin cytoskeleton, leading to the assembly of podosomes.

Inhibition of Myosin II Function Leads to Podosome Formation Both in Cells Expressing Wild-type and Kinase-dead TRPM7

Actomyosin contractility plays a central role in regulating the assembly and disassembly of adhesive contacts (Geiger and Bershadsky, 2002). Notably, the formation of podosomes requires a local inhibition of actomyosin contractility suggesting that TRPM7 may regulate podosome assembly by promoting relaxation of the actomyosin cytoskeleton (Burgstaller and Gimona, 2004). To further test this model, we investigated the effect of directly inhibiting myosin II function on cell adhesion in N1E-115 cells expressing TRPM7-WT and TRPM7-D1775A. Treatment of cells with blebbistatin (a selective inhibitor of myosin II ATPase activity) led to the transformation of focal adhesions into podosomes

Figure 5 Regulation of podosome formation by TRPM7 is kinase-dependent. N1E-115/TRPM7-WT and N1E-115/TRPM7-D1775A cells were stimulated with BK (10 nM; 30 min) and stained for actin (red), myosin IIA heavy chain (green) and vinculin (blue). Cells were analyzed by confocal microscopy. Scale bar = 20 μ m.

independently of TRPM7 kinase activity (Fig. 6A). These effects of blebbistatin do not require exogenous TRPM7 as parental N1E-115 cells also produce podosomes in response to myosin II inhibition (supplementary Fig. S4). Notably, the adhesive structures formed upon blebbistatin treatment are remarkably similar to those formed after BK stimulation of N1E-115/TRPM7-WT cells (Fig. 6B). These findings indicate that the transformation of normal focal adhesions into podosomes by BK-mediated activation of TRPM7 is due to a kinase-dependent inhibition of myosin II function.

TRPM7 Interacts with the Actomyosin Cytoskeleton

Since TRPM7 is a member of the α -kinase family, we hypothesized that it may affect actomyosin remodeling and podosome assembly by directly coupling to and potentially

phosphorylating components present within the actomyosin cytoskeleton. Therefore, we precipitated TRPM7 complexes with anti-HA antibodies and detected the presence of associated proteins by Western blotting. Both β -actin and the myosin IIA heavy chain were present in a complex with TRPM7 (Fig. 7A). Importantly, endogenous TRPM7 also associates with the actomyosin cytoskeleton (Fig. 7B). This interaction indeed suggests that TRPM7 can affect actomyosin function through a direct association with cytoskeletal proteins.

Bradykinin Causes Calcium and Kinase-dependent Association of the TRPM7 COOH-terminus with Myosin IIA

Since TRPM7 activation by BK results in Ca^{2+} -influx and a kinase-dependent remodeling of the actomyosin cytoskeleton, we investigated whether the interaction between TRPM7 and the actomyosin cytoskeleton is subject to regulation. BK stimulation of N1E-115/TRPM7 cells led to a transient increase in the amount of TRPM7-associated myosin IIA (Fig. 7C,D). Its kinetics closely correlate with those of calcium influx in response to TRPM7 activation (refer to Fig. 1) with maximal association observed at about 2 min after agonist addition. Moreover, chelation of Ca^{2+} , using BAPTA or EDTA, abrogated the association between TRPM7 and myosin IIA (Fig. 7E). These results indicate that the interaction between TRPM7 and myosin IIA is strictly Ca^{2+} -dependent, and suggest that TRPM7-mediated Ca^{2+} influx enhances the TRPM7/myosin IIA interaction. However, in addition to Ca^{2+} , the association of TRPM7 with the cytoskeleton requires an active kinase domain since the kinase-dead TRPM7-D1775A mutant did not interact with myosin IIA (Fig. 7F). A soluble COOH-terminus variant containing the kinase domain of TRPM7 also interacted with myosin IIA (Fig. 7G). We conclude therefore that the interaction between TRPM7 and myosin IIA is mediated by the COOH-terminus and requires an active kinase domain.

TRPM7 Phosphorylates Myosin IIA Heavy Chain

Hitherto, we have shown that TRPM7 associates with myosin IIA in a regulated manner and that both activation of TRPM7-WT (but not TRPM7-D1775A) and inhibition of myosin II




Figure 7 TRPM7 activation promotes its association via the COOH-terminus with the actomyosin cytoskeleton in a calcium- and kinase-dependent manner. In all experiments (except panel B), N1E-115 control or TRPM7-transduced cells were lysed and incubated with anti-HA-coupled protein G beads for 3 h at 4°C. Proteins in the complex were detected by Western blotting. **(A)** Coimmunoprecipitation of TRPM7 with β -actin and myosin IIA. Top, detection of TRPM7 with anti-HA antibodies; middle and bottom, detection of myosin IIA and β -actin, respectively, in the immune complexes (IP, left) and total lysates to control for protein levels (TL, right). **(B)** Endogenous TRPM7 associates with the actomyosin cytoskeleton. PC12 lysates were incubated with pre-immune serum or anti-TRPM7 serum and protein complexes were isolated using protein G-sepharose. The presence of myosin IIA was detected by Western blotting. Top, detection of associated myosin IIA heavy chain in immunoprecipitates; and Bottom, in total lysates. **(C)** Coimmunoprecipitation of TRPM7 with myosin IIA heavy chain in the pre- and post-BK stimulation (10 nM, 1 min). Top, detection of coimmunoprecipitated myosin IIA. Middle, detection of TRPM7 with anti-HA antibodies. Bottom, detection of myosin IIA heavy chain in total lysates to control for protein levels. **(D)** Kinetics of BK induced TRPM7 association with myosin IIA heavy chain. Cells were stimulated with BK (10 nM) for the indicated times and TRPM7 was immunoprecipitated using anti-HA antibodies. Top, Western blot showing coimmunoprecipitated myosin IIA heavy chain. Bottom, detection of myosin IIA heavy chain in total lysates. **(E)** Interaction between TRPM7 and myosin IIA heavy chain is Ca^{2+} -dependent. N1E-115/TRPM7 cells were pre-incubated with 10 mM BAPTA or EDTA for 1 min prior to cell lysis and TRPM7 was immunoprecipitated using anti-HA antibodies. **(F)** TRPM7-D1775A does not interact with myosin IIA heavy chain. TRPM7-WT and -D1775A were immunoprecipitated with anti-HA antibodies and associating myosin IIA heavy chain was detected by Western blotting. **(G)** The COOH-terminus of TRPM7 interacts with myosin IIA. The soluble COOH-terminus of TRPM7 (a.a. 1158-1864) was immunoprecipitated from N1E-115/TRPM7-C cells with anti-HA antibodies and associated myosin IIA was detected by Western blotting. Top, detection of associated myosin IIA heavy chain; bottom, detection of TRPM7-C using anti-HA antibodies in immunoprecipitation (IP, left) and in total lysates (TL, right).

function produce podosomes suggesting that TRPM7 activation promotes actomyosin relaxation. How does TRPM7 mediate the inhibition of myosin II function? By analogy to regulation of *Dictyostelium* myosin II function, we have previously proposed that Ca^{2+} -dependent MHC phosphorylation contributes to cytoskeletal relaxation (van Leeuwen *et al*, 1999). The finding that TRPM7 coprecipitates with myosin IIA, particularly after BK stimulation, enabled us to test whether TRPM7 can phosphorylate myosin IIA heavy chain. Since TRPM7 and myosin IIA heavy chain co-migrate on SDS-PAGE gels, TRPM7 was

immunoprecipitated both under conditions that favor association and dissociation (low vs. high stringency) of the cytoskeletal complex allowing

us to distinguish between TRPM7 autophosphorylation and myosin phosphorylation. By *in vitro* kinase assays, TRPM7 underwent autophosphorylation in agreement with earlier findings (Runnels *et al*, 2001; Ryazanova *et al*, 2004; Schmitz *et al*, 2003) (Fig. 8A; bottom panel). Strikingly, clear phosphorylation of associated myosin IIA heavy chain (which runs slightly faster than TRPM7) was also observed in these kinase reactions (Fig. 8A; bottom panel). In contrast, actin was not phosphorylated (data not shown).

To demonstrate that TRPM7 directly phosphorylates myosin IIA heavy chain, a myosin tail fragment was expressed as a GST-fusion protein in *E. coli* to serve as substrate whereas TRPM7 was immunoaffinity purified from




Figure 8 TRPM7 phosphorylates myosin IIA heavy chain.

(A) *In vitro* kinase assay detecting the phosphorylation of associated myosin IIA heavy chain by TRPM7. TRPM7/myosin IIA heavy chain complexes were isolated before and after stimulation of N1E-115 control and N1E-115/TRPM7 cells with BK (10 nM, 2 min) under low (1% triton X-100) and high (1% triton X-100/ 0.5% deoxycholate/ 0.1% SDS) stringency conditions. Substrates of TRPM7 were detected by labeling proteins with γ - ^{32}P -ATP, separating products of the *in vitro* kinase assay by SDS-PAGE (6% gel) followed by autoradiography. Top, Coomassie staining of precipitated proteins. Bottom, autoradiogram of phosphorylated proteins. Note that myosin association is lost under high stringency conditions (1% triton X-100/ 0.5% deoxycholate/ 0.1% SDS) whereas TRPM7 autophosphorylation is unaffected. (B) TRPM7 phosphorylates recombinant myosin IIA. TRPM7-WT and -D1775A were immunoaffinity purified from HEK293 cells using anti-HA antibodies under high stringency and mixed with 2 μg of GST or GST-myosin IIA in kinase buffer. To detect phosphorylated myosin IIA, the proteins were separated by SDS-PAGE (12 % gel) followed by autoradiography. Top, Coomassie staining of GST-fusion proteins. Note that GST co-migrates with the antibody light chain (Ig LC). Middle, autoradiogram of phosphorylated GST-fusion proteins. Bottom, autoradiogram showing autophosphorylation of WT but not kinase-dead TRPM7. (C) Recombinant TRPM7 kinase phosphorylates myosin IIA. The TRPM7 kinase domain (1403-1864 aa) was produced in *E. coli* as a fusion with maltose binding protein and purified on an amylose column. The purified kinase was incubated with 2 μg of GST or GST-myosin IIA in kinase buffer. The proteins were resolved on a 12% SDS-PAGE gel and detected by coomassie. Phosphorylated proteins were visualized by autoradiography. Left, coomassie staining of gel; right, autoradiogram.

mammalian cells or the COOH-terminus of TRPM7 was purified from *E. coli*. In *in vitro* kinase assays, immunoaffinity purified WT but not kinase-dead TRPM7 efficiently phosphorylated recombinant myosin IIA (Fig. 8B). Notably, GST was not phosphorylated by TRPM7 (Fig. 8B). Similar results were obtained when using the soluble COOH-terminus of TRPM7 (data not shown). Importantly, contaminating actin in the purified TRPM7 fractions is below detection levels and disruption of the actin cytoskeleton prior to TRPM7 purification had no effect on the level of myosin II phosphorylation arguing against the presence of contaminating kinases (supplementary Fig. S5). Finally, recombinant TRPM7 purified from *E. coli* lysates also efficiently phosphorylated myosin IIA, in the absence of other sources of eukaryotic proteins (Fig. 8C). Thus, TRPM7 itself and not an associated kinase is responsible for myosin IIA heavy chain phosphorylation. Collectively, our results demonstrate that TRPM7 associates with the actomyosin cytoskeleton in a Ca^{2+} - and kinase-dependent manner to regulate myosin II activity and consequently, actomyosin contractility. Moreover, our *in vitro* kinase data suggest that phosphorylation of the myosin IIA heavy chain by TRPM7 serves as a regulatory mechanism.

Discussion

In this study, we have tested the hypothesis that TRPM7, by analogy to its α -kinase family members from *Dictyostelium*, affects actomyosin contractility. We provide evidence that TRPM7 promotes relaxation of the actomyosin cytoskeleton via a kinase-dependent inhibition of myosin II, potentially involving myosin IIA heavy chain phosphorylation. The evidence includes: i) electrophysiological measurement of TRPM7 channel opening after BK stimulation; ii) biochemical analyses of the TRPM7 complex showing that TRPM7 interacts in a Ca^{2+} - and kinase-dependent manner with an actomyosin protein complex; iii) *in vitro* kinase reactions demonstrating that TRPM7 phosphorylates myosin IIA heavy chain; and iv) cell biological studies revealing that TRPM7 promotes a loss of cortical tension leading to podosome formation by a kinase-dependent mechanism. Collectively, our data indicate

that TRPM7 plays a role in linking receptor-mediated signals to actomyosin remodeling and cell adhesion. Furthermore, our observations reveal for the first time that TRP channels affect the cytoskeleton by directly associating with cytoskeletal proteins in a highly regulated manner.

What is the relationship between channel opening and the kinase domain? Several studies have suggested that kinase activity regulates TRPM7 channel opening (Runnels *et al*, 2001; Schmitz *et al*, 2003). However, in our model system, TRPM7 channel opening was independent of kinase activity, since the responses to BK stimulation were identical between cells expressing WT and kinase-dead TRPM7. While TRPM7 kinase activity does not directly affect channel opening, it cannot be excluded that actomyosin remodeling serves to regulate TRPM7 function in an indirect manner as reported for other TRP channels (Itagaki *et al*, 2004; Lockwich *et al*, 2001). Based on our observations, rather than the kinase domain regulating TRPM7-channel function, the reverse relationship exists. In this model (Fig. 9), phosphorylation of downstream targets (myosin II, annexin I; (Dorovkov and Ryazanov, 2004)) by the TRPM7-kinase domain is tightly regulated by ion-influx (Ca^{2+} , Mg^{2+}) through the TRPM7-channel. In support of this model, we observed that Ca^{2+} is strictly required for the association of TRPM7 with the cytoskeleton, that increased Ca^{2+} -influx through TRPM7 upon agonist-induced activation enhances its association with myosin IIA and that MHC phosphorylation is strictly Ca^{2+} -dependent (van Leeuwen *et al*, 1999). Recently, Dorovkov and Ryazanov (2004) showed that the kinase activity of TRPM7 is potentiated *in vitro* by the addition of Ca^{2+} and conversely, is inhibited by chelation of Ca^{2+} . Therefore, we propose that ion influx through the channel pore regulates the TRPM7 kinase domain by activating the kinase and controlling the recruitment of its substrates.

Kinases are maintained in an active state by intramolecular and intermolecular protein-protein or protein-lipid interactions (Roskoski, 2004; Van Etten, 2003). Activation of kinases

generally involves autophosphorylation or transphosphorylation at regulatory sites inducing conformational remodeling to expose the catalytic domain to its substrate. We postulate that TRPM7 is similarly regulated, since its kinase activity appears to be essential for its association with the actomyosin cytoskeleton. Autophosphorylation of TRPM7, which occurs on serine and threonine residues (Ryazanova *et al*, 2004; Schmitz *et al*, 2003), may release a cryptic binding site for myosin IIA heavy chain (Fig. 9). Thus, autophosphorylation of TRPM7 may serve as an important regulatory mechanism to modulate the interactions between TRPM7 and its substrates.

A functional consequence of TRPM7 activation is an increase in cell adhesion and spreading, which involves both kinase-dependent and -independent pathways. Expression of a TRPM7 kinase-dead mutant still promotes cell adhesion and spreading but does not lead to the formation of podosomes upon BK stimulation. The fact that the global effects of TRPM7 on cell morphology are independent of kinase activity or its association with the cytoskeleton may reflect the increase in cytosolic Ca^{2+} concentrations observed in TRPM7-transduced cells. Indeed, reducing the internal Ca^{2+} concentration with BAPTA-AM impaired cell spreading (data not shown). Ca^{2+} may act directly at the level of integrin activation or affect the actomyosin cytoskeleton. The activation of integrins can lead to the remodeling of the actomyosin cytoskeleton to promote cell spreading via outside-in signaling pathways (DeMali *et al*, 2003). Alternatively, Ca^{2+} is an important second messenger in actin remodeling including polymerization, severing of filaments and F-actin-membrane interactions (Forscher, 1989; Sun *et al*, 1999). Moreover, since TRPM7 can directly interact with PLC-isoforms, it may influence local concentrations of PIP₂, and thus affect actin polymerization (van Rheenen and Jalink, 2002).

The kinase-dependent effects of TRPM7 on cell adhesion are consistent with a role in regulating myosin IIA-dependent contractile responses. Notably, both BK stimulation and myosin II inhibition of N1E-115/TRPM7 cells

induces the formation of large podosomes. Spatial and temporal regulation of actomyosin contractility is central to modulating the assembly and disassembly of focal adhesions and podosomes (Burgstaller and Gimona, 2004; Burrige and Wennerberg, 2004; DeMali *et al*, 2003; Geiger and Bershadsky, 2002; Linder and Aepfelbacher, 2003). While local regulation of contractility appears to be controlled by Rho GTPases (Burgstaller and Gimona, 2004), α -kinases may also fulfill a role in spatial and temporal regulation of myosin II activity. Indeed, TRPM7, which is expressed at the cell surface, was found enriched in cell adhesion structures. By similarity, *Dictyostelium* myosin II function is precisely regulated by a family of MHC kinases with each member possessing a unique spatial distribution (Liang *et al*, 2002). Therefore, we postulate that an increase in TRPM7-mediated MHC phosphorylation contributes to local relaxation of the cortical cytoskeleton causing the transformation of focal adhesions into podosomes. Ablation of TRPM7 kinase activity maintains myosin-based cytoskeletal contractility preventing the reorganization of focal adhesions. Hence, TRPM7 regulates cell adhesion by modulating actomyosin contractility in a kinase-dependent manner and it is tempting to speculate that this phenomenon operates through MHC phosphorylation.

How to reconcile the small increase in TRPM7 expression (2-fold) with dramatic changes in cell morphology and response to BK? Although the kinase activity of TRPM7 is a linear relationship with concentration *in vitro*, it is most likely not linear *in vivo* as the protein will be subject to regulation. In support of this hypothesis, we have found that TRPM7 channels have increased activity as measured by patch-clamp and Ca^{2+} fluorometric experiments in N1E-115/TRPM7 cells. It should be noted that the basal Ca^{2+} levels in the cells are significantly higher upon TRPM7 overexpression, which could influence kinase activity as reported by Dorovkov and Ryazanov (2004). Moreover, TRPM7-mediated Ca^{2+} influx affects the recruitment of substrates for the kinase, by associating with the actomyosin cytoskeleton. Therefore, it can be expected that the activity of the TRPM7 complex in the N1E-115/TRPM7 *in vivo* is significantly more than the two-fold increase in kinase activity that is observed *in vitro*. Finally, if TRPM7 is targeted to a particular part of the cytoskeleton including cell adhesion structures, it is likely that the local activity of TRPM7 kinase increases

significantly more than 2-fold in those areas of the cell.

Bradykinin stimulation of N1E-115 cells leads to cell spreading and focal adhesion formation whereas it promotes the transformation of focal adhesions into podosomes in N1E-115/TRPM7 cells. Although N1E-115 cells express endogenous TRPM7, these cells do not produce podosomes in response to BK. We believe these results reflect the contractile state of the cell and that it is necessary to overexpress TRPM7 to achieve complete relaxation of the actomyosin cytoskeleton required for BK-mediated podosome formation (see supplementary Fig. S6). In parental N1E-115 cells, myosin II is very active leading to a contractile phenotype. BK stimulation of these cells relaxes the cytoskeleton but since focal adhesions are formed, myosin II must be at least partly active as tension is required for focal adhesion biogenesis (Geiger and Bershadsky, 2002). Overexpression of TRPM7 mimics these effects and BK stimulation of these cells leads to further inhibition of myosin II and relaxation of the actomyosin cytoskeleton, which allows for the formation of podosomes (adhesion structures which require an inhibition of myosin II for their formation (Burgstaller and Gimona, 2004)). In support of this model, inhibition of myosin II by blebbistatin in parental N1E-115 cells similarly induces podosomes (supplementary Fig. S4). Thus, since TRPM7 has a basal activity, its expression levels dictate myosin II activity, and thereby cellular phenotype.

Although mammalian nonmuscle actomyosin contractility appears to be primarily regulated by phosphorylation of the myosin regulatory light chain, a role for myosin heavy chain phosphorylation is emerging. Several studies have demonstrated that the MHC tail is phosphorylated upon stimulation of mammalian cells (van Leeuwen *et al*, 1999; Wilson *et al*, 1998). Based on homology to *Dictyostelium* MHC kinases, we hypothesized that TRPM7 may regulate myosin II function by phosphorylating the MHC α -helical tail. Indeed, *in vitro* kinase assays demonstrated that WT but not kinase-dead TRPM7 directly phosphorylates the MHC tail indicating broad evolutionary conservation with respect to both substrate and function. An unexpected result since other α -kinases, α -kinase 1 and EFK-2, regulate protein transport and protein synthesis, respectively (Heine *et al*, 2005; Ryazanov, 2002).

Myosin II forms bipolar thick filaments through electrostatic interactions of the MHC tail




Figure 9 Regulation of actomyosin contractility by TRPM7. (A) In the resting state, TRPM7 is not associated with the actomyosin cytoskeleton (1). (B) Stimulation of cells with PLC-activating agonists induces TRPM7-mediated Ca^{2+} influx (2) as well as TRPM7 kinase activity. Autophosphorylation (3) promotes a conformational change within TRPM7 allowing for Ca^{2+} -dependent association with myosin IIA (4). (C) Subsequently, TRPM7 phosphorylates the myosin IIA heavy chain, presumably leading to dissociation of myosin filaments and cytoskeletal remodeling (5).

(Hostetter *et al*, 2004). In *Dictyostelium*, phosphorylation of the α -helical MHC tail leads to filament disassembly thereby releasing cortical tension (Egelhoff *et al*, 1993). Whether phosphorylation of mammalian myosin II heavy chain by TRPM7 will have a similar effect on filament assembly remains to be determined (Fig. 9C). Importantly, the region of myosin II phosphorylated by TRPM7 corresponds to the domain responsible for regulating myosin filament assembly (Nakasawa *et al*, 2005). Moreover, phosphorylation by PKC and casein kinase as well as point mutations affecting electrostatic charges (E1841K) within this domain affect myosin II function (Dulyaninova *et al*, 2005; Franke *et al*, 2005). Therefore, we propose that phosphorylation of this region of the myosin IIA tail affects filament formation rather than ATPase activity of myosin II. Future investigations aimed at mapping the TRPM7 target residues within the myosin IIA heavy chain and defining the physiological relevance of MHC phosphorylation will provide further insight into the mechanisms underlying actomyosin contractility in mammalian cells.

In conclusion, we identified TRPM7 as a novel regulator of actomyosin contractility and cell adhesion in response to agonist stimulation. Moreover, we have demonstrated that TRP channels can interact with the actomyosin cytoskeleton to affect cell adhesion. Future experiments will be aimed at defining the role of TRPM7-containing protein assemblies in regulating actomyosin function, and establishing how these cytoskeletal changes affect cell adhesion and/or TRPM7 function.

Materials & Methods

Constructs

Mouse TRPM7 in pTracer-CMV2, and GFP-myosin IIA in pEGFP-C3 were kind gifts from David Clapham (Harvard, Boston, USA) and Robert Adelstein (NIH, Bethesda, USA), respectively. An HA-tag has been inserted at the COOH-terminus of TRPM7 (Runnels *et al*, 2001). To generate retroviral expression vectors, the TRPM7 cDNA was inserted as a *XhoI*-*NotI* fragment into the LZRS-ires-neomycin vector. The kinase-dead mutant TRPM7-D1775A was produced by site-directed mutagenesis using the Quickchange kit (Stratagene). The soluble TRPM7 COOH-terminus construct encodes for amino acids 1158-1864 and contains a HA-tag at the NH₂-

terminus. The cDNA was amplified by PCR and inserted as a *XhoI*-*NotI* fragment into the LZRS-ires-neomycin vector. GFP- β -actin cDNA was inserted as an *EcoRI*-*NotI* fragment into the LZRS-ires-neomycin vector. Mouse myosin IIA heavy chain tail (aa 1795-1960) was amplified by RT-PCR from N1E-115 cells and cloned in frame in pGEX-1N using *BamHI*-*EcoRI* sites. The TRPM7 kinase domain (aa 1403-1864) in pMAL-p2x was described previously (Ryazanova *et al*, 2004). All constructs were verified by DNA sequencing.

Cell Culture

Mouse N1E-115, HEK293 and Phoenix packaging cells were cultured in DMEM medium with 10% FCS while PC12 cells were grown in RPMI medium with 10% horse serum and 5% FCS. Stable cell lines expressing the various TRPM7 constructs and GFP- β -actin were generated by retroviral transduction (van Leeuwen *et al*, 1999). N1E-115 cells transduced with the empty vector served as a control for all experiments. Cells were selected by the addition of 0.8 mg/ml G418 to the media and the selection was complete within 7 days. For transient expression, cells were transfected using Fugene 6 (Roche) according to the manufacturer's recommendations.

Intracellular Calcium Determinations

Calcium measurements were performed by confocal ratiometric imaging of cells loaded with Oregon Green 488 BAPTA-1 AM and Fura Red AM (Molecular Probes) essentially as described (Schild *et al*, 1994). Experiments were performed at 37°C in a HEPES/bicarbonate-buffered saline under 5% CO₂, pH 7.3 (140 mM NaCl, 23 mM NaHCO₃, 5 mM KCl, 2 mM MgCl₂, 1 mM CaCl₂, 10 mM HEPES and 10 mM glucose), and calibration was with ionomycin (Calbiochem). Shown are representative traces from experiments performed at least in 10-fold; data are presented as mean \pm SEM.

Adhesion Assay

Cells (5×10^5) were seeded in a T25 flask and allowed to adhere for 16 h. To quantitate cell adhesion, non-adherent cells were washed away with PBS. The adherent cells were fixed in 3.7% formaldehyde in PBS for 10 min, washed with PBS and subsequently stained with 2% crystal

violet for 50 min. Excess stain was removed by extensive washing in PBS. Cells were lysed in PBS containing 1% NP40 and the absorbance at 570 nm of the lysate was measured. Maximal adhesion as measured on poly-L-lysine coated dishes was set to 100%. Values reported are representative of two independent experiments performed in triplicate.

Microscopy

Cells seeded on glass coverslips, were serum starved (0.1% FCS) overnight prior to stimulation with BK or blebbistatin (Calbiochem). For fixed specimens, cells were stained as previously described (van Leeuwen *et al*, 1997). Antibodies used for immunofluorescence were rat anti-HA (3F10; 1:100; Roche), mouse anti-vinculin (1:400; Sigma), rabbit anti-myosin IIA (1:100; BTI), alexa 488-conjugated anti-rabbit and anti-rat IgG, and alexa 647-conjugated anti-mouse IgG (1:200; Molecular Probes). F-actin was detected using either texas-red or alexa 546-labelled phalloidin (1:50; Molecular Probes). Cells were viewed using either a Zeiss-LSM 510-meta microscope equipped with a Plan-Apochromat 63X, 1.4 NA oil immersion lens or a Leica-DMRA microscope with a 63X, 1.32 NA oil immersion lens. For videomicroscopy, cells were transferred to a bicarbonate-buffered medium as described above and maintained at 37°C on a temperature- and CO₂-controlled stage. Cells were imaged with a DM-IRE2 inverted microscope fitted with a TCS-SP2 scanhead and a 63X, 1.32 NA oil immersion lens (Leica). Images were collected at a 30 s time interval and surface area was calculated by quantifying the amount of pixels under a digital mask constructed by using the binary-fill operation after a thresholding step.

Generation of Anti-TRPM7 Antibodies

To generate anti-TRPM7 antibodies, a GST-fusion protein encoding amino acids 1447 to 1555 of TRPM7 was expressed in *E. coli* and purified on a glutathione-sepharose column. Rabbits were injected with the antigen mixed with Freund's adjuvant and serum was collected 10 days after every vaccination. Serum collected prior to immunization (pre-immune) was used as a negative control in all experiments.

Immunoprecipitations

Cells were serum starved (0.1% FCS) overnight prior to stimulation as indicated in the figure legends. After stimulation, cells were lysed on ice for 20 min in lysis buffer (50 mM Tris pH 7.5, 300 mM NaCl, 0.5 mM DTT, 1.5 mM MgCl₂, 0.2 mM EDTA, 1% triton x-100 supplemented with protease inhibitors) and the extract was cleared by centrifugation. For the immunoprecipitation of exogenously expressed TRPM7, protein G-sepharose beads, which were blocked with 0.5% BSA and pre-coupled with anti-HA antibody (clone 12CA5), were added to the lysate of N1E-115/ empty vector (control) or N1E-115/TRPM7 cells. The samples were incubated at 4°C for 3 h. Endogenous TRPM7 was immunoprecipitated by incubating cellular lysates with anti-TRPM7 antibodies at 4°C for 3 h followed by the addition of protein G-sepharose beads at 4°C for 15 min. Subsequently, the beads were washed 3 times with lysis buffer, protein complexes were solubilized in Laemmli sample buffer and separated by SDS-PAGE. Proteins were detected by immunoblotting using the following antibodies: anti-HA (clone 3F10; 1:1000), anti-myosin IIA (1:500), anti-β actin (1:20000; Sigma).

In Vitro Kinase Assays

Recombinant TRPM7 kinase domain (amino acids 1402-1864) and myosin tail fragment (amino acids 1795-1960) were expressed, respectively, as a maltose binding protein- and GST-fusion protein in *Escherichia coli* and purified by standard methods. Immunocomplexes containing TRPM7 with or without associated myosin or TRPM7 mixed with 2 μg of GST-myosin were solubilized in kinase buffer (50 mM HEPES pH 7.0, 4 mM MnCl₂, 5 mM DTT) without ATP. The kinase reaction was initiated by adding 0.1 mM ATP in combination with 5 μCi γ-³²P-ATP and proceeded for 30 min at 30°C. The products of the kinase reaction were resolved by SDS-PAGE and detected by autoradiography.

Statistical Analysis

All data are representative of at least 3 independent experiments. Quantitative data are presented as the mean ± SEM. Statistical significance of differences between experimental groups was assessed with Student's *t* test.

Differences in means were considered significant if $p < 0.05$.

Supplemental Material

Fig. S1 shows that mutation of aspartate 1775 to alanine abrogates the ability of TRPM7 to undergo autophosphorylation. Fig. S2 shows the relative TRPM7 kinase activity in the different N1E-115 cell lines. Fig. S3 shows that both TRPM7-WT and TRPM7-D1775A are glycosylated and properly localized to the plasma membrane. Fig. S4 depicts the induction of podosomes in parental N1E-115 cells upon blebbistatin treatment. Fig. S5 shows that depletion of actin from TRPM7 immunoprecipitation by latrunculin A treatment has no effect on myosin IIA phosphorylation. Fig. S6 provides a model explaining the correlation between cell phenotype, TRPM7 expression and myosin II activity. Movie 1 depicts the dynamics of cell spreading and focal adhesion formation in N1E-115/TRPM7 cells in response to BK stimulation.

Acknowledgements

We are grateful to David Clapham, Loren Runnels and Robert Adelstein for providing DNA constructs. We thank Nannette Teunissen for technical assistance and Frank de Lange for help with confocal microscopy. We also thank Wiljan Hendriks and Erik Danen for critically reading the manuscript. This work was supported by grants from the Dutch Cancer Society (W.H.M., F.L., C.F.; grant- NKB 2002-2593).

References

- Burgstaller G, Gimona M (2004) Actin cytoskeleton remodelling via local inhibition of contractility at discrete microdomains. *J Cell Sci* **117**: 223-231
- Burridge K, Wennerberg K (2004) Rho and Rac take center stage. *Cell* **116**: 167-179
- De la Roche MA, Smith JL, Betapudi V, Egelhoff TT, Cote GP (2002) Signaling pathways regulating Dictyostelium myosin II. *J Muscle Res Cell Motil* **23**: 703-718
- DeMali KA, Wennerberg K, Burridge K (2003) Integrin signaling to the actin cytoskeleton. *Curr Opin Cell Biol* **15**: 572-582
- Dorovkov MV, Ryazanov AG (2004) Phosphorylation of annexin I by TRPM7 channel-kinase. *J Biol Chem* **279**: 50643-50646
- Drennan D, Ryazanov AG (2004) Alpha-kinases: analysis of the family and comparison with conventional protein kinases. *Prog Biophys Mol Biol* **85**: 1-32
- Dulyaninova NG, Malashkevich VN, Almo SC, Bresnick AR (2005) Regulation of myosin-IIA assembly and Mts1 binding by heavy chain phosphorylation. *Biochemistry* **44**: 6867-6876
- Egelhoff TT, Lee RJ, Spudich JA (1993) Dictyostelium myosin heavy chain phosphorylation sites regulate myosin filament assembly and localization in vivo. *Cell* **75**: 363-371
- Forscher P (1989) Calcium and polyphosphoinositide control of cytoskeletal dynamics. *Trends Neurosci* **12**: 468-474
- Franke JD, Dong F, Rickoll WL, Kelley MJ, Kiehart DP (2005) Rod mutations associated with MYH9-related disorders disrupt nonmuscle myosin-IIA assembly. *Blood* **105**: 161-169
- Geiger B, Bershadsky A (2002) Exploring the neighborhood: adhesion-coupled cell mechanosensors. *Cell* **110**: 139-142
- Heid PJ, Wessels D, Daniels KJ, Gibson DP, Zhang H, Voss E, Soll DR (2004) The role of myosin heavy chain phosphorylation in Dictyostelium motility, chemotaxis and F-actin localization. *J Cell Sci* **117**: 4819-4835
- Heine M, Cramm-Behrens CI, Ansari A, Chu HP, Ryazanov AG, Naim HY, Jacob R (2005) Alpha-kinase 1, a new component in apical protein transport. *J Biol Chem* **280**: 25637-25643
- Hostetter D, Rice S, Dean S, Altman D, McMahon PM, Sutton S, Tripathy A, Spudich JA (2004) Dictyostelium myosin bipolar thick filament formation: importance of charge and specific domains of the myosin rod. *PLoS Biol* **2**: 1880-1892
- Itagaki K, Kannan KB, Singh BB, Hauser CJ (2004) Cytoskeletal reorganization internalizes multiple transient receptor potential channels and blocks calcium entry into human neutrophils. *J Immunol* **172**: 601-607
- Jalink K, van Corven EJ, Hengeveld T, Morii N, Narumiya S, Moolenaar WH (1994) Inhibition of lysophosphatidate- and thrombin-induced neurite retraction and neuronal cell rounding by ADP ribosylation of the small GTP-binding protein Rho. *J Cell Biol* **126**: 801-810
- Kolman MF, Futey LM, Egelhoff TT (1996) Dictyostelium myosin heavy chain kinase A regulates myosin localization during growth and development. *J Cell Biol* **132**: 101-109
- Kozak JA, Cahalan MD (2003) MIC channels are inhibited by internal divalent cations but not ATP. *Biophys J* **84**: 922-927
- Liang W, Licate L, Warrick H, Spudich J, Egelhoff T (2002) Differential localization in cells of myosin II heavy chain kinases during cytokinesis and polarized migration. *BMC Cell Biol* **3**: 19
- Linder S, Aepfelbacher M (2003) Podosomes: adhesion hot-spots of invasive cells. *Trends Cell Biol* **13**: 376-385

- Lockwich T, Singh BB, Liu X, Ambudkar IS (2001) Stabilization of cortical actin induces internalization of transient receptor potential 3 (Trp3)-associated caveolar Ca²⁺ signaling complex and loss of Ca²⁺ influx without disruption of Trp3-inositol trisphosphate receptor association. *J Biol Chem* **276**: 42401-42408
- Nadler MJ, Hermosura MC, Inabe K, Perraud AL, Zhu Q, Stokes AJ, Kurosaki T, Kinet JP, Penner R, Scharenberg AM, Fleig A (2001) LTRPC7 is a Mg.ATP-regulated divalent cation channel required for cell viability. *Nature* **411**: 590-595
- Nakasawa T, Takahashi M, Matsuzawa F, Aikawa S, Togashi Y, Saitoh T, Yamagishi A, Yazawa M (2005) Critical regions for assembly of vertebrate nonmuscle myosin II. *Biochemistry* **44**: 174-183
- Rico M, Egelhoff TT (2003) Myosin heavy chain kinase B participates in the regulation of myosin assembly into the cytoskeleton. *J Cell Biochem* **88**: 521-532
- Roskoski R, Jr. (2004) Src protein-tyrosine kinase structure and regulation. *Biochem Biophys Res Commun* **324**: 1155-1164
- Runnels LW, Yue L, Clapham DE (2001) TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science* **291**: 1043-1047
- Runnels LW, Yue L, Clapham DE (2002) The TRPM7 channel is inactivated by PIP(2) hydrolysis. *Nat Cell Biol* **4**: 329-336
- Ryazanov AG (2002) Elongation factor-2 kinase and its newly discovered relatives. *FEBS Lett* **514**: 26-29
- Ryazanova LV, Dorovkov MV, Ansari A, Ryazanov AG (2004) Characterization of the protein kinase activity of TRPM7/ChaK1, a protein kinase fused to TRP ion channel. *J Biol Chem* **279**: 3708-3716
- Schild D, Jung A, Schultens HA (1994) Localization of calcium entry through calcium channels in olfactory receptor neurones using a laser scanning microscope and the calcium indicator dyes Fluo-3 and Fura-Red. *Cell Calcium* **15**: 341-348
- Schmitz C, Perraud AL, Johnson CO, Inabe K, Smith MK, Penner R, Kurosaki T, Fleig A, Scharenberg AM (2003) Regulation of vertebrate cellular Mg²⁺ homeostasis by TRPM7. *Cell* **114**: 191-200
- Sun HQ, Yamamoto M, Mejillano M, Yin HL (1999) Gelsolin, a multifunctional actin regulatory protein. *J Biol Chem* **274**: 33179-33182
- Van Etten RA (2003) c-Abl regulation: a tail of two lipids. *Curr Biol* **13**: R608-610
- van Leeuwen FN, Kain HE, Kammen RA, Michiels F, Kranenburg OW, Collard JG (1997) The guanine nucleotide exchange factor Tiam1 affects neuronal morphology; opposing roles for the small GTPases Rac and Rho. *J Cell Biol* **139**: 797-807
- van Leeuwen FN, van Delft S, Kain HE, van der Kammen RA, Collard JG (1999) Rac regulates phosphorylation of the myosin-II heavy chain, actinomyosin disassembly and cell spreading. *Nat Cell Biol* **1**: 242-248
- van Rheenen J, Jalink K (2002) Agonist-induced PIP(2) hydrolysis inhibits cortical actin dynamics: regulation at a global but not at a micrometer scale. *Mol Biol Cell* **13**: 3257-3267
- Wilson JR, Ludowyke RI, Biden TJ (1998) Nutrient stimulation results in a rapid Ca²⁺-dependent threonine phosphorylation of myosin heavy chain in rat pancreatic islets and RINm5F cells. *J Biol Chem* **273**: 22729-22737

Supplemental Material




Figure S1 Characterization of TRPM7 kinase-dead mutant. Mutation of aspartate 1775 to alanine abrogates the ability of TRPM7 to undergo autophosphorylation. WT and kinase-dead TRPM7 constructs were transiently transfected into HEK293 cells and immuno-affinity purified complexes were subjected to *in vitro* kinase assays. Top, *In vitro* autophosphorylation of TRPM7 detected by ^{32}P labeling and autoradiography. Bottom, expression levels of TRPM7 constructs in HEK293 cells detected using anti-HA antibodies.




Figure S2 Quantification of TRPM7 kinase activity in the different N1E-115 cell lines. (A) Autoradiogram of TRPM7 autophosphorylation levels in the different N1E-115 cell lines. N1E-115 cells transduced with either the empty vector, TRPM7-WT or TRPM7-D1775A were lysed in RIPA buffer. Protein concentration was determined using the BCA kit and equal amounts of protein (400 μ g) were transferred to each tube. TRPM7 was immunoprecipitated using a rabbit polyclonal TRPM7 antibody or rabbit pre-immune serum as a negative control. To assess the level of kinase activity, TRPM7 was autophosphorylated in the presence of 5 μ Ci of γ - 32 P-ATP, separated on an 8% SDS-PAGE gel and detected by autoradiography. (B) Quantification of TRPM7 kinase activity. Autophosphorylation of TRPM7 was quantified using phospho-imager analysis. The level of kinase activity in control N1E-115 cells was set to 1 and the data from the other cell lines is expressed as the kinase activity in relation to control cells (n=6).




Figure S3 TRPM7 is glycosylated. (A) Cells were treated with vehicle control or 2 $\mu\text{g/ml}$ tunicamycin for 24 h. TRPM7 was immunoprecipitated and detected by Western blotting using anti-HA antibodies. Note that while TRPM7 runs as a doublet at 216 kDa, only the fast migrating band is observed upon treatment with tunicamycin. (B) Glycosylation is independent of kinase activity. TRPM7-WT and TRPM7-D1775A were detected by immunoprecipitation followed by Western blotting using anti-HA antibodies. Note that both TRPM7-WT and TRPM7-D1775A migrate as doublets on SDS-PAGE gels indicative of glycosylation. (C) Expression of TRPM7-WT and TRPM7-D1775A at the cell surface. N1E-115 cells were transfected with YFP-TRPM7-WT or CFP-TRPM7-D1775A. Live cells were visualized by confocal microscopy 48 h post-transfection. Cell surface expression was also detected in N1E-115/TRPM7-WT and N1E-115/TRPM7-D1775A using anti-HA antibodies (Fig. 4 and data not shown).




Figure S4 Inhibition of myosin II function in N1E-115 cells leads to cell spreading and podosome formation. N1E-115 cells were seeded on fibronectin-coated coverslips to increase their adherence. Cells were treated with 50 μM blebbistatin for 30 min, fixed and stained for vinculin (green) and F-actin (red). (A) Untreated N1E-115 cells. Note the appearance of focal adhesions. (B) and (C) Blebbistatin treated N1E-115 parental cells. Scale bar =10 μm .




Figure S5 Disruption of the actin cytoskeleton prior to TRPM7 purification does not affect the phosphorylation of myosin IIA. (A) Effect of latrunculin A on the phosphorylation of myosin IIA by TRPM7. TRPM7-C WT and KD were immunopurified using anti-HA antibodies under high stringency conditions before and after latrunculin A treatment (1 μ M; 30 min). The kinase was incubated with 2 μ g of GST or GST-myosin IIA in kinase buffer. Phosphorylated proteins were detected by SDS-PAGE followed by autoradiography. Top, coomassie staining of GST-fusion proteins; Middle, autoradiogram showing that phosphorylation of GST-myosin IIA is unaffected by prior treatment with latrunculin A; Bottom, autophosphorylation of TRPM7-C constructs. (B) Contaminating actin is below detection levels. An aliquot of the purified TRPM7-C WT and KD from experiment in section A was run on gel and probed for β -actin (Top) or HA-tagged proteins (Middle). The bottom panel shows β -actin in total lysates as a control for protein levels.




Figure S6 Relationship between TRPM7 expression, myosin activity and cellular phenotype. Parental N1E-115 cells have high myosin II activity which is translated into a contractile phenotype where cells are round and poorly adherent. Stimulation of these cells with BK leads to a loss of contractility as observed by the spreading of cells. However, myosin II is not fully inhibited as focal adhesions form at the periphery of the cell. Overexpression of TRPM7 mimics the effect of BK on N1E-115 cells as it has basal activity. Stimulation of N1E-115/TRPM7 cells leads to a further inhibition of myosin II activity leading to the formation of podosomes. Directly inhibiting all myosin II activity leads to the formation of podosomes regardless of whether TRPM7 is present or not. Thus, the level of TRPM7 expression and activation dictates the extent of myosin II activity and thereby the phenotype of N1E-115 cells.