



Universiteit
Leiden
The Netherlands

Immune evasion by varicelloviruses : the identification of a new family of TAP-inhibiting proteins

Koppers-Lalić, D.

Citation

Koppers-Lalić, D. (2007, September 11). *Immune evasion by varicelloviruses : the identification of a new family of TAP-inhibiting proteins*. Retrieved from <https://hdl.handle.net/1887/12381>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/12381>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER

1

Introduction

The families of herpesviruses

The general outlook in the herpesvirus field has gradually changed the last 5 years, predominantly due to the discovery of a number of herpesviruses in fish, amphibians and mollusks (1-3). This has led to the recent proposal to create an order of *Herpesvirales* divided into three families: the *Herpesviridae*, the *Alloherpesviridae* and the *Malacoherpesviridae* (4). The *Alphaherpesvirinae*, the *Betaherpesvirinae* and the *Gammaherpesvirinae* still remain the only subfamilies of the *Herpesviridae*.

Besides their large double stranded DNA genome (ranging from 124 kbp to 295 kbp), all members of the *Herpesvirales* share another common property, namely their virion architecture. The DNA genome is found in an icosahedral (T=16) capsid of 115 to 130 nm with a specific portal complex. The capsid is surrounded by a highly ordered structure named tegument that supports the surrounding envelope (5). Beside structural proteins, the tegument contains such molecules as the virion host shutoff (vhs) protein, which functions as soon as the virus enters the cell (6).

At present, 56 complete herpesvirus genomes have been published. The genomes have 70 to 165 genes (but this number may go up to 200 when the larger viruses have been fully sequenced). At least one gene, encoding a DNA packaging enzyme, has a homolog in all families (2). Within the family of *Herpesviridae*, 43 genes are conserved in all members and were obviously inherited from a common ancestor (7). These conserved genes are named 'core' genes and they encode proteins that function predominantly in DNA replication, capsid formation and capsid egress, or are part of the tegument or envelope. The UL49.5 or UL49A gene that encodes homologs of glycoprotein N (gN) belongs to the 'core' genes and is thus found in all presently sequenced members of the *Herpesviridae*. Core genes are thought to play a fundamental role in the replication of the *Herpesviridae*, in contrast to the 'non-core' genes that have a more accessory function (2). The UL41 gene encoding the vhs protein is an example of a 'non-core' gene that is only found in *Alphaherpesvirinae*.

The last 10 years it has become clear that herpesviruses have developed many mechanisms to counteract the host's innate and acquired immune responses. For example, the gD protein of alphaherpesviruses can postpone apoptosis before the virus enters the cell (8). The vhs protein present in the virion not only down-regulates the host gene expression, it can also specifically counteract the interferon type I response of the host cell (9). For example, the product of the immediate-early gene of bovine herpesvirus 1 ICP0 (infected cell protein 0; BICP0) not only acts on viral promoters, but also inhibits interferon-dependent promoters (10). Herpesviruses have developed many additional mechanisms to evade the acquired immune response, as will be discussed later on. The fact that immune evasion genes belong to the 'non-core' or 'accessory' genes may reflect a more extensive form of adaptation to the host defense mechanisms.

Co-evolution of herpesviruses with their hosts

Viruses co-evolve with their hosts through adaptation to the host's cellular processes/pathways that they exploit. The best adapted virus will successfully maintain co-existence with its host (reservoir) and it will consequently ensure the survival of both. For example, the ability to escape their host's immune response represents an advantageous evolutionary adaptation from the virus's perspective.

The common ancestor of all the *Herpesvirales* is thought to have existed 570-505 million years ago (mya), around the time the first vertebrates appeared (4). The most recent common ancestor of the *Herpesviridae* is estimated to have emerged 400 mya (11). This is

long before the appearance of the most recent ancestor of the mammals (around 220 mya). The phylogenetic trees of the *Herpesviridae* and their hosts have roughly the same form, implying a co-evolution between viruses and hosts. This is particularly convincing for clades of the *Alphaherpesvirinae* and their hosts (12). This synchronous development implies a co-adaptation between the virus and its host.

Co-evolving with their hosts, herpesviruses have acquired elaborate strategies for evading the host's immune response. For this purpose, herpesviruses have exploited multiple cellular pathways, including the MHC class I antigen-processing pathway. This appears to be the principal means by which herpesviruses interfere with CD8⁺ T cell function, and it is certainly the best characterized until now. Numerous herpesviruses-encoded proteins that interfere with MHC class I antigen presentation have been described and it appears that these viral proteins are able to interfere at essentially every step of this pathway (reviewed in 13 and 14).

The MHC class I antigen processing and presentation pathway

T-lymphocytes or T-cells recognize antigenic peptides presented on the surface of host cells by MHC molecules (reviewed in 15). T-cells owe their specificity to the T-cell receptor (TCR). The TCR interacts with two types of MHC molecules: MHC class I and MHC class II. TCR-mediated recognition of class I and class II molecules works in combination with one of two co-receptors, CD8 or CD4, respectively.

MHC class I molecules are type I membrane glycoproteins associated with the small subunit β 2microglobulin (β 2m) found on all nucleated cells in vertebrates. MHC class I molecules mainly present peptides of endogenous proteins to CD8⁺ cytotoxic T-cells that scan for displayed peptide repertoire (figure 1). Detection of non-self peptides derived from endogenous pathogens or mutated proteins will trigger the elimination of antigen-presenting cells.




Fig. 1. Model of the MHC class I antigen presentation pathway. MHC class I molecules folding and complex formation and assembly of peptide loading complex (see text for further description). *Adopted from Abele & Tampe FEBS Lett. 2006 580:1156-1163*

MHC class II molecules exist as heterodimers of two transmembrane glycoprotein subunits, the α and β chain and are found only on the cell surface of specialized immune cells (dendritic cells, monocytes, macrophages, B-cells and thymic epithelial cells). They present peptide fragments of exogenous proteins. CD4⁺ T helper cells recognize peptide-MHC class II complexes and stimulate B-cells to produce antibodies.

Generation and processing of antigenic peptides

Most MHC class I molecules acquire their peptide cargo in the ER, and most of these peptides are generated from cytosolic substrates. Proteins can reach the cytosol either by being synthesized by ribosomes or by penetrating the plasma membrane through internalization. Endogenous peptides appear to originate primarily from defective forms of newly synthesized proteins (termed DRiPs, for defective ribosomal products) (16) that are rapidly degraded by the ubiquitin-proteasome pathway (17-19). In general, ubiquitination targets misfolded proteins as well as DRiPs for degradation by the proteasome (20). Proteasomes are large multi-subunit complexes with protease activity (21, 22). They are responsible for the bulk turnover of proteins in the cell, therefore maintaining homeostasis between anabolic and catabolic pathways (15). A direct function for proteasomes in the generation of antigens for class I presentation has been demonstrated to be essential for this process (18, 23, 24). Additional cytosolic proteases can further cut and trim proteasome-generated peptides (e.g. tripeptidyl peptidase II – TPPII; 25). Treatment of cells with interferon- γ or its production due to inflammation stimulates the formation of immuno-proteasomes, where the catalytically active subunits are replaced by LMP2, LMP7, and MECL1 (low molecular mass polypeptides 2 and 7; multi-catalytic endopeptidase complex-like 1) (26, 27). The resulting exchange gives a higher quantity of antigenic peptides with suitable carboxyl-termini and of the approximate length (6 to 11 amino acids) preferred by MHC class I molecules (28). However, only a fraction of these peptides makes its way to the lumen of the ER (29). Through the Transporter associated with Antigenic Processing (TAP)-mediated transport (30), peptides reach the endoplasmic reticulum (ER) (figure 1) where they subsequently face the aminopeptidase associated with antigen processing (ERAAP, also named ER aminopeptidase 1 ERAP1), which is capable of trimming the amino-termini of certain peptides. This increases the binding affinity of these peptides to MHC class I and the fraction of ER luminal peptides available for stable presentation (31, 32).

MHC class I/peptide-loading complex assembly

Proper folding and assembly of MHC class I/ β 2m molecules with peptides is supposed to be controlled by a number of accessory proteins and chaperones that are, to a varying degree, dedicated to this process (figure 1). The present model is that MHC class I heavy-chain molecules (approximately 45 kDa) are translated into the ER and glycosylated co-translationally. Addition of the single N-linked glycan on class I heavy-chain molecules serves as a point of recognition for the ER membrane chaperone calnexin (the ER-resident lectin) (33). Together with calnexin, the immunoglobulin-binding protein (BiP) interacts transiently with the newly synthesized heavy chain. Both chaperones promote folding and assembly of the class I heavy-chain with soluble β 2m (12 kDa). Calnexin is replaced by the ER-resident lectin calreticulin (34) after the heterodimer of heavy chain and β 2m has achieved its proper conformation (figure 1).

Another ER chaperone, the thiol-oxidoreductase ERp57, was found to be involved in the maturation of class I heavy-chain/ β_2m molecules. ERp57 assists with the disulfide bond formation probably at the early stage of class I heavy-chain folding (34). At the later stages of assembly, MHC class I-associated ERp57 can be found as part of the peptide-loading complex where ERp57 is covalently bound to tapasin. The ERp57-tapasin conjugate is dependent on the presence of MHC class I (35, 36).

Peptide loading of class I molecules does not proceed efficiently unless the calreticulin/class I heavy-chain/ β_2m /ERp57 complex binds to tapasin (figure 2), an MHC class I chaperone




Fig. 2. Model of peptide-loading complex (PLC) (see text for further description). Adapted from Abele & Tampe *FEBS Lett.* 2006 580:1156-1163

required for the association of the complex with TAP (37-40). Tapasin was found to be in concurrent association with the class I heavy-chain and both subunits of TAP (TAP1 and 2). This multi-component association results in the formation of the MHC class I peptide-loading complex (PLC; figure 2) that facilitates the binding of the antigenic peptide within the peptide-binding groove of the heavy chain. Stable MHC class I/ β_2m /peptide complexes are formed and released from the loading complex to be further translocated via the Golgi apparatus to the cell surface (figure 1) (41, 42).

Key players of peptide-loading complex assembly: tapasin and TAP

Essential building elements of the peptide-loading complex are represented by two TAP subunits (TAP1 and 2) associated with four tapasin and four MHC class I molecules (43).

Tapasin

Tapasin, an ER-resident type I membrane glycoprotein, is the central stabilizer of the peptide-loading complex and interacts with almost all proteins involved in the formation of this multimeric structure (44). The N-terminal domain of tapasin (the ER luminal part) is necessary for the interaction with MHC class I, calreticulin and ERp57, whereas the C-terminal domain (mainly the transmembrane region) of tapasin is sufficient for the interaction with TAP (45). By having discrete binding sites at opposite ends of the molecule, tapasin bridges TAP and class I molecules, thereby facilitating an increase of the local concentration of peptides for MHC class

I loading (46, 47). The MHC class I molecules deprived of peptides will remain associated with tapasin, which retains unloaded molecules within the ER (48). In the TAP-deficient human cell line T2, the association of tapasin with class I molecules appears to be independent of tapasin/TAP interaction. Furthermore, the MHC class I molecules expressed in the tapasin-deficient human cell line 721.220 (37, 38) or in tapasin knockout mice (49, 50) do not interact with TAP. In these cell lines, there is a reduced surface expression of peptide-loaded MHC class I (51) as well as a decrease in cell surface stability of peptide-bound class I molecules (38, 52). These findings signify an important role of tapasin in MHC class I-peptide assembly (53). Surprisingly, the binding affinities of the MHC class I-associated peptides are not altered in the presence or absence of tapasin (46), but the variety of bound peptides is increased when tapasin is associated with MHC class I molecules. Thus, cells containing functional tapasin exhibit improved MHC class I-peptide complex formation and stability, which in its turn provides enhanced levels of MHC class I complexes at the cell surface.

Transporter associated with Antigen Processing (TAP)

TAP belongs to the large family of ABC transporters (54, 55). The family of ABC transporters is defined by their homology within the ATP-binding cassette (ABC) region (56). This region (also known as the nucleotide-binding domain) contains three highly conserved motifs called Walker A and B motifs (figure 3B) as well as the C loop (also known as the ABC signature motif). The Walker A and B motifs are present in many ATP-binding proteins, whereas the C loop is characteristic for ABC transporters. Generally, all ABC transporters consist of a four-domain structure with two hydrophobic transmembrane domains (TMDs) and two hydrophilic nucleotide-binding domains (NBDs). The two NBDs located in the cytoplasm are involved in ATP binding and/or hydrolysis. The NBDs act as “motor domains”, since they convert the chemical energy of ATP hydrolysis into mechanical work. This process induces conformational changes within the TMDs to enable the passage of substrates.




Fig. 3. Model of transporter associated with antigen processing TAP. A Schematic model of the TAP complex. TAP forms a heterodimer composed of TAP1 and TAP2. Each subunit comprises an N-terminal transmembrane domain and a C-terminal, cytosolic NBD (red). The transmembrane domain can be subdivided into a six helices containing core domain and an NH₂-terminal extensions (tapasin-binding domains) of four and three helices for TAP1 and TAP2, respectively. Beside the translocation pathway, the TMDs also form the peptide-binding region (orange). *Adapted from Abele & Tampe FEBS Lett. 2006 580:1156-1163.* B TAP forms a transmembrane pore in the ER. The pore is followed by a peptide binding domain located at the cytoplasmic side. The structure is concluded by the two ATP-binding domains (see text for further description). *Adapted from Reits et al. Immunol Today 2000 21:518-600.*

Structural organization of the TAP complex

The peptide transporter TAP is a heterodimer composed of TAP1 (human ABCB2; calculated mol. mass 81 kDa) and TAP2 (human ABCB3; calculated mol. mass 75 kDa), each consisting of an N-terminal transmembrane domain (TMD) and a C-terminal, cytosolic nucleotide-binding domain (NBD) (figure 3A) (57, 58). Both subunits are essential and sufficient for antigen processing (59, 60). TAP is localized in the ER and *cis*-Golgi (61). Neither an N-terminal signal peptide for ER import nor a retention signal for any subunit is known as yet.

The topology of TAP is still under study. Hydrophobicity analysis and sequence comparison with members of the ABC-B subfamily imply that TAP1 and TAP2 contain 10 and 9 transmembrane segments (TMs), respectively (figure 3A). Only the last six transmembrane segments of both subunits show sequence homology to other ABC transporters. Interestingly, a peptide transporter missing the N-terminal domains of four and three putative transmembrane segments of TAP1 and TAP2, respectively, is still active with respect to peptide binding and transport (62). The N-terminal extensions of both subunits are required for the interaction with the transmembrane domain of tapasin (44, 63). However, the precise binding site(s) for tapasin is (are) still unknown (58).

Tapasin is usually present within a functional TAP1 and TAP2 complex and plays an important role in the assembly and stability of TAP. However, it has also been found to bind TAP1 or TAP2 subunits individually. Generally, tapasin stabilizes the TAP1 protein, resulting in a higher steady-state protein expression level of TAP (64). In tapasin-deficient cells or cells expressing tapasin mutants, which are unable to bind to TAP, the steady-state level of TAP is decreased up to 100-fold (44). Therefore, by promoting the TAP heterodimer assembly and stabilizing the peptide-binding site of TAP, tapasin enhances the peptide translocation rate acting as a “chaperone” for TAP (44, 45, 51, 64, 65).

The peptide-binding region of TAP is localized in the last cytosolic loop and a 15-amino acid extension of the last putative transmembrane segments of TAP1 and TAP2 (66). The NBDs of TAP convert the energy of ATP hydrolysis to peptide transport. For such multi-domain machinery, a multi-step process is plausible, where peptide and ATP binding, ATP hydrolysis, and peptide transport occur in an ordered manner. As known from biochemical studies on isolated NBDs, ATP-binding induces a conformational change of the NBD, enabling the dimerization of the catalytic domain (figure 3B), which is a prerequisite for ATP hydrolysis and substrate transport (67-69).

TAP also fulfils the function of a chaperone for the other members of the loading complex. The assembly of class I molecules, ERp57, calreticulin, and tapasin is less efficient in the absence of TAP than in its presence (70). Moreover, class I adopts a different conformation in the absence of TAP than in its presence (71). TAP may also serve as a relay to transmit a signal for dissociation of peptide-loaded MHC class I complexes (72).

How does TAP translocate peptides across the ER membrane?

As already stated, TAP is composed of two subunits, TAP1 and TAP2. Both subunits contain a multi-membrane spanning domain forming the actual pore and also encompassing the peptide-binding site. In addition, TAP1 and TAP2 each carry a hydrophilic nucleotide-binding domain (figure 3B) (reviewed in 73-75). The actual position of TAP1 and TAP2 regarding their orientation within the functional heterodimer still remains questionable, although a head-head/tail-tail orientation has been proposed (58, 76). Studies on tapasin interaction with TAP indicated that the first four and three TMDs of TAP1 and TAP2, respectively, are not essential

for the formation of the functional peptide translocation pore and most likely provide two separate platforms for tapasin/class I association (63).

Although the TAP transporter has been the subject of numerous studies, it has proved very difficult to assess the sequence of events during the peptide transport cycle. The following model for peptide transport by TAP could be proposed (figure 4 modified after 74; reviewed in 73 - 75): Peptide and ATP bind independently to TAP (step 1). ATP binding enables TAP1 and TAP2 NBDs to dimerize. It has been suggested that the binding of ATP first occurs at the TAP2 NBD site (NBD2) which is also the first to be hydrolyzed. Also, ATP hydrolysis in NBD2 is sufficient for peptide transport by weakening the dimer interface. However, ATP hydrolysis in TAP1 NBD (NBD1) alone does not interrupt the dimer interface. ATP can only be hydrolyzed at NBD1 after ATP hydrolysis has occurred at NBD2. Peptide binding (step 2) induces




Fig. 4. Proposed peptide transport cycle of TAP. See text for description of each step of the peptide transportation cycle. P-phosphate. Modified after Abele and Tampe *Physiology* 2004 19:216-224.

structural rearrangements in the TMDs of both subunits (step 3), which weakens rigid interactions within NBDs. As a result, ATP-loaded NBDs form a dimer, which subsequently induces peptide release into the ER lumen (step 4). The peptide is released from the altered peptide-binding site, and subsequent translocation through the opened pore (TMDs) into the ER lumen occurs. ATP hydrolysis and phosphate release, first at NBD2 (step 5) and then at

NBD1 (step 6) weaken the interaction of the NBDs. ATP hydrolysis at NBD1 may be required to close the pore, restore the peptide-binding domain and complete the translocation cycle. After the dimer dissociation (step 7), ADP is released and replaced by ATP.

Even though the transport activity of TAP can be directly measured with model substrates (77), conformational changes that occur during transport are difficult to study. However, the conformational changes associated with activity of TAP can, to a certain extent, be visualized in living cells by measuring the lateral mobility of fluorescent GFP-tagged TAP over the ER membrane (78). For this purpose, fluorescence recovery after photo-bleaching (FRAP) technique was used to visualize the lateral mobility and dynamics of TAP complexes in living cells (79). By using this technique, it has been shown that the lateral mobility of TAP molecules is inversely proportional to TAP activity, as peptide transporting TAP molecules diffuse at a slower rate than inactive, closed TAP complexes. Thus, the mobility of inactive TAP is higher when compared to TAP actively pumping peptides.

Apparently, the decrease in mobility is likely due to conformational changes within the TAP complex during pore opening and peptide transfer. In addition, the effect of chemical and viral TAP inhibitors can be visualized through confocal microscopy, the FRAP technique and the measurement of lateral mobility of TAP, provided they affect the conformational changes within the TAP complex (78, 80).

Herpesviruses: masters of disguise

To establish a lifelong infection in the presence of an active (immunocompetent) host immune system, herpesviruses have acquired an impressive array of immune modulatory strategies that contribute to their successful long-term persistence. The success of these strategies relies on virus-encoded proteins that either mimic specific components of the host's immune system or target a variety of host immune surveillance mechanisms. The effectiveness of these viral proteins in escaping from host immune surveillance, ultimately contributes to lifelong infection and pathogenesis of herpesviruses. The number of identified virus-encoded proteins with immune evasion properties has increased dramatically in recent years. In this thesis, only a subset of immune evasion molecules will be covered with the accent on proteins encoded by herpesviruses that target the MHC class I antigen presentation pathway (summarized in table 1).

The virion host shut-off (vhs) mechanism

The shut-off of cellular protein synthesis by the virion host shut-off (vhs) protein is a characteristic feature of alphaherpesviruses. The vhs protein of herpes simplex virus-1 (HSV-1), the prototype alphaherpesvirus, is the best characterized vhs protein. Vhs homologs have been identified in the genome of all the alphaherpesviruses sequenced thus far. However, only a few have been analyzed for host protein synthesis shut-off properties (81-83). The vhs protein, encoded by the HSV-1 UL41 gene, is a tegument protein, which enters the host cell as a component of the infectious virion. Upon entry, the virion vhs protein (but also the vhs proteins that is expressed during the early stage of infection) accelerate the degradation of mRNA, contributing to an overall decrease in host protein synthesis (84-86). It has been shown that during the early stages of BHV-1 infection, the BHV-1 UL41 gene product also strongly affects the synthesis of host proteins, among which the MHC class I molecules. These findings indicate that the BHV-1 vhs protein plays a role in MHC class I down-regulation. In addition to the vhs protein activity, other BHV-1 gene products specifically down-regulate MHC class I

molecules. Thus, the suppression of host gene expression seems to support immune evasion in the early stage of infection (81, 87-89). Apparently, most herpesviruses do not concentrate on a single immuno-evasion strategy but utilize several strategies in parallel.

Table 1

Alphaherpesviruses	Protein	Target	Mechanism (References)
HSV-1& -2	ICP47	TAP	Inhibition of peptide binding (137-138)
BHV-1	UL49.5	TAP	Inhibition and degradation of the PLC (80, 168)
PRV	UL49.5(gN)	TAP	Inhibition of peptide transport (80)
EHV-1& -4	UL49.5	TAP	Inhibition of ATP binding (80, *)
VZV	ORF66	MHC I	Retention in Golgi (159, 170)
Betaherpesviruses			
HCMV	US2	MHC I	Dislocation and proteasomal degradation (91-95)
	US3	MHC I/tapasin	ER retention/ Inhibition of peptide loading (101-105)
	US6	TAP	Inhibition of ATP binding (143-148)
	US10	MHC I	Retention in the ER (106)
	US11	MHC I	Dislocation and proteasomal degradation (91-95)
MCMV	m04	MHC I	Binding to MHC/display at cell surface (110-111)
	m06	MHC I	Degradation in lysosomes (112)
	m152	MHC I	Retention in the ER (108-109)
Gammaherpesviruses			
EBV	EBNA-1	Proteasome	Prevention of proteasomal degradation (156,157)
	vIL-10	TAP	Inhibition of TAP1 expression (154)
	BNLF2a	TAP	Inhibition of peptide transport (155; Hislop, Rensing, Rickinson, Wiertz et al., Pers. comm.)
HHV-7	U21	MHC I	Degradation in lysosomes (127,128)
KSHV	K3	MHC I	Ubiquitinylation/endocytosis in lysosomes (114-119)
	K5	MHC I	Ubiquitinylation/endocytosis in lysosomes (114,115)
MHV-68	mK3	MHC I	Ubiquitinylation/proteasomal degradation (121-126, 149-151)
		TAP/tapasin	Proteasomal degradation (122,149-151)

* Chapter 5 of this thesis

Targeting of the MHC class I complex by viral proteins

The use of multiple strategies to subvert antigen presentation has been clearly illustrated for human cytomegalovirus (HCMV) (90). The US2-US11 (US, unique short region) of HCMV potentially encodes 8-10 membrane glycoproteins, four of which, US2, US3, US10 and US11, are known to target MHC class I molecules and one, US6, specifically inhibits TAP-mediated peptide transport.

US2 and US11, both type-I membrane glycoproteins, induce rapid proteasome-dependent degradation of nascent MHC class I molecules. Located in the ER, they both bind directly to MHC class I heavy chains and cause them to be removed from the ER back to the cytosol through a process termed "dislocation" (retro-translocation) (91-95). It is known that dislocation is a central component of ER associated protein degradation (ERAD), an important mechanism used by cells to degrade improperly assembled or misfolded ER proteins (96). Whereas the ER-luminal domain of US11 is involved in the binding of MHC class I heavy chains, its transmembrane domain is essential for their dislocation (97). The transmembrane domain of US11 mediates the recruitment of MHC class I molecules to Derlin-1, a protein essential for the degradation of misfolded ER-luminal proteins. US11 bridges Derlin-1 with MHC class I, probably delivering MHC class I molecules to the dislocation pore (98, 99). Interestingly, Derlin-1 is not required for US2-mediated degradation of MHC class I molecules, indicating different mechanisms of action (98). Recently, signal peptide peptidase (SPP) has been identified as a partner for the active form of US2, implicating SPP in the US2-mediated MHC class I degradation pathway. Apparently, a decrease in SPP levels inhibits heavy-chain dislocation by US2 but not by US11 (100).

The HCMV-encoded US3 and US10 proteins retain MHC complexes in the ER. US3, the only immediate early US gene, interacts transiently with MHC molecules and prevents their egress to the cell surface (101-103). The transmembrane and the ER luminal domain of US3 are both required for the retention of MHC class I molecules in the ER (102, 104). However, substantial amounts of MHC class I molecules still escape US3-mediated ER arrest. Apparently, US3 down-regulates the cell surface expression of MHC class I molecules in an allele-specific manner. US3 directly binds tapasin, thereby affecting only MHC class I alleles that are dependent on tapasin for their surface expression. Through its interactions, US3 reduces the association between TAP and tapasin. While US3 binds to both tapasin and TAP, it was evident that US3 did not affect TAP function for peptide transport (105).

Like US3, US10 also inhibits the export of MHC class I out of the ER (106). US10 is an ER-localized type I membrane glycoprotein that delays but does not block the transport of MHC class I molecules from the ER. However, the mechanism of interaction between US10 and class I heavy chains remains to be determined.

Thus, HCMV encodes at least 5 proteins with specific tasks to obstruct MHC class I assembly and presentation. Interestingly, two of these, US2 and US3, appear to be multifunctional as they both also target MHC class II molecules, preventing their egress to the cell surface (107).

Murine cytomegalovirus (MCMV) carries three genes encoding type-I membrane glycoproteins known to interfere with the intracellular trafficking of MHC class I molecules: m152/gp40, m04/gp34 and m06/gp48. These viral proteins are expressed sequentially during viral infection and they act synergistically. Whereas gp40 can be detected within a few hours after infection, gp34 and gp48 are expressed later. Although the direct interaction of gp40 with MHC class I could not be detected, the expression of gp40 protein triggers the retention of MHC class I complexes in the ER-Golgi intermediate compartment (ERGIC)/*cis*-Golgi (108, 109). These MHC class I complexes are stable, whereas gp40 is re-directed and degraded in endosomal/lysosomal compartments (109). In contrast to gp40, gp34 and gp48 are both found

in direct association with MHC class I molecules. gp34 does not alter MHC class I surface expression, but it is found in association with MHC class I molecules at the cell surface (110, 111), whereas gp48-MHC class I complexes are diverted to late endosomes and lysosomes for rapid proteolysis. The lysosomal targeting of gp48-bound MHC class I molecules is mediated by the cytosolic domain of gp48 that contains a di-leucine-motif (LL) (112).

Herpesvirus 8 or Kaposi's sarcoma associated virus (KSHV) encodes two proteins, K3 and K5, which are, until now, unique for KSHV. K3 and K5 are expressed during the early lytic cycle of viral replication and they exhibit 40% of amino acid sequence identity to each other. K3 and K5 are also known as modulator of immune recognition 1 and 2 (MIR1 and MIR2) respectively (113). In their presence, MHC class I molecules exit the ER and reach the cell surface, but are then rapidly endocytosed and degraded in lysosomal compartments (114, 115). K3 and K5 are predominantly associated with the ER. They are type III membrane proteins containing two transmembrane helices. The N-terminus and C-terminus of K3 and K5 are placed in the cytosol. K3 and K5 have an N-terminal RING finger motif that shares structural and sequence homology with the RING finger domain that is found in a subset of E3 ubiquitin ligases. Thus, both proteins belong to the family of type 3 (E3) ubiquitin ligases (116, 117) and can regulate the last step of the ubiquitination cascade (114). The N-terminal domain and the central sorting motif are involved in triggering the internalization of the MHC class I molecules and in re-routing them to the trans-Golgi network (TGN). The C-terminal cluster is engaged in targeting MHC class I molecules to lysosomal compartments (118). K3-mediated ubiquitination of MHC class I molecules signals for both internalization and sorting to late endosomes (119, 120). K3 and K5 both down-regulate MHC class I molecules in a similar way, but they differ in their specificity for MHC class I alleles (114, 115).

Murine γ -herpesvirus 68 (MHV-68) is a γ 2-herpesvirus of small rodents that has a homology to the Kaposi's sarcoma-associated herpesvirus (KSHV) in humans. MHV-68 encodes K3 protein (mK3), a zinc binding RING finger protein, that down-regulates MHC class I molecules expression. mK3 is a type III membrane protein localized in the ER membrane where it binds to the cytoplasmic tail of newly synthesized MHC class I molecules. mK3 catalyses ubiquitination of MHC class I through its N-terminal RING-CH domain and causes their rapid, proteasome-dependent degradation (121-126, 150, 169).

A human herpesvirus-7 (HHV-7)-encoded type I membrane glycoprotein, U21, employs a different strategy to stop MHC class I molecules on their journey to the cell surface membrane. U21 binds to MHC class I shortly after its synthesis in the ER and targets MHC class I complexes to lysosomes for degradation (127). Apparently, the ER-luminal domain of U21 is sufficient for the induction of lysosomal degradation (128). U21 shares no significant sequence homology with any other presently known protein, except for U21 of HHV-6.

TAP under attack

Viral proteins can prevent MHC class I presentation at the cell surface by targeting MHC class I molecules directly or through inhibiting TAP-mediated peptide transport. The blocking of MHC class I antigen presentation is a strategy employed by different herpesviruses, and several of them have chosen TAP as the target (figure 5). Peptide transport into the ER can be impaired by different mechanisms, including down-regulation of TAP expression, proteasomal degradation of TAP mediated by viral proteins, and blocking peptide transport by direct interaction of viral proteins with TAP complex (reviewed in 14, 58, 129 and 130).

TAP is a target of ICP47 (Infected Cell Protein 47) of HSV-1 and -2. Although the ICP47

protein (encoded by immediate-early gene US12) of both serotypes share an overall sequence identity of only 42%, they do not differ significantly in their effect on TAP (131). HSV-1 and 2 encoded ICP47s share no relevant sequence homology with other proteins, except with homologues from simian agent 8 (SA8; 132), monkey B virus (Cercopithecine herpesvirus 1; CeHV-1), and baboon herpesvirus 2 (herpesvirus papio 2; HVP-2). ICP47 is expressed as cytosolic protein, of which its N-terminal domain of 32 residues has been identified as the minimal active region required for TAP inhibition (133, 134). This domain appears to be unstructured in an aqueous solution, but an alpha-helical structure is induced




Fig. 5. Modulation of TAP functions by viral factors. The mK3 protein of MHV-68 binds directly to TAP and induces polyubiquitination and proteasomal degradation. The interaction of UL49.5 of BHV-1 arrests TAP in a transport-incompetent conformation, in which binding of ATP and peptides is not affected. Ultimately, UL49.5 mediates the degradation of TAP by proteasomes. ICP47 of the HSV inhibits peptide binding from the cytosolic side of TAP. The association of US6 of the HCMV to the ER-luminal transmembrane core of TAP blocks ATP binding to the cytosolic NBDs. (for the description of the peptide-loading complex components see figure 2) *Adapted from Abele & Tampe FEBS Lett. 2006 580:1156-1163*

at a membrane interface (135, 136). By binding to TAP from its cytosolic side, ICP47 blocks peptide-binding. Although it does not affect ATP binding, the subsequent ATP hydrolysis is abolished (137-140). Additionally, binding of ICP47 seems to disturb stabilization of the TAP1 and TAP2 complex (137). ICP47 is partially species-specific, since it inhibits human, monkey, canine and bovine TAP function, whereas TAP of small rodents is not affected (131, 137, 141). At present, it is unknown which regions of ICP47 and TAP specifically interact with each other (142).

The unique short region of the HCMV genome encodes yet another immuno-evasion protein called US6. This type I transmembrane glycoprotein blocks TAP-dependent peptide transport. The ER luminal domain of US6 has been shown to be essential and sufficient for TAP inhibition (143, 144). Interestingly, US6 inhibits ATP binding and hydrolysis of TAP at its cytosolic side through interaction with TAP at the ER luminal side. This interaction does not affect peptide binding or the stability of TAP complex (143-147). Although US6 binds to TAP1 and TAP2 independently, the US6-mediated inactivation of the transporter requires simultaneous interaction with conformation-dependent ER-luminal loops of both transporter subunits. The US6 luminal domain is enriched with cysteine residues (8 cysteines) that are likely to form a complex intra- and intermolecular network of disulfide bridges. Indeed, US6 forms oligomers that are apparently involved in the bridging of TAP1 and 2 subunits (148). It is likely that such a compact multimeric formation is crucial for inducing an aberrant structural conformation of the transporter, which in its turn prevents acquisition of ATP.

The murine γ -herpesvirus 68 (MHV-68)-encoded mK3 protein is another example of a viral protein with the ability to interfere with the function of peptide-loading complex. mK3 interacts with MHC class I molecules causing their rapid degradation by proteasomes (121-126, 149, 169, 170). In addition, by association with the peptide-loading complex through its C-terminal tail (150), mK3 initiates the degradation of TAP and tapasin (122, 151). To induce TAP degradation, mK3 requires the transmembrane/cytoplasmic part of tapasin (122). Interestingly, mK3-mediated TAP/tapasin degradation appears to be restricted to certain cell types (lymphoid cell lines). This may have an implication in vivo, as the MHV-68 virus establishes latency in B lymphocytes, macrophages and dendritic cells. Indeed, the major function of mK3 appears to be CTL (cytotoxic T lymphocyte) evasion during viral latency expansion (152).

Epstein-Barr virus (EBV; γ 1-herpesvirus) is another gammaherpesvirus with the ability to inhibit antigen presentation at the level of TAP (153). During the late lytic phase, EBV encodes an IL-10 homolog (viral interleukin-10; vIL-10), involved in down-regulation of TAP1 expression in EBV-infected B cells (154). Moreover, during the early lytic phase of EBV gene expression, specific inhibition in TAP-dependent transport has been observed (155). Recently, the product of the early lytic gene BNLF2a has been identified as an inhibitor of peptide transport by TAP (Hislop, Rensing, Rickinson, Wiertz et al., personal communication).

The list of herpesviruses-encoded “immuno-evaders” has increased dramatically in recent years. Yet there are many more to identify and explore. The group of viruses that belong to the subfamily of *Alphaherpesvirinae*, genus *Varicellovirus*, also specifically target MHC class I antigen presentation. Varicella-zoster virus (VZV), bovine herpesvirus 1 (BHV-1), pseudorabies virus (PRV), equine herpesvirus 1 and 4 (EHV-1 and 4) and many others are members of the genus *Varicellovirus*. For a few members their immuno-evasive strategies have been elucidated only recently (80). The strategies of other members of this genus remain to be further analyzed.

VZV interferes with MHC class I presentation (158) by retaining MHC class I molecules in Golgi compartments of infected cells. However, the mechanism employed for this strategy is still unknown (159). Marek’s disease virus (MDV; genus *Mardivirus*) of chickens is the only known oncogenic alphaherpesvirus inducing T-cell tumors (160). MDV is recognized by virus-specific CTLs (161); however this response appears to be weak, suggesting potential interference with the MHC class I antigen-presentation pathway. Indeed, MDV shows the ability to down-regulate the surface expression of MHC class I molecules during the early phase of virus replication (162, 163).

The studies reported in this thesis focused on bovine herpesvirus 1, pseudorabies virus, and equine herpesvirus 1, which specifically target TAP function (81, 164-166). The aim was to identify the viral gene(s) responsible for the observed TAP inhibition and to characterize the mechanisms employed.

Outline of this thesis

At the start of this study it was unclear whether the down-regulation of the MHC class I cell surface expression observed after infection by BHV-1 is the result of a general or a specific mechanism. The viral host shut-off (vhs) mechanism largely affects host protein expression, and a BHV-1 vhs deletion mutant was constructed and studied for its ability to down-regulate MHC class I. **Chapter 2** describes that the vhs protein was indeed responsible for part of the down-regulation of the MHC class I cell surface expression. On the other hand, the MHC class I molecules that accumulated in the endoplasmic reticulum (ER) after infection with a BHV-1 vhs deletion mutant did not mature into a stable MHC class I-peptide complexes. This observation led to the hypothesis that BHV-1 could also specifically inhibit peptide transport by TAP, which was shown to be the case.

However, studies aiming at the identification of BHV-1-encoded TAP inhibitor have been hampered by the lack of bovine-specific immunological reagents. The experiments described in **Chapter 3** were performed to investigate whether BHV-1 can infect human cells and interfere with the function of human TAP. Interestingly, a human melanoma cell line appeared to be permissive for BHV-1 infection. BHV-1 mediated inhibition of human TAP and down-regulation of MHC class I molecules at the surface of infected cells. The data indicated that the BHV-1-encoded molecule(s) that block bovine TAP are able to interact with homologous components of the MHC class I peptide-loading complex in human cells.

Indeed, the fact that immune evasion by BHV-1 can be studied in human cells facilitated the identification of a new TAP inhibiting protein, UL49.5, encoded by BHV-1. The results reported in **Chapter 4** establish a novel immune evasion strategy for the inactivation of TAP. The BHV-1 encoded UL49.5 protein blocks TAP through a mechanism that is distinct from those employed by other viral inhibitors described above. The effect of BHV-1 UL49.5 on the TAP complex is striking and relies on two events: inhibition of peptide transport by arresting the TAP complex in a translocation-incompetent state and destabilization of both TAP subunits causing their proteasome-dependent degradation.

Furthermore, a characterization of the UL49.5 gene product encoded by various varicelloviruses as a potent inhibitor of TAP is described in **Chapter 5**. PRV and EHV1-encoded UL49.5 homologs have been identified in association with the peptide-loading complex and are responsible for the inhibition of its function. The UL49.5 proteins of BHV-1, PRV and EHV-1 appear to be relatively homologous based on their amino acid sequences identity and structural organization. Remarkably, experimental data revealed seemingly diverse mechanisms employed by UL49.5 homologs to block TAP. Thus, the results described in chapter 5 classify a number of varicellovirus-encoded UL49.5 homologs as members of a novel family of TAP-inhibiting proteins.

BHV-1 UL49.5, a type-I membrane protein of only 9 kDa, is found as a monomer, homodimer and heterodimer associated with the viral glycoprotein M (gM) in infected cells and virion envelopes. In **Chapter 6** it is shown that UL49.5 can combine the interactions with gM and the TAP complex in the context of natural infection. Although UL49.5/gM association

in isolated system can interfere with the inhibition of TAP, a sufficient amount of non-gM bound UL49.5 protein is available to exceed this interference during productive BHV-1 infection. Thus, BHV-1 UL49.5 inhibits TAP despite complex formation with glycoprotein M.

The results presented above already suggested that the BHV-1 UL49.5 and other varicelloviruses encoded homologs scrutinized for the inhibition of TAP represent potential immune suppressive agents for use in humans. **Chapter 7** shows that the varicellovirus-encoded TAP-inhibitor UL49.5 represents a new tool to study and modulate endogenous antigen-presentation. UL49.5 down-regulates HLA class I expression and inhibits minor (mHag) and major Histocompatibility antigen-specific CTL responses more efficiently than the other two known TAP inhibitors, US6 and ICP47. UL49.5 also reduces alloHLA-reactivity, affecting both peptide-dependent and peptide-independent antigen-presentation. Interestingly, the empty MHC class I molecules that appear at the surface of UL49.5-expressing cells can be loaded with peptides exogenously added to the cells. The resulting MHC class I-peptide complexes are stably expressed and can serve to activate T cells of a desired specificity.

In **Chapter 8**, inhibition of TAP by UL49.5 is described to induce the presentation of a new category of peptides, T cell Epitopes associated with Impaired Peptide Processing (TEIPP) (167). Although these peptides are derived from widely distributed self proteins, they are not presented by normal, antigen processing-proficient cells and therefore the immune system recognizes them as immunogenic neo-antigens. TEIPP have been found to be presented by tumor cells that have antigen processing defects. UL49.5 can be used to specifically induce TEIPP-specific T lymphocytes directed against tumor cells, thereby creating a new opportunity for immunotherapy against cancer. Thus, UL49.5 provides a powerful new tool for the study of fundamental aspects of antigen processing and allows specific modulation of the peptide repertoire presented by MHC class I molecules.

In **Chapter 9** the molecular interaction between the UL49.5 protein and the peptide loading complex are discussed together with the possible therapeutic application of the UL49.5 protein and the use of UL49.5 deletion mutants as vaccines.

References

1. **Davison AJ** (1998) The genome of salmonid herpesvirus 1. *J Virol* 72:1974-1982.
2. **Davison AJ** (2002) Evolution of the herpesviruses. *Vet Microbiol* 86:69-88.
3. **Davison AJ, Trus BL, Cheng N, Steven AC, Watson MS, Cunningham C, Le Deuff RM, Renault T** (2005) A novel class of herpesvirus with bivalve hosts. *J Gen Virol* 86:41-53.
4. **McGeoch DJ, Rixon FJ, Davison AJ** (2006) Topics in herpesvirus genomics and evolution. *Virus Res* 117:90-104.
5. **Mettenleiter TC** (2006) Intriguing interplay between viral proteins during herpesvirus assembly or: the herpesvirus assembly puzzle. *Vet Microbiol* 113:163-169.
6. **Strand SS, Leib DA** (2004) Role of the VP16-binding domain of vhs in viral growth, host shutoff activity, and pathogenesis. *J Virol* 78:13562-13572.
7. **McGeoch DJ, Davison AJ** (1999) The molecular evolutionary history of the herpesviruses. In: Domingo E, Webster R, Holland J (Eds.) *Origin and Evolution of Viruses*. Academic Press, London, pp. 441-465.
8. **Zhou G, Avitabile E, Campadelli-Fiume G, Roizman B** (2003) The domains of glycoprotein D required to block apoptosis induced by herpes simplex virus 1 are largely distinct from those involved in cell-cell fusion and binding to nectin 1. *J Virol* 77:3759-3767.
9. **Duerst RJ, Morrison LA** (2004) Herpes simplex virus 2 virion host shutoff protein interferes with type I interferon production and responsiveness. *Virology* 322:158-167.
10. **Henderson G, Zhang Y, Jones C** (2005) The Bovine herpesvirus 1 gene encoding infected cell protein 0 (bICP0) can inhibit interferon-dependent transcription in the absence of other viral genes. *J Gen Virol* 86:2697-2702.
11. **McGeoch DJ, Gatherer D** (2005) Integrating reptilian herpesviruses into the family Herpesviridae. *J Virol* 79:725-731.
12. **McGeoch DJ, Dolan A, Ralph AC** (2000) Toward a comprehensive phylogeny for mammalian and avian herpesviruses. *J Virol* 74:10401-10406.
13. **Ploegh HL** (1998) Viral strategies of immune evasion. *Science* 280:248-253.
14. **Loch S, Tampe R** (2005) Viral evasion of the MHC class I antigen-processing machinery. *Eur J Physiol* 451:409-417. Review.
15. **Paulsson KM** (2004) Evolutionary and functional perspectives of the major histocompatibility complex class I antigen-processing machinery. *Cell Mol Life Sci* 61:2446-2460.
16. **Yewdell JW, Nicchitta CV** (2006) The DRiP hypothesis decennial: support, controversy, refinement and extension. *Trends Immunol* 27:368-373.
17. **Kloetzel PM** (2001) Antigen processing by the proteasome. *Nat Rev Mol Cell Biol* 2:179-187. Review.
18. **Rock KL, Goldberg AL** (1999) Degradation of cell proteins and the generation of MHC class I-presented peptides. *Annu Rev Immunol* 17:739-779. Review.
19. **Reits EA, Vos JC, Gromme M, Neefjes J** (2000) The major substrates for TAP in vivo are derived from newly synthesized proteins. *Nature* 404:774-778.
20. **Schubert U, Anton LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR** (2000) Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* 404:770-774.
21. **Coux O, Tanaka K, Goldberg AL** (1996) Structure and functions of the 20S and 26S proteasomes. *Annu Rev Biochem* 65:801-847. Review.
22. **Baumeister W, Walz J, Zuhl F, Seemuller E** (1998) The proteasome - paradigm of a self-compartmentalizing protease. *Cell* 92:367-380.
23. **Lehner PJ, Cresswell P** (1996) Processing and delivery of peptides presented by MHC class I molecules. *Curr Opin Immunol* 8:59-67. Review.
24. **Pamer E, Cresswell P** (1998) Mechanisms of MHC class I-restricted antigen processing. *Annu Rev Immunol* 16:323-358. Review.

25. **Reits E, Neijssen J, Herberts C, Benckhuijsen W, Janssen L, Drijfhout JW, Neeffjes J** (2004) A major role for TPP11 in trimming proteasomal degradation products for MHC class I antigen presentation. *Immunity* 20:495-506.
26. **Brooks P, Murray RZ, Mason GG, Hendil KB, Rivett AJ** (2000) Association of immunoproteasomes with the endoplasmic reticulum. *Biochem J* 352:611-615.
27. **Goldberg AL, Cascio P, Saric T, Rock KL** (2002) The importance of the proteasome and subsequent proteolytic steps in the generation of antigenic peptides. *Mol Immunol* 39:147-164. Review.
28. **Cascio P, Hilton C, Kisselev AF, Rock KL, Goldberg AL** (2001) 26S proteasomes and immunoproteasomes produce mainly N-extended versions of an antigenic peptide. *EMBO J* 20:2357-2366.
29. **Princiotta MF, Finzi D, Qian SB, Gibbs J, Schuchmann S, Buttgerit F, Bennink JR, Yewdell JW** (2003) Quantitating protein synthesis, degradation, and endogenous antigen processing. *Immunity* 18:343-354.
30. **Neeffjes JJ, Momburg F, Hammerling GJ** (1993) Selective and ATP-dependent translocation of peptides by the MHC-encoded transporter. *Science* 261:769-771.
31. **York IA, Chang SC, Saric T, Keys JA, Favreau JM, Goldberg AL, Rock KL** (2002) The ER aminopeptidase ERAP1 enhances or limits antigen presentation by trimming epitopes to 8-9 residues. *Nat Immunol* 3:1177-1184.
32. **Serwold T, Gonzalez F, Kim J, Jacob R, Shastri N** (2002) ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* 419:480-483.
33. **Molinari M, Eriksson KK, Calanca V, Galli C, Cresswell P, Michalak M, Helenius A** (2004) Contrasting functions of calreticulin and calnexin in glycoprotein folding and ER quality control. *Mol Cell* 13:125-135.
34. **Dick TP** (2004) Assembly of MHC class I peptide complexes from the perspective of disulfide bond formation. *Cell Mol Life Sci* 61:547-556. Review.
35. **Peaper DR, Wearsch PA, Cresswell P** (2005) Tapasin and ERp57 form a stable disulfide-linked dimer within the MHC class I peptide-loading complex. *EMBO J* 24:3613-3623.
36. **Garbi N, Hammerling G, Tanaka S** (2007) Interaction of ERp57 and tapasin in the generation of MHC class I-peptide complexes. *Curr Opin Immunol* 19:99-105. Review.
37. **Sadasivan B, Lehner PJ, Ortmann B, Spies T, Cresswell P** (1996) Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* 5:103-114.
38. **Grande AG 3rd, Lehner PJ, Cresswell P, Spies T** (1997) Regulation of MHC class I heterodimer stability and interaction with TAP by tapasin. *Immunogenetics* 46:477-483.
39. **Lehner PJ, Trowsdale J** (1998) Antigen presentation: coming out gracefully. *Curr Biol* 8:605-608. Review.
40. **Cresswell P, Bangia N, Dick T, Diedrich G** (1999) The nature of the MHC class I peptide loading complex. *Immunol Rev* 172:21-28.
41. **Cox JH, Yewdell JW, Eisenlohr LC, Johnson PR, Bennink JR** (1990) Antigen presentation requires transport of MHC class I molecules from the endoplasmic reticulum. *Science* 247:715-718.
42. **Spiliotis ET, Osorio M, Zuniga MC, Eddin M** (2000) Selective export of MHC class I molecules from the ER after their dissociation from TAP. *Immunity* 13:841-851.
43. **Ortmann B, Copeman J, Lehner PJ, Sadasivan B, Herberg JA, Grande AG, Riddell SR, Tampe R, Spies T, Trowsdale J, Cresswell P** (1997) A critical role for tapasin in the assembly and function of multimeric MHC class I-TAP complexes. *Science* 277:1306-1309.
44. **Tan P, Kropshofer H, Mandelboim O, Bulbuc N, Hammerling GJ, Momburg F** (2002) Recruitment of MHC class I molecules by tapasin into the transporter associated with antigen processing-associated complex is essential for optimal peptide loading. *J Immunol* 168:1950-1960.
45. **Bangia N, Lehner PJ, Hughes EA, Surman M, Cresswell P** (1999) The N-terminal region of tapasin is required to stabilize the MHC class I loading complex. *Eur J Immunol* 29:1858-1870.
46. **Zarling AL, Luckey CJ, Marto JA, White FM, Brame CJ, Evans AM, Lehner PJ, Cresswell P, Shabanowitz J, Hunt DF, Engelhard VH** (2003) Tapasin is a facilitator, not an editor, of class I MHC peptide binding. *J Immunol* 171:5287-5295.
47. **Bangia N, Cresswell P** (2005) Stoichiometric tapasin interactions in the catalysis of major histocompatibility complex class I molecule assembly. *Immunology* 114:346-353.

48. **Schoenhals GJ, Krishna RM, Granda AG 3rd, Spies T, Peterson PA, Yang Y, Fruh K** (1999) Retention of empty MHC class I molecules by tapasin is essential to reconstitute antigen presentation in invertebrate cells. *EMBO J* 18:743-753.
49. **Granda AG 3rd, Golovina TN, Hamilton SE, Sriram V, Spies T, Brutkiewicz RR, Harty JT, Eisenlohr LC, Van Kaer L** (2000) Impaired assembly yet normal trafficking of MHC class I molecules in Tapasin mutant mice. *Immunity* 13:213-222.
50. **Garbi N, Tan P, Diehl AD, Chambers BJ, Ljunggren HG, Momburg F, Hammerling GJ** (2000) Impaired immune responses and altered peptide repertoire in tapasin-deficient mice. *Nat Immunol* 1:234-238.
51. **Lehner PJ, Surman MJ, Cresswell P** (1998) Soluble tapasin restores MHC class I expression and function in the tapasin-negative cell line .220. *Immunity* 8:221-231.
52. **Purcell AW, Gorman JJ, Garcia-Peydro M, Paradelo A, Burrows SR, Talbo GH, Laham N, Peh CA, Reynolds EC, Lopez De Castro JA, McCluskey J** (2001) Quantitative and qualitative influences of tapasin on the class I peptide repertoire. *J Immunol* 166:1016-1027.
53. **Granda AG, Van Kaer L** (2001) Tapasin: an ER chaperone that controls MHC class I assembly with peptide. *Trends Immunol* 22:194-199.
54. **Higgins CF** (1992) ABC transporters: from microorganisms to man. *Annu Rev Cell Biol* 8:67-113. Review.
55. **Dean M, Rzhetsky A, Allikmets R** (2001) The human ATP-binding cassette (ABC) transporter superfamily. *Genome Res* 11:1156-1166. Review.
56. **Walker JE, Saraste M, Runswick MJ, Gay NJ** (1982) Distantly related sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J* 18:945-951.
57. **Kelly A, Powis SH, Kerr LA, Mockridge I, Elliott T, Bastin J, Uchanska-Ziegler B, Ziegler A, Trowsdale J, Townsend A** (1992) Assembly and function of the two ABC transporter proteins encoded in the human major histocompatibility complex. *Nature* 355:641-644.
58. **Koch J, Tampe R** (2006) The macromolecular peptide-loading complex in MHC class I-dependent antigen presentation. *Cell Mol Life Sci* 63:653-662.
59. **Powis SJ, Townsend AR, Deverson EV, Bastin J, Butcher GW, Howard JC** (1991) Restoration of antigen presentation to the mutant cell line RMA-S by an MHC-linked transporter. *Nature* 354:528-531.
60. **Spies T, DeMars R** (1991) Restored expression of major histocompatibility class I molecules by gene transfer of a putative peptide transporter. *Nature* 351:323-324.
61. **Kleijmeer MJ, Kelly A, Geuze HJ, Slot JW, Townsend A, Trowsdale J** (1992) Location of MHC-encoded transporters in the endoplasmic reticulum and *cis*-Golgi. *Nature* 357:342-344.
62. **Koch J, Guntrum R, Tampe R** (2005) Exploring the minimal functional unit of the transporter associated with antigen processing. *FEBS Lett* 579:4413-4416.
63. **Koch J, Guntrum R, Heintke S, Kyritsis C, Tampe R** (2004) Functional dissection of the transmembrane domains of the transporter associated with antigen processing (TAP). *J Biol Chem* 279:10142-10147.
64. **Raghuraman G, Lapinski PE, Raghavan M** (2002) Tapasin interacts with the membrane-spanning domains of both TAP subunits and enhances the structural stability of TAP1 x TAP2 Complexes. *J Biol Chem* 277:41786-41794.
65. **Garbi N, Tiwari N, Momburg F, Hammerling GJ** (2003) A major role for tapasin as a stabilizer of the TAP peptide transporter and consequences for MHC class I expression. *Eur J Immunol* 33:264-273.
66. **Nijenhuis M, Schmitt S, Armandola EA, Obst R, Brunner J, Hammerling GJ** (1996) Identification of a contact region for peptide on the TAP1 chain of the transporter associated with antigen processing. *J Immunol* 156:2186-2195.
67. **Moody JE, Millen L, Binns D, Hunt JF, Thomas PJ** (2002) Cooperative, ATP-dependent association of the nucleotide binding cassettes during the catalytic cycle of ATP-binding cassette transporters. *J Biol Chem* 277:21111-21114.
68. **van der Does C, Presenti C, Schulze K, Dinkelaker S, Tampe R** (2006) Kinetics of the ATP hydrolysis cycle of the nucleotide-binding domain of Mdl1 studied by a novel site-specific labeling technique. *J Biol Chem* 281:5694-5701.
69. **Zaitseva J, Jenewein S, Oswald C, Jumpertz T, Holland IB, Schmitt L** (2005) A molecular understanding of the catalytic cycle of the nucleotide-binding domain of the ABC transporter HlyB. *Biochem Soc Trans* 33:990-995. Review.

70. **Momburg F, Tan P** (2002) Tapasin—the keystone of the loading complex optimizing peptide binding by MHC class I molecules in the endoplasmic reticulum. *Mol Immunol* 39:217-233. Review.
71. **Owen BA, Pease LR** (1999) TAP association influences the conformation of nascent MHC class I molecules. *J Immunol* 162:4677-4684.
72. **Knittler MR, Alberts P, Deverson EV, Howard JC** (1999) Nucleotide binding by TAP mediates association with peptide and release of assembled MHC class I molecules. *Curr Biol* 9:999-1008.
73. **Reits EA, Griekspoor AC, Neeffjes J** (2000) How does TAP pump peptides? insights from DNA repair and traffic ATPases. *Immunol Today* 21:598-600. Review.
74. **Abele R, Tampe R** (2004) The ABCs of immunology: structure and function of TAP, the transporter associated with antigen processing. *Physiology* 19:216-24.
75. **van Enderp PM, Saveanu L, Hewitt EW, Lehner P** (2002) Powering the peptide pump: TAP crosstalk with energetic nucleotides. *Trends Biochem Sci* 27:454-61.
76. **Vos JC, Reits EA, Wojcik-Jacobs E, Neeffjes J** (2000) Head-head/tail-tail relative orientation of the pore-forming domains of the heterodimeric ABC transporter TAP. *Curr Biol* 10:1-7.
77. **Neumann L, Tampe R** (1999) Kinetic analysis of peptide binding of the TAP transporter complex: evidence for structural rearrangements induced by substrate binding. *J Mol Biol* 294:1203-1213.
78. **Reits EA, Vos JC, Gromme M, Neeffjes J** (2000) The major substrates for TAP in vivo are derived from newly synthesized proteins. *Nature* 404:774-778.
79. **Reits EA, Neeffjes JJ** (2001) From fixed to FRAP: measuring protein mobility and activity in living cells. *Nat Cell Biol* 3:145-147. Review.
80. **Koppers-Lalic D, Reits EA, Rensing ME, Lipińska AD, Abele R, Koch J, Marcondes Rezende M, Admiraal P, van Leeuwen D, Bieńkowska-Szewczyk K, Mettenleiter TC, Rijsewijk FA, Tampe R, Neeffjes J, Wiertz EJ** (2005) Varicelloviruses avoid T cell recognition by UL49.5-mediated inactivation of the transporter associated with antigen processing. *Proc Natl Acad Sci U S A* 102:5144-5149.
81. **Koppers-Lalic D, Rijsewijk FA, Verschuren SB, van Gaans-Van den Brink JA, Neisig A, Rensing ME, Neeffjes J, Wiertz EJ** (2001) The UL41-encoded virion host shutoff (vhs) protein and vhs-independent mechanisms are responsible for down-regulation of MHC class I molecules by bovine herpesvirus 1. *J Gen Virol* 82:2071-2081.
82. **Sato H, Callanan LD, Pesnicak L, Krogmann T, Cohen JI** **Varicella-zoster virus (VZV) ORF17 protein induces RNA cleavage and is critical for replication of VZV at 37 degrees C but not 33 degrees C.** (2002) *J Virol* 76:11012-11023.
83. **Ambagala AP, Gopinath RS, Srikumaran S.** (2003) Inhibition of TAP by pseudorabies virus is independent of its vhs activity. *Virus Res* 96:37-48.
84. **Jones FE, Smibert CA, Smiley JR** (1995) Mutational analysis of the herpes simplex virus virion host shutoff protein: evidence that vhs functions in the absence of other viral proteins. *J Virol* 69:4863-4871.
85. **Read GS, Karr BM, Knight K** (1993) Isolation of a herpes simplex virus type 1 mutant with a deletion in the virion host shutoff gene and identification of multiple forms of the vhs (UL41) polypeptide. *J Virol* 67:7149-7160.
86. **Smiley JR, Elgadi MM, Saffran HA** (2001) Herpes simplex virus vhs protein. *Methods Enzymol* 342:440-451.
87. **Tigges MA, Leng S, Johnson DC, Burke RL** (1996) Human herpes simplex virus (HSV)-specific CD8⁺ CTL clones recognize HSV-2-infected fibroblasts after treatment with IFN-gamma or when virion host shutoff functions are disabled. *J Immunol* 156:3901-3910.
88. **Smiley JR** (2004) Herpes simplex virus virion host shutoff protein: immune evasion mediated by a viral RNase? *J Virol* 78:1063-1068. Review.
89. **Gopinath RS, Ambagala AP, Hinkley S, Srikumaran S** (2002) Effects of virion host shut-off activity of bovine herpesvirus 1 on MHC class I expression. *Viral Immunol* 15:595-608
90. **Hengel H, Brune W, Koszinowski UH** (1998) Immune evasion by cytomegalovirus—survival strategies of a highly adapted opportunist. *Trends Microbiol* 6:190-197.
91. **Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL** (1996) The human cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the endoplasmic reticulum to the cytosol. *Cell* 84:769-779.

92. **Wiertz EJ, Tortorella D, Bogyo M, Yu J, Mothes W, Jones TR, Rapoport TA, Ploegh HL** (1996) Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature* 384:432–438.
93. **Shamu CE, Story CM, Rapoport TA, Ploegh HL** (1999) The pathway of US11-dependent degradation of MHC class I heavy chains involves a ubiquitin-conjugated intermediate. *J Cell Biol* 147:45–58.
94. **Shamu CE, Flierman D, Ploegh HL, Rapoport TA, Chau V** (2001) Polyubiquitination is required for US11-dependent movement of MHC class I heavy chain from endoplasmic reticulum into cytosol. *Mol Biol Cell* 12:2546–2555.
95. **Blom D, Hirsch C, Stern P, Tortorella D, Ploegh HL** (2004) A glycosylated type I membrane protein becomes cytosolic when peptide: N-glycanase is compromised. *EMBO J* 23:650–658.
96. **McCracken AA, Brodsky JL** (2003) Evolving questions and paradigm shifts in endoplasmic-reticulum-associated degradation (ERAD). *Bioessays* 25:868–877. Review.
97. **Lilley BN, Tortorella D, Ploegh HL** (2003) Dislocation of a type I membrane protein requires interactions between membrane-spanning segments within the lipid bilayer. *Mol Biol Cell* 14:3690–3698.
98. **Lilley BN, Ploegh HL** (2004) A membrane protein required for dislocation of misfolded proteins from the ER. *Nature* 429:834–840.
99. **Ye Y, Shibata Y, Yun C, Ron D, Rapoport TA** (2004) A membrane protein complex mediates retro-translocation from the ER lumen into the cytosol. *Nature* 429:841–847.
100. **Loureiro J, Lilley BN, Spooner E, Noriega V, Tortorella D, Ploegh HL** (2006) Signal peptide peptidase is required for dislocation from the endoplasmic reticulum. *Nature* 441:894–897.
101. **Gruhler A, Peterson PA, Fruh K** (2000) Human cytomegalovirus immediate early glycoprotein US3 retains MHC class I molecules by transient association. *Traffic* 1:318–325.
102. **Misaghi S, Sun Z-YJ, Stern P, Gaudet R, Wagner G, Ploegh H** (2004) Structural and functional analysis of human cytomegalovirus US3 protein. *J Virol* 78:413–423.
103. **Park B, Lee S, Kim E, Cho K, Riddell SR, Cho S, Ahn K** (2006) Redox regulation facilitates optimal peptide selection by MHC class I during antigen processing. *Cell* 127:369–382.
104. **Lee S, Yoon J, Park B, Jun Y, Jin M, Sung HC, Kim I-H, Kang S, Choi E-J, Ahn BY, Ahn K** (2000) Structural and functional dissection of human cytomegalovirus US3 in binding major histocompatibility complex Class I molecules. *J Virol* 74:11262–11269.
105. **Park B, Kim Y, Shin J, Lee S, Cho K, Fruh K, Lee S, Ahn K** (2004) Human cytomegalovirus inhibits tapasin-dependent peptide loading and optimization of the MHC class I peptide cargo for immune evasion. *Immunity* 20:71–85.
106. **Furman MH, Dey N, Tortorella D, Ploegh HL** (2002) The human cytomegalovirus US10 gene product delays trafficking of major histocompatibility complex class I molecules. *J Virol* 76:11753–11756
107. **Johnson DC, Hegde NR** (2002) Inhibition of the MHC class II antigen presentation pathway by human cytomegalovirus. *Curr Top Microbiol Immunol* 269:101–115. Review.
108. **Ziegler H, Thale R, Lucin P, Muranyi W, Flohr T, Hengel H, Farrell H, Rawlinson W, Koszinowski UH** (1997) A mouse cytomegalovirus glycoprotein retains MHC class I complexes in the ERGIC/*cis*-Golgi compartments. *Immunity* 6:57–66.
109. **Ziegler H, Muranyi W, Burgert HG, Kremmer E, Koszinowski UH** (2000) The luminal part of the murine cytomegalovirus glycoprotein gp40 catalyzes the retention of MHC class I molecules. *EMBO J* 19:870–881.
110. **Kavanagh DG, Gold MC, Wagner M, Koszinowski UH, Hill AB** (2001) The multiple immune-evasion genes of murine cytomegalovirus are not redundant: m4 and m152 inhibit antigen presentation in a complementary and cooperative fashion. *J Exp Med* 194:967–978.
111. **Kleijnen MF, Huppa JB, Lucin P, Mukherjee S, Farrell H, Campbell AE, Koszinowski UH, Hill AB, Ploegh HL** (1997) A mouse cytomegalovirus glycoprotein, gp34, forms a complex with folded class I MHC molecules in the ER which is not retained but is transported to the cell surface. *EMBO J* 16:685–694.
112. **Reusch U, Muranyi W, Lucin P, Burgert HG, Hengel H, Koszinowski UH** (1999) A cytomegalovirus glycoprotein reroutes MHC class I complexes to lysosomes for degradation. *EMBO J* 18:1081–1091.
113. **Coscoy L, Sanchez DJ, Ganem D** (2001) A novel class of herpesvirus-encoded membrane-bound E3 ubiquitin ligases regulates endocytosis of proteins involved in immune recognition. *J Cell Biol* 155:1265–1273.

114. **Coscoy L, Ganem D** (2000) Kaposi's sarcoma-associated herpesvirus encodes two proteins that block cell surface display of MHC class I chains by enhancing their endocytosis. *Proc Natl Acad Sci USA* 97:8051–8056.
115. **Ishido S, Wang C, Lee BS, Cohen GB, Jung JU** (2000) Downregulation of major histocompatibility complex class I molecules by Kaposi's sarcoma-associated herpesvirus K3 and K5 proteins. *J Virol* 74:5300–5309.
116. **Lehner PJ, Cresswell P** (2004) Recent developments in MHC-class-I-mediated antigen presentation. *Curr Opin Immunol* 16:82–89. Review.
117. **Lehner PJ, Hoer S, Dodd R, Duncan LM** (2005) Downregulation of cell surface receptors by the K3 family of viral and cellular ubiquitin E3 ligases. *Immunol Rev* 207:112–125. Review.
118. **Means RE, Ishido S, Alvarez X, Jung JU** (2002) Multiple endocytic tracking pathways of MHC class I molecules induced by a Herpesvirus protein. *EMBO J* 21:1638–1649.
119. **Hewitt EW, Duncan L, Mufti D, Baker J, Stevenson PG, Lehner PJ** (2002) Ubiquitylation of MHC class I by the K3 viral protein signals internalization and TSG101-dependent degradation. *EMBO J* 21:2418–2429.
120. **Duncan LM, Piper S, Dodd RB, Saville MK, Sanderson CM, Luzio JP, Lehner PJ** (2006) Lysine-63-linked ubiquitination is required for endolysosomal degradation of class I molecules. *EMBO J* 25:1635–1645.
121. **Boname JM, Stevenson PG** (2001) MHC class I ubiquitination by a viral PHD/LAP finger protein. *Immunity* 15:627–636.
122. **Boname JM, May JS, Stevenson PG** (2004) The murine gamma-herpesvirus-68 MK3 protein causes TAP degradation independent of MHC class I heavy chain degradation. *Eur J Immunol* 35:171–179.
123. **Stevenson PG, Efstathiou S, Doherty PC, Lehner PJ** (2000) Inhibition of MHC class I-restricted antigen presentation by gamma 2-herpesviruses. *Proc Natl Acad Sci USA* 97:8455–8460.
124. **Yu YY, Harris MR, Lybarger L, Kimpler LA, Myers NB, Virgin HW 4th, Hansen TH** (2002) Physical association of the K3 protein of gamma-2 herpesvirus 68 with major histocompatibility complex class I molecules with impaired peptide and beta(2)-microglobulin assembly. *J Virol* 76:2796–2803.
125. **Wang X, Ye Y, Lencer W, Hansen TH** (2006) The viral E3 ubiquitin ligase mK3 uses the Derlin/p97 endoplasmic reticulum-associated degradation pathway to mediate down-regulation of major histocompatibility complex class I proteins. *J Biol Chem* 281:8636–8644.
126. **Wang X, Connors R, Harris MR, Hansen TH, Lybarger L** (2005) Requirements for the selective degradation of endoplasmic reticulum-resident major histocompatibility complex class I proteins by the viral immune evasion molecule mK3. *J Virol* 79:4099–4108.
127. **Hudson AW, Howley PM, Ploegh HL** (2001) A human herpesvirus 7 glycoprotein, U21, diverts major histocompatibility complex class I molecules to lysosomes. *J Virol* 75:12347–12358.
128. **Hudson AW, Blom D, Howley PM, Ploegh HL** (2003) The ER luminal domain of the HHV-7 immunoevasin U21 directs class I MHC molecules to lysosomes. *Traffic* 4:824–837.
129. **Bauer D, Tampe R** (2002) Herpes viral proteins blocking the transporter associated with antigen processing TAP—from genes to function and structure. *Curr Top Microbiol Immunol* 269:87–99.
130. **Abele R, Tampe R** (2006) Modulation of the antigen transport machinery TAP by friends and enemies. *FEBS Lett* 580:1156–1163. Review.
131. **Tomazin R, van Schoot NEG, Goldsmith K, Jugovic P, Sempe P, Fruh K, Johnson DC** (1998) Herpes simplex virus type 2 ICP47 inhibits human TAP but not mouse TAP. *J Virol* 72:2560–2563.
132. **Bigger JE, Martin DW** (2004) Identification of an ICP47 homologue in simian agent 8 (SA8). *Virus Genes* 28:223–225.
133. **Neumann L, Kraas W, Uebel S, Jung G, Tampe R** (1997) The active domain of the herpes simplex virus protein ICP47: a potent inhibitor of the transporter associated with antigen processing. *J Mol Biol* 272:484–492.
134. **Galocha B, Hill A, Barnett BC, Dolan A, Raimondi A, Cook RF, Brunner J, McGeoch DJ, Ploegh HL** (1997) The Active Site of ICP47, a Herpes Simplex Virus-encoded Inhibitor of the Major Histocompatibility Complex (MHC)-encoded Peptide Transporter Associated with Antigen Processing (TAP), Maps to the NH2-terminal 35 Residues. *J Exp Med* 185:1565–1572.
135. **Beinert D, Neumann L, Uebel S, Tampe R** (1997) Structure of the viral TAP-inhibitor ICP47 induced by membrane association. *Biochemistry* 36:4694–4700.

136. **Pfänder R, Neumann L, Zweckstetter M, Seger C, Holak TA, Tampe R** (1999) Structure of the active domain of the herpes simplex virus protein ICP47 in water/sodium dodecyl sulfate solution determined by nuclear magnetic resonance spectroscopy. *Biochemistry* 38:13692–13698.
137. **Ahn K, Meyer TH, Uebel S, Sempe P, Djaballah H, Yang Y, Peterson PA, Fruh K, Tampe R** (1996) Molecular mechanism and species specificity of TAP inhibition by herpes simplex virus ICP47. *EMBO J* 15:3247–3255.
138. **Fruh K, Ahn K, Djaballah H, Sempe P, van Endert PM, Tampe R, Peterson PA, Yang Y** (1995) A viral inhibitor of peptide transporters for antigen presentation. *Nature* 375:415–418.
139. **Gorbulev S, Abele R, Tampe R** (2001) Allosteric crosstalk between peptide-binding, transport, and ATP hydrolysis of the ABC transporter TAP. *Proc Natl Acad Sci USA* 98:3732–3737.
140. **Hill A, Jugovic P, York I, Russ G, Bennink J, Yewdell J, Ploegh H, Johnson D** (1995) Herpes simplex virus turns-off the TAP to evade host immunity. *Nature* 375:411–415.
141. **Jugovic P, Hill AM, Tomazin R, Ploegh H, Johnson DC** (1998) Inhibition of major histocompatibility complex class I antigen presentation in pig and primate cells by herpes simplex virus type 1 and 2 ICP47. *J Virol* 72:5076–5084.
142. **Tomazin R, Hill AB, Jugovic P, York I, van Endert P, Ploegh HL, Andrews DW, Johnson DC** (1996) Stable binding of the herpes simplex virus ICP47 protein to the peptide binding site of TAP. *EMBO J* 15:3256–3266.
143. **Kyritsis C, Gorbulev S, Hutschenreiter S, Pawlitschko K, Abele R, Tampe R** (2001) Molecular mechanism and structural aspects of transporter associated with antigen processing inhibition by the cytomegalovirus protein US6. *J Biol Chem* 276:48031–48039.
144. **Ahn K, Gruhler A, Galocha B, Jones TR, Wiertz EJ, Ploegh HL, Peterson PA, Yang Y, Fruh K** (1997) The ER-luminal domain of the HCMV glycoprotein US6 inhibits peptide translocation by TAP. *Immunity* 6:613–621.
145. **Hengel H, Koopmann JO, Flohr T, Muranyi W, Goulmy E, Hammerling GJ, Koszinowski UH, Momburg F** (1997) A viral ER-resident glycoprotein inactivates the MHC-encoded peptide transporter. *Immunity* 6:623–632.
146. **Hewitt EW, Gupta SS, Lehner PJ** (2001) The human cytomegalovirus gene product US6 inhibits ATP binding by TAP. *EMBO J* 20:387–396.
147. **Lehner PJ, Karttunen JT, Wilkinson GW, Cresswell P** (1997) The human cytomegalovirus US6 glycoprotein inhibits transporter associated with antigen processing-dependent peptide translocation. *Proc Natl Acad Sci USA* 94:6904–6909.
148. **Halenius A, Momburg F, Reinhard H, Bauer D, Lobigs M, Hengel H** (2006) Physical and functional interactions of the cytomegalovirus US6 glycoprotein with the transporter associated with antigen processing. *J Biol Chem* 281:5383–5390.
149. **Lybarger L, Wang X, Harris MR, Virgin HWT, Hansen TH** (2003) Virus subversion of the MHC class I peptide-loading complex. *Immunity* 18:121–130.
150. **Wang X, Lybarger L, Connors R, Harris MR, Hansen TH** (2004) Model for the interaction of gammaherpesvirus 68 RING-CH finger protein mK3 with major histocompatibility complex class I and the peptide-loading complex. *J Virol* 78:8673–8686.
151. **Boname JM, de Lima BD, Lehner PJ, Stevenson PG** (2004) Viral degradation of the MHC class I peptide loading complex. *Immunity* 20:305–317.
152. **Stevenson PG, May JS, Smith XG, Marques S, Adler H, Koszinowski UH, Simas JP, Efstathiou S** (2002) K3-mediated evasion of CD8(+) T cells aids amplification of a latent gamma-herpesvirus. *Nat Immunol* 3:733–740.
153. **Hahn AM, Huye LE, Ning S, Webster-Cyriaque J, Pagano JS** (2005) Interferon regulatory factor 7 is negatively regulated by the Epstein-Barr virus immediate-early gene, BZLF-1. *J Virol* 79:10040–10052.
154. **Zeidler R, Eissner G, Meissner P, Uebel S, Tampe R, Lazis S, Hammerschmidt W** (1997) Downregulation of TAP1 in B lymphocytes by cellular and Epstein-Barr virus-encoded interleukin-10. *Blood* 90:2390–2397.
155. **Resing ME, Keating SE, van Leeuwen D, Koppers-Lalic D, Pappworth IY, Wiertz EJ, Rowe M** (2005) Impaired transporter associated with antigen processing-dependent peptide transport during productive EBV infection. *J Immunol* 174:6829–6838.
156. **Levitskaya J, Sharipo A, Leonchiks A, Ciechanover A, Masucci MG** (1997) Inhibition of ubiquitin/proteasome-dependent protein degradation by the Gly-Ala repeat domain of the Epstein-Barr virus nuclear antigen 1. *Proc Natl Acad Sci USA* 94:12616–12621.

157. **Sharipo A, Imreh M, Leonchiks A, Imreh S, Masucci MG** (1998) A minimal glycine-alanine repeat prevents the interaction of ubiquitinated I kappaB alpha with the proteasome: a new mechanism for selective inhibition of proteolysis. *Nat Med* 4:939–944.
158. **Cohen JI** (1998) Infection of cells with varicella-zoster virus down-regulates surface expression of class I major histocompatibility complex antigens. *J Infect Dis* 177:1390-1393.
159. **Abendroth A, Lin I, Slobedman B, Ploegh H, Arvin AM** (2001) Varicella-zoster virus retains major histocompatibility complex class I proteins in the Golgi compartment of infected cells. *J Virol* 75:4878–4888.
160. **Nazerian K, Witter RL** (1975) Properties of a chicken lymphoblastoid cell line from Marek's disease tumor. *J Natl Cancer Inst* 54:453-458.
161. **Omar AR, Schat KA** (1997) Characterization of Marek's disease herpesvirus-specific cytotoxic T lymphocytes in chickens inoculated with a non-oncogenic vaccine strain of MDV. *Immunology* 90:579-585.
162. **Hunt HD, Lupiani B, Miller MM, Gimeno I, Lee LF, Parcels MS** (2001) Marek's disease virus down-regulates surface expression of MHC (B Complex) Class I (BF) glycoproteins during active but not latent infection of chicken cells. *Virology* 282:198-205.
163. **Levy AM, Davidson I, Burgess SC, Dan Heller E** Major histocompatibility complex class I is downregulated in Marek's disease virus infected chicken embryo fibroblasts and corrected by chicken interferon. *Comp Immunol Microbiol Infect Dis* 26:189-198.
164. **Hinkley S, Hill AB, Srikumaran S** (1998) Bovine herpesvirus-1 infection affects the peptide transport activity in bovine cells. *Virus Res* 53:91-96.
165. **Ambagala AP, Hinkley S, Srikumaran S** (2000) An early pseudorabies virus protein down-regulates porcine MHC class I expression by inhibition of transporter associated with antigen processing (TAP). *J Immunol* 164:93–99.
166. **Ambagala AP, Gopinath RS, Srikumaran S** (2004) Peptide transport activity of the transporter associated with antigen processing (TAP) is inhibited by an early protein of equine herpesvirus-1. *J Gen Virol* 85: 349–353.
167. **van Hall T, Wolpert EZ, van Veelen P, Laban S, van der Veer M, Roseboom M, Bres S, Grufman P, de Ru A, Meiring H, de Jong A, Franken K, Teixeira A, Valentijn R, Drijfhout JW, Koning F, Camps M, Ossendorp F, Karre K, Ljunggren HG, Melief CJ, Offringa R** (2006) Selective cytotoxic T-lymphocyte targeting of tumor immune escape variants. *Nat Med* 12:417-424.
168. **Lipińska AD, Koppers-Lalic D, Rychłowski M, Admiraal P, Rijsewijk FA, Bienkowska-Szewczyk K, Wiertz E** (2006) Bovine Herpesvirus 1 UL49.5 Protein Inhibits the Transporter Associated with Antigen Processing despite Complex Formation with Glycoprotein M. *J Virol* 80:5822-5832.
169. **Wang X, Herr RA, Chua W-J, Lybarger L, Wiertz EJHJ, Hansen TH** (2007) Ubiquitination of serine or lysine residues on the cytoplasmic tail can induce ERAD of MHC-I by viral E3 ligase mK3. *J Cell Biol* doi:10.1083/jcb.200611063
170. **Eisfeld AJ, Yee MB, Erazo A, Abendroth A, Kinchington PR** (2007) Downregulation of class I Major Histocompatibility Complex surface expression by varicella-zoster virus involves Open Reading Frame 66 protein kinase dependent and independent mechanisms doi:10.1128/jvi.00711-07

