



Universiteit
Leiden
The Netherlands

GEP-NET : rare tumour connections. Pathophysiological aspects of gastroenteropancreatic neuroendocrine tumours

Kuiper, P.

Citation

Kuiper, P. (2011, April 21). *GEP-NET : rare tumour connections. Pathophysiological aspects of gastroenteropancreatic neuroendocrine tumours*. Retrieved from <https://hdl.handle.net/1887/17584>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/17584>

Note: To cite this publication please use the final published version (if applicable).

GEP-NET: rare tumour connections
Pathophysiological aspects of gastroenteropancreatic neuroendocrine tumours

Patricia Kuiper

GEP-NET: rare tumour connections
Pathophysiological aspects of gastroenteropancreatic neuroendocrine tumours

Patricia Kuiper

Thesis, University of Leiden
ISBN: 9789090260433

Cover design by Wouter Kanselaar (photograph) and Patricia Kuiper (art work).
Photograph of Seven Mile Bridge, Florida Keys, United States (2008).

Printed by Ridderprint, Ridderkerk

© 2011 P. Kuiper, Leiden, the Netherlands

All rights reserved. No part of this thesis may be reproduced or transmitted in any form, by any means, electronic or mechanical, without prior written permission of the author.

Printing of this thesis was financially supported by the Section Experimental Gastroenterology (SEG) of the Netherlands society of Gastroenterology (NvGE), the Netherlands society of Gastroenterology (NvGE), AstraZeneca B.V., Dr. Falk Pharma Benelux B.V., Ipsen Farmaceutica B.V., Novartis Oncology, and Stichting voor Patienten met Kanker aan het Spijsverteringskanaal (SPKS).

GEP-NET: rare tumour connections
*Pathophysiological aspects of gastroenteropancreatic
neuroendocrine tumours*

PROEFSCHRIFT

ter verkrijging van de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus Prof. mr. P.F. van der Heijden,
volgens besluit van het College voor Promoties
te verdedigen op donderdag 21 april 2011

klokke 15.00 uur

door

Patricia Kuiper

geboren te 's Gravenhage

in 1986

Promotiecommissie

Promotor:

Prof. dr. C.B.H.W. Lamers

Co-Promotores:

Dr. ir. H.W. Verspaget

Dr. ir. I. Biemond

Overige leden:

Prof. dr. A.A.M. Masclee, Maastricht Universitair Medisch Centrum

Prof. dr. J.B. Jansen, Universitair Medisch Centrum St Radboud Nijmegen,
tevens Elkerliek Ziekenhuis Helmond

Dr. B. G. Taal, Nederlands Kanker Instituut - Antoni van Leeuwenhoek

Ziekenhuis Amsterdam

Prof. dr. J.H. Bolk

Prof. dr. H.F.A. Vasen

Contents

| | |
|--|-----|
| Abbreviations | 7 |
| Chapter 1 General Introduction and Outline of the Thesis | 9 |
| Chapter 2 An overview of the current diagnosis and recent developments in neuroendocrine tumours of the gastroenteropancreatic tract: the diagnostic approach <i>Neth J Med</i> 2011; 69: 14-20 | 31 |
| Chapter 3 Pathological incidence of duodenopancreatic neuroendocrine tumours in the Netherlands: a Pathologisch Anatomisch Landelijk Geautomatiseerd Archief study <i>Pancreas</i> 2010; 39: 1134-1139 | 49 |
| Chapter 4 A case of recurrent gastrinoma in the liver with a review of "primary" hepatic gastrinomas <i>BMJ Case Reports</i> [published online 11 June 2009] | 67 |
| Chapter 5 Diagnostic efficacy of the secretin stimulation test for the Zollinger-Ellison syndrome: an intra-individual comparison using different dosages in patients and controls <i>Pancreatology</i> 2010; 10: 14-18 | 81 |
| Chapter 6 Expression and ligand binding of bombesin receptors in pulmonary and intestinal carcinoids: The role of bombesin in carcinoids <i>J Endocrinol Invest</i> 2010 [Epub ahead of print] | 97 |
| Chapter 7 Angiogenic markers endoglin and vascular endothelial growth factor in gastroenteropancreatic neuroendocrine tumours <i>World J Gastroenterol</i> 2011; 17: 219-225 | 115 |
| Chapter 8 The IGF-matrilysin network in gastroenteropancreatic neuroendocrine tumours <i>Submitted</i> | 133 |

| | |
|-------------------------------------|-----|
| Chapter 9 | 149 |
| Summarizing discussion | |
| Chapter 10 | 159 |
| Nederlandse samenvatting voor leken | |
| List of publications | 169 |
| Curriculum Vitae | 171 |
| Nawoord | 173 |
| Full-colour illustrations | 175 |

Abbreviations

| | |
|-------------|--|
| 5-HIAA | 5-hydroxyindoleacetic acid |
| ACTH | Adrenocorticotrophic hormone |
| APUDoma | Amine precursor uptake decarboxylase tumour |
| BB1R, BRS-1 | Bombesin-receptor type 1, Neuromedin B receptor |
| BB2R, BRS-2 | Bombesin-receptor type 2, Gastrin-releasing peptide receptor |
| BB3R, BRS-3 | Bombesin receptor subtype 3 |
| BB4R, BRS-4 | Bombesin receptor subtype 4 |
| BBS | Bombesin |
| BLP | Bombesin-like peptide |
| BSA | Bovine serum albumin |
| CBS | Central Bureau for Statistics |
| CCK | Cholecystokinin |
| CD105 | Endoglin |
| CgA | Chromogranin A |
| CNS | Central nerve system |
| CRC | Colorectal cancer |
| CT | Computed tomography |
| DAB | 3,3'-diaminobenzidine |
| DNET | Duodenal neuroendocrine tumour |
| EC | Enterochromaffin |
| ECL | Enterochromaffin-like |
| ECM | Extracellular matrix |
| ELISA | Enzyme linked immuno-sorbent assay |
| EUS | Endoscopic ultrasonography |
| Flt-1 | VEGF receptor 1 |
| F-PNET | Functioning/functional pancreatic neuroendocrine tumour |
| FSG | Fasting serum gastrin |
| GEP-NET | Gastroenteropancreatic neuroendocrine tumour |
| GI | Gastrin increase |
| GI-NET | Gastrointestinal neuroendocrine tumour |
| GIST | Gastrointestinal stromal tumour |
| GRF | Growth-hormone releasing factor |
| GRP | Gastrin releasing peptide |
| GRPR | Gastrin releasing peptide receptor |
| HRP | Horseradish peroxidase |
| IGF | Insulin-like growth factor |
| kDa | Kilo Dalton |
| KDR | VEGF receptor 2 |
| MEN-1 | Multiple endocrine neoplasia syndrome type 1 |
| MMP | Matrix metalloproteinases |
| MMP-7 | Matrilysin, matrix metalloproteinase-7 |
| MRI | Magnetic resonance imaging |
| mRNA | Messenger ribonucleic acid |
| MVD | Microvessel density |

| | |
|--------------|---|
| NEC | Neuroendocrine carcinoma |
| NET | Neuroendocrine tumour |
| NF-1 | Neurofibromatosis type 1 |
| NF-PNET | Non-functioning/non-functional pancreatic neuroendocrine tumour |
| NMB | Neuromedin B |
| NMBR | Neuromedin B receptor |
| NME | Necrolytic migratory erythema |
| NSE | Neuron specific enolase |
| PALGA | Nationwide network and registry for histo- and cytopathology in the Netherlands |
| PBS | Phosphate buffered saline |
| PET | Pancreatic endocrine tumour |
| PNET | Pancreatic neuroendocrine tumour |
| PP | Pancreas polypeptide |
| PPI | Proton pump inhibitor |
| ROC | Receiver operating characteristic |
| RT | Room temperature |
| s.e. | Standard error |
| SEER | Surveillance, Epidemiology and End Results database |
| SEM | Standard error of the mean |
| sEndoglin | Soluble endoglin |
| SRS | Somatostatin receptor scintigraphy |
| SSPS | Statistical Package for Social Sciences |
| St. dev. | Standard deviation |
| TBS | Tris-buffered saline |
| TGF- β | Transforming growth factor beta |
| TNM | Tumour node metastasis |
| VEGF | Vascular endothelial growth factor |
| vHLD | Von Hippel-Lindau disease |
| VIP | Vasoactive intestinal peptide |
| WDHA | Watery diarrhea hypokalemia achlorhydria |
| WHO | World Health Organization |
| ZES | Zollinger-Ellison syndrome |