



Universiteit
Leiden
The Netherlands

Peripheral nerve graft architecture affects regeneration

Vleggeert-Lankamp, Carmen Lia Anne-Marie

Citation

Vleggeert-Lankamp, C. L. A. -M. (2006, December 14). *Peripheral nerve graft architecture affects regeneration*. Retrieved from <https://hdl.handle.net/1887/5566>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/5566>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 9

Adhesion and proliferation of human Schwann cells on adhesive coatings

C.L.A.M. Vleggeert-Lankamp^a, A.P. Pêgô^b, E.A.J.F. Lakke^a, M. Deenen^a, E. Marani^a,
R.T.W.M. Thomeer^a

^aNeuroregulation group, Department of Neurosurgery, ^cDepartment of Neurophysiology,
Leiden University Medical Centre (LUMC), P.O. Box 9600, NL-2300 RC Leiden, the Netherlands

^bInstitute for Biomedical Technology (BMTI), Department of Polymer Chemistry and
Biomaterials, University of Twente, P.O. Box 217, NL-7500 AE Enschede, The Netherlands

Biomaterials, 2004; 25: 2741-2751

*If it's true that no gentleman is a hero to his valet, and no writer is a genius to his typist, then
certainly no promovendus is a genius to her co-promotor.*

Abstract

Attachment to and proliferation on the substrate are deemed important considerations when Schwann cells are to be seeded in synthetic nerve grafts. Attachment is a prerequisite for the Schwann cells to survive and fast proliferation will yield large numbers of Schwann cells in a short time, which appears promising for stimulation of peripheral nerve regeneration. The aim of the present study was to compare the adhesion and proliferation of human Schwann cells on different substrates. The following were selected for their suitability as an internal coating of synthetic nerve grafts; the extracellular matrix proteins fibronectin, laminin and collagen type I and the poly-electrolytes poly(D-lysine) and poly(ethylene-imine). On all coatings attachment of human Schwann cells was satisfactory and comparable, indicating that this factor is not a major consideration in choosing a suitable coating.

9
200

Proliferation was best on fibronectin, laminin and poly(D-lysine), and worst on collagen type I and poly(ethylene-imine). Since nerve regeneration is enhanced by laminin and/or fibronectin, these are preferred as coatings for synthetic nerve grafts seeded with Schwann cells.

Introduction

Schwann cells (SCs) play an important role in mediating peripheral nerve regeneration [1, 2]. After nerve transection Schwann cells in the distal nerve stump align to form the so called 'bands of Bungner', which conduct the outgrowing regenerating axons to the target organ [3]. In the presence of axons, SCs assemble a complete extracellular matrix (ECM) [4, 5], and express cellular adhesion molecules (CAMs; N-CAM, MAG, L1/Ng-CAM and N-cadherin) on their surface. These CAMs interact with the CAMs, integrins, and integrin-related ECM receptors on the neuronal growth cones [2].

Introduction of cultured Schwann cells into the lumen of a synthetic nerve graft enhances peripheral nerve regeneration [6-11]. For SCs to survive in the graft, attachment is mandatory, since attachment is a prerequisite for survival and proliferation of SCs [12, 13]. In the absence of axons, it seems necessary to supply the SCs with additives, i.e. an adhesive coating, to allow them to assemble basal-lamina like structures, and thus to adhere to the grafts' interior. Preferably such an adhesive coating should also stimulate proliferation, because the few studies that evaluated the influence of the number of SCs in artificial nerve grafts on nerve regeneration, advocate the addition of a large number of cells [6, 9, 11]. The success of predegenerated autologous nerve grafts seems to be at least partially based on Schwann cell proliferation subsequent to Wallerian degeneration [14, 15].

Fibronectin, laminin and collagen type I [16-18], interact with the integrins on the SC surface [13, 19], and support Schwann cell attachment and proliferation. The positively charged polyelectrolytes poly(D-lysine) (PDL) [20, 21] and poly(ethylene-imine) (PEI) [22] allow attachment of negatively charged Schwann cells.

Although the adhesive and proliferation stimulating properties of these coatings are important in deciding what coating to apply to the synthetic nerve guide, no previous study, to the best of our knowledge, systemically quantified and compared attachment and proliferation on these coatings before. In the present study we compared both attachment and proliferation of human Schwann cells (HSCs) on fibronectin, laminin, collagen, PDL, and PEI amongst each other and against customary coatings in Schwann cell culturing, like gelatin, gelatin crosslinked with glutaraldehyde [23] and uncoated glass.

Materials and methods

Isolation and culture of human Schwann cells

Small pieces of human sural nerve that remained after nerve transplantation were used. All material was obtained from the Department of Neurosurgery at the Leiden University Medical Center. Patients agreed by informed consent to the use of this material.

After stripping the epineurium and connective tissue, the sural nerve was cut into pieces of ca. 1 mm³. These were placed in gelatin-coated (see below) culture flasks (Greiner, The Neth

erlands) and covered with a thin layer of LAK culture medium [24], consisting of Dulbecco's modified Eagle's medium (DMEM; Bio-Whittaker Europe, Belgium), 10% lymphokine activated killer cells conditioned medium (LAK) [25], 5% fetal calf serum (FCS; Gibco BRL, Life Technologies, Germany), 0.25 µl/ml phytohaem-agglutinin (PHA; Difco Laboratories, USA), 100 IU/ml penicillin (Gist-Brocades, The Netherlands), and 50 µg/ml streptomycin (Gist-Brocades).

To obtain highly purified human Schwann cell cultures, a sequential explantation technique was used [26]. Fibroblasts migrate faster out of the nerve pieces and after three to five explantation cycles, only Schwann cells emerge. Cultures were regularly sampled, and the presence of fibroblasts in these samples was assessed with immunostainings. If more than 2% of the sampled cells (rough estimate) were fibroblasts, 10 ml of $1 \cdot 10^{-3}$ M arabinoside-C was added to the culture medium [26]. After two days, arabinoside-C was removed and cells were thoroughly rinsed with fresh LAK culture medium. In the subsequent weeks, Schwann cells were allowed to recover.

Immunocytochemistry of human Schwann cells

Standard immunocytochemical stainings were used to determine the identity of the cultured cells. Cell cultures were made on crosslinked gelatin coated coverslips (see below), fixed with Cryofix® (Merck, Darmstadt, Germany), rinsed three times with phosphate buffered saline (PBS; 0.1 M, pH 7.2) and incubated with antibodies appropriately diluted in PBS containing 0.1% bovine serum albumin (BSA; Sigma, St. Louis, Missouri) and 1% normal goat serum (NGS; CLB, Amsterdam, the Netherlands). To identify Schwann cells, antibodies directed against S100 (1:1000; Sigma) and GFAP (1:500; Boehringer Mannheim Biochemica, Germany) were added. For fibroblast identification, antibodies directed against fibroblasts (1:100; S5 clone, Sigma) and Thy 1.1 (1:10,000; Serotec, Oxford, United Kingdom) were used. After overnight incubation at 4°C in a moist chamber, cultures were again rinsed 3 times with PBS and subsequently stained with GAM/FITC (Molecular Probes, The Netherlands) (anti-S100, anti-fibroblast, and anti-Thy 1.1) and GAR/FITC (GFAP) appropriately diluted in PBS containing 0.1% BSA and 1% NGS. After extensive rinsing with PBS, sections were mounted, dehydrated, and coverslipped with Fluoromount® (Merck) and viewed with a fluorescence microscope.

Schwann cell identity was further confirmed with a Reverse Transcriptase Polymeric Chain Reaction (RT-PCR) protocol [27]. In short, cDNA was generated from total RNA isolated from trypsinized Schwann cell cultures. As a control, cDNA was also generated from total RNA isolated from fibroblasts. PCR reactions were allowed to take place between the generated cDNA and primers for 2',3'-cyclic nucleotide-3'-phosphatase (CNPase), S100β and GFAP. After amplification, the PCR products were applied to an agarose gel and visualized under UV.

As a final control, human Schwann cells were collected from the gelatin-coated culture flasks by incubation with a solution of 0.25% trypsin (Difco Laboratories, USA) and 10 mM ethylenediamine tetra-acetic acid (EDTA) in PBS [24] for approximately 2 minutes at room temperature, followed by the addition of an equal amount of culture medium. The cell number was estimated using a Bürker-Türk chamber. The cell suspension was centrifuged for 5 minutes at

1,600 rpm and subsequently the cells were incubated in 2 ml 40 μ M Dil in culture medium for 30 minutes. After a second identical centrifugation step the cells were resuspended in fresh culture medium to a final concentration of ca. $2 \cdot 10^3$ cells/ml and seeded on cultures of DRG neurons [28]. After 7 days, the ability of the HSCs cells to align along neurites was evaluated with a fluorescence microscope.

Coatings

Round glass coverslips with a diameter of 15 mm (176.7 mm^2) were washed in a 10% g/v potassiumdichromate in 10% H_2SO_4 solution, rinsed and sterilized. All glass coverslips were first placed in a 24-well plate and subsequently coated. For each timepoint 3 glass coverslips were prepared with each coating cf. [18]. Glass coverslips coated with gelatin and gelatin crosslinked with glutaraldehyde, and uncoated glass coverslips were used as controls.

9
203

Fibronectin and Laminin:

A 50 μ g/ml solution of either fibronectin or laminin (Boehringer Mannheim, Almere, The Netherlands) in PBS (200 μ l) was applied to glass coverslips. After 45 minutes of incubation (37°C and humidified air/5% CO_2), the surplus of fluid was removed. In order to prevent the surface from drying out, HSCs were seeded on the surface immediately.

Collagen type I:

Collagen type I from bovine dermis (Vitrogen 100, Collagen Corporation, Fremont, CA) was applied (500 μ l) to glass coverslips. After overnight incubation at 37°C and humidified air/5% CO_2 , the surface was rinsed thoroughly with PBS. In order to prevent the surface from drying out, HSCs were seeded on the surface immediately.

PDL and PEI:

A 0.1% solution (w/v) of either poly-D-lysine or poly-(ethylene-imine) (Sigma) was applied to glass coverslips. After 2 hours the coverslips were rinsed thoroughly with PBS. The surfaces were dried overnight in a flow chamber.

Gelatin:

A 0.5% solution (w/v) of gelatin (400 μ l; Difco Laboratories, USA) in PBS was applied to glass coverslips. After 45 minutes, the surplus of gelatin was removed and the surface was rinsed with PBS. The surfaces were dried overnight in a flow chamber.

Crosslinked gelatin:

A 0.5% solution (w/v) of gelatin (400 μ l) in PBS was applied to glass coverslips. After 45 minutes the surplus of gelatin was removed and the surface was treated for 15 minutes with a 0.5% solution (w/v) of glutaraldehyde (400 μ l). This was done to prevent the gelatin from dissolving

in the culture medium [23]. Then the surface was thoroughly rinsed with PBS. The surfaces were dried overnight in a flow chamber.

Human Schwann cell seeding

HSCs were collected from the gelatin-coated culture flasks, by incubation for approximately 2 minutes at room temperature with a solution of 0.25% trypsin and 10 mM EDTA in PBS, followed by the addition of an equal amount of culture medium. A sample of the suspension was stained with True Blue (Janssen Chemica, Belgium) to evaluate cell viability. The cell suspension was centrifuged for 5 minutes at 1,600 rpm, and subsequently the cell pellet was washed with LAK culture medium. The cell number was estimated using a Bürker-Türk chamber. After a second identical centrifugation step the cells were resuspended in fresh culture medium to a final concentration of $2 \cdot 10^3$ cells/ml for the attachment study and to a final concentration of $3.8 \cdot 10^3$ cells/ml for the proliferation study. The concentration of HSCs was lower in the attachment study to reduce attachment of HSCs to each other. Aliquots of the final cell suspension (500 μ l) were added to the (coated) glass slides in the wells and incubated at 37°C in humidified air/5% CO₂. Medium was refreshed three times a week.

9
204

Human Schwann cell adhesion and proliferation

For evaluation at the indicated times, the (coated) glass coverslips in the 24-well plate were gently rinsed with PBS to remove the non-attached cells. Cryofix® was added for fixation and gently rinsed away after 20 minutes. Coverslips were taken out of the 24-well plate and stained with 0.25 wt/vol% Coomassie blue solution in methanol:water:acetic acid (5:5:1). Subsequently (coated) glass coverslips with adhering stained HSCs were coverslipped with Aquamount® (Merck) and examined under a light microscope (Olympus) at 100 or 200x magnification. The morphology of the cells was described and pictures of representative cells on all slides were made. All cells present on the coatings and glass were counted and the logarithms of the number of cells were compared. When a confluent cell layer was present, the total number of cells was calculated from representative samples.

Attachment of HSCs was evaluated 1, 3, 6 and 24 hours after seeding [29]. The attachment ratio was calculated as the lognumber of cells counted after 24 hours, divided by the lognumber of cells seeded.

Proliferation of HSCs was evaluated 3, 6, 9, 12 and 15 days after seeding. The proliferation rate was calculated as the slope of the line resulting from applying a linear regression analysis to the logarithms of the cell numbers on each coating or glass at the studied timepoints. First, this was done for the overall proliferation (time interval day 3 to 12), and subsequently for the time intervals day 0 to 3, day 3 to 6, day 6 to 9, and day 9 to 12, separately.

Statistical data analysis

The lognumber of cells in the attachment and proliferation studies and the attachment ratios are presented as mean \pm standard deviation (SD) and were analyzed using a one-way ANOVA,

followed by a Tukey's least significant differences multiple comparisons test if there was a difference between the groups beyond a significance level of $p=0.05$. Proliferation rates were presented as mean \pm standard deviation (SD) and compared using a univariate general linear model with a difference contrast for time. The SPSS statistical program, version 11.0, was used to perform all calculations. P-values of less than 0.05 were regarded as significant.

Results


Purification and characterization of Schwann cell cultures

Purification by the sequential explantation method resulted in cultures that stained almost exclusively with S100 and GFAP antibodies, and only sporadically with fibroblast and Thy1.1 antibodies, indicating that highly purified HSC cultures were obtained. RT-PCR demonstrated that HSC cultures expressed the Schwann cell markers CNPase, S-100 β , and GFAP. Control fibroblast cultures were negative in RT-PCR for all Schwann cell markers. Virtually all cells in the samples stained with True Blue turned blue, indicating high HSC viability. Dil labelled HSCs, cultured on rat DRG neurons, were observed to align along the neurites, thus expressing a specific Schwann cell property (fig. 1).

Human Schwann cell adhesion on adhesive coatings


One hour after seeding, most HSCs adhering to the (coated) coverslips demonstrated small lamellipodial extensions and resembled 'fried eggs' (fig. 2). After 3 hours, most cells stretched out, although there was also a considerable number of cells that appeared partially folded, indicating that the cells were not yet fully attached. Except on PEI, cells were stretched out or transformed to a spindle shape and formed clusters after 6 hours. On PEI, after 24 hours a lot

Figure 1. Human Schwann cells stained with Dil seeded on DRG neurites.



The left figure is a phase contrast micrograph of HSCs aligning along DRG neurites. The neurites are easily recognizable. The fluorescence micrograph (right figure) clearly demonstrates the alignment of the fluorescent HSCs along the DRG neurites.

Figure 2. Attachment of human Schwann cells on coatings - qualitative



Morphological appearances of human Schwann cells on fibronectin, laminin, collagen type I, PDL, PEI, gelatine, crosslinked gelatine, and glass, 1, 3, 6 and 24 hours after seeding. Historically, Schwann cells were described to be spindle shaped [52, 53]. However, in later years it was demonstrated that, although many of cultured Schwann cells have this characteristic spindle-shaped morphology, some have a more flattened (fibroblast-like) morphology [32, 54-58].


of cells were still not fully stretched. Mitotic figures were absent throughout the cell cultures on all coatings at 24 hours, indicating that multiplication of cells did not influence attachment ratios.

Initially, the lognumber of adhering cells differed among the coatings, but these differences disappeared progressively during the first 6 hours of culturing (fig. 3A). One hour after seeding the lognumbers of cells adhering to collagen, PDL, gelatin and glass were lower compared to crosslinked gelatin, and the lognumbers of cells on collagen and glass were also lower compared to fibronectin and PEI. After 3 hours the lognumber of cells on collagen was lower compared to gelatin and crosslinked gelatin, and the lognumber of adhering cells to glass was lower compared to all other coatings, except collagen. Six and twenty-four hours after seeding, no significant differences could be detected in the lognumbers of adhering cells to the coatings and glass. Moreover, the attachment ratio was equal ($p=0.09$; fig. 3B).

Human Schwann cell proliferation on adhesive coatings

HSCs proliferated on all coatings and on uncoated glass, between 3 and 15 days of culturing (fig. 4). Except on PEI HSCs appeared stretched out and demonstrated a typical bi- or tripolar


Figure 3. Attachment of human Schwann cells on coatings – quantitative



A: The lognumber of HSCs adhering to different coatings and glass at 1, 3, 6 and 24 hours after seeding. The number of cells seeded was 1000 cells/well (log 1000 = 3). Data are represented as mean \pm SD. The seven coatings and glass were compared among each other: * lower compared to crosslinked gelatin at the corresponding time point; # lower compared to fibronectin and PEI at the corresponding time point; § lower compared to gelatin and crosslinked gelatin at the corresponding time point; † lower compared to all surfaces, except collagen at the corresponding time point.

B: Attachment ratios calculated as the lognumber of cells present after 24 hours divided by the log number of cells seeded. Data are represented as mean \pm SD.

Figure 4. Proliferation of human Schwann cells on coatings – qualitative



Morphological appearances of human Schwann cells on coatings, 3, 6, 9, 12 and 15 days after seeding. Schwann cells can appear either as spindle-shaped or as flat cells.

morphology with oval nuclei [30]. After six days of culturing, cell layers reached confluency. On PEI it took the cells more than six days to fully recover from the seeding procedure; it was not until the ninth day of culturing that HSCs on PEI were fully stretched out and reached confluency. The increase in number of cells on the coatings and on glass was accompanied

Table 1. Proliferation rate of human Schwann cells on coatings – timeinterval 3 – 12 days

	Correlation coefficient of linear fit (R)	Proliferation rate (3-12 days)
Fibronectin	0.820	0.10 ± 0.02
Laminin	0.950	0.09 ± 0.01
Collagen	0.799	0.13 ± 0.03
PDL	0.900	0.09 ± 0.02
PEI	0.798	0.09 ± 0.02
Crossl. gelatin	0.901	0.07 ± 0.01
Gelatin	0.859	0.14 ± 0.03
Glass	0.853	0.12 ± 0.02

The proliferation rate of the cells on the coatings for the whole culturing period (day 3 to 12) was calculated as the slope of the line resulting from applying a linear regression analysis to the lognumbers of cells on each surface on all timepoints.

by a decrease in cell size. This is clearly illustrated by comparing the cells on day 12 and 15 (enlargement 200x) to the cells on day 3, 6 and 9 (enlargement 100x). On day 15, the number of cells increased to such an extent that the cell layers started to detach from the coated surfaces and from the glass. Mitotic figures were present throughout the cell cultures on all coatings.

The proliferation rates on all coatings and glass in the time interval 3 to 12 days were in the same range, but the accompanying correlation coefficients were relatively low (table 1). For this reason we wanted to gain insight in the trends of proliferation within the 0-12 days period. Figures 5A and B demonstrate that the proliferation rate during the first three days after seeding was highest on laminin and crosslinked gelatin, and also high on fibronectin and PDL. As a result, the lognumber of cells at day 3 on collagen, PEI, gelatin and glass was lower compared to laminin and crosslinked gelatin. Moreover, the logarithm of cell numbers on gelatin and collagen was also lower compared to PDL.

From three to six days of culturing, proliferation rates changed considerably. On PEI, the proliferation rate increased more than 20 times, on fibronectin, collagen, gelatin, and glass, the proliferation rate increased 6 to 10 times. The proliferation rate on laminin increased only a little, in contrast to a threefold increase in the cell number during the first three days. On crosslinked gelatin the proliferation rate even decreased. Consequently, the lognumber of cells was approximately the same at six days after seeding, except on crosslinked gelatin and PEI, which displayed a lower lognumber of cells.


From six to nine days, the lognumber of cells on the coatings and glass remained constant, with the exception of PEI, that demonstrated a considerable proliferation rate, though also lower in comparison with the previous interval (fig. 5B). After nine days, no significant differences could be detected in the lognumber of cells on the adhesive coatings or on uncoated glass (fig. 5A).

Between nine and twelve days, proliferation rates again increased on all coatings and glass, except on collagen and PEI. On collagen, the number of cells remained approximately the same, and on PEI the number of cells even decreased. Consequently, after 12 days the log

number of cells on collagen became smaller compared to laminin and the lognumber of cells on PEI became smaller compared to all other coatings, except collagen (fig. 5A).

Figure 5. Proliferation of human Schwann cells on coatings – quantitative


A



9

210

B



The lognumber of HSCs adhering to different coatings and glass at 3, 6, 9 and 12 days after seeding. The initial amount of cells was 1900 cells/well ($\log 1900 = 3.3$). Data are represented a mean \pm SD. The seven coatings and glass were compared among each other: * lower compared to laminin and crosslinked gelatin at the corresponding time point; ‡ lower compared to PDL at the corresponding time point; § lower compared to collagen and glass at the corresponding time point; # lower compared to fibronectin, laminin, PDL, and gelatin at the corresponding time point; & lower compared to laminin at the corresponding time point; † lower compared to all surfaces, but collagen at the corresponding time point.

B: Proliferation rates calculated as the slope of the linear fit calculated from the logarithm of the number of cells over the time intervals: 0 to 3 days, 3 to 6 days, 6 to 9 days and 9 to 12 days. For all coatings, the comparison of proliferation rates differed significantly when two time intervals were compared, except for collagen when the proliferation rates of the time interval 6 to 9 days was compared with the time interval 9 to 12 days (indicated with an *).

After fifteen days of culturing, the number of cells on all coatings and on glass increased to such an extent that the cell layer detached from the underlying surface. It was therefore not possible to count and to compare the cells quantitatively. Qualitatively however, the space between the cells was larger on collagen and PEI (fig. 4).

Discussion

Using a relatively simple explantation technique, we were able to obtain confluent human Schwann cell cultures from which fibroblasts were virtually absent. Purity was assessed by positive staining of S100- and GFAP-antibodies, and negative staining with fibroblast- and Thy 1.1-antibodies. The positive staining results were confirmed with RT-PCR Schwann cell identification. We demonstrated that the cultured cells aligned along neurites, a highly specific Schwann cell property. Therefore, we qualified this explantation technique as adequate to obtain sufficient numbers of human Schwann cells for subsequent seeding on the coatings and glass.

Attachment of human Schwann cells

The attachment study demonstrated that within six hours after seeding, human Schwann cells attached as well to ECM proteins, poly-electrolytes, gelatin and crosslinked gelatin as to glass. On PEI however, complete stretching of HSCs took more than 24 hours.

It was demonstrated in several studies that Schwann cells could satisfactorily attach to ECM proteins and poly-electrolytes [16, 18, 20, 21, 31]. However, attachment to these substrata was never studied comparatively. The present results demonstrate that the number of cells adhering to these coatings is comparable. Moreover, no differences could be detected in Schwann cell morphology, with the exception of HSCs on PEI which still demonstrated partial folding after 24 hours in culture. Fibronectin was previously demonstrated to encourage cell spreading [32], and laminin to stimulate SCs to elongate [17], but we did not observe more spreading or elongation of the HSCs on fibronectin or laminin compared to other coatings (fig. 2).

In a previous paper we described Schwann cell attachment to different biomaterials, and likewise demonstrated that within six hours after seeding, attachment was the same [33]. Apparently Schwann cells easily attach to almost any culture substratum. Thus, contrary to the prevailing opinion, attachment is not an important consideration in choosing a suitable material for coating the interior of a synthetic nerve guide.

Proliferation of human Schwann cells

Proliferation rates of human Schwann cells varied largely during the first six days of culture. The first increase in proliferation (3 – 6 days of culture) was most profound on fibronectin, laminin, PDL and crosslinked gelatin, and less on collagen type I, PEI, gelatin and glass. Proliferation

diminished whenever a density of ca. 10 cells / mm² (1850 cells / 176.7 mm²) was reached, but after nine days, a second increase in cell proliferation was observed.

The initial increase in proliferation was least prominent on collagen type I, PEI, gelatin and glass. The attachment ratios were also low on these surfaces. It seems plausible that the proliferation rate is partially dependent on the attachment ratio [17]. Newly emerged cells must attach prior to the next proliferation cycle, and delayed attachment (low attachment ratio) may thus slow down the proliferation rate. Insignificant differences in attachment may thus be magnified to significant differences in proliferation.

We observed a decrease in proliferation rate upon reaching a density of approximately 10 cells / mm². Cessation of proliferation upon increase of HSC density has been observed before, and was recently ascribed to a contact-mediated mechanism of growth-regulation [34].

Contrary to the other coatings and glass, the second increase in proliferation rate after 9 days was not observed on collagen type I and PEI. The number of HSCs on collagen remained equal, while the number of HSCs on PEI even declined.

In conclusion, the proliferation rate is lowest on collagen type I and on PEI because (i) proliferation during the first 9 days of culturing is slower; (ii) the second burst in proliferation does not occur. Moreover, on PEI cells eventually die. Thus when a coating is needed to encourage HSC proliferation, we do not recommend choosing collagen type I or PEI from our panel. The results on collagen are remarkable, since it is frequently used as a matrix or coating to culture Schwann cells [20]. However, systematic comparison of HSC proliferation on different coatings has not been performed before, and it was therefore not known that other coatings enhance HSC proliferation more.

Previous studies in which the SC number seeded in an artificial nerve graft was varied, demonstrated that the total number of axons and the number of myelinated axons was larger in those groups in which a large number of Schwann cells was seeded [6, 9]. It is to be expected that Schwann cell proliferation ceases upon arrival of axons, since axonal signals control SC differentiation and myelination by regulation of the Oct-6 Schwann cell-specific enhancer [35], and cause SCs to transform into a non-proliferating cell state [19]. Regenerating axons will start ingrowth into an artificial nerve guide approximately two to three weeks after implantation [36]. It is therefore necessary that the coating chosen enhances Schwann cell proliferation especially during this period. The coating on which Schwann cells proliferate the most is therefore preferable. Fibronectin, laminin and PDL show maximal proliferation of human Schwann cells during the first 15 days of culturing, and are therefore the most suitable candidates from our panel for coating of a synthetic nerve graft.

The extent to which the coating enhances neurite extension is also of great importance. Poly-electrolytes allow neurite extension from peripheral and central nervous system neurons *in vitro* [37-39], but, to the best of our knowledge, PDL was never used to coat a synthetic nerve graft *in vivo*. Fibronectin and laminin to the contrary, were repeatedly demonstrated not only to enhance neurite extension *in vitro* [38, 40, 41], but to also stimulate nerve regeneration *in vivo* in a synthetic nerve graft [3, 42-44]. It was even demonstrated that laminin and

fibronectin act synergistically with respect to the enhancement of neurite outgrowth, both *in vitro* [16] and *in vivo* [45-47].

The enhancement of neurite regeneration by laminin and fibronectin can be explained by the observation that regenerating axons upregulate receptors for ECM molecules in order to establish an integrin mediated interaction with these proteins [48, 49]. Recently, the $\alpha 7 \beta 1$ integrin, a specific receptor for the laminin 1, 2 and 4 isoforms, was demonstrated to be strongly upregulated after axotomy, indicating an important role for this integrin in axonal regeneration [50].

Combining the HSC proliferation promotive capacities that we demonstrated in this study, and the neurite outgrowth promotive capacities described in literature, we would choose to coat a synthetic nerve guide with a combination of laminin and fibronectin. It would even be better to disperse these ECMs in a three dimensional matrix as to allow the outgrowing neurites to exploit the entire volume of the synthetic nerve graft. Results obtained with Matrigel®, a matrix consisting of 80% laminin, were contradictory [6, 42-44, 51], while results obtained with a lower concentration of laminin (50 $\mu\text{g}/\text{ml}$; 5% m/v) were uniformly positive [3, 38, 40, 45-47].

At such low concentrations, however, laminin does not aggregate into a gel. Thus we need another matrix protein to constitute the bulk of the gel. Although we demonstrated that collagen type I did not promote the proliferation of HSCs, it did not inhibit proliferation either. Therefore, collagen gel is suitable to serve as a matrix into which low concentrations of laminin and fibronectin (50 $\mu\text{g}/\text{ml}$; 5% m/v) can be dispersed. In a follow up study we will evaluate whether the introduction of this gel into a synthetic nerve graft, indeed improves peripheral nerve regeneration.

Conclusion

Attachment and proliferation of human Schwann cells on various culture substrata were compared, with the aim to select among these substrata the most suitable ones for coating the interior of synthetic nerve guides. The attachment study demonstrated that the number of Schwann cells adhering to the coatings was comparable, as was their morphology. We concluded that human Schwann cells easily attach to almost any culture substratum, and that attachment is not an important consideration in the selection of suitable materials for coating the interior of synthetic nerve guides.

In the first six days after seeding the proliferation rate was high on fibronectin, laminin, PDL and crosslinked gelatin, but only moderate on collagen type I, PEI, gelatin and glass. On all coatings and on glass the proliferation rate diminished whenever a density of ca. 10 cells / mm^2 was reached. After nine days, on all coatings, except on collagen type I and on PEI, Schwann cell proliferation increased again, which was ascribed to a contact-mediated mechanism of

growth-regulation. The number of Schwann cells on collagen, however, remained equal, while the number of Schwann cells on PEI even declined.

Considering the relative poor performance of collagen type I or PEI with respect to Schwann cell proliferation, we do not recommend the application of these materials to synthetic nerve guides. Fibronectin, laminin and PDL demonstrated the best proliferation rates during the 15 day culture period. Of these fibronectin and laminin have a proven ability to enhance neurite extension. Fibronectin and laminin are therefore the most suitable candidates from our panel for coating the interior of a synthetic nerve guide.

Acknowledgements

9

214

We wish to thank Dr. P. Eilers for help with the statistical analysis of the data.

References

1. Fawcett JW, Keynes RJ. Peripheral nerve regeneration. *Annu Rev Neurosci* 1990; 13: 43-60.
2. Bixby JL, Lilien J, Reichardt LF. Identification of the major proteins that promote neuronal process outgrowth on Schwann cells in vitro. *J Cell Biol* 1988; 107: 353-61.
3. Longo FM, Hayman EG, Davis GE, Ruoslahti E, Engvall E, Manthorpe M, Varon S. Neurite-promoting factors and extracellular matrix components accumulating in vivo within nerve regeneration chambers. *Brain Res* 1984; 309: 105-17.
4. Bunge MB, Williams AK, Wood PM, Uitto J, Jeffrey JJ. Comparison of nerve cell and nerve cell plus Schwann cell cultures, with particular emphasis on basal lamina and collagen formation. *J Cell Biol* 1980; 84: 184-202.
5. Bunge MB, Williams AK, Wood PM. Neuron-Schwann cell interaction in basal lamina formation. *Dev Biol* 1982; 92: 449-60.
6. Guenard V, Kleitman N, Morrissey TK, Bunge RP, Aebischer P. Syngeneic Schwann cells derived from adult nerves seeded in semipermeable guidance channels enhance peripheral nerve regeneration. *J Neurosci* 1992; 12: 3310-20.
7. Kim DH, Connolly SE, Kline DG, Voorhies RM, Smith A, Powell M, Yoes T, Daniloff JK. Labeled Schwann cell transplants versus sural nerve grafts in nerve repair. *J Neurosurg* 1994; 80: 254-60.
8. Bryan DJ, Wang KK, Chakalis-Haley DP. Effect of Schwann cells in the enhancement of peripheral nerve regeneration. *J Reconstr Microsurg* 1996; 12: 439-46.
9. Ansellin AD, Fink T, Davey DF. Peripheral nerve regeneration through nerve guides seeded with adult Schwann cells. *Neuropathol Appl Neurobiol* 1997; 23: 387-98.
10. Hadlock T, Elisseeff J, Langer R, Vacanti J, Cheney M. A tissue-engineered conduit for peripheral nerve repair. *Arch Otolaryngol Head Neck Surg* 1998; 124: 1081-6.
11. Rutkowski GE, Heath CA. Development of a Bioartificial Nerve Graft. II. Nerve Regeneration in Vitro. *Biotechnol Prog* 2002; 18: 373-9.
12. Kleinman HK, Klebe RJ, Martin GR. Role of collagenous matrices in the adhesion and growth of cells. *J Cell Biol* 1981; 88: 473-85.
13. Keilhoff G, Fansa H, Smalla KH, Schneider W, Wolf G. Neuroma: a donor-age independent source of human Schwann cells for tissue engineered nerve grafts. *Neuroreport* 2000; 11: 3805-9.
14. Shen ZL, Lassner F, Becker M, Walter GF, Bader A, Berger A. Viability of cultured nerve grafts: An assessment of proliferation of Schwann cells and fibroblasts. *Microsurgery* 1999; 19: 356-63.
15. Keilhoff G, Fansa H, Schneider W, Wolf G. In vivo predegeneration of peripheral nerves: an effective technique to obtain activated Schwann cells for nerve conduits. *J Neurosci Methods* 1999; 89: 17-24.
16. Baron-van Evercooren A. Fibronectin promotes rat Schwann cell growth and motility. *J Cell Biol* 1982; 93: 211-6.
17. McGarvey ML, Baron-Van Evercooren A, Kleinman HK, Dubois-Dalcq M. Synthesis and effects of basement membrane components in cultured rat Schwann cells. *Dev Biol* 1984; 105: 18-28.
18. Chi H, Horie H, Hikawa N, Takenaka T. Isolation and age-related characterization of mouse Schwann cells from dorsal root ganglion explants in type I collagen gels. *J Neurosci Res* 1993; 35: 183-7.
19. Chernousov MA, Carey DJ. Schwann cell extracellular matrix molecules and their receptors. *Histol Histopathol* 2000; 15: 593-601.
20. Eccleston PA, Mirsky R, Jessen KR. Type I collagen preparations inhibit DNA synthesis in glial cells of the peripheral nervous system. *Exp Cell Res* 1989; 182: 173-85.
21. Hu M, Sabelman EE, Tsai C, Tan J, Hentz VR. Improvement of Schwann cell attachment and proliferation on modified hyaluronic acid strands by polylysine. *Tissue Eng* 2000; 6: 585-93.

22. Ruegg UT, Hefti F. Growth of dissociated neurons in culture dishes coated with synthetic polymeric amines. *Neurosci Lett* 1984; 49: 319-24.
23. Reinders JH, De Groot PG, Gonsalves MD, Zandbergen J, Loesberg C, Van Mourik JA. Isolation of a storage and secretory organelle containing Von Willebrand protein from cultured human endothelial cells. *Biochem Biophys Acta* 1984; 804: 361-9.
24. Van den Berg LH, Bar PR, Sodaar P, Mollee I, Wokke JH, Logtenberg T. Selective expansion and long-term culture of human Schwann cells from sural nerve biopsies. *Ann Neurol* 1995; 38: 674-8.
25. Lamers CH, van de Griend RJ, Braakman E, Ronteltap CP, Benard J, Stoter G, Gratama JW, Bolhuis RL. Optimization of culture conditions for activation and large-scale expansion of human T lymphocytes for bispecific antibody-directed cellular immunotherapy. *Int J Cancer* 1992; 51: 973-9.
26. Morrissey TK, Kleitman N, Bunge RP. Isolation and functional characterization of Schwann cells derived from adult peripheral nerve. *J Neurosci* 1991; 11: 2433-42.
27. Spierings E, De Boer T, Zulianello L, Ottenhoff TH. Novel mechanisms in the immunopathogenesis of leprosy nerve damage: the role of Schwann cells, T cells and *Mycobacterium leprae*. *Immunol Cell Biol* 2000; 78: 349-55.
28. van Dorp R, Jalink K, Oudega M, Marani E, Ypey DL, Ravestloot JH. Morphological and functional properties of rat dorsal root ganglion cells cultured in a chemically defined medium. *Eur J Morphol* 1990; 28: 430-44.
29. van Wachem PB, Beugeling T, Feijen J, Bantjes A, Detmers JP, van Aken WG. Interaction of cultured human endothelial cells with polymeric surfaces of different wettabilities. *Biomaterials* 1985; 6: 403-8.
30. Levi ADO. Characterization of the technique involved in isolating Schwann cells from adult human peripheral nerve. *J Neurosci Meth* 1996; 68: 21-6.
31. Miller C, Shanks H, Witt A, Rutkowski G, Mallapragada S. Oriented Schwann cell growth on micropatterned biodegradable polymer substrates. *Biomaterials* 2001; 22: 1263-9.
32. Kleinman HK, Cannon FB, Laurie GW, Hassell JR, Aumailley M, Terranova VP, Martin GR, DuBois-Dalcq M. Biological activities of laminin. *J Cell Biochem* 1985; 27: 317-25.
33. Pego AP, Vleggeert-Lankamp CLAM, Deenen M, Lakke EAJF, Grijpma DW, Poot AA, Marani E, Feijen J. Adhesion and growth of human Schwann cells on trimethylene carbonate (co)polymers. *J Biomed Mater Res* 2003; accepted for publication.
34. Casella GT, Wieser R, Bunge RP, Margitich IS, Katz J, Olson L, Wood PM. Density dependent regulation of human Schwann cell proliferation. *Glia* 2000; 30: 165-77.
35. Mandemakers W, Zwart R, Jaegle M, Walbeehm E, Visser P, Grosveld F, Meijer D. A distal Schwann cell-specific enhancer mediates axonal regulation of the Oct-6 transcription factor during peripheral nerve development and regeneration. *Embo J* 2000; 19: 2992-3003.
36. Williams LR, Longo FM, Powell HC, Lundborg G, Varon S. Spatial-temporal progress of peripheral nerve regeneration within a silicone chamber: parameters for a bioassay. *J Comp Neurol* 1983; 218: 460-70.
37. Yavin Z, Yavin E. Survival and maturation of cerebral neurons on poly(L-lysine) surfaces in the absence of serum. *Dev Biol* 1980; 75: 454-9.
38. Rogers SL, Letourneau PC, Palm SL, McCarthy J, Furcht LT. Neurite extension by peripheral and central nervous system neurons in response to substratum-bound fibronectin and laminin. *Dev Biol* 1983; 98: 212-20.
39. Mouveroux JM, Lakke EA, Marani E. Intrinsic properties inhibit axonal outgrowth from neonatal rat spinal cord explant. *Arch Physiol Biochem* 2002; 110: 177-85.
40. Manthorpe M, Engvall E, Ruoslahti E, Longo FM, Davis GE, Varon S. Laminin promotes neuritic regeneration from cultured peripheral and central neurons. *J Cell Biol* 1983; 97: 1882-90.

41. Sephel GC, Burrous BA, Kleinman HK. Laminin neural activity and binding proteins. *Dev Neurosci* 1989; 11: 313-31.
42. Madison RD, da Silva CF, Dikkes P. Entubulation repair with protein additives increases the maximum nerve gap distance successfully bridged with tubular prostheses. *Brain Res* 1988; 447: 325-34.
43. Madison R, da Silva CF, Dikkes P, Chiu TH, Sidman RL. Increased rate of peripheral nerve regeneration using bioresorbable nerve guides and a laminin-containing gel. *Exp Neurol* 1985; 88: 767-72.
44. Kljavin IJ, Madison RD. Peripheral nerve regeneration within tubular prostheses; effects of laminin and collagen matrices on cellular ingrowth. *Cells Materials* 1991; 1: 17-28.
45. Woolley AL, Hollowell JP, Rich KM. Fibronectin-laminin combination enhances peripheral nerve regeneration across long gaps. *Otolaryngol Head Neck Surg* 1990; 103: 509-18.
46. Bailey SB, Eichler ME, Villadiego A, Rich KM. The influence of fibronectin and laminin during Schwann cell migration and peripheral nerve regeneration through silicone chambers. *J Neurocytol* 1993; 22: 176-84.
47. Chen YS, Hsieh CL, Tsai CC, Chen TH, Cheng WC, Hu CL, Yao CH. Peripheral nerve regeneration using silicone rubber chambers filled with collagen, laminin and fibronectin. *Biomaterials* 2000; 21: 1541-7.
48. Lefcort F, Venstrom K, McDonald JA, Reichardt LF. Regulation of expression of fibronectin and its receptor, alpha 5 beta 1, during development and regeneration of peripheral nerve. *Development* 1992; 116: 767-82.
49. Kloss CU, Werner A, Klein MA, Shen J, Menuz K, Probst JC, Kreutzberg GW, Raivich G. Integrin family of cell adhesion molecules in the injured brain: regulation and cellular localization in the normal and regenerating mouse facial motor nucleus. *J Comp Neurol* 1999; 411: 162-78.
50. Werner A, Willem M, Jones LL, Kreutzberg GW, Mayer U, Raivich G. Impaired axonal regeneration in alpha7 integrin-deficient mice. *J Neurosci* 2000; 20: 1822-30.
51. Valentini RF, Aebischer P, Winn SR, Galletti PM. Collagen- and laminin-containing gels impede peripheral nerve regeneration through semipermeable nerve guidance channels. *Exp Neurol* 1987; 98: 350-6.
52. Cravioto H, Lockwood R. The behaviour of normal peripheral nerve in tissue culture. *Z Zellforsch* 1968; 90: 186-201.
53. Murray MR, Stout AP. Characteristics of human Schwann cells in vitro. *Anat Rec* 1942; 84: 275-94.
54. Brockes JP, Fields KL, Raff MC. A surface antigenic marker for rat Schwann cells. *Nature* 1977; 266: 364-6.
55. Bolin LM, Iismaa TP, Shooter EM. Isolation of activated adult Schwann cells and a spontaneously immortal Schwann cell clone. *J Neurosci Res* 1992; 33: 231-8.
56. Scarpini E, Meola G, Baron P, Beretta S, Velicogna M, Scarlato G. S-100 protein and laminin: immunocytochemical markers for human Schwann cells in vitro. *Exp Neurol* 1986; 93: 77-83.
57. Fields KL, Raine CS. Ultrastructure and immunocytochemistry of rat Schwann cells and fibroblasts in vitro. *J Neuroimm* 1982; 2: 155-66.
58. Cornbrooks CJ, Carey DJ, McDonald JA, Timpl R, Bunge RP. In vivo and in vitro observations on laminin production by Schwann cells. *Proc natl Acad Sci USA* 1983; 80: 3850-4.

