



Universiteit  
Leiden  
The Netherlands

## **Gangliosides and anti-ganglioside antibodies in neuromuscular synaptic function**

Zitman, F.M.P.

### **Citation**

Zitman, F. M. P. (2010, January 20). *Gangliosides and anti-ganglioside antibodies in neuromuscular synaptic function*. Retrieved from <https://hdl.handle.net/1887/14568>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14568>

**Note:** To cite this publication please use the final published version (if applicable).

---

# CHAPTER 3.5

## The neuropathic potential of anti-GM1 autoantibodies is regulated by the local glycolipid environment in mice

Kay N. Greenshields <sup>a</sup>, Susan K. Halstead <sup>a</sup>, Femke M.P. Zitman <sup>b,c</sup>, Simon Rinaldi <sup>a</sup>, Kathryn M. Brennan <sup>a</sup>, Colin O'Leary <sup>a</sup>, Luke H. Chamberlain <sup>d</sup>, Alistair Easton <sup>a</sup>, Jennifer Roxburgh <sup>a</sup>, John Padiani <sup>d</sup>, Koichi Furukawa <sup>e</sup>, Keiko Furukawa <sup>e</sup>, Carl S. Goodyear <sup>a</sup>, Jaap J. Plomp <sup>b,c</sup>, and Hugh J. Willison <sup>a</sup>

<sup>a</sup>Division of Clinical Neurosciences, Glasgow Biomedical Research Centre, University of Glasgow, Glasgow G12 8TA, UK.

Departments of <sup>b</sup>Neurology and <sup>c</sup>Molecular Cell Biology – Group Neurophysiology, Leiden University Medical Centre, PO Box 9600, NL-2300 RC Leiden, The Netherlands.

<sup>d</sup>Henry Wellcome Laboratory of Cell Biology, Division of Biochemistry and Molecular Biology, Faculty of Biomedical and Life Sciences, University of Glasgow, Glasgow G12 8QQ, UK.

<sup>e</sup>Department of Biochemistry II, Nagoya University School of Medicine, Nagoya, Japan.

*Published in the Journal of Clinical Investigation 119(3) (2009) 595 - 610*

## ***Abstract***

Anti-GM1 ganglioside autoantibodies are used as diagnostic markers for motor axonal peripheral neuropathies and are believed to be the primary mediators of such diseases. However, their ability to bind and exert pathogenic effects at neuronal membranes is highly inconsistent. Using human and mouse monoclonal anti-GM1 antibodies to probe the GM1-rich motor nerve terminal membrane in mice, we here show that the antigenic oligosaccharide of GM1 in the live plasma membrane is cryptic, hidden on surface domains that become buried for a proportion of anti-GM1 antibodies, due to a masking effect of neighbouring gangliosides. The cryptic GM1 binding domain was exposed by sialidase treatment that liberated sialic acid from masking gangliosides including GD1a or by disrupting the live membrane by freezing or fixation. This cryptic behaviour was also recapitulated in solid-phase immunoassays. These data show that certain anti-GM1 antibodies exert potent complement activation-mediated neuropathogenic effects, including morphological damage at living terminal motor axons, leading to a block of synaptic transmission. This occurred only when GM1 was topologically available for antibody binding, but not when GM1 was cryptic. This revised understanding of the complexities in ganglioside membrane topology provides a mechanistic account for wide variations in the neuropathic potential of anti-GM1 antibodies.

## ***Acknowledgements***

This work was sponsored by grants from the Wellcome Trust (#077041/Z/05/Z, to HJW) and the Prinses Beatrix Fonds (#MAR04-0213 to JJP). KNG was supported by the Medical Research Council doctoral training account awarded to the University of Glasgow.

## Introduction

The sialic acid-containing glycosphingolipids known as gangliosides are concentrated in plasma membrane microdomains where they modulate the topological organization and function of membrane proteins (Sandhoff and Kolter, 2003; Simons and Ikonen, 1997). Their oligosaccharide head groups protrude from the lipid bilayer into the extracellular environment to act as (co)receptors for a diverse range of glycan binding proteins including autoantibodies, sialic acid binding Ig-like lectins (siglecs), microbial toxins and viral components (Varki, 2007; Bullens et al., 2002; Willison, 2005; Crocker et al., 2007; Lencer and Tsai, 2003; Tsai et al., 2003).

In a subset of autoimmune peripheral nerve diseases, including Guillain-Barré syndrome (GBS) and multifocal motor neuropathy, autoantibody-ganglioside interactions are believed to be a critical pathogenic factor (Willison and Yuki, 2002; Yuki, 2001). Serum anti-GM1, -GD1b -GQ1b and -GD1a ganglioside antibodies are associated with nerve injury in both human clinical studies and animal models (Goodfellow et al., 2005; Kusunoki et al., 1996; Yuki et al., 2001; Plomp et al., 1999), with anti-GM1 antibodies being highly associated with motor neuropathy variants (Willison and Yuki, 2002). With respect to the antibody *induction* phase of the illness, it is clearly established that anti-GM1 antibodies can arise through molecular mimicry with structurally homologous *Campylobacter jejuni* lipooligosaccharides (LOS) (Bowes et al., 2002; Yuki et al., 1993; Yuki et al., 2004; Ang et al., 2002). In contrast, examination of the *effector* pathways through which anti-GM1 antibodies selectively bind to and induce injury in motor nerve membranes, while avoiding damage to other neural and non-neural plasma membranes containing abundant GM1, is confounded by inconsistent and often counterintuitive data (Willison and Yuki, 2002; Arasaki et al., 1993; Buchwald et al., 2007; Nores et al., 2007; Paparounas et al., 1999). In particular, the sensitivity or resistance of the membrane toward undergoing anti-GM1 antibody-mediated injury cannot be fully explained by the presence and density of plasma membrane GM1.

One reason for the uncertainties surrounding anti-GM1 effector pathways may be that protein-ganglioside interactions are typically identified by in vitro solid phase binding studies using immobilized gangliosides or structurally related natural and synthetic glycans. The translation of this in vitro binding data to physiologically and pathophysiologically relevant protein-glycan binding behaviour in intact membranes in vivo is where the complexities and inconsistencies arise. For example, an antibody that binds a specific glycan by immunoassay may apparently be unable to bind the same ganglioside when present in an intact membrane (Schwarz and Futerman, 1996). Furthermore, different anti-GM1-specific antibodies can have very different binding patterns in the CNS (Iwamoto et al., 1990; Laev and Mahadik, 1989). In addition to differences in antibody affinities, one explanation for such discrepancies might be that, within the complex environment of glycolipid-enriched microdomains, the interacting oligosaccharide headgroup is masked from the protein binding partner by surrounding molecules. Furthermore, fixation procedures might influence the antibody binding characteristics of gangliosides (Schwarz and Futerman, 1996). However, the detailed mechanisms underlying these determinants of antibody-ganglioside binding are unknown.

In the current study we addressed these issues by investigating a group of mouse and human anti-GM1 monoclonal antibodies (mAbs) for their potential neuropathogenic effects at mouse motor nerve terminals and by studying in detail the underlying

topological requirements for their binding to GM1 in neuronal membrane. Previously we showed that anti-GQ1b and anti-GD1a antibodies bind to the presynaptic motor nerve ending and activate complement, leading to membrane attack complex (MAC, C5b-9) formation, which causes intense neurotransmitter release and ultrastructural destruction, thereby blocking synaptic transmission at the neuromuscular junction (NMJ) (Goodfellow et al., 2005; Plomp et al., 1999; Goodyear et al., 1999; Halstead et al., 2004). We here show that mouse and human anti-GM1 mAbs also bring about these destructive neuropathic effects but that antigenic GM1 in the living neuronal membrane is cryptic for a proportion of the mAbs, thereby rendering them harmless, due to a masking effect of neighbouring gangliosides including GD1a. These findings indicate that anti-GM1 antibodies are pathogenically relevant in motor neuropathy and that masking and unmasking of GM1, such as may occur in membranes undergoing dynamic changes associated with normal axonal maintenance or during axonal regeneration, might have a profound influence on antibody pathogenicity (Yang et al., 2006).

## **Methods**

### **Anti-ganglioside mAbs**

Mice lacking complex gangliosides (GM2s-KO) were used to generate anti-ganglioside antibodies in response to inoculation of ganglioside liposomes or ganglioside mimicking LOS (Bowes et al., 2002; Townson et al., 2007; Aspinall et al., 1994). Hybridoma cells were maintained to generate stocks of mAbs, which were affinity purified by protein G sepharose chromatography (Amersham BioSciences), desalted and stored at -70 °C. Their binding properties are shown in the Table 3.2. Human mAbs were cloned as previously described (Townson et al., 2008) and their binding properties are shown in the Table 3.2.

### **Mice**

Male and female GD3s-KO (Okada et al., 2002), GM2s-KO (Takamiya et al., 1996) and wildtype (WT) mice were used at 4-8 weeks of age. All mice were killed by a rising concentration of CO<sub>2</sub> in accordance with UK Home Office, Dutch law and Leiden University guidelines.

### **Immunohistochemistry on frozen tissue sections**

*Topical mAb binding studies.* Diaphragm was snap-frozen and cryostat sectioned at 8 µm onto APES (Sigma-Aldrich) coated slides. For staining, mAbs were applied at 20 µg/ml and CTB-FITC (Sigma-Aldrich) at 1 µg/ml diluted in phosphate-buffered saline (PBS), along with BTx-TRITC (Molecular Probes, 1.3 µg/ml). mAbs/CTB were incubated for 3.5 h at 4 °C. For secondary detection of mAbs, slides were rinsed in PBS and anti-mouse IgG-FITC (Southern Biotech, 1.3 µg/ml in PBS) was applied for 3 h at 4 °C. All slides were mounted in Citifluor (Citifluor) following a final rinse in PBS. For CTB blocking of mAbs, sections were firstly incubated in unlabelled CTB (4 µg/ml in PBS, Sigma) or PBS as a control, for 1.5 h at 4 °C, rinsed thoroughly in PBS and mAbs applied as before.

*Ig, C3, MAC and neurofilament quantification.* Diaphragm (previously exposed *ex vivo* to mAbs and normal human serum as complement source) was cryostat cut at 8

and 20  $\mu\text{m}$  onto APES coated slides (thicker sections were designated for neurofilament staining). All solutions were made up in PBS, and incubations done at 4°C. BTx-TRITC was used at 1.3  $\mu\text{g}/\text{ml}$  throughout. To detect mAb binding, FITC subtype specific anti-mouse IgG or anti-human IgM (Southern Biotech) was applied at 3.3  $\mu\text{g}/\text{ml}$  for 3 h at 4 °C, along with BTx-TRITC. For detection of the activated complement component C3, slides were incubated in FITC-Goat anti-human C3 (Dako, 33  $\mu\text{g}/\text{ml}$ ) and BTx-TRITC for 2 h at 4 °C. MAC staining was performed with mouse anti-human C5b-9 (Dako, 363  $\mu\text{g}/\text{ml}$ ), applied to slides for 3.5 h with BTx-TRITC. After rinsing in PBS, MAC was detected with FITC goat anti-mouse IgG2a applied at 5  $\mu\text{g}/\text{ml}$  for 3.5 h. Sections for neurofilament analysis were firstly stained for 1 h in BTx-TRITC, rinsed in PBS, and placed for 20 min in freezing ethanol (-20 °C) prior to overnight application of 1211 (rabbit polyclonal, diluted 1:750, Affinity BioReagents). Slides were rinsed and incubated in FITC goat anti-rabbit IgG (3.3  $\mu\text{g}/\text{ml}$ ) for 3.5 h. Slides were finally rinsed and mounted in Citifluor.

### **Immunostaining of live tissue**

For triangularis sternae neuromuscular preparations, the ribcage was removed and pinned in a Sylgard (Sigma-Aldrich) lined dish with standard Ringer's medium (119 mM NaCl, 4.5 mM KCl, 2 mM  $\text{CaCl}_2$ , 1 mM  $\text{MgSO}_4$ , 1 mM  $\text{NaH}_2\text{PO}_4$ , 23 mM  $\text{NaHCO}_3$ , 11 mM glucose, pH 7.4 pre-gassed with 95%  $\text{O}_2$  / 5%  $\text{CO}_2$ ) and triangularis sternae was exposed. Preparations were incubated in mAbs (100  $\mu\text{g}/\text{ml}$  in Ringer's) or CTB (2  $\mu\text{g}/\text{ml}$ ) and BTx-TRITC (2  $\mu\text{g}/\text{ml}$ ) for 2 h at 32 °C, followed by 30 min at 4 °C. Tissue was thoroughly rinsed in Ringer's and fixed for 20 min at room temperature in 4% PFA (Sigma-Aldrich). Tissue was rinsed again in Ringer's and incubated for 10 min in 0.1 M glycine. For detection of mAbs, tissue was incubated overnight in subtype-specific FITC anti-mouse or anti-human IgM secondary antibodies (3.3  $\mu\text{g}/\text{ml}$  in PBS). For neurofilament staining, 1211 (Affinity BioReagents) was applied overnight (rabbit polyclonal, diluted 1 in 200) in 1% Triton-X100/PBS and following 3 rinses in PBS, was detected with cy-5 anti-rabbit IgG (3.3  $\mu\text{g}/\text{ml}$  in PBS) at 4 °C for 8 h. Following a final rinse in PBS, all immunostained muscles were whole-mounted in Citifluor.

### ***Ex vivo* pathogenesis: hemi-diaphragm**

Mouse hemi-diaphragms with attached phrenic nerves were dissected out and pinned under tension in Sylgard lined dishes with oxygenated Ringer's medium. The dorsal portion of each was removed and immediately snap frozen for baseline immunohistological measurements. Hemi-diaphragms were incubated in either Ringer's (as a control) or mAbs (100  $\mu\text{g}/\text{ml}$ ) for 2 h at 32 °C, followed by 30 min at 4 °C and 10 min at room temperature. mAb solutions were removed and retained, then preparations rinsed in Ringer's.

Normal human serum (NHS) from a single donor stock was diluted to 40% in Ringer's, and added as a source of complement. After 1 h at room temperature, NHS was rinsed off and the diaphragm tissue snap-frozen before being processed for analysis.

### ***Ex vivo* electrophysiology at the NMJ**

Left and right hemi-diaphragms were dissected with their phrenic nerve attached and mounted in standard Ringer's medium (as before) at room temperature. mAbs and normal human serum were applied as described above.

Intracellular recordings of MEPPs at the NMJ were made from muscle fibres at room temperature (20-22 °C) using a glass micro-electrode (10-20 M $\Omega$ , filled with 3 M KCl) connected to a Geneclamp 500B (Axon Instruments/Molecular devices) for amplifying and filtering (1 Hz high-pass and 10 kHz low-pass) of the signal. The signal was digitized using a Digidata 1322A interface (Axon Instruments/Molecular Devices) using Mini Analysis 6.0.3 (Synaptosoft).

Electrophysiological data is presented as group mean  $\pm$  SEM of the mean muscle values calculated from the mean NMJ values. At least 10 NMJs were sampled per muscle per experimental condition.

### **Microscopy**

For quantification of immunofluorescence over the NMJ, each experimental repeat (n of 3 animals/experiments for statistical analysis) was stained in a separate staining run, and at least 120 NMJs were imaged using constant acquisition settings for each marker. Dual colour images were acquired with a Zeiss LSM 5 Pascal microscope (Carl Zeiss). All samples were analysed observer-blinded. Semi-quantification was based on the  $\alpha$ BTx-TRITC signal to define the NMJ, and was done with Scion Image analysis software (Scion Corporation), either by analysing the percentage of immunofluorescence over the NMJ (as previously described, (O'Hanlon et al., 2001)) or analysing the intensity of fluorescence over the NMJ. For illustrative images of whole mount triangularis sternae, 3D reconstructions were generated from z-stacks using the software package Voxx-2 (<http://www.nephrology.iupui.edu/imaging/voxx>).

### **Membrane treatments**

For ex-vivo neuromuscular preparations, 1-5 units/ml of Neuraminidase (*Clostridium perfringens*, Sigma) was applied to tissue for 1-1.5 h at 32 °C. The enzyme was rinsed off and tissues incubated in mAb and/or complement as already described. For CTB pre-incubation of tissue prior to neuraminidase, unconjugated CTB (4  $\mu$ g/ml) was applied to tissue for 1.5 h at 32 °C, and thoroughly rinsed off with Ringer's prior to neuraminidase treatment. For GPI-anchored protein removal, PC12 cells were incubated at 37 °C for 1 h in 2 units/ml of PI-PLC (Sigma-Aldrich). *Ex vivo* neuromuscular preparations were treated for 1.5 h at 32 °C in 3.5 units of PI-PLC. For cholesterol depletion, 5 mM methyl- $\mu$ -cyclodextrin (Sigma-Aldrich) was incubated for 45 min at 37 °C. Cellular cholesterol was measured using Infinity Cholesterol reagent (Alpha Laboratories).

### **GD1a/GM1 co-localization**

MOG35 (IgG2b) was primary labelled using a Zenon Mouse IgG Labelling Kit (Invitrogen) according to manufacturer's instructions. Co-localization of DG2 and MOG35 was determined using the Metamorph (version 6.3.3; Molecular Devices Corp.) "correlation plot" application to compare the fluorescence in each pixel of the FITC and TRITC channels after subtracting background fluorescence.

### **PC12 cell staining**

Cells were grown on poly-L-lysine (Sigma-Aldrich) coated coverslips, and mAbs applied (12.5  $\mu$ g/ml in PBS) for 30 min at 4 °C. Following 3 rinses in PBS, mAbs were detected with FITC anti-mouse IgG (3.3  $\mu$ g/ml) for 45 min at 4 °C. Coverslips were rinsed again, fixed for 15 min in 4% PFA, and finally rinsed before mounting in Citiflour. For "Live/Dead staining" (Molecular Probes Inc.) a mixture of Calcein green (2  $\mu$ M) and ethidium homodimer-1 (EthD-1, 2  $\mu$ M) was applied to cells in PBS

for 30 min at 37 °C. Coverslips were rinsed in PBS, mounted in PBS and immediately imaged.

#### **Raft isolation, affinity purification and Western blotting**

Approximately  $100 \times 10^6$  PC12 cells were suspended in either MOG35 or control mAb EG1 (anti-GD3, GT1a and GQ1b) at 15 µg/ml for 1 h at 4 °C. Cells were washed in chilled PBS, and kept on ice for subsequent steps. Cells were resuspended in 500 µl of 25 mM MES/150 mM NaCl, pH 6.5 (MBS), 0.5% Brij-96 (Sigma-Aldrich) and protease inhibitors (Roche Applied Science). Following incubation at 4 °C for 20 min, solubilised cells were homogenised with 10 strokes of a Dounce homogenizer. Of the lysate, 0.4 ml was mixed with 80% sucrose and overlaid successively with 2.2 ml of 30% sucrose and 1.4 ml of 5% sucrose. Following overnight centrifugation at 240,000 g, fraction 1 was collected by removing 390 µl from the top of the gradient and successive fractions through to fraction 12 were collected in this way. Protease inhibitors were added immediately upon fraction collection.

Fractions 3-5 from MOG 35 incubated cells were pooled following retention of an equal aliquot from each (for Western blotting along with other fractions). This was also done for control mAb-incubated cells. Pooled samples were halved; half for affinity purification and half retained as starting material. For affinity purification, anti-mouse IgG-coated Dynabeads (Invitrogen) were incubated with the samples for 30 min at 4 °C on a rotating mixer. Tubes were exposed to a magnet and the unbound material removed and retained. Beads were rinsed 4x in PBS, Novex loading buffer (Invitrogen) added to the rinsed beads and boiled to release the bound fractions for electrophoresis alongside unbound material and starting material.

For electrophoresis, samples were reduced by addition of 25 mM dithiothreitol (DTT, Amersham) and run on Novex 16% Tricine gels (Invitrogen). Gels were blotted overnight in a transfer cell according to manufacturer's instructions (BioRad) onto nitrocellulose membranes. Membranes were blocked in 5% non-fat milk in PBS/0.01% Tween-20 (PBST) and incubated in anti-ganglioside mAbs (20 µg/ml) or 0.1 µg/ml horseradish-peroxidase conjugated CTB (HRP-CTB, Invitrogen). After rinsing, and if required, secondary antibody (HRP-conjugated anti-mouse IgG, Sigma) was added at 0.38 µg/ml in 1% non-fat milk/PBST. ECL (Amersham) was used to develop the membranes. Fractions 1-12 were also subject to Western blotting to confirm success of the raft isolation using antibodies against flotillin (BD Biosciences), SNAP 25 (Chamberlain et al., 2001) and transferrin receptor (Invitrogen), all at 10 µg/ml.

#### **Immunoassays**

Gangliosides were purchased from Sigma Chemicals or Accurate Chemical and Scientific. ELISA was performed as previously described (Willison et al., 1999). In brief, Immunolon 2HB plates (Thermo) were coated with 100 µl of ganglioside (2 µg/ml) diluted in methanol. To create ganglioside complexes, a 50:50 (v/v) mixture of the two component gangliosides in methanol was sonicated for 3 min. A volume of 100 µl of this solution, (containing 100 ng of each ganglioside) was then applied to the appropriate wells. Control wells were coated with methanol alone. The average OD reading from control wells was subtracted from all other wells to correct for background. The mAbs DG1 and DG2, used at their GM1 half maximal binding concentrations of 0.5 µg/ml and 0.4µg/ml respectively, and the appropriate secondary antibodies (peroxidase conjugated goat anti-mouse IgG, 1/3000, Sigma-Aldrich) were



diluted in 1% BSA (bovine serum albumin). The assay was conducted at 4 °C. For the PVDF glycoarray, gangliosides and ganglioside complexes (100 µg/ml) were prepared as described for ELISA. PVDF (polyvinylidene difluoride) membrane on glass slides was spotted (100 nl per spot) with each target ganglioside or ganglioside complex in duplicate at predefined locations on each membrane using an automated TLC sampler (Camag) as previously described (Kanter et al., 2006). Membranes were blocked with 2% BSA, and then probed with the same primary (1 µg/ml) and secondary antibodies (diluted 1/30000) as for ELISA. Chemiluminescent detection was performed using ECL plus (Amersham). Digitised film images were analysed and quantified by the array analysis component of ImageQuant TL software (Amersham).

### *Campylobacter jejuni*

Bacteria were cultured on Skirrow agar medium (E & O Laboratories) and maintained under microaerophilic conditions. For inactivation, bacteria were suspended in PBS and placed in a UV crosslinker for 30 min or suspended in 4% PFA for 25 min.

### FACS analysis

For FACS analysis, anti-ganglioside mAbs were used at 10 µg/ml, CTB-FITC at 1 µg/ml and Thy-1 (CD90, AbCam) at 7.8 µg/ml in PBS. Secondary antibodies (anti-mouse IgG-FITC) were used at 3.3 µg/ml, and cells were rinsed by pelleting in PBS. FACS analysis was performed using a FACScan flow cytometer (Becton Dickinson).

mAb	Isotype	Immunogen	Half maximal binding <sup>A</sup>				Affinity	
			GM1	GA1	GD1b	GD1a	K <sub>D</sub> <sup>B</sup>	Ligand
Mouse								
DG1	IgG2b	HS19 LOS <sup>C</sup> (GM1, GT1a <sup>+</sup> )	2.0x10 <sup>3</sup>	-	-	-	1.8 x10 <sup>-6</sup>	GM1
DG2	IgG3	GM1 liposome	2.5x10 <sup>3</sup>	6.7x10 <sup>2</sup>	2.0x10 <sup>2</sup>	-	3.5 x10 <sup>-7</sup>	GA1
MOG1	IgG3	GD1b liposome	-	-	5.0x10 <sup>3</sup>	-	1.0 x10 <sup>-8</sup>	GD1b
MOG35	IgG2b	O:19 LOS <sup>C</sup> (GM1, GD1a <sup>+</sup> )	-	-	-	1.0x10 <sup>4</sup>	9.5 x10 <sup>-7</sup>	GD1a
Human <sup>D</sup>								
SM1	IgM	Unknown	6.5x10 <sup>3</sup>	50	-	-	Unknown	
DO1	IgM	Unknown	5.5x10 <sup>3</sup>	4.3x10 <sup>3</sup>	5.7x10 <sup>3</sup>	-	Unknown	

<sup>A</sup> Half-maximal binding values represent the reciprocal of the antibody concentration (in mg/ml) giving half maximal binding as determined by ELISA. <sup>B</sup> BiaCore analysis of antibody Fab fragments was used to determine the binding affinity (K<sub>D</sub>) of the antibodies to the shown ligands (Townson et al., 2007). <sup>C</sup> HS19 and O:19 are the *C. jejuni* strains from which the LOS species were isolated and used to immunize GM2s-KO mice (Bowes et al., 2002). <sup>D</sup> Cloned from human patients with acute and chronic motor neuropathy (Paterson et al., 1995).

**Table 3.2. Mouse and human mAb ganglioside binding characteristics**

### Statistics

The Mann-Whitney test was used for statistical analyses of non-parametric immunohistological data, otherwise, a one way analysis of variance was used to determine statistical significance, both employing a 5% level of significance. Data are represented as box and whisker plots, with the horizontal line representing the median. One quarter of the data lying above the median, and one quarter of the data lying below the median are represented by the large box, and the vertical lines on either side show the remaining quarters of the data lying above and below the interquartile ranges. Outliers have not been shown for clarity, but were included in the statistical analysis. To assess whether binding to GM1 complexes was statistically different to GM1 alone on ELISA and PVDF array, General Linear Model ANOVA was used. Dunnett's method was employed to maintain a family error rate of <0.05 for multiple comparisons to the control level. Differences between DG1 and DG2

binding to each complex were assessed by the two sample t-test. Bonferroni's correction was applied in view of the multiple comparisons being made, again to maintain a family error rate of <0.05. The normality of each data set was verified by the Kolmogorov-Smirnov test ( $p > 0.15$  indicating normality).


## **Results**

### **Anti-GM1 binding characteristics of mouse and human mAbs**


The origin and properties of the anti-GM1 mouse mAbs DG1, DG2, anti-GD1b mAb MOG1, anti-GD1a mAb MOG35, and human anti-GM1 mAbs SM1 and DO1 (derived from motor neuropathy patients) are shown in Table 3.2. By ELISA, both DG1 and DG2 react with GM1 with very similar half maximal binding values (0.5 and 0.4  $\mu\text{g/ml}$  respectively) as previously reported (Townson et al., 2007). Screening by ELISA against a panel of gangliosides (GM1, GM2, GM3, GD1a, GD1b, GT1a, GT1b, GQ1b, GD2, GD3, GA1) has demonstrated that DG1 only binds GM1, suggesting a binding epitope unique to GM1 that includes the internal sialic acid, whereas DG2 binds GM1, GA1, and more weakly to GD1b, suggesting the epitope principally comprises the non-reducing galactose- $\beta$  1-3 *N*-acetylgalactosamine [Gal( $\beta$ 1-3)GalNAc] moiety common to the 3 glycolipids. Supplemental figure 1 shows details of relevant gangliosides and epitopes. Antibody sequencing has previously determined the variable region gene usage of DG1 and DG2 and established that they contain different immunoglobulin heavy and light chain gene rearrangements and thus have distinct clonal origins (Townson et al., 2007). Human mAb SM1 binds principally to GM1, and mAb DO1 binds to the Gal( $\beta$ 1-3)GalNAc moiety common to GM1, GA1 and GD1b.

### **Both DG1 and DG2 bind to motor nerve terminals in frozen nerve-muscle tissue**

Prior to evaluating the ability of DG1 and DG2 to mediate any pathogenic effects on nerve, we first assessed the binding potency of DG1 and DG2 to NMJs in immunohistological analyses of unfixed frozen muscle-nerve preparation of diaphragm from WT, GD3synthase deficient (GD3s-KO) and GalNAc-transferase deficient (GM2s-KO) mice. The ganglioside composition of the glycosyltransferase deficient mice is shown in supplemental figure 1. These mice are used as they are powerful tools for analysing the relevance of specific ganglioside targets, with GM1 being absent in GM2s-KO mice and overrepresented, along with GD1a and other a-series gangliosides, in GD3s-KO mice as previously reported (Okada et al., 2002). Frozen muscle sections were immunostained with DG1 or DG2 or, as positive control, with cholera toxin B-subunit (CTB), a highly specific GM1 ligand (Merritt et al., 1994). Fluorescence intensity and coverage of the NMJ area (defined by the area stained with fluorescent  $\alpha$ -bungarotoxin [ $\alpha$ BTx], labelling acetylcholine receptors at the NMJ) was quantitated. Under these conditions, all 3 ligands bound over nerve terminals in WT and GD3s-KO tissue (Figure 3.17). The intensity of DG1 and DG2 immunostaining at NMJs was similar in GD3s-KO tissue ( $\sim 100$  arbitrary units, AU), but the level was somewhat lower for DG1 than DG2 in WT tissue. As expected, in view of the GM1 deficiency, no DG1, DG2 or CTB staining was observed at NMJs of GM2s-KO mice.



**Figure 3.17.** Topical immunostaining of frozen diaphragm sections from WT, GD3s-KO (GD3s<sup>-/-</sup>) and GM2s-KO (GalNAcT<sup>-/-</sup>) mice. Fluorescence was quantified as either the area of the NMJ (stained with  $\alpha$ BTx-TRITC) that is covered with by anti-GM1 ligand (i.e. FITC fluorescence), or as the intensity of the FITC signal overlying the NMJ. All ligands bound most abundantly at the GD3s-KO NMJ compared to the WT NMJ. \*  $p < 0.05$ , WT versus GD3s-KO. All images were acquired at x 40 magnification. Scale bars: 60 $\mu$ m. **A. & B.** CTB fluorescence is strong at the WT and GD3s-KO, but absent at the GM2s-KO NMJ. **C.** Examples of CTB binding at NMJs of each strain, taken under constant microscope settings to represent the quantifications shown in **A** and **B**. **D. & E.** Anti-GM1 mAb DG1 stains both the WT and GD3s-KO, and is absent in the GM2s-KO NMJs. **F.** DG1 topical staining. **G. & H.** DG2 binding profile is evident at WT and GD3s-KO, but absent at the GM2s-KO NMJs. **I.** DG2 binding at the WT, GD3s-KO and GM2s-KO NMJs.



**Figure 3.18. Pathophysiological effects of anti-GM1 antibodies in living nerve-muscle preparations *ex vivo***

**A.** Reconstructed confocal images of triangularis sterni NMJs from GD3s-KO (GD3s<sup>-/-</sup>) mice, following *ex vivo* incubation in CTB, DG2 and DG1. For CTB and DG2, axonal staining is present along with staining of the parajunctional fibroblast. DG1 binding is undetectable.

**B.** Electrophysiology in *ex vivo* hemi-diaphragm of WT and GD3s-KO mice. mAbs were applied, followed by a source of complement, and under such conditions only DG2 caused an increase in MEPP frequency of each strain. Con, control.

**C.** DG1 effect in *ex vivo* hemi-diaphragm (of WT and GD3s-KO). Following *ex vivo* muscle nerve incubations in mAb (or Ringer's medium alone [R] as control), tissue was snap-frozen and sectioned for quantitative analysis. Graphs show IgG, C3c and MAC deposition (quantified as the signal intensity over the NMJ). Neurofilament (NF) loss is quantified as area coverage over the NMJ. DG1 binding is undetectable in the WT NMJ, and only weakly present in the GD3s-KO NMJ under maximal detection settings. MAC and C3c deposition do not translate into NF loss.

**D.** The experiment was performed as described for **C.**, but with the mAb DG2. DG2 is detectable in WT and GD3s-KO NMJs and activates complement to cause a NF loss compared with Ringer's medium alone-incubated control tissue. For both **C** and **D** note that the Y axes have been adjusted for clarity. \* p<0.05 compared with the Ringer's control.

Scale bars: 20µm.

**DG2 and DG1 have different capacities to bind to live mouse motor nerve terminals and to exert complement-mediated neuropathogenic effects**


We next examined motor nerve terminal binding and possible pathophysiological effects of DG1 and DG2 at living NMJs in dissected triangularis sternae and diaphragm nerve-muscle preparations maintained as live preparations in oxygenated Ringer's medium. In positive control experiments conducted to demonstrate the presence and level of GM1 at nerve terminals, CTB bound strongly to both GD3s-KO (Figure 3.18A, left panel) and WT (not shown) nerve terminals, with strong staining of the pre-synaptic axon of the NMJ. Since GM1 is widely distributed throughout the body, albeit at low levels, staining is also seen elsewhere, including connective tissue and blood vessels; thus CTB staining over other GM1 bearing structures is also seen in the upper region of Figure 3.18A (left panel). DG2 (Figure 3.18A, middle panel) showed a similar staining pattern to CTB, with particularly pronounced staining in this image of the para-junctional fibroblast, an extralaminar cell that caps the NMJ and also bears the GM1 ligand (Court et al., 2008). GM2s-KO nerve terminals and other structures were not stained under these conditions, as expected (data not shown). DG2 strongly bound to GD3s-KO NMJs (~150 arbitrary units) and to a lesser extent (20-50 arbitrary units) to WT NMJs (Figure 3.20D). Unexpectedly, and in marked contrast to DG2 and to the situation observed in frozen sections as described above, DG1 binding was undetectable at all living WT NMJs and bound only very weakly at GM1-rich GD3s-KO NMJs (undetectable at >50% of NMJ and only at <10 arbitrary units at the most strongly positive NMJs, Figure 3.18A, C). As expected, neither of the two mAbs bound at GM2s-KO NMJs (supplemental figure 2A). Thus, whereas both DG1 and DG2 are able to bind well to GM1 at NMJs in frozen muscle-nerve preparation sections, only DG2 is capable of binding GM1 at live NMJs in physiological preparations, indicating that the topographical conformation of GM1 and/or adjacent molecules in the neuronal membrane is substantially different under these two conditions.

In order to study the ability of bound anti-GM1 antibody at live NMJs to activate complement and induce neuropathological effects, we exposed DG1- and DG2-pretreated live nerve-muscle diaphragm to normal human serum (NHS; as a complement source) and electrophysiologically monitored the occurrence of miniature endplate potentials (MEPPs, the spontaneous unquantal acetylcholine release at NMJs) with intracellular microelectrode measurement. In the presence of human serum, DG2-pretreated NMJs showed a dramatic rise in MEPP frequency in WT and GD3s-KO preparations (Figure 3.18B and supplemental figure 2B), which caused asynchronous twitching of muscle fibres which eventually subsided and was followed by complete synaptic "silence". This resulted in paralysis by block of transmission at the NMJ as judged from the inability of the diaphragm muscle to contract following phrenic nerve stimulation. As expected, in view of the finding that DG1 lacks binding activity to live WT NMJs, no such NHS-induced electrophysiological effects were observed at DG1-pretreated preparations (Figure 3.18B and supplemental figure 2C).

We screened for immunohistological evidence of antibody-mediated complement activation by assessing C3c and MAC deposition at NMJs (Figure 3.18C, D and supplemental figure 2). Both components were clearly deposited at GD3s-KO and WT nerve terminals in conjunction with DG2 binding, although with less intensity in WT. At DG1 pre-treated nerve terminals from GD3s-KO mice in which very low levels of DG1 antibody were deposited, only very low levels of C3c (median <5 arbitrary units) and MAC (median <5 arbitrary units) were present. WT NMJs pre-treated with DG1 showed no antibody binding and lacked any complement


deposition. We next examined the structural integrity of the motor nerve terminal after MAC formation at the presynaptic membrane by quantifying neurofilament staining. DG2 pre-treated NMJs from WT and GD3s-KO mice that showed MAC deposition also displayed neurofilament loss (Figure 3.20D), whereas the very low level of DG1-dependent complement activation at GD3s-KO NMJs was insufficient to induce such loss (Figure 3.18C).

Taken together, these results in living nerve-muscle preparations show that of the two anti-GM1 mAbs DG1 and DG2, only DG2 binds to motor nerve terminals and induces complement-mediated neuropathophysiological and structural effects at motor nerve terminals. Apparently, DG1 is inhibited from binding GM1, despite its abundant presence, by unknown factors in the local environment of the living neuronal membrane. The observed neuropathic effects of DG2 are identical to the complement-mediated effects we have seen previously for anti-GQ1b and anti-GD1a antibodies at NMJs (Goodfellow et al., 2005; Plomp et al., 1999; Goodyear et al., 1999; O’Hanlon et al., 2001).



**Figure 3.19. Reactivity of anti-GM1 mAbs DG1 and DG2 to ganglioside complexes containing GM1 in solid-phase**

The ganglioside complex at each location is established by combining the row and column labels. Thus, coordinates 1,4 and 4,1 represent GM1/GD1a complex. Wells labelled x are negative controls (methanol only). **A.** ELISA. DG1 (left) and DG2 (right) both bind GM1 alone, with no difference in average OD. The binding of both antibodies to complexes of GM1 and GM2 or GD1a is reduced as compared with GM1 alone. No difference was observed with other combinations investigated (GM1:GM3, GM1:GD1b and GM1:GD3). DG1 binding to complexes GM1/GM2 and GM1/GD1a was less than that of DG2. Mean results  $\pm$  SEM from 3 experiments are shown. **B.** PVDF glycoarrays. DG1 was the primary antibody on the left membrane, DG2 on the right. Mean results  $\pm$  SEM for 3 experiments are shown. No significant difference in GM1 binding was observed for the two antibodies. DG1 binding to GM1 complexes was significantly reduced compared to GM1 alone ( $p > 0.05$ , significance level for these comparisons are not indicated in graph). DG2 binding GM1 complexes was marginally different compared with GM1 alone, but significant for GM1/GM2 and GM1/GD1a. The inhibitory effect of complexes on antibody binding is greater for DG1 than for DG2. The average absolute reduction in signal intensity for GM1/GD1a complex, compared to GM1 alone, is 75.1% for DG1 and 18.2% for DG2 ( $p < 0.05$ ). Error bars indicate SEM. 1,GM1; 2,GM2; 3,GM3; 4,GD1a; 5,GD1b; 6,GD3. \*  $p < 0.05$ , \*\*  $p < 0.01$ .



**Figure 3.20. Effect of neuraminidase treatment on anti-GM1 antibody binding and pathogenic activity**  
**A.** Reconstructed confocal images of DG1 binding at *ex vivo* GD3s-KO triangularis sterni NMJs. Left: DG1 binding in living tissue is undetectable at the NMJ (stained with  $\alpha$ BTx-TRITC). Right: DG1 binding (FITC) following neuraminidase treatment of tissue. DG1 binding overlies the NMJ, and is co-localized with the axonal NF staining (Cy5), suggesting DG1 is binding to the presynaptic aspect (the axon terminal). **B.** *Ex vivo* GD3s-KO hemi-diaphragm preparations, as described in Figure 3.180. Neuraminidase (N'ase) treatment of tissue prior to incubation with DG1 enabled DG1 to bind the NMJ (as shown by IgG deposition), fix complement, and cause a neurofilament (NF) loss. \*  $p < 0.05$  compared with control tissue incubated in Ringer's (minus neuraminidase) followed by DG1. **C.** Electrophysiology. DG1 and NHS as a source of complement caused a massive increase in MEPP frequency in the *ex vivo* GD3s-KO NMJ only when applied to neuraminidase-pretreated tissue. Scale bars: 20  $\mu$ m. Error bars represent SEM for experiments performed in triplicate.

**The nerve terminal ligand for DG1 and DG2, and the immunopathological target for DG2 in live membranes is GM1**

The absence of nerve terminal staining in GM2s-KO tissue with CTB, DG1 or DG2 (Figure 3.17) demonstrates that any potential ganglioside cross-reactivity of these GM1 ligands should only occur with the complex ganglioside series that are missing in GM2s-KO mice (supplemental figure 1). However, in order to assess for certain whether DG1 and DG2 might be binding to any of these ganglioside antigens other than GM1 at the nerve terminal in either frozen or live tissue, we conducted blocking experiments using the GM1-specific, high affinity ligand, CTB (supplemental figure 3A). In frozen tissue experiments, CTB pretreatment completely inhibited binding of DG1 and DG2 to GD3s-KO NMJs. Furthermore, CTB also inhibited DG2 binding in WT NMJs, excluding possible cross reactivity with Gal( $\beta$ 1-3)GalNAc bearing epitopes. We next demonstrated that DG2 (which binds GM1, GA1 and GD1b in ELISA) does not bind GD1b in the living neuronal membrane as follows. We first showed the presence of GD1b at NMJs by incubating tissues with an anti-GD1b monospecific antibody, MOG1 (Boffey et al., 2005), and observing intense immunohistochemical staining at motor nerve terminals of WT mice (but neither GM2s-KO nor GD3s-KO mice that both lack GD1b). MOG1 also induced associated complement deposition and structural injury, as determined by neurofilament loss (supplemental figure 3B). We then showed that MOG1 binding was unaffected by CTB blockade (supplemental figure 3C). These data therefore exclude the possibility that DG2 is binding GD1b in the living tissue. They thus demonstrate that the Gal( $\beta$ 1-3)GalNAc epitope on GD1b is cryptically orientated for DG2 binding *in vivo*, whereas the MOG1-reactive GD1b epitope is available for MOG1 binding *in vivo*. Since GA1 (asialo-GM1) is absent from the mammalian nervous system (Seyfried et al., 1996), this potential cross-reactive antigen is not relevant in this context and potential binding to Gal( $\beta$ 1-3)GalNAc-bearing glycoproteins is excluded by the absence of staining in GM2s-KO mice. Furthermore, and consistent with the complete blocking seen in frozen tissue sections, CTB pre-incubation of *ex vivo* muscle tissue from WT mice inhibits the complement mediated actions of DG2, evidenced by abolition of C3c and MAC depositions over the NMJ (supplemental figure 3D). Collectively, these data indicate that DG2 is only binding GM1 in nerve terminal membranes *in vivo*. It should also be noted that the binding of DG2 to GD1b in ELISA is very weak, (half maximal binding,  $2 \times 10^2$ , see Table 3.2), further indicating that it is unlikely to bind avidly to GD1b in membranes.

**Inhibitory interactions between GM1 and adjacent gangliosides form under solid-phase conditions**

We speculated that the inability of DG1 to bind GM1 in the living plasma membrane was due to a masking effect arising from *cis* interactions with adjacent gangliosides and that these were disrupted following the death of tissue and subsequent processing to frozen sections. Since GD3s-KO mice are only able to synthesise a-series gangliosides, the most abundantly expressed species being GM1 and GD1a, we concluded that the masking effect in this mouse strain was most likely due to a GM1/GD1a *cis* interaction, notwithstanding the possibility that other gangliosides (e.g. GM2, GM3) could mediate similar blocking function in membranes with other ganglioside compositions. To investigate the potential inhibitory interaction between GM1 and GD1a for DG1 binding, we first attempted to recapitulate the *in vivo* interaction in solid phase binding assays using ELISA and then using newly developed PVDF membrane formats (Figure 3.19). In ELISA, conducted with



GM1:GD1a-coated at a 1:1 ratio by weight, DG1 was significantly inhibited and DG2 partially inhibited from binding GM1 in the presence of GD1a (Figure 3.19A). Both monoclonal antibodies were used at their GM1 half maximal binding concentrations of 0.5  $\mu\text{g/ml}$  and 0.4  $\mu\text{g/ml}$  respectively. Furthermore, DG1 was significantly more impaired than DG2 from binding the GM1/GD1a complex. This inhibitory effect was not unique to GD1a, also occurring to significant extent with GM1:GM2 and a mild extent (not significant) with GM1/GD1b. In marked contrast, complexes of GM1/GM3 and GM1/GD3 did not produce any inhibitory effects on DG1 or DG2 binding, excluding the possibility that non-specific displacement of GM1 by a second, non-reactive ganglioside is accounting for the lower signal seen for the inhibitory GM1/ganglioside complexes.

When isolated GM1 was displayed on PVDF membranes as dot blots, there was no significant difference in the binding of DG1 and DG2 (Figure 3.19B). When GM1 was complexed with a range of other gangliosides, DG1 was almost completely inhibited from binding GM1. In contrast, DG2 binding to GM1 was minimally affected, although the difference in binding did reach significance for GM1/GM2 and GM1/GD1a complexes (Figure 3.19B). Throughout these PVDF studies, the reduction in complex binding for DG1 was substantially and significantly greater than for DG2. Under the PVDF array and ELISA assay conditions used in this study, DG2 is not demonstrated to bind to GD1b because of the very low half-maximal binding value for the DG2/GD1b interaction.

#### **Inhibitory interactions between GM1 and GD1a are not present in *C. jejuni* LOS**

To investigate whether or not this inhibitory interaction between GM1 and GD1a was a ganglioside-specific phenomenon or might also occur in other situations when GM1 and GD1a oligosaccharides were juxtaposed, we exploited the neuropathy-associated *Campylobacter* species that bear GM1 and GD1a mimics on their LOS (it should be noted that DG1 was cloned following immunization with LOS). A Penner O:19 *Campylobacter jejuni* isolate bearing both GM1 and GD1a mimics (50:50 ratio) in the core oligosaccharide was assessed for binding to DG1 (anti-GM1) and MOG35 (anti-GD1a) mAbs and CTB by FACS, in comparison with a non-ganglioside mimicking O:3 *C. jejuni* strain. DG1 was able to bind well to both live and killed (ultraviolet light or paraformaldehyde) O:19 isolate, indicating that an inhibitory complex between the GM1 and GD1a mimicking moieties of LOS is not present (supplemental figure 4). MOG35 and CTB also bound well to the O:19 isolate, confirming the presence of both the GM1 and GD1a mimics, and no ligands bound the control O:3 isolate (data shown for CTB only).

#### **Access of DG1 to GM1 in the living membrane is prevented by a masking effect of GD1a**

Based on the above data, we considered that the inability of DG1 to bind GM1 on the living neuronal membrane was due to closely adjacent gangliosides, including GD1a, which might *cis*-interact with GM1 to create a masking effect. Using live nerve-muscle tissue preparations from GD3s-KO mice (which highly overexpress GM1 and GD1a), we cleaved the terminal sialic acid from GD1a with neuraminidase treatment thereby converting GD1a to GM1, and subsequently assessed the live tissue for DG1 binding (Figure 3.20). This pretreatment massively promoted DG1 binding at the NMJ (Figure 3.20A, B), which subsequently activated complement, as shown by C3c and MAC deposition (Figure 3.22B). It thereby induced the previously described pathophysiological cascade (Figure 3.20C), characterized by increased MEPP

frequency, muscle fibre twitching, and paralysis, described above for DG2. This unmasking effect for DG1 was also evident in neuraminidase-treated tissue from WT C57Bl/6 and Balb/c mice (supplemental figure 5A).

In control experiments, we verified that neuraminidase treatment was able to digest most of the GD1a present at motor nerve terminals as assessed by a substantial reduction in the binding of the anti-GD1a mAb, MOG35 after treatment, in comparison with strong binding before treatment (supplemental figure 5B). DG1 binding following neuraminidase treatment was also completely blocked by pre-incubation with CTB, indicating that binding of DG1 was GM1-specific (supplemental figure 5C). Since neuraminidase treatment creates de novo GM1 from GD1a, it is very likely that this de novo GM1 would also be capable of acting as a ligand for DG1. Such binding confounds (without contradicting) the conclusion that masked “native” GM1 is released from its *cis* inhibition with GD1a by neuraminidase treatment to become available for DG1 binding. In order to investigate the extent to which the DG1 binding was directed to de novo GM1 generated from GD1a (rather than the pre-existing, native GM1), we first blocked native GM1 in the tissue preparation with CTB and then treated the tissue with neuraminidase, followed by probing with DG1 (supplemental figure 5D). Under these conditions DG1 binding to CTB-pretreated tissue was greatly reduced compared with binding to CTB-untreated tissue, indicating that a proportion of DG1 binding was to native GM1, unmasked from GD1a by neuraminidase treatment. After neuraminidase incubation, DG2 bound equally to both CTB pre-treated and untreated tissue, indicating that abundant de novo GM1 produced from GD1a was available for DG2 binding. The relatively weak binding of DG1 in comparison to DG2 (supplemental figure 5D) to de novo GM1 (i.e. CTB-blocked, neuraminidase-treated tissue; supplemental figure 5D), suggests that not all the de novo GM1 is available for DG1 binding. One explanation for this would be that de novo GM1 might itself form inhibitory *cis* interactions with remaining a-series gangliosides (e.g. GM2, GM3) in the plasma membrane of GD3s-KO mice, although this was not addressed experimentally beyond the solid-phase PVDF membrane data shown in Figure 3.19.

### **GM1 and GD1a co-localize to form an inhibitory environment for DG1 antibody binding in living PC12 cells and in lipid rafts**

To investigate the molecular environment that might account for the masking effects of GD1a we used rat neuroendocrine PC12 cells which, in their undifferentiated state, express gangliosides including GM1 and GD1a (Townson et al., 2008). DG1 was entirely unable to bind GM1 in living PC12 cells, in contrast to strong binding with CTB, DG2 and the anti-GD1a mAb MOG35 (Figure 3.21), as predicted from the live nerve terminal findings. Neuraminidase treatment resulted in intense DG1 binding to PC12 cells with complete concomitant loss of MOG35 binding (Figure 3.21B) under conditions in which cell viability was maintained, as monitored by ethidium homodimer-1 staining (data not shown). In co-localization studies of GM1 and GD1a, performed by comparing the binding of the mAbs DG2 and MOG35, the pixel by pixel correlation coefficient was 0.96, indicating very close proximity at the light microscopy level (Figure 3.21C). These findings in PC12 cells indicate a similar PC12 membrane interaction between GM1 and GD1a similar to that occurring in the mouse motor nerve terminal membrane and thereby suggest that this *cis*-inhibitory behaviour may be common other plasma membranes.


Since gangliosides are enriched in lipid rafts (Simons and Ikonen, 1997), we considered that the raft microenvironment might favour interactions between GM1


and GD1a that were inhibitory for DG1 binding. First we disrupted rafts in live PC12 cells with the cholesterol-sequestering agent methyl- $\beta$ -cyclodextrin to establish whether GM1 became available for DG1 binding. This was not observed; indeed a general attenuation of ligand binding was seen (supplemental figure 6A). We next considered that raft-localized (GPI-anchored) proteins might exert a masking effect on GM1 for DG1 binding and investigated this by cleaving them with phosphatidylinositol-specific PLC (PI-PLC). PI-PLC treatment enhanced DG2 and MOG35 binding, suggesting that GPI anchored proteins can exert a general and non-specific masking effect on both GM1 and GD1a, but had no significant effect on DG1 binding which remained negligible or absent (supplemental figure 6B). We also conducted the PI-PLC study at live NMJs and observed the same trend as seen in PC12 cells (supplemental figure 6C).

In order to support a molecular interaction hypothesis between GM1 and GD1a, it would be necessary to show that they co-localize in the same raft environment. To investigate this, we incubated PC12 cells with the anti-GD1a mAb MOG35 and an irrelevant isotype control, EG1 (which reacts with GQ1b, GT1a and GD3 but does not bind to PC12 cells; data not shown) and examined GD1a affinity-purified raft fractions by Western blot for GM1 content (Figure 3.21D). This demonstrated that at least a proportion of GM1 was present in the GD1a-enriched raft compartment. In control studies, EG1-affinity purification of raft fractions did not isolate any raft material (as assessed by raft protein markers) or yield any GM1 or GD1a signal. The raft fraction containing GD1a and GM1 also contained the raft marker flotillin, whereas SNAP25 was undetectable, indicating that a heterogeneous population of rafts exist with respect to protein and ganglioside composition in PC12 cells. This latter point was not investigated further.

**Figure 3.21. Localization of GM1 and GD1a to raft fractions in PC12 cells (next page).**

**A.** PC12 cell immunostaining (original magnification, x40). DG1 binding to PC12 cells (shown in phase contrast) is not detectable despite the presence of GM1, as shown by CTB staining of cells. Scale bars: 15  $\mu$ m. **B.** Effect of neuraminidase on DG1 and MOG35 binding to PC12 cells. Double staining reveals that control cells are positively stained with MOG35 (TRITC), with no binding of DG1 (FITC). Following neuraminidase (N<sup>d</sup>ase) treatment, MOG35 staining is diminished alongside a concomitant increase in DG1 binding. Scale bars: 15  $\mu$ m (images acquired at x40 magnification). **C.** GM1 and GD1a pixel-by-pixel co-localization. FITC and TRITC (anti-GM1 and anti-GD1a respectively) images (x 63 magnification) from double-stained PC12 cells, with co-localization appearing as yellow overlap. Scale bar represents 15  $\mu$ m. Plane by plane co-localization (linear scatter plot) shows strong co-localization. **D.** Western blot of raft immunoprecipitation based on MOG35 binding, allowing isolation of GD1a-positive rafts by anti-mouse IgG-coated beads. In irrelevant antibody-incubated cells, no rafts were isolated by anti-mouse IgG-coated beads. Bound sample was concentrated (x10) to amplify any potentially weak signal. In MOG35-incubated cells, a population of rafts was isolated by the beads. Isolated fractions contained the raft-associated protein flotillin, but not SNAP25, which was taken as evidence that the raft extraction procedure did not lead to coalescence of the heterogeneous raft population. Bound fractions also contained both the light chain (LC) and heavy chain (HC) of the anti-GD1a antibody, and the isolated rafts were positive for both GM1 and GD1a. SM, starting material, UB, unbound, B, bound, Bc, bound concentrated.





**Figure 3.22. Binding and neuropathophysiological effects of human anti-GM1 mAbs SM1 and DO1**

**A.** *Ex vivo* hemi-diaphragm preparations from GD3s-KO mice, as described for Figure 3.18. SM1 only binds NMJs following neuraminidase treatment and then causes a complement-mediated neurofilament (NF) loss. **B.** The experiment was performed as described for **A.**, but with DO1. **C.** Absence of SM1 staining at GD3s-KO triangularis sterni NMJs (left), and presence of staining when the tissue was pre-incubated in neuraminidase (N'ase; right). **D.** Electrophysiological analysis of the effects of SM1 and DO1 at GD3s-KO hemi-diaphragm NMJs, as described in Figure 3.20 for mouse mAb DG1. Following neuraminidase treatment, both SM1 and DO1 bind at NMJs and cause a complement-dependent increase in MEPP frequency.

Scale bars: 20  $\mu$ m. Error bars represent SEM for 3 experiments performed in triplicate.

### Human neuropathy-associated anti-GM1 mAbs also exert complement-mediated neuropathophysiological effects at the mouse motor nerve terminal, provided GM1 is unmasked

In order to bring clinical relevance to the findings observed with mouse mAbs, we screened human anti-GM1 antibodies cloned from motor neuropathy cases for similar behaviour. Human anti-GM1 IgM antibodies SM1 and DO1 (Paterson et al., 1995) were examined for their binding and complement-mediated neuropathophysiological effects at the mouse NMJ and we assessed a possible inhibitory effect of interaction between GM1 and GD1a (Figure 3.22). Neither mAb bound to GD3s-KO motor nerve terminals, and consequently they did not induce complement deposition (C3c and MAC) or neurofilament loss, as assessed with immunohistochemistry, and did not cause increased MEPP frequency, the electrophysiological hallmark of

neuropathogenicity. However, following neuraminidase pre-treatment of the nerve-muscle preparations, deposits of IgM were massively present. Consequently, the complement cascade was activated, as shown by C3 and MAC deposits, with resultant neurofilament loss and electrophysiological injury as manifested by an increased MEPP frequency.

## **Discussion**

This study has two principle findings. First, we demonstrate that under specific conditions, both human and mouse anti-GM1 antibodies have potent complement activation-mediated neuropathogenic effects at mouse terminal motor axons, including electrophysiological and morphological injury, leading to block of synaptic transmission at the NMJ, thereby resulting in paralysis of the nerve-muscle preparation. Second, we discovered that GM1 ganglioside in living neuronal membranes is masked from binding certain classes of anti-GM1 autoantibody by an interaction with GD1a (and presumably other gangliosides) that is able to shield a critical domain of the oligosaccharide head group involved in antibody binding. Consequently, this masking prevented the neuropathophysiological effects of these anti-GM1 antibodies from occurring. These findings have major implications for our understanding of the pathogenic potential of anti-ganglioside antibodies in neuropathy, their use and relevance as autoimmune disease diagnostics, and as ganglioside detection reagents. The epitope masking effect of this interaction is not a feature concerning all anti-GM1 antibodies, but is dependent upon subtle features of the antibody-antigen interaction that confer these characteristics, and needs to be assessed for each antibody on a case by case basis. Importantly, our results demonstrate that the ability of mouse and human anti-GM1 antibodies to bind isolated GM1 in solid phase assays or in frozen tissue sections cannot be used to predict their pathogenic behaviour *in vivo*. Our findings are likely to account for the markedly discrepant effects of anti-GM1 antibodies observed in many previously published models.

Anti-GM1 antibodies are mainly associated with the acute motor axonal variants of GBS and chronic multifocal motor neuropathies (Willison and Yuki, 2002) and our immunohistochemical and electrophysiological findings at motor nerve terminals may provide a neuropathophysiological explanation for a component of the paralytic symptoms in such patients. In support of this as a clinically relevant site, motor nerve terminal degeneration has been shown at NMJs in a motor point muscle biopsy of an acute motor axonal neuropathy variant GBS patient with anti-GM1 antibodies, while sural nerve biopsy was normal (Ho et al., 1997). The terminal portion of motor axons may be relatively vulnerable for autoimmune attack because it is not protected by a blood-nerve barrier, in contrast to more proximal parts of the axon. GM1 is also present at nodes of Ranvier throughout the length of the nerve. Our results certainly do not exclude anti-GM1 binding and complement-mediated neuropathogenic effects at the more proximal parts of the axon, as has been observed in a rabbit neuropathy model (Susuki et al., 2003).

The complement-mediated pathogenic effects of anti-GM1 mAbs at the NMJ shown here are identical to those observed by us in earlier (*ex vivo* and *in vivo*) mouse studies on mAbs against GQ1b ganglioside and on anti-GQ1b positive patient sera (Plomp et

al., 1999; Halstead et al., 2004; O'Hanlon et al., 2001). Furthermore, anti-GD1a antibodies produce these effects, provided the density of GD1a is high, as in neuronal membranes of GD3s-KO mice (Goodfellow et al., 2005). Thus, this neuropathophysiological effect at murine NMJs is a common feature of certain classes of anti-GM1, -GD1b, -GQ1b and -GD1a antibodies. The heterogeneous distribution of clinical effects observed in patients with these anti-ganglioside antibody specificities does not correlate with the common paralytic effect of these antibodies at mouse NMJs. It is thus likely that regional differences in ganglioside expression, co-expression and density in different neural and glial membranes from different anatomical regions in different species are critical determining factors.

The differences in binding and pathogenic properties of individual anti-glycolipid antibodies may in part be attributable to differences in their affinity or avidity, in addition to their precise specificity. If the affinity of an antibody is high enough to compete effectively for a critical oligosaccharide domain that may otherwise be masked by complex formation with a *cis*-interacting glycolipid, then the pathophysiological consequence of any complex formation might be annulled. Whether this range of affinities can be achieved for anti-glycolipid antibodies is unknown. However, in the case of the widely used high-affinity GM1 ligand CTB, the affinity for GM1 is reported as being in the  $10^{-10}$  to  $10^{-12}$  M range (Kuziemko et al., 1996), which is  $10^4$ - to  $10^6$ -fold higher than the mAb DG1, used in the current study. Structural studies have demonstrated that CTB binds GM1 through a 2-fingered pinch interaction involving the nonreducing terminal galactose epitope of GM1 and the internal sialic acid, the latter being masked in our complex formation model (Kuziemko et al., 1996). DG1 also requires the presence of this internal sialic acid for binding to GM1. The implication is therefore that the very high affinity of CTB for GM1 is sufficient to disrupt an inhibitory complex that disguises the sialic acid from its binding domain on the toxin. In contrast, the affinity of DG1 for GM1 is substantially lower than CTB which most likely accounts for DG1's inability to compete effectively for the cryptic sialic acid. Thus affinity may be a relevant factor in disrupting glycolipid-glycolipid complex interactions to allow ligand binding in certain situations. The affinities of DG2 and DG1 for GM1 and GA1 were within fivefold of each other and the half maximal binding values for GM1 were very similar, yet they were completely polarized in terms of their pathogenic potential. It is therefore improbable that a simple explanation of affinity or avidity differences, regardless of dependence upon any complex formation, is the critical factor influencing binding in the current setting, but more likely that the pathogenic differences are due to differences in the fine specificity of the 2 antibodies, specifically their dependence upon the internal sialic acid for binding. Other experimental evidence discussed below also strongly supports this view.

Our study underscores the complexities surrounding the topographical organization of glycolipids in living membranes and how these relate to topography in other types of detection systems such as are used in laboratory immunoassays. Considerable effort is currently focused on achieving optimal platforms for assessing protein-glycan interactions and the issues surrounding complex formation need to be considered in this context (Paulson et al., 2006). The cryptic behaviour of glycolipid headgroups in the plasma membrane, as assessed by accessibility to antibodies is a long recognized phenomenon (Hakomori et al., 1968; Lloyd et al., 1992), but its structural basis has never been demonstrated experimentally. This behaviour may in part be dependent upon ganglioside organisation in lipid rafts. In this study, raft disruption with a cholesterol sequestering agent did not allow the DG1 antibody to bind GM1,

indicating that the interaction of GM1 and GD1a is not cholesterol-dependent, as we also observed by solid phase immunoassay. Rather, it appears to be an intrinsic property of the interacting ganglioside species, possibly localized in a glycosphingolipid signalling sub-domain (Iwabuchi et al., 1998). The ganglioside clustering property may be due to the glycosphingolipid hydroxyl groups acting as hydrogen bond donors, allowing side-by-side interactions within a microdomain (Hakomori, 2002). However, gangliosides require substantial surface area to host their oligosaccharide chain and its associated hydration water, inferring that complex headgroups such as GM1 and GD1a will have large space requirements in the membrane (Sonnino et al., 2007), the consequence of which is unknown in terms of influencing their *cis*-interaction.

Our observations in the GD3s-KO mouse implicate GD1a as the ganglioside that masks GM1 because in this artificial context these are the two predominant gangliosides. Our data on WT mice and in solid-phase assays also strongly suggest that other gangliosides (but not necessarily all gangliosides) may be capable of mediating this masking effect. In the GD3s-KO mouse, the GM1 concentration is substantially higher than in wild-type mice, thereby adding support to the view that the failure of DG1 to bind sialidase-naive WT or GD3s-KO tissue cannot be simply ascribed to limited availability of GM1 ligand.

The heterogeneity of ganglioside and protein content in lipid rafts means there will be a wide range of raft-associated molecules which could potentially interact with GM1 or inhibit binding of anti-GM1 antibodies through steric hindrance. Our observation that cleavage of GPI anchored proteins resulted in increased DG2 and CTB binding suggests that these proteins have a general masking effect for GM1, although this was not the particular masking effect that prevented the binding of DG1, because that remained at the same low level.

Anti-GM1 mAbs cloned from neuropathy patients were also masked from binding GM1 and therefore unable to bind nerve terminals in *ex vivo* nerve-muscle preparations in these studies. This indicates that the observations with the mouse monoclonal anti-GM1 antibody DG1 is likely to be clinically relevant in man. Further studies are needed to explore the diversity of anti-GM1 antibodies seen in different patient populations to determine the extent of diversity in the context of ganglioside-interaction mediated inhibition. In recent experiments, we have identified anti-GM1 positive neuropathy patient sera that are capable of inducing the neuropathophysiological effects at NMJs as observed here, but only after neuraminidase treatment (J.J. Plomp et al., unpublished results). This further shows that clinically relevant anti-GM1 antisera exists that only bind to unmasked GM1. Such human antibodies may only become clinically important in conditions where the topological organization of gangliosides is destabilized, as for instance during nerve regeneration and the mechanisms by which they may or may not induce immunopathology in normal nerve are currently unknown. In regenerating nerve fibres (following experimental transection injury), it has been shown that growth cones have enhanced activity of a membrane sialidase that acts to reduce complex gangliosides into GM1 that may thereby render GM1 available for complex attenuated antibody binding. Targeted delivery of sialidase to experimental nerve lesions in the rat enhances axon outgrowth, and this may in part be a result of the increased level of GM1 in the regenerating axon (Yang et al., 2006). In addition, the sialidase-mediated decrease of GD1a interrupts the neuronal adhesion of myelin-associated glycoprotein, which normally inhibits regeneration (Vyas et al., 2002), causing disinhibited axonal regeneration. Based on our observations with neuraminidase, it is possible that



unmasking of cryptic GM1 could allow previously harmless DG1-like antibodies to bind to GM1 rich growth cones and therefore become pathogenic through their ability to impair regeneration. There is experimental evidence, at least for anti-GD1a antibodies, that anti-ganglioside antibodies may influence neuronal regeneration (Lehmann et al., 2007). The existence of anti-GM1 antibodies like DG1 that specifically bind regenerating neural tissue and cause complement-mediated injury would have significant clinical consequences in that it could be the start of recurrent cycles of axonal injury and regeneration. Therefore, rapid and persistent clearance of such antibodies should be an important therapeutic goal.

It is possible that GM1 masking occurs by interaction with gangliosides other than GD1a. Indeed, in our control experiments the Gal( $\beta$ 1-3)GalNAc moiety of GD1b was apparently shielded from binding DG2, whereas MOG1 - which presumably binds at least in part the internal disialosyl structure of GD1b as it is GD1b monospecific - was able to bind GD1b. Studies showing neuropathic effects restricted to sensory ataxia due to mono-specific anti-GD1b antibodies in a rabbit model are consistent with this finding (Kusunoki et al., 1999), as are recent studies using human ataxic GBS sera (Kaida et al., 2008a). We did not explore further the nature of this antibody-binding inhibitory interaction of GD1b with other gangliosides.

With regard to the induction of anti-GM1 antibodies by *C. jejuni* infection in GBS, our findings illuminate the protective role of antibody in microbial defence, as balanced against the development of autoimmune injury. Thus, DG1 is able to bind the GM1 mimic in LOS on live *C. jejuni* that bears also a GD1a mimic in a 50:50 ratio, indicating that a masking interaction does not occur. This shows that DG1-like antibodies could be bactericidal without binding to neuronal GM1, and thus (at least initially) would not exert an autoimmune neuropathic effect. Furthermore, DG1-like-producing B-cell clones do not recognize self-GM1 in living membranes. Therefore, breaking tolerance to GM1 might not be required in order to generate protective immunity against self-mimicking microbial molecules.

We were able to reproduce a masking effect of GD1a and other gangliosides on GM1 in immunoassays conducted either on polystyrene plates or PVDF membranes, recapitulating the discordant findings observed in living membranes between DG1 and DG2. Interestingly, the inhibitory complexes between GM1 and GD1a form on both polystyrene and PVDF immobilization platforms and thus the molecular interactions between GM1 and other gangliosides must spontaneously form when they are admixed, in addition to being organized or disorganized in living or dead membranes, respectively. Furthermore, these solid phase immunoassays demonstrated that interaction is not unique to GM1 and GD1a but can also occur between GM1 and other gangliosides. In addition, they form important support for the conclusion drawn from our neuraminidase experiments in live tissues - that the enabling factor for DG1 binding to the neuronal membrane under these conditions is an unmasking of pre-existing GM1 by the removal of the external neuraminic acid of GD1a, as opposed to simply an increase in GM1 density by creation of de novo GM1 from GD1a.

Based on these experimental findings, we conclude that the micro-heterogeneity of ganglioside binding by anti-GM1 antibodies substantially influences their neuropathogenic potential, and that the basis for these differences is likely due to subtle conformational arrangements within the membrane, leading to complexes of GM1 with other gangliosides that inhibit binding of some types of anti-GM1 antibodies. These ganglioside interactions appear to be stably formed in living membranes, but are highly susceptible to disintegration (and thus epitope exposure) in frozen or otherwise altered tissue. The changes in the molecular architecture of the

living and dead/disrupted membrane that underlie these properties are unknown and difficult to investigate at the molecular biophysical level. In solid-phase immunoassays, these gangliosides interactions are also able to form. Recent studies showed that complexes of two different gangliosides can form neo-epitopes that specifically bind certain neuropathy-associated antibodies which do not bind to the single ganglioside species (Willison, 2005; Kaida et al., 2004b; Kaida et al., 2006). Together with the data provided here this leads to the conclusion that ganglioside interaction may either enable or inhibit antibody binding to the neuronal membrane, or be neutral, depending of the type of circulating antibody. To describe these 3 conditions, we propose the terms *complex-enhanced*, *complex-attenuated*, and *complex-independent*. In the light of these findings, we need to critically evaluate any individual antibody-ganglioside interaction detected by the wide range of techniques available in order to be able to interpret the neuropathogenic potency of such an interaction in the living neuronal membrane.

**Supplemental material is available online with this article;  
doi:10.1172/JCI37338DS1**

