



Universiteit
Leiden
The Netherlands

Gangliosides and anti-ganglioside antibodies in neuromuscular synaptic function

Zitman, F.M.P.

Citation

Zitman, F. M. P. (2010, January 20). *Gangliosides and anti-ganglioside antibodies in neuromuscular synaptic function*. Retrieved from <https://hdl.handle.net/1887/14568>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14568>

Note: To cite this publication please use the final published version (if applicable).

List of References

-
- Akyildiz B, Gumus H, Kumandas S, Coskun A, Baykan A, Yikilmaz A, Kara I, Okur A. Guillain-Barré syndrome associated with Legionella infection. *J Trop Pediatr* 2008; (54): 275-277.
- Albert L J, Inman R D. Molecular mimicry and autoimmunity. *N Engl J Med* 1999; (341): 2068-2074.
- Allende M L, Proia R L. Lubricating cell signaling pathways with gangliosides. *Curr Opin Struct Biol* 2002; (12): 587-592.
- Ando S. Gangliosides in the nervous system. *Neurochem Int* 1983; (5): 507-537.
- Ando S, Tanaka Y, Waki H, Kon K, Iwamoto M, Fukui F. Gangliosides and sialylcholesterol as modulators of synaptic functions. *Ann N Y Acad Sci* 1998; (845): 232-239.
- Ando S, Tanaka Y, Kobayashi S, Fukui F, Iwamoto M, Waki H, Tai T, Hirabayashi Y. Synaptic function of cholinergic-specific Chol-1alpha ganglioside. *Neurochem Res* 2004; (29): 857-867.
- Ang C W, Endtz H P, Jacobs B C, Laman J D, de Klerk M A, van der Meche F G, van Doorn P A. *Campylobacter jejuni* lipopolysaccharides from Guillain-Barré syndrome patients induce IgG anti-GM1 antibodies in rabbits. *J Neuroimmunol* 2000; (104): 133-138.
- Ang C W, Laman J D, Willison H J, Wagner E R, Endtz H P, de Klerk M A, Tio-Gillen A P, Van den B N, Jacobs B C, van Doorn P A. Structure of *Campylobacter jejuni* lipopolysaccharides determines antiganglioside specificity and clinical features of Guillain-Barré and Miller Fisher patients. *Infect Immun* 2002; (70): 1202-1208.
- Ang C W, Jacobs B C, Laman J D. The Guillain-Barré syndrome: a true case of molecular mimicry. *Trends Immunol* 2004; (25): 61-66.
- Arasaki K, Kusunoki S, Kudo N, Kanazawa I. Acute conduction block in vitro following exposure to antiganglioside sera. *Muscle Nerve* 1993; (16): 587-593.
- Arimura K, Watanabe O, Kitajima I, Suehara M, Minato S, Sonoda Y, Higuchi I, Takenaga S, Maruyama I, Osame M. Antibodies to potassium channels of PC12 in serum of Isaacs' syndrome: Western blot and immunohistochemical studies. *Muscle Nerve* 1997; (20): 299-305.
- Asano A, Selvaraj V, Buttke D E, Nelson J L, Green K M, Evans J E, Travis A J. Biochemical characterization of membrane fractions in murine sperm: identification of three distinct sub-types of membrane rafts. *J Cell Physiol* 2009; (218): 537-548.
- Aspinall G O, Fujimoto S, McDonald A G, Pang H, Kurjanczyk L A, Penner J L. Lipopolysaccharides from *Campylobacter-jejuni* associated with Guillain-Barré syndrome patients mimic human gangliosides in structure. *Infect Immun* 1994; (62): 2122-2125.
- Aydin M, Cengiz S, Agachan B, Yilmaz H, Isbir T. Age-related changes in GM1, GD1a, GT1b components of gangliosides in Wistar albino rats. *Cell Biochem Funct* 2000; (18): 41-45.
- Bach J F. Infections and autoimmune diseases. *J Autoimmun* 2005; (25 Suppl): 74-80.
- Badalamente M A, Stracher A. Delay of muscle degeneration and necrosis in mdx mice by calpain inhibition. *Muscle Nerve* 2000; (23): 106-111.
- Baldwin M R, Kim J J, Barbieri J T. Botulinum neurotoxin B-host receptor recognition: it takes two receptors to tango. *Nat Struct Mol Biol* 2007; (14): 9-10.
- Barrantes F J. The lipid environment of the nicotinic acetylcholine receptor in native and reconstituted membranes. *Crit Rev Biochem Mol Biol* 1989; (24): 437-478.
- Barrier L, Ingrand S, Damjanac M, Bilan A R, Hugon J, Page G. Genotype-related changes of ganglioside composition in brain regions of transgenic mouse models of Alzheimer's disease. *Neurobiol Aging* 2007; (28): 1863-1872.
- Benatar M G, Willison H J, Vincent A. Lack of effect of Miller Fisher sera/plasmas on transmitter release from PC12 cells. *J Neuroimmunol* 1997; (80): 1-5.
- Bergelson L D. Serum gangliosides as endogenous immunomodulators. *Immunol Today* 1995; (16): 483-486.
- Boffey J, Odaka M, Nicoll D, Wagner E R, Townson K, Bowes T, Conner J, Furukawa K, Willison H J. Characterisation of the immunoglobulin variable region gene usage encoding the murine anti-ganglioside antibody repertoire. *J Neuroimmunol* 2005; (165): 92-103.
- Bowes T, Wagner E R, Boffey J, Nicholl D, Cochrane L, Benboubetra M, Conner J, Furukawa K, Furukawa K, Willison H J. Tolerance to self gangliosides is the major factor restricting the antibody response to lipopolysaccharide core oligosaccharides in *Campylobacter jejuni* strains associated with Guillain-Barre syndrome. *Infect Immun* 2002; (70): 5008-5018.
- Breathnach C S. Johann Ludwig Wilhelm Thudichum 1829-1901, bane of the Protagonisers. *Hist Psychiatry* 2001; (12): 283-296.
- Breckenridge W C, Gombos G, Morgan I G. The lipid composition of adult rat brain synaptosomal plasma membranes. *Biochim Biophys Acta* 1972; (266): 695-707.
- Brodsky R A. How I treat paroxysmal nocturnal hemoglobinuria. *Blood* 2009.
- Brody A J, Sternbach G, Varon J. Octave Landry: Guillain-Barré syndrome. *J Emerg Med* 1994; (12): 833-837.
- Brunetti-Pierri N, Scaglia F. GM1 gangliosidosis: review of clinical, molecular, and therapeutic aspects. *Mol Genet Metab* 2008; (94): 391-396.
- Brunger A T, Rummel A. Receptor and substrate interactions of clostridial neurotoxins. *Toxicon* 2009.
- Buccoliero R, Futerman A H. The roles of ceramide and complex sphingolipids in neuronal cell function. *Pharmacol Res* 2003; (47): 409-419.

- Buccoliero R, Bodenne J, Echten-Deckert G, Sandhoff K, Futerman A H. Phospholipid synthesis is decreased in neuronal tissue in a mouse model of Sandhoff disease. *J Neurochem* 2004; (90): 80-88.
- Buchwald B, Weishaupt A, Toyka K V, Dudel J. Immunoglobulin G from a patient with Miller-Fisher syndrome rapidly and reversibly depresses evoked quantal release at the neuromuscular junction of mice. *Neurosci Lett* 1995; (201): 163-166.
- Buchwald B, Bufler J, Carpo M, Heidenreich F, Pitz R, Dudel J, Nobile-Orazio E, Toyka K V. Combined pre- and postsynaptic action of IgG antibodies in Miller Fisher syndrome. *Neurology* 2001; (56): 67-74.
- Buchwald B, Ahangari R, Toyka K V. Differential blocking effects of the monoclonal anti-GQ1b IgM antibody and alpha-latrotoxin in the absence of complement at the mouse neuromuscular junction. *Neurosci Lett* 2002; (334): 25-28.
- Buchwald B, Zhang G, Vogt-Eisele A K, Zhang W, Ahangari R, Griffin J W, Hatt H, Toyka K V, Sheikh K A. Anti-ganglioside antibodies alter presynaptic release and calcium influx. *Neurobiol Dis* 2007; (28): 113-121.
- Bullens R W, O'Hanlon G M, Goodyear C S, Molenaar P C, Conner J, Willison H J, Plomp J J. Anti-GQ1b antibodies and evoked acetylcholine release at mouse motor endplates. *Muscle Nerve* 2000; (23): 1035-1043.
- Bullens R W, O'Hanlon G M, Wagner E, Molenaar P C, Furukawa K, Furukawa K, Plomp J J, Willison H J. Complex gangliosides at the neuromuscular junction are membrane receptors for autoantibodies and botulinum neurotoxin but redundant for normal synaptic function. *J Neurosci* 2002; (22): 6876-6884.
- Cantu' L, Corti M, Brocca P, Del Favero E. Structural aspects of ganglioside-containing membranes. *Biochim Biophys Acta* 2009; (1788): 202-208.
- Cao Z, Qiu J, Domeniconi M, Hou J, Bryson J B, Mellado W, Filbin M T. The inhibition site on myelin-associated glycoprotein is within Ig-domain 5 and is distinct from the sialic acid binding site. *J Neurosci* 2007; (27): 9146-9154.
- Caporale C M, Papola F, Fioroni M A, Aureli A, Giovannini A, Notturmo F, Adorno D, Caporale V, Uncini A. Susceptibility to Guillain-Barre syndrome is associated to polymorphisms of CD1 genes. *J Neuroimmunol* 2006; (177): 112-118.
- Chamberlain L H, Burgoyne R D, Gould G W. SNARE proteins are highly enriched in lipid rafts in PC12 cells: implications for the spatial control of exocytosis. *Proc Natl Acad Sci U S A* 2001; (98): 5619-5624.
- Chen F, Qian L, Yang Z H, Huang Y, Ngo S T, Ruan N J, Wang J, Schneider C, Noakes P G, Ding Y Q, Mei L, Luo Z G. Rapsyn interaction with calpain stabilizes AChR clusters at the neuromuscular junction. *Neuron* 2007b; (55): 247-260.
- Chen Y W, Pedersen J W, Wandall H H, Levery S B, Pizette S, Clausen H, Cohen S M. Glycosphingolipids with extended sugar chain have specialized functions in development and behavior of *Drosophila*. *Dev Biol* 2007a; (306): 736-749.
- Cheng Q, Wang D S, Jiang G X, Han H, Zhang Y, Wang W Z, Fredrikson S. Distinct pattern of age-specific incidence of Guillain-Barré syndrome in Harbin, China. *J Neurol* 2002; (249): 25-32.
- Cheng Q, Wang D S, Jiang G X, Han H, Zhang Y, Wang W Z, Fredrikson S. A case-control study of Guillain-Barre syndrome in Harbin, China. *Eur J Neurol* 2006; (13): 953-957.
- Chiavegatto S, Sun J, Nelson R J, Schnaar R L. A functional role for complex gangliosides: motor deficits in GM2/GD2 synthase knockout mice. *Exp Neurol* 2000; (166): 227-234.
- Chiba A, Kusunoki S, Shimizu T, Kanazawa I. Serum IgG antibody to ganglioside GQ1b is a possible marker of Miller Fisher syndrome. *Ann Neurol* 1992; (31): 677-679.
- Chiba A, Kusunoki S, Obata H, Machinami R, Kanazawa I. Serum anti-GQ1b IgG antibody is associated with ophthalmoplegia in Miller Fisher syndrome and Guillain-Barre syndrome: clinical and immunohistochemical studies. *Neurology* 1993; (43): 1911-1917.
- Chiba A, Kusunoki S, Obata H, Machinami R, Kanazawa I. Ganglioside composition of the human cranial nerves, with special reference to pathophysiology of Miller Fisher syndrome. *Brain Research* 1997; (745): 32-36.
- Choi H J, Chung T W, Kim S J, Cho S Y, Lee Y S, Lee Y C, Ko J H, Kim C H. The AP-2alpha transcription factor is required for the ganglioside GM3-stimulated transcriptional regulation of a PTEN gene. *Glycobiology* 2008; (18): 395-407.
- Cosi V, Versino M. Guillain-Barré syndrome. *Neurol Sci* 2006; (27 Suppl 1): S47-S51.
- Court FA, Gillingwater T H, Melrose S, Sherman D L, Greenshields K N, Morton A J, Harris J B, Willison H J, Ribchester R R. Identity, developmental restriction and reactivity of extralaminar cells capping mammalian neuromuscular junctions. *J Cell Sci* 2008; (121): 3901-3911.
- Crespo P M, Zurita A R, Giraudo C G, Maccioni H J, Daniotti J L. Ganglioside glycosyltransferases and newly synthesized gangliosides are excluded from detergent-insoluble complexes of Golgi membranes. *Biochem J* 2004; (377): 561-568.
- Crocker P R, Paulson J C, Varki A. Siglecs and their roles in the immune system. *Nat Rev Immunol* 2007; (7): 255-266.
- Csurhes P A, Sullivan A A, Green K, Greer J M, Pender M P, McCombe P A. Increased circulating T cell reactivity to GM1 ganglioside in patients with Guillain-Barré syndrome. *J Clin Neurosci* 2005; (12): 409-415.
- d'Azzo A, Tessitore A, Sano R. Gangliosides as apoptotic signals in ER stress response. *Cell Death Differ* 2006; (13): 404-414.
- De Maria R, Lenti L, Malisan F, d'Agostino F, Tomassini B, Zeuner A, Rippon M R, Testi R. Requirement for GD3 ganglioside in. *Science* 1997; (277): 1652-1655.

- Debuch H, Ernst Klenk 1896-1971. *J Neurochem* 1973; (21): 725-727.
- Desplats P A, Denny C A, Kass K E, Gilmartin T, Head S R, Sutcliffe J G, Seyfried T N, Thomas E A. Glycolipid and ganglioside metabolism imbalances in Huntington's disease. *Neurobiol Dis* 2007; (27): 265-277.
- Dobrowsky R T, Carter B D. Coupling of the p75 neurotrophin receptor to sphingolipid signaling. *Ann N Y Acad Sci* 1998; (845): 32-45.
- Dolly J O, Aoki K R. The structure and mode of action of different botulinum toxins. *Eur J Neurol* 2006; (13 Suppl 4): 1-9.
- Donev R M, Sivasankar B, Mizuno M, Morgan B P. The mouse complement regulator CD59b is significantly expressed only in testis and plays roles in sperm acrosome activation and motility. *Mol Immunol* 2008; (45): 534-542.
- Duan J, Zhang J, Zhao Y, Yang F, Zhang X. Ganglioside GM2 modulates the erythrocyte Ca²⁺-ATPase through its binding to the calmodulin-binding domain and its 'receptor'. *Arch Biochem Biophys* 2006; (454): 155-159.
- Durand M C, Porcher R, Orlikowski D, Aboab J, Devaux C, Clair B, Annane D, Gaillard J L, Lofaso F, Raphael J C, Sharshar T. Clinical and electrophysiological predictors of respiratory failure in Guillain-Barré syndrome: a prospective study. *Lancet Neurol* 2006; (5): 1021-1028.
- Ebanks R O, Isenman D E. Mouse complement component C4 is devoid of classical pathway C5 convertase subunit activity. *Mol Immunol* 1996; (33): 297-309.
- Egorushkina N V, Ratushnyak A S, Egorushkin I V. The influence of exogenous gangliosides on the dynamics of the development of prolonged posttetanic potentiation. *Neurosci Behav Physiol* 1993; (23): 435-438.
- Fearon D T. Regulation by membrane sialic acid of beta1H-dependent decay-dissociation of amplification C3 convertase of the alternative complement pathway. *Proc Natl Acad Sci U S A* 1978; (75): 1971-1975.
- Feasby T E, Gilbert J J, Brown W F, Bolton C F, Hahn A F, Koopman W F, Zochodne D W. An acute axonal form of Guillain-Barré polyneuropathy. *Brain* 1986; (109 (Pt 6)): 1115-1126.
- Feng G, Mellor R H, Bernstein M, Keller-Peck C, Nguyen Q T, Wallace M, Nerbonne J M, Lichtman J W, Sanes J R. Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of GFP. *Neuron* 2000; (28): 41-51.
- Fiekers J F. Concentration-dependent effects of neostigmine on the endplate acetylcholine receptor channel complex. *J Neurosci* 1985; (5): 502-514.
- Fisher M. An unusual variant of acute idiopathic polyneuritis (syndrome of ophthalmoplegia, ataxia and areflexia). *N Engl J Med* 1956; (255): 57-65.
- Fishman P H. Role of membrane gangliosides in the binding and action of bacterial toxins. *J Membr Biol* 1982; (69): 85-97.
- Flachenecker P. Epidemiology of neuroimmunological diseases. *J Neurol* 2006; (253 Suppl 5): V2-V8.
- Forsberg A, Ahlstrom G, Holmqvist L W. Falling ill with Guillain-Barré syndrome: patients' experiences during the initial phase. *Scand J Caring Sci* 2008; (22): 220-226.
- Fujita A, Cheng J, Hirakawa M, Furukawa K, Kusunoki S, Fujimoto T. Gangliosides GM1 and GM3 in the living cell membrane form clusters susceptible to cholesterol depletion and chilling. *Mol Biol Cell* 2007; (18): 2112-2122.
- Funakoshi K, Koga M, Takahashi M, Hirata K, Yuki N. *Campylobacter coli* enteritis and Guillain-Barré syndrome: no evidence of molecular mimicry and serological relationship. *J Neurol Sci* 2006; (246): 163-168.
- Furiya Y, Hirano M, Kusunoki S, Ueda M, Sugie K, Nishiwaki T, Ueno S. Complete recovery of an aged patient with Guillain-Barré syndrome associated with multiple IgM anti-ganglioside antibodies. *Muscle Nerve* 2008; (38): 1630-1633.
- Furuse H, Waki H, Kaneko K, Fujii S, Miura M, Sasaki H, Ito K I, Kato H, Ando S. Effect of the mono- and tetrasialogangliosides, GM1 and GQ1b, on long-term potentiation in the CA1 hippocampal neurons of the guinea pig. *Exp Brain Res* 1998; (123): 307-314.
- Futerman A H, Hannun Y A. The complex life of simple sphingolipids. *EMBO Rep* 2004; (5): 777-782.
- Garssen M P, Schillings M L, van Doorn P A, Van Engelen B G, Zwartz M J. Contribution of central and peripheral factors to residual fatigue in Guillain-Barre syndrome. *Muscle Nerve* 2007; (36): 93-99.
- Geleijns K, Brouwer B A, Jacobs B C, Houwing-Duistermaat J J, van Duijn C M, van Doorn P A. The occurrence of Guillain-Barré syndrome within families. *Neurology* 2004; (63): 1747-1750.
- Geleijns K, Emonts M, Laman J D, van Rijs W, van Doorn P A, Hermans P W, Jacobs B C. Genetic polymorphisms of macrophage-mediators in Guillain-Barré syndrome. *J Neuroimmunol* 2007; (190): 127-130.
- Gilhus N E. Autoimmune myasthenia gravis. *Expert Rev Neurother* 2009; (9): 351-358.
- Gillard B K, Thomas J W, Nell L J, Marcus D M. Antibodies against ganglioside GT3 in the sera of patients with type I diabetes mellitus. *J Immunol* 1989; (142): 3826-3832.
- Ginzburg L, Li S C, Li Y T, Futerman A H. An exposed carboxyl group on sialic acid is essential for gangliosides to inhibit calcium uptake via the sarco/endoplasmic reticulum Ca²⁺-ATPase: relevance to gangliosidoses. *J Neurochem* 2008; (104): 140-146.
- Giraud C G, Maccioni H J. Ganglioside glycosyltransferases organize in distinct multienzyme complexes in CHO-K1 cells. *J Biol Chem* 2003; (278): 40262-40271.
- Godschalk P C, Heikema A P, Gilbert M, Komagamine T, Ang C W, Glerum J, Brochu D, Li J, Yuki N, Jacobs B C, van Belkum A, Endtz H P. The crucial role of *Campylobacter jejuni* genes in anti-ganglioside antibody induction in Guillain-Barre syndrome. *J Clin Invest* 2004; (114): 1659-1665.

- Godschalk P C, Gilbert M, Jacobs B C, Kramers T, Tio-Gillen A P, Ang C W, Van den B N, Li J, Verbrugh H A, van Belkum A, Endtz H P. Co-infection with two different *Campylobacter jejuni* strains in a patient with the Guillain-Barre syndrome. *Microbes Infect* 2006; (8): 248-253.
- Godschalk P C R, Kuijff M L, Li J J, St Michael F, Ang C W, Jacobs B C, Karwaski M F, Brochu D, Moteras A, Endtz H P, van Belkum A, Gilbert M. Structural characterization of *Campylobacter jejuni* lipooligosaccharide outer cores associated with Guillain-Barré and Miller Fisher syndromes. *Infection and Immunity* 2007; (75): 1245-1254.
- Gong Y, Tagawa Y, Lunn M P, Laroy W, Heffer-Lauc M, Li C Y, Griffin J W, Schnaar R L, Sheikh K A. Localization of major gangliosides in the PNS: implications for immune neuropathies. *Brain* 2002; (125): 2491-2506.
- Goodfellow J A, Bowes T, Sheikh K, Odaka M, Halstead S K, Humphreys P D, Wagner E R, Yuki N, Furukawa K, Furukawa K, Plomp J J, Willison H J. Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy. *J Neurosci* 2005; (25): 1620-1628.
- Goodyear C S, O'Hanlon G M, Plomp J J, Wagner E R, Morrison I, Veitch J, Cochrane L, Bullens R W, Molenaar P C, Conner J, Willison H J. Monoclonal antibodies raised against Guillain-Barre syndrome-associated *Campylobacter jejuni* lipopolysaccharides react with neuronal gangliosides and paralyze muscle-nerve preparations. *J Clin Invest* 1999; (104): 697-708.
- Green W N, Andersen O S. Surface charges and ion channel function. *Annu Rev Physiol* 1991; (53): 341-359.
- Greenshields K N, Halstead S K, Zitman F M, Rinaldi S, Brennan K M, O'Leary C, Chamberlain L H, Easton A, Roxburgh J, Pediani J, Furukawa K, Furukawa K, Goodyear C S, Plomp J J, Willison H J. The neuropathic potential of anti-GM1 autoantibodies is regulated by the local glycolipid environment in mice. *J Clin Invest* 2009; (119): 595-610.
- Groshong J S, Spencer M J, Bhattacharyya B J, Kudryashova E, Vohra B P, Zayas R, Wollmann R L, Miller R J, Gomez C M. Calpain activation impairs neuromuscular transmission in a mouse model of the slow-channel myasthenic syndrome. *J Clin Invest* 2007; (117): 2903-2912.
- Guillain G, Barré J A, Strohl A. Sur un syndrome de radiculo-nevrite avec hyperalbuminose du liquide cephalo-rachidien sans reaction cellulaire: remarques sur les caracteres cliniques et graphiques des reflexes tendineux. *Bull Soc Med Hop Paris* 1916; (40): 1462-1470.
- Haber P, DeStefano F, Angulo F J, Iskander J, Shadomy S V, Weintraub E, Chen R T. Guillain-Barré syndrome following influenza vaccination. *JAMA* 2004; (292): 2478-2481.
- Hafer-Macko C, Hsieh S T, Li C Y, Ho T W, Sheikh K, Cornblath D R, McKhann G M, Asbury A K, Griffin J W. Acute motor axonal neuropathy: an antibody-mediated attack on axolemma. *Ann Neurol* 1996b; (40): 635-644.
- Hafer-Macko C E, Sheikh K A, Li C Y, Ho T W, Cornblath D R, McKhann G M, Asbury A K, Griffin J W. Immune attack on the Schwann cell surface in acute inflammatory demyelinating polyneuropathy. *Ann Neurol* 1996a; (39): 625-635.
- Hakomori S. Glycosylation defining cancer malignancy: new wine in an old bottle. *Proc Natl Acad Sci U S A* 2002; (99): 10231-10233.
- Hakomori S. Structure, organization, and function of glycosphingolipids in membrane. *Curr Opin Hematol* 2003; (10): 16-24.
- Hakomori S I, Teather C, Andrews H. Organizational difference of cell surface "hematoside" in normal and virally transformed cells. *Biochem Biophys Res Commun* 1968; (33): 563-568.
- Hakomori S I. Inaugural Article: The glycosynapse. *Proc Natl Acad Sci U S A* 2002; (99): 225-232.
- Hakomori S I. Structure and function of glycosphingolipids and sphingolipids: Recollections and future trends. *Biochim Biophys Acta* 2007.
- Halpern J L, Loftus A. Characterization of the receptor-binding domain of tetanus toxin. *J Biol Chem* 1993; (268): 11188-11192.
- Halstead S K, O'Hanlon G M, Humphreys P D, Morrison D B, Morgan B P, Todd A J, Plomp J J, Willison H J. Anti-disialoside antibodies kill perisynaptic Schwann cells and damage motor nerve terminals via membrane attack complex in a murine model of neuropathy. *Brain* 2004; (127): 2109-2123.
- Halstead S K, Humphreys P D, Goodfellow J A, Wagner E R, Smith R A, Willison H J. Complement inhibition abrogates nerve terminal injury in Miller Fisher syndrome. *Ann Neurol* 2005a; (58): 203-210.
- Halstead S K, Humphreys P D, Zitman F M, Hamer J, Plomp J J, Willison H J. C5 inhibitor rEV576 protects against neural injury in an in vitro mouse model of Miller Fisher syndrome. *J periph nerv syst* 2008b; (13): 228-235.
- Halstead S K, Morrison I, O'Hanlon G M, Humphreys P D, Goodfellow J A, Plomp J J, Willison H J. Anti-disialosyl antibodies mediate selective neuronal or Schwann cell injury at mouse neuromuscular junctions. *Glia* 2005b; (52): 177-189.
- Halstead S K, Zitman F M, Humphreys P D, Greenshields K, Verschuuren J J, Jacobs B C, Rother R P, Plomp J J, Willison H J. Eculizumab prevents anti-ganglioside antibody-mediated neuropathy in a murine model. *Brain* 2008a; (131): 1197-1208.
- Handa Y, Ozaki N, Honda T, Furukawa K, Tomitia Y, Inoue M, Furukawa K, Okada M, Sugiura Y. GD3 synthase gene knockout mice exhibit thermal hyperalgesia and mechanical allodynia but decreased response to formalin-induced prolonged noxious stimulation. *Pain* 2005; (117): 271-279.

-
- Hansson H A, Holmgren J, Svennerholm L. Ultrastructural localization of cell membrane GM1 ganglioside by cholera toxin. *Proc Natl Acad Sci U S A* 1977; (74): 3782-3786.
- Harris J B, Ribchester R R. The relationship between end-plate size and transmitter release in normal and dystrophic muscles of the mouse. *J Physiol* 1979; (296): 245-265.
- Hartung H P, Schwenke C, Bitter-Suermann D, Toyka K V. Guillain-Barré syndrome: activated complement components C3a and C5a in CSF. *Neurology* 1987; (37): 1006-1009.
- Hashiramoto A, Mizukami H, Yamashita T. Ganglioside GM3 promotes cell migration by regulating MAPK and c-Fos/AP-1. *Oncogene* 2006; (25): 3948-3955.
- Havelaar A H. Campylobacterios in Nederland. Risico's en interventiemogelijkheden. National institute for the public health and the environment, Bilthoven, The Netherlands 2002; (Report number: 250911001).
- Hayashi K, Muhleisen M, Probst W, Rahmann H. Binding of Ca²⁺ to phosphoinositols, phosphatidylserines and gangliosides. *Chem Phys Lipids* 1984; (34): 317-322.
- Hayes S. Use of ganglioside affinity filters to identify toxigenic strains of *Clostridium botulinum* types C and D. *Infect Immun* 1979; (26): 150-156.
- Hepburn N J, Williams A S, Nunn M A, Chamberlain-Banoub J C, Hamer J, Morgan B P, Harris C L. In vivo characterization and therapeutic efficacy of a C5-specific inhibitor from the soft tick *Ornithodoros moubata*. *J Biol Chem* 2007; (282): 8292-8299.
- Hettmer S, Ladisch S, Kaucic K. Low complex ganglioside expression characterizes human neuroblastoma cell lines. *Cancer Lett* 2005; (225): 141-149.
- Hill A, Richards S J, Hillmen P. Recent developments in the understanding and management of paroxysmal nocturnal haemoglobinuria. *Br J Haematol* 2007; (137): 181-192.
- Hillmen P, Hall C, Marsh J C, Elebute M, Bombara M P, Petro B E, Cullen M J, Richards S J, Rollins S A, Mojcik C F, Rother R P. Effect of eculizumab on hemolysis and transfusion requirements in patients with paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2004; (350): 552-559.
- Hillmen P, Young N S, Schubert J, Brodsky R A, Socie G, Muus P, Roth A, Szer J, Elebute M O, Nakamura R, Browne P, Risitano A M, Hill A, Schrezenmeier H, Fu C L, Maciejewski J, Rollins S A, Mojcik C F, Rother R P, Luzzatto L. The complement inhibitor eculizumab in paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2006; (355): 1233-1243.
- Ho T W, Hsieh S T, Nachamkin I, Willison H J, Sheikh K, Kiehlauch J, Flanigan K, McArthur J C, Cornblath D R, McKhann G M, Griffin J W. Motor nerve terminal degeneration provides a potential mechanism for rapid recovery in acute motor axonal neuropathy after *Campylobacter* infection. *Neurology* 1997; (48): 717-724.
- Houliston R S, Jacobs B C, Tio-Gillen A P, Verschuuren J J, Khieu N H, Gilbert M, Jarrell H C. STD-NMR used to elucidate the fine binding specificity of pathogenic anti-ganglioside antibodies directly in patient serum. *Biochemistry* 2009; (48): 220-222.
- Hu Z T, Zhao P, Liu J, Wu Z X, Xu T. Alpha-latrotoxin triggers extracellular Ca²⁺-dependent exocytosis and sensitizes fusion machinery in endocrine cells. *Acta Biochim Biophys Sin (Shanghai)* 2006; (38): 8-14.
- Hua S Y, Teylan M A, Cimenser A. An antibody to synaptotagmin I facilitates synaptic transmission. *Eur J Neurosci* 2007; (25): 3217-3225.
- Hughes R A, Cornblath D R. Guillain-Barré syndrome. *Lancet* 2005; (366): 1653-1666.
- Hughes R A, Raphael J C, Swan A V, van Doorn P A. Intravenous immunoglobulin for Guillain-Barré syndrome. *Cochrane Database Syst Rev* 2006;CD002063.
- Hughes R A, Swan A V, Raphael J C, Annane D, van Koningsveld R, van Doorn P A. Immunotherapy for Guillain-Barre syndrome: a systematic review. *Brain* 2007; (130): 2245-2257.
- Ilyas A A, Chen Z W. Lewis rats immunized with GM1 ganglioside do not develop peripheral neuropathy. *J Neuroimmunol* 2007; (188): 34-38.
- Inoue M, Fujii Y, Furukawa K, Okada M, Okumura K, Hayakawa T, Furukawa K, Sugiura Y. Refractory skin injury in complex knock-out mice expressing only the GM3 ganglioside. *Journal of Biological Chemistry* 2002; (277): 29881-29888.
- Isaev D, Isaeva E, Shatskih T, Zhao Q, Smits N C, Shworak N W, Khazipov R, Holmes G L. Role of extracellular sialic acid in regulation of neuronal and network excitability in the rat hippocampus. *J Neurosci* 2007; (27): 11587-11594.
- Ito M, Kuwabara S, Odaka M, Misawa S, Koga M, Hirata K, Yuki N. Bickerstaff's brainstem encephalitis and Fisher syndrome form a continuous spectrum: clinical analysis of 581 cases. *J Neurol* 2008; (255): 674-682.
- Itoh M, Fukumoto S, Iwamoto T, Mizuno A, Rokutanda A, Ishida H K, Kiso M, Furukawa K. Specificity of carbohydrate structures of gangliosides in the activity to regenerate the rat axotomized hypoglossal nerve. *Glycobiology* 2001; (11): 125-130.
- Iwabuchi K, Handa K, Hakomori S. Separation of "glycosphingolipid signaling domain" from caveolin-containing membrane fraction in mouse melanoma B16 cells and its role in cell adhesion coupled with signaling. *J Biol Chem* 1998; (273): 33766-33773.
- Iwamoto N, Suzuki Y, Makino Y, Haga C, Kosaka K, Iizuka R. Cell membrane changes in brains manifesting senile plaques: an immunohistochemical study of GM1 membranous ganglioside. *Brain Res* 1990; (522): 152-156.
- Izumi T, Ogawa T, Koizumi H, Fukuyama Y. Low levels of CSF gangliotetraose-series gangliosides in West syndrome: implication of brain maturation disturbance. *Pediatr Neurol* 1993; (9): 293-296.

- Jacobs B C, Vandoorn P A, Schmitz P I M, Tio-Gillen A P, Herbrink P, Visser L H, Hooijkaas H, Vandermeché F G A. *Campylobacter jejuni* infections and anti-GM1 antibodies in Guillain-Barré syndrome. *Ann Neurol* 1996; (40): 181-187.
- Jacobs B C, Bullens R W, O'Hanlon G M, Ang C W, Willison H J, Plomp J J. Detection and prevalence of alpha-latrotoxin-like effects of serum from patients with Guillain-Barré syndrome. *Muscle Nerve* 2002; (25): 549-558.
- Jacobs B C, O'Hanlon G M, Bullens R W, Veitch J, Plomp J J, Willison H J. Immunoglobulins inhibit pathophysiological effects of anti-GQ1b-positive sera at motor nerve terminals through inhibition of antibody binding. *Brain* 2003; (126): 2220-2234.
- Jennemann R, Sandhoff R, Wang S, Kiss E, Gretz N, Zuliani C, Martin-Villalba A, Jager R, Schorle H, Kenzelmann M, Bonrouhi M, Wiegandt H, Grone H J. Cell-specific deletion of glucosylceramide synthase in brain leads to severe neural defects after birth. *Proc Natl Acad Sci U S A* 2005; (102): 12459-12464.
- Jung W R, Kim H G, Kim K L. Ganglioside GQ1b improves spatial learning and memory of rats as measured by the Y-maze and the Morris water maze tests. *Neurosci Lett* 2008; (439): 220-225.
- Kaida K, Morita D, Kanzaki M, Kamakura K, Motoyoshi K, Hirakawa M, Kusunoki S. Ganglioside complexes as new target antigens in Guillain-Barré syndrome. *Ann Neurol* 2004b; (56): 567-571.
- Kaida K, Kusunoki S, Kanzaki M, Kamakura K, Motoyoshi K, Kanazawa I. Anti-GQ1b antibody as a factor predictive of mechanical ventilation in Guillain-Barré syndrome. *Neurology* 2004a; (62): 821-824.
- Kaida K, Kanzaki M, Morita D, Kamakura K, Motoyoshi K, Hirakawa M, Kusunoki S. Anti-ganglioside complex antibodies in Miller Fisher syndrome. *J Neurol Neurosurg Psychiatry* 2006; (77): 1043-1046.
- Kaida K, Morita D, Kanzaki M, Kamakura K, Motoyoshi K, Hirakawa M, Kusunoki S. Anti-ganglioside complex antibodies associated with severe disability in GBS. *J Neuroimmunol* 2007; (182): 212-218.
- Kaida K, Sonoo M, Ogawa G, Kamakura K, Ueda-Sada M, Arita M, Motoyoshi K, Kusunoki S. GM1/GalNAc-GD1a complex: a target for pure motor Guillain-Barré syndrome. *Neurology* 2008b; (71): 1683-1690.
- Kaida K, Kamakura K, Ogawa G, Ueda M, Motoyoshi K, Arita M, Kusunoki S. GD1b-specific antibody induces ataxia in Guillain-Barré syndrome. *Neurology* 2008a; (71): 196-201.
- Kaja S, van de Ven R C, Broos L A, Veldman H, van Dijk J G, Verschuuren J J, Frants R R, Ferrari M D, van den Maagdenberg A M, Plomp J J. Gene dosage-dependent transmitter release changes at neuromuscular synapses of CACNA1A R192Q knockin mice are non-progressive and do not lead to morphological changes or muscle weakness. *Neuroscience* 2005; (135): 81-95.
- Kaja S, van de Ven R C, van Dijk J G, Verschuuren J J, Arahata K, Frants R R, Ferrari M D, van den Maagdenberg A M, Plomp J J. Severely impaired neuromuscular synaptic transmission causes muscle weakness in the Cacna1a-mutant mouse rolling Nagoya. *Eur J Neurosci* 2007a; (25): 2009-2020.
- Kaja S, van de Ven R C, Broos L A, Frants R R, Ferrari M D, van den Maagdenberg A M, Plomp J J. Characterization of acetylcholine release and the compensatory contribution of non-Ca_v2.1 channels at motor nerve terminals of leaner Ca_v2.1-mutant mice. *Neuroscience* 2007b; (144): 1278-1287.
- Kaminski D A, Stavnezer J. Antibody class switching differs among SJL, C57BL/6 and 129 mice. *Int Immunol* 2007; (19): 545-556.
- Kaminski H J, Kusner L L, Richmonds C, Medof M E, Lin F. Deficiency of decay accelerating factor and CD59 leads to crisis in experimental myasthenia. *Exp Neurol* 2006; (202): 287-293.
- Kappel T, Anken R H, Hanke W, Rahmann H. Gangliosides affect membrane-channel activities dependent on ambient temperature. *Cell Mol Neurobiol* 2000; (20): 579-590.
- Kato K, Iwamori M, Hirabayashi Y. Increase of GQ1b in the hippocampus of mice following kindled-seizures. *Neurosci Lett* 2008; (441): 286-290.
- Kawai H, Allende M L, Wada R, Kono M, Sango K, Deng C, Miyakawa T, Crawley J N, Werth N, Bierfreund U, Sandhoff K, Proia R L. Mice expressing only monosialoganglioside GM3 exhibit lethal audiogenic seizures. *J Biol Chem* 2001; (276): 6885-6888.
- Kawano M. Complement regulatory proteins and autoimmunity. *Arch Immunol Ther Exp (Warsz)* 2000; (48): 367-372.
- Kazatchkine M D, Fearon D T, Austen K F. Human alternative complement pathway: membrane-associated sialic acid regulates the competition between B and beta1 H for cell-bound C3b. *J Immunol* 1979; (122): 75-81.
- Kieran D, Greensmith L. Inhibition of calpains, by treatment with leupeptin, improves motoneuron survival and muscle function in models of motoneuron degeneration. *Neuroscience* 2004; (125): 427-439.
- Kieseier B C, Hartung H P. Therapeutic strategies in the Guillain-Barré syndrome. *Semin Neurol* 2003; (23): 159-168.
- Kitamura M, Iwamori M, Nagai Y. Interaction between *Clostridium botulinum* neurotoxin and gangliosides. *Biochim Biophys Acta* 1980; (628): 328-335.
- Klenk E. Über die Ganglioside, eine neue Gruppe von zuckerhaltigen Gehirnlipoiden. *Z Physiol Chem* 1942; (273): 76-86.
- Koga M, Takahashi M, Masuda M, Hirata K, Yuki N. *Campylobacter* gene polymorphism as a determinant of clinical features of Guillain-Barré syndrome. *Neurology* 2005; (65): 1376-1381.
- Kolter T, Sandhoff K. Glycosphingolipid degradation and animal models of GM2-gangliosidosis. *J Inherit Metab Dis* 1998; (21): 548-563.

-
- Kolter T, Sandhoff K. Principles of lysosomal membrane digestion: stimulation of sphingolipid degradation by sphingolipid activator proteins and anionic lysosomal lipids. *Annu Rev Cell Dev Biol* 2005; (21): 81-103.
- Kolter T, Sandhoff K. Sphingolipid metabolism diseases. *Biochim Biophys Acta* 2006; (1758): 2057-2079.
- Koski C L, Sanders M E, Swoveland P T, Lawley T J, Shin M L, Frank M M, Joiner K A. Activation of terminal components of complement in patients with Guillain-Barré syndrome and other demyelinating neuropathies. *J Clin Invest* 1987; (80): 1492-1497.
- Kozaki S, Kamata Y, Watarai S, Nishiki T, Mochida S. Ganglioside GT1b as a complementary receptor component for Clostridium botulinum neurotoxins. *Microb Pathog* 1998; (25): 91-99.
- Kuijf M L, Godschalk P C, Gilbert M, Endtz H P, Tio-Gillen A P, Ang C W, van Doorn P A, Jacobs B C. Origin of ganglioside complex antibodies in Guillain-Barré syndrome. *J Neuroimmunol* 2007; (188): 69-73.
- Kuijf M L, Geleijns K, Ennaji N, van Rijs W, van Doorn P A, Jacobs B C. Susceptibility to Guillain-Barré syndrome is not associated with CD1A and CD1E gene polymorphisms. *J Neuroimmunol* 2008; (205): 110-112.
- Kumagai K, Yasuda S, Okemoto K, Nishijima M, Kobayashi S, Hanada K. CERT mediates intermembrane transfer of various molecular species of ceramides. *J Biol Chem* 2005; (280): 6488-6495.
- Kusunoki S, Shimizu J, Chiba A, Ugawa Y, Hitoshi S, Kanazawa I. Experimental sensory neuropathy induced by sensitization with ganglioside GD1b. *Ann Neurol* 1996; (39): 424-431.
- Kusunoki S, Hitoshi S, Kaida K, Arita M, Kanazawa I. Monospecific anti-GD1b IgG is required to induce rabbit ataxic neuropathy. *Ann Neurol* 1999; (45): 400-403.
- Kusunoki S, Kaida K I, Ueda M. Antibodies against gangliosides and ganglioside complexes in Guillain-Barré syndrome: New aspects of research. *Biochim Biophys Acta* 2007; (1780): 441-444.
- Kuwabara S, Misawa S, Takahashi H, Sawai S, Kanai K, Nakata M, Mori M, Hattori T, Yuki N. Anti-GQ1b antibody does not affect neuromuscular transmission in human limb muscle. *J Neuroimmunol* 2007; (189): 158-162.
- Kuziemko G M, Stroh M, Stevens R C. Cholera toxin binding affinity and specificity for gangliosides determined by surface plasmon resonance. *Biochemistry* 1996; (35): 6375-6384.
- Laev H, Mahadik S P. Topography of monosialoganglioside (GM1) in rat brain using monoclonal antibodies. *Neurosci Lett* 1989; (102): 7-14.
- Lang T, Bruns D, Wenzel D, Riedel D, Holroyd P, Thiele C, Jahn R. SNAREs are concentrated in cholesterol-dependent clusters that define docking and fusion sites for exocytosis. *EMBO J* 2001; (20): 2202-2213.
- Lange D J, DeAngelis T, Sivak M A. Single-fiber electromyography shows terminal axon dysfunction in Miller Fisher syndrome: a case report. *Muscle Nerve* 2006; (34): 232-234.
- Lasky T, Terracciano G J, Magder L, Koski C L, Ballesteros M, Nash D, Clark S, Haber P, Stolley P D, Schonberger L B, Chen R T. The Guillain-Barré syndrome and the 1993 and 1993-1994 influenza vaccines. *N Engl J Med* 1998; (339): 1797-1802.
- Ledeer R W. Gangliosides of the neuron. *Trends in Neurosciences* 1985; (8): 169-174.
- Ledeer R W, Diebler M F, Wu G, Lu Z H, Varoqui H. Ganglioside composition of subcellular fractions, including pre- and postsynaptic membranes, from Torpedo electric organ. *Neurochem Res* 1993; (18): 1151-1155.
- Ledeer R W, Wu G S. Ganglioside function in calcium homeostasis and signaling. *Neurochemical Research* 2002; (27): 637-647.
- Ledeer R W, Wu G S. Sphingolipids of the nucleus and their role in nuclear signaling. *Biochim Biophys Acta-Molecular and Cell Biology of Lipids* 2006b; (1761): 588-598.
- Ledeer R W, Wu G S. GM1 ganglioside: another nuclear lipid that modulates nuclear calcium. GM1 potentiates the nuclear sodium-calcium exchanger. *Canadian Journal of Physiology and Pharmacology* 2006a; (84): 393-402.
- Lee M S, Lee J H, Lee H S, Chang H, Kim Y S, Cho K H, Ahn S H, Song J H, Yoo M, Han J K, Park H Y. Scrub typhus as a possible aetiology of Guillain-Barré syndrome: two cases. *Ir J Med Sci* 2008.
- Lehmann H C, Lopez P H, Zhang G, Ngyuen T, Zhang J, Kieseier B C, Mori S, Sheikh K A. Passive immunization with anti-ganglioside antibodies directly inhibits axon regeneration in an animal model. *J Neurosci* 2007; (27): 27-34.
- Lencer W I, Tsai B. The intracellular voyage of cholera toxin: going retro. *Trends Biochem Sci* 2003; (28): 639-645.
- Lennon V A, Kryzer T J, Griesmann G E, Osuilleabhain P E, Windebank A J, Woppmann A, Miljanich G P, Lambert E H. Calcium-channel antibodies in the Lambert-Eaton syndrome and other paraneoplastic syndromes. *N Engl J Med* 1995; (332): 1467-1474.
- Leung S S. Guillain-Barré syndrome in elderly people. *J Am Geriatr Soc* 2008; (56): 381-382.
- Li H, Turley S D, Liu B, Repa J J, Dietschy J M. GM2/GD2 and GM3 gangliosides have no effect on cellular cholesterol pools or turnover in normal or NPC1 mice. *J Lipid Res* 2008; (49): 1816-1828.
- Lin F, Kaminski H J, Conti-Fine B M, Wang W, Richmonds C, Medof M E. Markedly enhanced susceptibility to experimental autoimmune myasthenia gravis in the absence of decay-accelerating factor protection. *J Clin Invest* 2002; (110): 1269-1274.
- Lin M S, Chen L Y, Wang S S, Chang Y, Chen W Y. Examining the levels of ganglioside and cholesterol in cell membrane on attenuation the cytotoxicity of beta-amyloid peptide. *Colloids Surf B Biointerfaces* 2008; (65): 172-177.

- Lin W, Burgess R W, Dominguez B, Pfaff S L, Sanes J R, Lee K F. Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse. *Nature* 2001; (410): 1057-1064.
- Lisman J E, Raghavachari S, Tsien R W. The sequence of events that underlie quantal transmission at central glutamatergic synapses. *Nat Rev Neurosci* 2007; (8): 597-609.
- Liu Y, Wada R, Kawai H, Sango K, Deng C, Tai T, McDonald M P, Araujo K, Crawley J N, Bierfreund U, Sandhoff K, Suzuki K, Proia R L. A genetic model of substrate deprivation therapy for a glycosphingolipid storage disorder. *J Clin Invest* 1999; (103): 497-505.
- Liu Y, Su Y, Wiznitzer M, Epifano O, Ladisch S. Ganglioside depletion and EGF responses of human GM3 synthase-deficient fibroblasts. *Glycobiology* 2008; (18): 593-601.
- Lloyd K O, Gordon C M, Thampoe I J, DiBenedetto C. Cell surface accessibility of individual gangliosides in malignant melanoma cells to antibodies is influenced by the total ganglioside composition of the cells. *Cancer Res* 1992; (52): 4948-4953.
- Lo Y L, Chan L L, Pan A, Ratnagopal P. Acute ophthalmoparesis in the anti-GQ1b antibody syndrome: electrophysiological evidence of neuromuscular transmission defect in the orbicularis oculi. *J Neurol Neurosurg Psychiatry* 2004; (75): 436-440.
- Lo Y L, Leoh T H, Dan Y F, Lim L L, Seah A, Fook-Chong S, Ratnagopal P. Presynaptic neuromuscular transmission defect in the Miller Fisher syndrome. *Neurology* 2006; (66): 148-149.
- Lopez P H, Zhang G, Bianchet M A, Schnaar R L, Sheikh K A. Structural requirements of anti-GD1a antibodies determine their target specificity. *Brain* 2008; (131): 1926-1939.
- Lu J L, Sheikh K A, Wu H S, Zhang J, Jiang Z F, Cornblath D R, McKhann G M, Asbury A K, Griffin J W, Ho T W. Physiologic-pathologic correlation in Guillain-Barre syndrome in children. *Neurology* 2000; (54): 33-39.
- Lunn M P, Willison H J. Diagnosis and treatment in inflammatory neuropathies. *J Neurol Neurosurg Psychiatry* 2009; (80): 249-258.
- Luongo L, Sajic M, Grist J, Clark A K, Maione S, Malcangio M. Spinal changes associated with mechanical hypersensitivity in a model of Guillain-Barre syndrome. *Neurosci Lett* 2008; (437): 98-102.
- Ma Q, Kobayashi M, Sugiura M, Ozaki N, Nishio K, Shiraishi Y, Furukawa K, Furukawa K, Sugiura Y. Morphological study of disordered myelination and the degeneration of nerve fibers in the spinal cord of mice lacking complex gangliosides. *Arch Histol Cytol* 2003; (66): 37-44.
- Maccioni H J. Glycosylation of glycolipids in the Golgi complex. *J Neurochem* 2007; (103 Suppl 1): 81-90.
- Maegawa G H, Stockley T, Tropak M, Banwell B, Blaser S, Kok F, Giugliani R, Mahuran D, Clarke J T. The natural history of juvenile or subacute GM2 gangliosidosis: 21 new cases and literature review of 134 previously reported. *Pediatrics* 2006; (118): e1550-e1562.
- Magleby K L, Stevens C F. A quantitative description of end-plate currents. *J Physiol* 1972; (223): 173-197.
- Malisan F, Testi R. GD3 in cellular ageing and apoptosis. *Exp Gerontol* 2002; (37): 1273-1282.
- Manganelli F, Pisciotta C, Iodice R, Calandro S, Dubbioso R, Ranieri A, Santoro L. Case of acute motor conduction block neuropathy (AMCBN). *Muscle Nerve* 2009; (39): 224-226.
- Mangano K, Dati G, Quattrocchi C, Proietti L, Mazzarino C, Di Marco R, Bendtzen K, Greco B, Zaratin P, Nicoletti F. Preventive and curative effects of cyclophosphamide in an animal model of Guillain Barre syndrome. *J Neuroimmunol* 2008; (196): 107-115.
- Mantipragada S B, Horvath L I, Arias H R, Schwarzmann G, Sandhoff K, Barrantes F J, Marsh D. Lipid-protein interactions and effect of local anesthetics in acetylcholine receptor-rich membranes from *Torpedo marmorata* electric organ. *Biochemistry* 2003; (42): 9167-9175.
- Marcheselli V, Daniotti J L, Vidal A C, Maccioni H, Marsh D, Barrantes F J. Gangliosides in acetylcholine receptor-rich membranes from *Torpedo marmorata* and *Discopyge tschudii*. *Neurochem Res* 1993; (18): 599-603.
- Marconi S, Acler M, Lovato L, De Toni L, Tedeschi E, Anghileri E, Romito S, Cordioli C, Bonetti B. Anti-GD2-like IgM autoreactivity in multiple sclerosis patients. *Mult Scler* 2006; (12): 302-308.
- Marengo F D, Wang S Y, Wang B, Langer G A. Dependence of cardiac cell Ca²⁺ permeability on sialic acid-containing sarcolemmal gangliosides. *J Mol Cell Cardiol* 1998; (30): 127-137.
- Maselli R A, Leung C. Analysis of Anticholinesterase-Induced Neuromuscular Transmission Failure. *Muscle Nerve* 1993; (16): 548-553.
- Matà S, Galli E, Amantini A, Pinto F, Sorbi S, Lolli F. Anti-ganglioside antibodies and elevated CSF IgG levels in Guillain-Barré syndrome. *Eur J Neurol* 2006; (13): 153-160.
- McGrogan A, Madle G C, Seaman H E, de Vries C S. The Epidemiology of Guillain-Barré Syndrome Worldwide. A Systematic Literature Review. *Neuroepidemiology* 2008; (32): 150-163.
- Meir A, Ginsburg S, Butkevich A, Kachalsky S G, Kaiserman I, Ahdut R, Demirgoren S, Rahamimoff R. Ion channels in presynaptic nerve terminals and control of transmitter release. *Physiol Rev* 1999; (79): 1019-1088.
- Merritt E A, Sarfaty S, van den A F, L'Hoir C, Martial J A, Hol W G. Crystal structure of cholera toxin B-pentamer bound to receptor GM1 pentasaccharide. *Protein Sci* 1994; (3): 166-175.
- Meyer zu Hörste G, Hu W, Hartung H P, Lehmann H C, Kieseier B C. The immunocompetence of Schwann cells. *Muscle Nerve* 2008; (37): 3-13.
- Middlebrook J L, Dorland R B. Bacterial toxins: cellular mechanisms of action. *Microbiol Rev* 1984; (48): 199-221.

-
- Morales A, Lee H, Goni F M, Kolesnick R, Fernandez-Checa J C. Sphingolipids and cell death. *Apoptosis* 2007; (12): 923-939.
- Moran A P, Prendergast M M. Molecular Mimicry in *Campylobacter jejuni* and *Helicobacter pylori* Lipopolysaccharides: Contribution of Gastrointestinal Infections to Autoimmunity. *Journal of Autoimmunity* 2001; (16): 241-256.
- Moran A P, Annuk H, Prendergast M M. Antibodies induced by ganglioside-mimicking *Campylobacter jejuni* lipooligosaccharides recognise epitopes at the nodes of Ranvier. *J Neuroimmunol* 2005; (165): 179-185.
- Morgan B P. Mechanisms of tissue damage by the membrane attack complex of complement. *Complement Inflamm* 1989; (6): 104-111.
- Morgan B P, Berg C W, Harris C L. "Homologous restriction" in complement lysis: roles of membrane complement regulators. *Xenotransplantation* 2005; (12): 258-265.
- Morgan B P, Chamberlain-Banoub J, Neal J W, Song W, Mizuno M, Harris C L. The membrane attack pathway of complement drives pathology in passively induced experimental autoimmune myasthenia gravis in mice. *Clin Exp Immunol* 2006; (146): 294-302.
- Mori M, Kuwabara S, Fukutake T, Yuki N, Hattori T. Clinical features and prognosis of Miller Fisher syndrome. *Neurology* 2001; (56): 1104-1106.
- Motomura M, Lang B, Johnston I, Palace J, Vincent A, Newsomdavis J. Incidence of serum anti-P/Q-type and anti-N-type calcium channel autoantibodies in the Lambert-Eaton myasthenic syndrome. *J Neurol Sci* 1997; (147): 35-42.
- Nachamkin I, Shadomy S V, Moran A P, Cox N, Fitzgerald C, Ung H, Corcoran A T, Iskander J K, Schonberger L B, Chen R T. Anti-ganglioside antibody induction by swine (A/NJ/1976/H1N1) and other influenza vaccines: insights into vaccine-associated Guillain-Barré syndrome. *J Infect Dis* 2008; (198): 226-233.
- Nagashima T, Koga M, Odaka M, Hirata K, Yuki N. Continuous spectrum of pharyngeal-cervical-brachial variant of Guillain-Barre syndrome. *Arch Neurol* 2007; (64): 1519-1523.
- Nakatani Y, Nagaoka T, Hotta S, Utsunomiya I, Yoshino H, Miyatake T, Hoshi K, Taguchi K. IgG anti-GalNAc-GD1a antibody inhibits the voltage-dependent calcium channel currents in PC12 pheochromocytoma cells. *Exp Neurol* 2007a; (204): 380-386.
- Nakatani Y, Kawakami K, Nagaoka T, Utsunomiya I, Tanaka K, Yoshino H, Miyatake T, Hoshi K, Taguchi K. Ca channel currents inhibited by serum from select patients with Guillain-Barré syndrome. *Eur Neurol* 2007b; (57): 11-18.
- Nakatani Y, Hotta S, Utsunomiya I, Tanaka K, Hoshi K, Ariga T, Yu R K, Miyatake T, Taguchi K. Cav2.1 voltage-dependent Ca²⁺ channel current is inhibited by serum from select patients with Guillain-Barré syndrome. *Neurochem Res* 2009; (34): 149-157.
- Ngamukote S, Yanagisawa M, Ariga T, Ando S, Yu R K. Developmental changes of glycosphingolipids and expression of glycogenes in mouse brains. *J Neurochem* 2007.
- Nguyen T, Mehta N R, Conant K, Kim K J, Jones M, Calabresi P A, Melli G, Hoke A, Schnaar R L, Ming G L, Song H, Keswani S C, Griffin J W. Axonal protective effects of the myelin-associated glycoprotein. *J Neurosci* 2009; (29): 630-637.
- Nores G A, Lardone R D, Comin R, Alaniz M E, Moyano A L, Irazoqui F J. Anti-GM1 antibodies as a model of the immune response to self-glycans. *Biochim Biophys Acta* 2007.
- Notturmo F, Caporale C M, Uncini A. Acute sensory ataxic neuropathy with antibodies to GD1b and GQ1b gangliosides and prompt recovery. *Muscle Nerve* 2008; (37): 265-268.
- Notturmo F, Luciani M, Caporale C M, Ciarelli A, Uncini A. Antibodies to ganglioside complexes in Guillain-Barre syndrome: clinical correlates, fine specificity and complement activation. *Int J Immunopathol Pharmacol* 2009; (22): 437-445.
- Nunn M A, Sharma A, Paesen G C, Adamson S, Lissina O, Willis A C, Nuttall P A. Complement inhibitor of C5 activation from the soft tick *Ornithodoros moubata*. *J Immunol* 2005; (174): 2084-2091.
- O'Hanlon G M, Paterson G J, Veitch J, Wilson G, Willison H J. Mapping immunoreactive epitopes in the human peripheral nervous system using human monoclonal anti-GM1 ganglioside antibodies. *Acta Neuropathol* 1998; (95): 605-616.
- O'Hanlon G M, Plomp J J, Chakrabarti M, Morrison I, Wagner E R, Goodyear C S, Yin X, Trapp B D, Conner J, Molenaar P C, Stewart S, Rowan E G, Willison H J. Anti-GQ1b ganglioside antibodies mediate complement-dependent destruction of the motor nerve terminal. *Brain* 2001; (124): 893-906.
- O'Hanlon G M, Bullens R W, Plomp J J, Willison H J. Complex gangliosides as autoantibody targets at the neuromuscular junction in Miller Fisher syndrome: a current perspective. *Neurochem Res* 2002; (27): 697-709.
- O'Hanlon G M, Humphreys P D, Goldman R S, Halstead S K, Bullens R W, Plomp J J, Ushkaryov Y, Willison H J. Calpain inhibitors protect against axonal degeneration in a model of anti-ganglioside antibody-mediated motor nerve terminal injury. *Brain* 2003; (126): 2497-2509.
- Okada M, Itoh M M, Haraguchi M, Okajima T, Inoue M, Oishi H, Matsuda Y, Iwamoto T, Kawano T, Fukumoto S, Miyazaki H, Furukawa K, Aizawa S, Furukawa K. b-series Ganglioside deficiency exhibits no definite changes in the neurogenesis and the sensitivity to Fas-mediated apoptosis but impairs regeneration of the lesioned hypoglossal nerve. *J Biol Chem* 2002; (277): 1633-1636.
- Ong G L, Mattes M J. Mouse strains with typical mammalian levels of complement activity. *J Immunol Methods* 1989; (125): 147-158.

- Ong G L, Baker A E, Mattes M J. Analysis of high complement levels in *Mus hortulanus* and BUB mice. *J Immunol Methods* 1992; (154): 37-45.
- Ortiz N, Rosa R, Gallardo E, Illa I, Tomàs J, Aubry J, Sabater M, Santafé M. IgM monoclonal antibody against terminal moiety of GM2, GalNAc-GD1a and GalNAc-GM1b from a pure motor chronic demyelinating polyneuropathy patient: effects on neurotransmitter release. *J Neuroimmunol* 2001; (119): 114-123.
- Osmers I, Szalai A J, Tenner A J, Barnum S R. Complement in BuB/BnJ mice revisited: Serum C3 levels and complement opsonic activity are not elevated. *Molecular Immunol* 2006; (43): 1722-1725.
- Pangburn M K, Schreiber R D, Muller-Eberhard H J. Human complement C3b inactivator: isolation, characterization, and demonstration of an absolute requirement for the serum protein beta1H for cleavage of C3b and C4b in solution. *J Exp Med* 1977; (146): 257-270.
- Pangburn M K, Muller-Eberhard H J. Complement C3 convertase: cell surface restriction of beta1H control and generation of restriction on neuraminidase-treated cells. *Proc Natl Acad Sci U S A* 1978; (75): 2416-2420.
- Paparounas K, O'Hanlon G M, O'Leary C P, Rowan E G, Willison H J. Anti-ganglioside antibodies can bind peripheral nerve nodes of Ranvier and activate the complement cascade without inducing acute conduction block in vitro. *Brain* 1999; (122 (Pt 5)): 807-816.
- Paparounas K. Anti-GQ1b ganglioside antibody in peripheral nervous system disorders: pathophysiologic role and clinical relevance. *Arch Neurol* 2004; (61): 1013-1016.
- Park E J, Suh M, Ramanujam K, Steiner K, Begg D, Clandinin M T. Diet-induced changes in membrane gangliosides in rat intestinal mucosa, plasma and brain. *J Pediatr Gastroenterol Nutr* 2005; (40): 487-495.
- Paterson G, Wilson G, Kennedy P G E, Willison H J. Analysis of anti-GM1 ganglioside IgM antibodies cloned from motor neuropathy patients demonstrates diverse v-region gene usage with extensive somatic mutation. *J Immunol* 1995; (155): 3049-3059.
- Pato C, Stetzkowski-Marden F, Gaus K, Recouvreur M, Cartaud A, Cartaud J. Role of lipid rafts in agrin-elicited acetylcholine receptor clustering. *Chem Biol Interact* 2008; (175): 64-67.
- Paulson J C, Blixt O, Collins B E. Sweet spots in functional glycomics. *Nat Chem Biol* 2006; (2): 238-248.
- Paw B H, Kaback M M, Neufeld E F. Molecular basis of adult-onset and chronic GM2 gangliosidosis in patients of Ashkenazi Jewish origin: substitution of serine for glycine at position 269 of the alpha-subunit of beta-hexosaminidase. *Proc Natl Acad Sci U S A* 1989; (86): 2413-2417.
- Perera V N, Nachamkin I, Ung H, Patterson J H, McConville M J, Coloe P J, Fry B N. Molecular mimicry in *Campylobacter jejuni*: role of the lipo-oligosaccharide core oligosaccharide in inducing anti-ganglioside antibodies. *FEMS Immunol Med Microbiol* 2007; (50): 27-36.
- Perillo M A, Yu R K, Maggio B. Modulation of the activity of *Clostridium perfringens* neuraminidase by the molecular organization of gangliosides in monolayers. *Biochim Biophys Acta* 1994; (1193): 155-164.
- Peters I, Igbavboa U, Schutt T, Haidari S, Hartig U, Rosello X, Bottner S, Copanaki E, Deller T, Kogel D, Wood W G, Muller W E, Eckert G P. The interaction of beta-amyloid protein with cellular membranes stimulates its own production. *Biochim Biophys Acta* 2009; (1788): 964-972.
- Peterson M E. Black widow spider envenomation. *Clin Tech Small Anim Pract* 2006; (21): 187-190.
- Petrov A M, Giniatullin A R, Sitdikova G F, Zefirov A L. The role of cGMP-dependent signaling pathway in synaptic vesicle cycle at the frog motor nerve terminals. *J Neurosci* 2008; (28): 13216-13222.
- Phongsisay V, Susuki K, Matsuno K, Yamahashi T, Okamoto S, Funakoshi K, Hirata K, Shinoda M, Yuki N. Complement inhibitor prevents disruption of sodium channel clusters in a rabbit model of Guillain-Barre syndrome. *J Neuroimmunol* 2008; (205): 101-104.
- Plomp J J, Molenaar P C, O'Hanlon G M, Jacobs B C, Veitch J, Daha M R, van Doorn P A, van der Meché F G, Vincent A, Morgan B P, Willison H J. Miller Fisher anti-GQ1b antibodies: alpha-latrotoxin-like effects on motor end plates. *Ann Neurol* 1999; (45): 189-199.
- Plomp J J, Willison H J. Pathophysiological actions of neuropathy-related anti-ganglioside antibodies at the neuromuscular junction. *J Physiol* 2009; (doi: 10.1113/jphysiol.2009.171702).
- Prendergast M M, Tribble D R, Baqar S, Scott D A, Ferris J A, Walker R I, Moran A P. In vivo phase variation and serologic response to lipooligosaccharide of *Campylobacter jejuni* in experimental human infection. *Infect Immun* 2004; (72): 916-922.
- Prinetti A, Loberto N, Chigorno V, Sonnino S. Glycosphingolipid behaviour in complex membranes. *Biochim Biophys Acta* 2009; (1788): 184-193.
- Pritchard J, Hughes R A. Guillain-Barré syndrome. *Lancet* 2004; (363): 2186-2188.
- Proia R L. Glycosphingolipid functions: insights from engineered mouse models. *Philos Trans R Soc Lond B Biol Sci* 2003; (358): 879-883.
- Putzu G A, Figarella-Branger D, Bouvier-Labit C, Liprandi A, Bianco N, Pellissier J F. Immunohistochemical localization of cytokines, C5b-9 and ICAM-1 in peripheral nerve of Guillain-Barré syndrome. *J Neurol Sci* 2000; (174): 16-21.
- Rahmann H, Kortje K H, Seybold V, Rosner H. Ultrastructural localization of gangliosides, calcium and a high-affinity Ca²⁺-ATPase in nerve terminals: a contribution to the possible functional role of gangliosides. *Indian J Biochem Biophys* 1990; (27): 420-424.
- Rahmann H, Schifferer F, Beiting H. Calcium-ganglioside interactions and synaptic plasticity: effect of calcium on specific ganglioside/peptide (valinomycin, gramicidin A)-complexes in mixed mono- and bilayers. *Neurochem Int* 1992; (20): 323-338.

-
- Rahmann H, Jonas U, Kappel T, Hilderbrandt H. Differential involvement of gangliosides versus phospholipids in the process of temperature adaptation in vertebrates. A comparative phenomenological and physicochemical study. *Ann N Y Acad Sci* 1998; (845): 72-91.
- Ramaglia V, King R H, Nourallah M, Wolterman R, de Jonge R, Ramkema M, Vigar M A, van der W S, Morgan B P, Troost D, Baas F. The membrane attack complex of the complement system is essential for rapid Wallerian degeneration. *J Neurosci* 2007; (27): 7663-7672.
- Ramaglia V, Wolterman R, de Kok M, Vigar M A, Wagenaar-Bos I, King R H, Morgan B P, Baas F. Soluble complement receptor 1 protects the peripheral nerve from early axon loss after injury. *Am J Pathol* 2008; (172): 1043-1052.
- Ramirez O A, Gomez R A, Carrer H F. Gangliosides improve synaptic transmission in dentate gyrus of hippocampal rat slices. *Brain Res* 1990; (506): 291-293.
- Raphaël J C, Chevret S, Hughes R A, Annane D. Plasma exchange for Guillain-Barré syndrome. *Cochrane Database Syst Rev* 2002;CD001798.
- Ravichandra B, Joshi P G. Regulation of transmembrane signaling by ganglioside GM1: interaction of anti-GM1 with Neuro2a cells. *J Neurochem* 1999; (73): 557-567.
- Rekand T, Gramstad A, Vedeler C A. Fatigue, pain and muscle weakness are frequent after Guillain-Barré syndrome and poliomyelitis. *J Neurol* 2009.
- Ribeiro-Resende V T, Oliveira-Silva A, Ouverney-Brandao S, Santiago M F, Hedin-Pereira C, Mendez-Otero R. Ganglioside 9-O-acetyl GD3 expression is upregulated in the regenerating peripheral nerve. *Neuroscience* 2007; (147): 97-105.
- Rice C E. The interchangeability of the complement components of different animal species; literature survey. *Can J Comp Med Vet Sci* 1950; (14): 369-379.
- Richards D A, Guatimosim C, Rizzoli S O, Betz W J. Synaptic vesicle pools at the frog neuromuscular junction. *Neuron* 2003; (39): 529-541.
- Ricklin D, Lambris J D. Complement-targeted therapeutics. *Nat Biotechnol* 2007; (25): 1265-1275.
- Rinaldi S, Willison H J. Ganglioside antibodies and neuropathies. *Curr Opin Neurol* 2008; (21): 540-546.
- Roberts M, Willison H, Vincent A, Newsom-Davis J. Serum factor in Miller-Fisher variant of Guillain-Barré syndrome and neurotransmitter release. *Lancet* 1994; (343): 454-455.
- Roberts M, Willison H J, Vincent A, Newsom-Davis J. Multifocal motor neuropathy human sera block distal motor nerve conduction in mice. *Ann Neurol* 1995; (38): 111-118.
- Rogozhin A A, Pang K K, Bukharaeva E, Young C, Slater C R. Recovery of mouse neuromuscular junctions from single and repeated injections of botulinum neurotoxin A. *J Physiol* 2008; (586): 3163-3182.
- Rohou A, Nield J, Ushkaryov Y A. Insecticidal toxins from black widow spider venom. *Toxicon* 2007; (49): 531-549.
- Roth J, Kempf A, Reuter G, Schauer R, Gehring W J. Occurrence of sialic acids in *Drosophila melanogaster*. *Science* 1992; (256): 673-675.
- Rother R P, Rollins S A, Mojcik C F, Brodsky R A, Bell L. Discovery and development of the complement inhibitor eculizumab for the treatment of paroxysmal nocturnal hemoglobinuria. *Nat Biotechnol* 2007; (25): 1256-1264.
- Rudolph T, Larsen J P, Farbu E. The long-term functional status in patients with Guillain-Barré syndrome. *Eur J Neurol* 2008; (15): 1332-1337.
- Saito M, Ito M, Sugiyama K. A specific loss of C-series gangliosides in pancreas of streptozotocin-induced diabetic rats. *Life Sci* 1999; (64): 1803-1810.
- Saito M, Sugiyama K. Major and c-series gangliosides in lenticular tissues: mammals to molluscs. *Comp Biochem Physiol B Biochem Mol Biol* 2001; (130): 313-321.
- Saito M, Mao R F, Wang R, Vadasz C, Saito M. Effects of gangliosides on ethanol-induced neurodegeneration in the developing mouse brain. *Alcohol Clin Exp Res* 2007; (31): 665-674.
- Salaün C, James D J, Chamberlain L H. Lipid rafts and the regulation of exocytosis. *Traffic* 2004; (5): 255-264.
- Salazar B C, Castano S, Sanchez J C, Romero M, Recio-Pinto E. Ganglioside GD1a increases the excitability of voltage-dependent sodium channels. *Brain Res* 2004; (1021): 151-158.
- Sanders M E, Koski C L, Robbins D, Shin M L, Frank M M, Joiner K A. Activated terminal complement in cerebrospinal fluid in Guillain-Barre syndrome and multiple sclerosis. *J Immunol* 1986; (136): 4456-4459.
- Sandhoff K, Kolter T. Biosynthesis and degradation of mammalian glycosphingolipids. *Philos Trans R Soc Lond B Biol Sci* 2003; (358): 847-861.
- Sandhoff R, Geyer R, Jennemann R, Paret C, Kiss E, Yamashita T, Gorgas K, Sijmonsma T P, Iwamori M, Finaz C, Proia R L, Wiegandt H, Grone H J. Novel class of glycosphingolipids involved in male fertility. *J Biol Chem* 2005; (280): 27310-27318.
- Sanes J R, Lichtman J W. Development of the vertebrate neuromuscular junction. *Annu Rev Neurosci* 1999; (22): 389-442.
- Santafé M M, Urbano F J, Lanuza M A, Uchitel O D. Multiple types of calcium channels mediate transmitter release during functional recovery of botulinum toxin type A-poisoned mouse motor nerve terminals. *Neuroscience* 2000; (95): 227-234.
- Santafé M M, Sabaté M M, Garcia N, Ortiz N, Lanuza M A, Tomàs J. Changes in the neuromuscular synapse induced by an antibody against gangliosides. *Ann Neurol* 2005; (57): 396-407.

- Santafé M M, Sabaté M M, Garcia N, Ortiz N, Lanuza M A, Tomàs J. Anti-GM2 gangliosides IgM paraprotein induces neuromuscular block without neuromuscular damage. *J Neuroimmunol* 2008; (204): 20-28.
- Sarkey J P, Richards M P, Stubbs E B, Jr. Lovastatin attenuates nerve injury in an animal model of Guillain-Barré syndrome. *J Neurochem* 2007; (100): 1265-1277.
- Sartucci F, Cafforio G, Borghetti D, Domenici L, Orlandi G, Murri L. Electrophysiological evidence by single fibre electromyography of neuromuscular transmission impairment in a case of Miller Fisher syndrome. *Neurol Sci* 2005; (26): 125-128.
- Schnaar R L, Lopez P H. Myelin-associated glycoprotein and its axonal receptors. *J Neurosci Res* 2009.
- Schwarz A, Futerman A H. The localization of gangliosides in neurons of the central nervous system: The use of anti-ganglioside antibodies. *Biochimica et Biophysica Acta - Reviews on Biomembranes* 1996; (1286): 247-267.
- Seyfried T N, el Abbadi M, Ecsedy J A, Bai H W, Yohe H C. Influence of host cell infiltration on the glycolipid content of mouse brain tumors. *J Neurochem* 1996; (66): 2026-2033.
- Shafqat S, Khealani B A, Awan F, Abedin S E. Guillain-Barré syndrome in Pakistan: similarity of demyelinating and axonal variants. *Eur J Neurol* 2006; (13): 662-665.
- Shahar E. Current therapeutic options in severe Guillain-Barré syndrome. *Clin Neuropharmacol* 2006; (29): 45-51.
- Sheikh K A, Sun J, Liu Y, Kawai H, Crawford T O, Proia R L, Griffin J W, Schnaar R L. Mice lacking complex gangliosides develop Wallerian degeneration and myelination defects. *Proc Natl Acad Sci U S A* 1999; (96): 7532-7537.
- Shevchuk N A, Hathout Y, Epifano O, Su Y, Liu Y, Sutherland M, Ladisch S. Alteration of ganglioside synthesis by GM3 synthase knockout in murine embryonic fibroblasts. *Biochim Biophys Acta* 2007; (1771): 1226-1234.
- Simons K, Ikonen E. Functional rafts in cell membranes. *Nature* 1997; (387): 569-572.
- Simpson M A, Cross H, Proukakis C, Priestman D A, Neville D C, Reinkensmeier G, Wang H, Wiznitzer M, Gurtz K, Verganelaki A, Pryde A, Patton M A, Dwek R A, Butters T D, Platt F M, Crosby A H. Infantile-onset symptomatic epilepsy syndrome caused by a homozygous loss-of-function mutation of GM3 synthase. *Nat Genet* 2004; (36): 1225-1229.
- Slomiany B L, Liu J, Fekete Z, Yao P, Slomiany A. Modulation of dihydropyridine-sensitive gastric mucosal calcium channels by GM1-ganglioside. *International Journal of Biochemistry* 1992; (24): 1289-1294.
- Sohn H, Kim Y S, Kim H T, Kim C H, Cho E W, Kang H Y, Kim N S, Kim C H, Ryu S E, Lee J H, Ko J H. Ganglioside GM3 is involved in neuronal cell death. *FASEB J* 2006; (20): 1248-1250.
- Sonnino S, Mauri L, Chigorno V, Prinetti A. Gangliosides as components of lipid membrane domains. *Glycobiology* 2007; (17): 1R-13R.
- Stangel M, Pul R. Basic principles of intravenous immunoglobulin (IVIg) treatment. *J Neurol* 2006; (253 Suppl 5): V18-V24.
- Stevens C F, Tsujimoto T. Estimates for the pool size of releasable quanta at a single central synapse and for the time required to refill the pool. *Proc Natl Acad Sci U S A* 1995; (92): 846-849.
- Sugiura Y, Furukawa K, Tajima O, Mii S, Honda T, Furukawa K. Sensory nerve-dominant nerve degeneration and remodeling in the mutant mice lacking complex gangliosides. *Neuroscience* 2005; (135): 1167-1178.
- Susuki K, Nishimoto Y, Yamada M, Baba M, Ueda S, Hirata K, Yuki N. Acute motor axonal neuropathy rabbit model: immune attack on nerve root axons. *Ann Neurol* 2003; (54): 383-388.
- Susuki K, Baba H, Tohyama K, Kanai K, Kuwabara S, Hirata K, Furukawa K, Furukawa K, Rasband M N, Yuki N. Gangliosides contribute to stability of paranodal junctions and ion channel clusters in myelinated nerve fibers. *Glia* 2007b; (55): 746-757.
- Susuki K, Rasband M N, Tohyama K, Koibuchi K, Okamoto S, Funakoshi K, Hirata K, Baba H, Yuki N. Anti-GM1 antibodies cause complement-mediated disruption of sodium channel clusters in peripheral motor nerve fibers. *J Neurosci* 2007a; (27): 3956-3967.
- Suzuki A. Genetic basis for the lack of N-glycolylneuraminic acid expression in human tissues and its implication to human evolution. *Proc Jpn Acad* 2008; (82): 93-103.
- Svennerholm L. The gangliosides. *J Lipid Res* 1964; (5): 145-155.
- Svennerholm L, Bostrom K, Fredman P, Mansson J E, Rosengren B, Rynmark B M. Human brain gangliosides: developmental changes from early fetal stage to advanced age. *Biochim Biophys Acta* 1989; (1005): 109-117.
- Svennerholm L. Designation and schematic structure of gangliosides and allied glycosphingolipids. *Prog Brain Res* 1994; (101): XI-XIV.
- Svennerholm L, Bostrom K, Jungbjer B. Changes in weight and compositions of major membrane components of human brain during the span of adult human life of Swedes. *Acta Neuropathol* 1997; (94): 345-352.
- Taboada E N, van Belkum A, Yuki N, Acedillo R R, Godschalk P C, Koga M, Endtz H P, Gilbert M, Nash J H. Comparative genomic analysis of Campylobacter jejuni associated with Guillain-Barré and Miller Fisher syndromes: neuropathogenic and enteritis-associated isolates can share high levels of genomic similarity. *BMC Genomics* 2007; (8): 359.
- Takamiya K, Yamamoto A, Furukawa K, Yamashiro S, Shin M, Okada M, Fukumoto S, Haraguchi M, Takeda N, Fujimura K, Sakae M, Kishikawa M, Shiku H, Furukawa K, Aizawa S. Mice with disrupted GM2/GD2 synthase gene lack complex gangliosides but exhibit only subtle defects in their nervous system. *Proc Natl Acad Sci U S A* 1996; (93): 10662-10667.

-
- Tanaka Y, Waki H, Kon K, Ando S. Gangliosides enhance KCl-induced Ca^{2+} influx and acetylcholine release in brain synaptosomes. *Neuroreport* 1997; (8): 2203-2207.
- Tatsumoto M, Koga M, Gilbert M, Odaka M, Hirata K, Kuwabara S, Yuki N. Spectrum of neurological diseases associated with antibodies to minor gangliosides GM1b and GalNAc-GD1a. *J Neuroimmunol* 2006; (177): 201-208.
- Taverna E, Saba E, Rowe J, Francolini M, Clementi F, Rosa P. Role of lipid microdomains in P/Q-type calcium channel ($Ca_v2.1$) clustering and function in presynaptic membranes. *J Biol Chem* 2004; (279): 5127-5134.
- Terrell J, Yadava P, Castro C, Hughes J. Liposome fluidity alters interactions between the ganglioside GM1 and cholera toxin B subunit. *J Liposome Res* 2008; (18): 21-29.
- Tettamanti G, Bonali F, Marchesini S, Zambotti V. A new procedure for the extraction, purification and fractionation of brain gangliosides. *Biochim Biophys Acta* 1973; (296): 160-170.
- Thomas T C, Rollins S A, Rother R P, Giannoni M A, Hartman S L, Elliott E A, Nye S H, Matis L A, Squinto S P, Evans M J. Inhibition of complement activity by humanized anti-C5 antibody and single-chain Fv. *Mol Immunol* 1996; (33): 1389-1401.
- Townson K, Boffey J, Nicholl D, Veitch J, Bundle D, Zhang P, Samain E, Antoine T, Bernardi A, Arosio D, Sonnino S, Isaacs N, Willison H J. Solid phase immunoabsorption for therapeutic and analytical studies on neuropathy-associated anti-GM1 antibodies. *Glycobiology* 2007; (17): 294-303.
- Townson K H, Speak A O, Greenshields K N, Goodyear C S, Willison H J, Platt F M. Glycosphingolipid depletion in PC12 cells using iminosugars protects neuronal membranes from anti-ganglioside antibody mediated injury. *J Neuroimmunol* 2008; (203): 33-38.
- Tsai B, Gilbert J M, Stehle T, Lencer W, Benjamin T L, Rapoport T A. Gangliosides are receptors for murine polyoma virus and SV40. *EMBO J* 2003; (22): 4346-4355.
- Uchitel O D, Protti D A, Sanchez V, Cherksey B D, Sugimori M, Llinas R. P-type voltage-dependent calcium channel mediates presynaptic calcium influx and transmitter release in mammalian synapses. *Proc Natl Acad Sci U S A* 1992; (89): 3330-3333.
- Uncini A, Lugaresi A. Fisher syndrome with tetraparesis and antibody to GQ1b: evidence for motor nerve terminal block. *Muscle Nerve* 1999; (22): 640-644.
- Urbano F J, Pagani M R, Uchitel O D. Calcium channels, neuromuscular synaptic transmission and neurological diseases. *J Neuroimmunol* 2008; (201-202): 136-144.
- Valaperta R, Valsecchi M, Rocchetta F, Aureli M, Prioni S, Prinetti A, Chigorno V, Sonnino S. Induction of axonal differentiation by silencing plasma membrane-associated sialidase Neu3 in neuroblastoma cells. *J Neurochem* 2007; (100): 708-719.
- van Beek J, van Meurs M, 't Hart B A, Brok H P, Neal J W, Chatagner A, Harris C L, Omidvar N, Morgan B P, Laman J D, Gasque P. Decay-accelerating factor (CD55) is expressed by neurons in response to chronic but not acute autoimmune central nervous system inflammation associated with complement activation. *J Immunol* 2005; (174): 2353-2365.
- van Doorn P A, Ruts L, Jacobs B C. Clinical features, pathogenesis, and treatment of Guillain-Barré syndrome. *Lancet Neurol* 2008; (7): 939-950.
- van Koningsveld R, Steyerberg E W, Hughes R A, Swan A V, van Doorn P A, Jacobs B C. A clinical prognostic scoring system for Guillain-Barré syndrome. *Lancet Neurol* 2007; (6): 589-594.
- van Sorge N M, Yuki N, Jansen M D, Nishimoto Y, Susuki K, Wokke J H, van de Winkel J G, van den Berg L H, van der Pol W L. Leukocyte and complement activation by GM1-specific antibodies is associated with acute motor axonal neuropathy in rabbits. *J Neuroimmunol* 2007a; (182): 116-123.
- van Sorge N M, Yuki N, Koga M, Susuki K, Jansen M D, van Kooten C, Wokke J H, van de Winkel J G, van der Pol W L, van den Berg L H. Ganglioside-specific IgG and IgA recruit leukocyte effector functions in Guillain-Barré syndrome. *J Neuroimmunol* 2007b; (182): 177-184.
- Varki A. Glycan-based interactions involving vertebrate sialic-acid-recognizing proteins. *Nature* 2007; (446): 1023-1029.
- Varoqueaux F, Sons M S, Plomp J J, Brose N. Aberrant morphology and residual transmitter release at the Munc13-deficient mouse neuromuscular synapse. *Mol Cell Biol* 2005; (25): 5973-5984.
- Verderio C, Rossetto O, Grumelli C, Frassoni C, Montecucco C, Matteoli M. Entering neurons: botulinum toxins and synaptic vesicle recycling. *EMBO Rep* 2006; (7): 995-999.
- Vincent A, Lang B, Kleopa K A. Autoimmune channelopathies and related neurological disorders. *Neuron* 2006; (52): 123-138.
- Vyas A A, Patel H V, Fromholt S E, Heffer-Laue M, Vyas K A, Dang J, Schachner M, Schnaar R L. Gangliosides are functional nerve cell ligands for myelin-associated glycoprotein (MAG), an inhibitor of nerve regeneration. *Proc Natl Acad Sci U S A* 2002; (99): 8412-8417.
- Waki H, Kon K, Tanaka Y, Ando S. Facile methods for isolation and determination of gangliosides in a small scale: age-related changes of gangliosides in mouse brain synaptic plasma membranes. *Anal Biochem* 1994; (222): 156-162.
- Walport M J. Complement. First of two parts. *N Engl J Med* 2001; (344): 1058-1066.
- Wang L H, Tu Y P, Yang X Y, Tsui Z C, Yang F Y. Effect of ganglioside GM3 on the activity and conformation of reconstituted Ca^{2+} -ATPase. *FEBS Letters* 1996; (388): 128-130.

- Wang Y L, Tsui Z C, Yang F Y. Mechanistic study of modulation of SR Ca²⁺-ATPase activity by gangliosides GM1 and GM3 through some biophysical measurements. *Glycoconjugate Journal* 1999b; (16): 781-786.
- Wang Y L, Tsui Z C, Yang F Y. Antagonistic effect of ganglioside GM1 and GM3 on the activity and conformation of sarcoplasmic reticulum Ca²⁺-ATPase. *Febs Letters* 1999a; (457): 144-148.
- Wang Z, Sun Z, Li A V, Yarema K J. Roles for UDP-GlcNAc 2-epimerase/ManNAc 6-kinase outside of sialic acid biosynthesis: modulation of sialyltransferase and BiP expression, GM3 and GD3 biosynthesis, proliferation, and apoptosis, and ERK1/2 phosphorylation. *J Biol Chem* 2006; (281): 27016-27028.
- Wanschitz J, Maier H, Lassmann H, Budka H, Berger T. Distinct time pattern of complement activation and cytotoxic T cell response in Guillain-Barré syndrome. *Brain* 2003; (126): 2034-2042.
- Wasser C R, Ertunc M, Liu X, Kavalali E T. Cholesterol-dependent balance between evoked and spontaneous synaptic vesicle recycling. *J Physiol* 2007; (579): 413-429.
- Weiler J M, Daha M R, Austen K F, Fearon D T. Control of the amplification convertase of complement by the plasma protein beta1H. *Proc Natl Acad Sci U S A* 1976; (73): 3268-3272.
- Wieraszko A, Seifert W. Evidence for a functional role of gangliosides in synaptic transmission: studies on rat brain striatal slices. *Neurosci Lett* 1984; (52): 123-128.
- Wieraszko A, Seifert W. The role of monosialoganglioside GM1 in the synaptic plasticity: in vitro study on rat hippocampal slices. *Brain Res* 1985; (345): 159-164.
- Willison H J, Paterson G, Veitch J, Inglis G, Barnett S C. Peripheral neuropathy associated with monoclonal IgM anti-Pr2 cold agglutinins. *J Neurol Neurosurg Psychiatry* 1993; (56): 1178-1183.
- Willison H J, Veitch J. Immunoglobulin subclass distribution and binding characteristics of anti-GQ1b antibodies in Miller Fisher syndrome. *J Neuroimmunol* 1994; (50): 159-165.
- Willison H J, O'Hanlon G M, Paterson G, Veitch J, Wilson G, Roberts M, Tang T, Vincent A. A somatically mutated human antiganglioside IgM antibody that induces experimental neuropathy in mice is encoded by the variable region heavy chain gene, V1-18. *J Clin Invest* 1996; (97): 1155-1164.
- Willison H J, O'Hanlon G M. The immunopathogenesis of Miller Fisher syndrome. *J Neuroimmunol* 1999; (100): 3-12.
- Willison H J, Veitch J, Swan A V, Baumann N, Comi G, Gregson N A, Lila I, Jacobs B C, Zielasek J, Hughes R A C. Inter-laboratory validation of an ELISA for the determination of serum anti-ganglioside antibodies. *European Journal of Neurology* 1999; (6): 71-77.
- Willison H J, Yuki N. Peripheral neuropathies and anti-glycolipid antibodies. *Brain* 2002; (125): 2591-2625.
- Willison H J. Ganglioside complexes: new autoantibody targets in Guillain-Barré syndromes. *Nat Clin Pract Neurol* 2005; (1): 2-3.
- Willison H J, Plomp J J. Anti-ganglioside antibodies and the presynaptic motor nerve terminal. *Ann N Y Acad Sci* 2008; (1132): 114-123.
- Winer J B, Hughes R A, Anderson M J, Jones D M, Kangro H, Watkins R P. A prospective study of acute idiopathic neuropathy. II. Antecedent events. *J Neurol Neurosurg Psychiatry* 1988; (51): 613-618.
- Winer J B. Guillain-Barré syndrome. *BMJ* 2008; (337): a671.
- Wirguin I, Ifergane G, Almog Y, Lieberman D, Bersudsky M, Herishanu Y O. Presynaptic neuromuscular transmission block in Guillain-Barre syndrome associated with anti-GQ1b antibodies. *Neuromuscul Disord* 2002; (12): 292-293.
- Wood S J, Slater C R. The contribution of postsynaptic folds to the safety factor for neuromuscular transmission in rat fast- and slow-twitch muscles. *J Physiol* 1997; (500 (Pt 1)): 165-176.
- Wood S J, Slater C R. Safety factor at the neuromuscular junction. *Prog Neurobiol* 2001; (64): 393-429.
- Wu G, Xie X, Lu Z H, Ledeen R W. Cerebellar neurons lacking complex gangliosides degenerate in the presence of depolarizing levels of potassium. *Proc Natl Acad Sci U S A* 2001; (98): 307-312.
- Wu G, Lu Z H, Xie X, Ledeen R W. Susceptibility of cerebellar granule neurons from GM2/GD2 synthase-null mice to apoptosis induced by glutamate excitotoxicity and elevated KCl: rescue by GM1 and LIGA20. *Glycoconj J* 2004; (21): 305-313.
- Wu G, Lu Z H, Wang J, Wang Y, Xie X, Meyenhofer M F, Ledeen R W. Enhanced susceptibility to kainate-induced seizures, neuronal apoptosis, and death in mice lacking gangliotetraose gangliosides: protection with LIGA 20, a membrane-permeant analog of GM1. *J Neurosci* 2005; (25): 11014-11022.
- Wu G, Lu Z H, Obukhov A G, Nowycky M C, Ledeen R W. Induction of calcium influx through TRPC5 channels by cross-linking of GM1 ganglioside associated with alpha5beta1 integrin initiates neurite outgrowth. *J Neurosci* 2007; (27): 7447-7458.
- Wu G S, Vaswani K K, Lu Z H, Ledeen R W. Gangliosides stimulate calcium flux in neuro-2A cells and require exogenous calcium for neuritogenesis. *J Neurochem* 1990; (55): 484-491.
- Xiang S L, Zhong M, Cai F C, Deng B, Zhang X P. The sialic acid residue is a crucial component of C. jejuni lipooligosaccharide ganglioside mimicry in the induction Guillain-Barré syndrome. *J Neuroimmunol* 2006; (174): 126-132.
- Yamamoto N, Matsubara T, Sato T, Yanagisawa K. Age-dependent high-density clustering of GM1 ganglioside at presynaptic neuritic terminals promotes amyloid beta-protein fibrillogenesis. *Biochim Biophys Acta* 2008; (1778): 2717-2726.

-
- Yamashita T, Wada R, Proia R L. Early developmental expression of the gene encoding glucosylceramide synthase, the enzyme controlling the first committed step of glycosphingolipid synthesis. *Biochim Biophys Acta* 2002; (1573): 236-240.
- Yamashita T, Hashiramoto A, Haluzik M, Mizukami H, Beck S, Norton A, Kono M, Tsuji S, Daniotti J L, Werth N, Sandhoff R, Sandhoff K, Proia R L. Enhanced insulin sensitivity in mice lacking ganglioside GM3. *Proc Natl Acad Sci U S A* 2003; (100): 3445-3449.
- Yamashita T, Wu Y P, Sandhoff R, Werth N, Mizukami H, Ellis J M, Dupree J L, Geyer R, Sandhoff K, Proia R L. Interruption of ganglioside synthesis produces central nervous system degeneration and altered axon-glia interactions. *Proc Natl Acad Sci U S A* 2005; (102): 2725-2730.
- Yang L J, Zeller C B, Shaper N L, Kiso M, Hasegawa A, Shapiro R E, Schnaar R L. Gangliosides are neuronal ligands for myelin-associated glycoprotein. *Proc Natl Acad Sci U S A* 1996; (93): 814-818.
- Yang L J, Lorenzini I, Vajn K, Mountney A, Schramm L P, Schnaar R L. Sialidase enhances spinal axon outgrowth in vivo. *Proc Natl Acad Sci U S A* 2006; (103): 11057-11062.
- Yates A J, Rampersaud A. Sphingolipids as receptor modulators. An overview. *Ann N Y Acad Sci* 1998; (845): 57-71.
- Yowler B C, Schengrund C L. Botulinum neurotoxin A changes conformation upon binding to ganglioside GT1b. *Biochemistry* 2004; (43): 9725-9731.
- Yu R K, Usuki S, Ariga T. Ganglioside molecular mimicry and its pathological roles in Guillain-Barré syndrome and related diseases. *Infect Immun* 2006; (74): 6517-6527.
- Yuki N, Taki T, Inagaki F, Kasama T, Takahashi M, Saito K, Handa S, Miyatake T. A bacterium lipopolysaccharide that elicits Guillain-Barré syndrome has a GM1 ganglioside-like structure. *J Exp Med* 1993; (178): 1771-1775.
- Yuki N. Molecular mimicry between gangliosides and lipopolysaccharides of *Campylobacter jejuni* isolated from patients with Guillain-Barré syndrome and Miller Fisher syndrome. *J Infect Dis* 1997; (176): S150-S153.
- Yuki N. Infectious origins of, and molecular mimicry in, Guillain-Barré and Fisher syndromes. *Lancet Infect Dis* 2001; (1): 29-37.
- Yuki N, Yamada M, Koga M, Odaka M, Susuki K, Tagawa Y, Ueda S, Kasama T, Ohnishi A, Hayashi S, Takahashi H, Kamijo M, Hirata K. Animal model of axonal Guillain-Barré syndrome induced by sensitization with GM1 ganglioside. *Ann Neurol* 2001; (49): 712-720.
- Yuki N, Susuki K, Koga M, Nishimoto Y, Odaka M, Hirata K, Taguchi K, Miyatake T, Furukawa K, Kobata T, Yamada M. Carbohydrate mimicry between human ganglioside GM1 and *Campylobacter jejuni* lipooligosaccharide causes Guillain-Barré syndrome. *Proc Natl Acad Sci U S A* 2004; (101): 11404-11409.
- Yuki N. *Campylobacter* sialyltransferase gene polymorphism directs clinical features of Guillain-Barré syndrome. *J Neurochem* 2007b; (103 Suppl 1): 150-158.
- Yuki N. Ganglioside mimicry and peripheral nerve disease. *Muscle Nerve* 2007a; (35): 691-711.
- Zamir O, Charlton M P. Cholesterol and synaptic transmitter release at crayfish neuromuscular junctions. *J Physiol* 2006; (571): 83-99.
- Zhang G, Lopez P H, Li C Y, Mehta N R, Griffin J W, Schnaar R L, Sheikh K A. Anti-ganglioside antibody-mediated neuronal cytotoxicity and its protection by intravenous immunoglobulin: implications for immune neuropathies. *Brain* 2004; (127): 1085-1100.
- Zhu D, Xiong W C, Mei L. Lipid rafts serve as a signaling platform for nicotinic acetylcholine receptor clustering. *J Neurosci* 2006; (26): 4841-4851.
- Zitman F M, Todorov B, Jacobs B C, Verschuuren J J, Furukawa K, Furukawa K, Willison H J, Plomp J J. Neuromuscular synaptic function in mice lacking major subsets of gangliosides. *Neuroscience* 2008; (156): 885-897.
- Zitman F M, Todorov B, Verschuuren J J, Jacobs B C, Furukawa K, Furukawa K, Willison H J, Plomp J J. Neuromuscular synaptic transmission in aged ganglioside-deficient mice. *Neurobiol Aging* 2009; (doi:10.1016/j.neurobiolaging.2009.01.007).
- Zuo Y, Lubischer J L, Kang H, Tian L, Mikesh M, Marks A, Scofield V L, Maika S, Newman C, Krieg P, Thompson W J. Fluorescent proteins expressed in mouse transgenic lines mark subsets of glia, neurons, macrophages, and dendritic cells for vital examination. *J Neurosci* 2004; (24): 10999-11009.
- Zwirner J, Wittig A, Kremmer E, Gotze O. A novel ELISA for the evaluation of the classical pathway of complement. *J Immunol Methods* 1998; (211): 183-190.