



Universiteit
Leiden
The Netherlands

Chemical analysis of organic molecules in carbonaceous meteorites

Torrao Pinto Martins, Zita Carla

Citation

Torrao Pinto Martins, Z. C. (2007, January 24). *Chemical analysis of organic molecules in carbonaceous meteorites*. Retrieved from <https://hdl.handle.net/1887/9450>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/9450>

Note: To cite this publication please use the final published version (if applicable).



CHAPTER 3

Extraterrestrial nucleobases in the Murchison meteorite

Carbon-rich meteorites, carbonaceous chondrites, contain many biologically relevant organic molecules and delivered prebiotic material to the young Earth. We present for the first time compound-specific carbon isotope data indicating that measured purine and pyrimidine compounds are indigenous components of the Murchison meteorite. Carbon isotope ratios for uracil and xanthine of $\delta^{13}\text{C} = +44.5\text{‰}$ and $+37.7\text{‰}$, respectively, indicate a non-terrestrial origin for these compounds. These new results demonstrate that organic compounds, which are components of the genetic code in modern biochemistry, were already present in the early solar system and may have played a key role in life's origin.

Z. Martins, O. Botta, M. L. Fogel, M. A. Sephton, D. P. Glavin, J. S. Watson, J. P. Dworkin, A. W. Schwartz and Pascale Ehrenfreund (2006)
Proceedings of the National Academy of Sciences, submitted.

3.1 Introduction

One of the most fundamental discoveries of modern science is the role of nucleobases as components of nucleic acids, the central molecules in the storage, transcription and translation of life's genetic code (Watson and Crick 1953). Since the genetic code is an ancient feature of life, it is likely that nucleobases were utilized by Earth's earliest living systems (Dworkin *et al.* 2003). One potentially important source of these compounds is the extraterrestrial delivery of material to Earth by comets, asteroids and their fragments as well as interplanetary dust particles (IDPs) (Chyba and Sagan 1992). Carbonaceous chondrites, which contain many biologically relevant organic compounds (for reviews see e.g. Botta and Bada 2002; Sephton 2002) have been analysed for nucleobases by several different research groups (Hayatsu 1964; Hayatsu *et al.* 1968, 1975; Folsome *et al.* 1971, 1973; Lawless *et al.* 1972; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981; Shimoyama *et al.* 1990). Hayatsu (1964) reported on the detection of N-heterocyclic compounds from an acetylated acid-hydrolysed Orgueil sample. The nucleobases adenine and guanine, as well as triazines (ammeline and melamine) were identified using paper chromatography. Later, Hayatsu *et al.* (1968) used HCl-hydrolysis but no acetylation in their analytical procedure to prevent the alteration or destruction of organic compounds during the acetylation step applied in their previous work. Guanylurea and the purine adenine were identified in the Orgueil meteorite, but no guanine was detected. Analysis by a different group (Folsome *et al.* 1971, 1973; Lawless *et al.* 1972) using gas chromatography-mass spectrometry (GC-MS) of trimethylsilyl (TMS) derivatives produced clearly different results, with no purines, triazines or guanylurea detected in water and formic acid extracts of the Orgueil, Murchison and Murray meteorites. However, Folsome *et al.* (1971, 1973) and Lawless *et al.* (1972) detected 4-hydroxypyrimidine in the same meteorite extracts. Hayatsu *et al.* (1975) analysed Murchison samples using the same extraction procedure as Folsome *et al.* (1971, 1973) and detected only aliphatic amines and alkylpyridines by direct sample volatilisation in a mass spectrometer. However, when more "drastic" extraction conditions were applied (3-6 M HCl or trifluoroacetic acid), Hayatsu *et al.* (1975) found the purines adenine and guanine, guanylurea and triazines, but no 4-hydroxypyrimidine, and this suggested that purines and triazines were released by acid hydrolysis from the meteorite macromolecular material. Two years later, van der Velden and Schwartz (1977) analysed a sample of the Murchison meteorite using high performance liquid chromatography (HPLC) with UV spectroscopy, a technique that did not require derivatisation or volatilisation prior to analysis. With this technique xanthine was detected at an abundance of 2.4 parts-per-million (ppm) in formic acid extracts; guanine and hypoxanthine were also tentatively identified (with a concentration of 0.1 ppm and 0.04 ppm, respectively), while no pyrimidines were found at levels higher than the background (0.01 ppm) in water or formic acid extracts. However, after silylation hydroxypyrimidines appeared in the water extracts leading to the suggestion that the compounds detected by Folsome *et al.* (1971, 1973) and Lawless *et al.* (1972) might have been produced from contaminants present in the silylation reagents (van der Velden and Schwartz 1977). Stoks and Schwartz (1979) re-analysed water and formic acid extracts of the Murchison, Murray and Orgueil meteorites using specific fractionation techniques (including charcoal columns) and ion exclusion chromatography with UV

spectroscopy and detected for the first time the nucleobase uracil, in the extracts of all these meteorites. Further re-analysis of the formic acid extracts of Murchison, Murray and Orgueil using GC, HPLC and mass spectrometry (MS) yielded in the detection of xanthine, adenine, hypoxanthine and guanine in concentrations ranging from 114 to 655 parts-per-billion (ppb) in all three meteorites (Stoks and Schwartz 1981). In contrast, hydroxypyrimidines and triazines were not identified above detection limit of 10 ppb and 50 ppb, respectively, suggesting that previous identifications of triazines by Hayatsu (1964) and Hayatsu *et al.* (1968, 1975) may have been artefacts synthesized during the experimental procedure (Stoks and Schwartz 1981). Shimoyama *et al.* (1990) detected guanine, and possible xanthine and hypoxanthine in the Antarctic meteorites Yamato (Y-) 74662 and Y-791198 using HPLC with UV spectroscopy. The concentrations of guanine ranged from 30 to 420 ppb, and no pyrimidines were found. Two other Antarctic meteorites, Y-793321 and Belgica (B-) 7904, also yielded no nucleobases (Shimoyama *et al.* 1990).

Several circumstantial lines of evidence suggested that the origin of the nucleobases present in carbonaceous meteorites is extraterrestrial (e.g. high relative xanthine abundance, low thymine to uracil ratio, and low abundance of cytosine) (van der Velden and Schwartz 1977). However, significant quantitative and qualitative variations even between different fragments of the same meteorite (Folsome *et al.* 1971, 1973; Hayatsu *et al.* 1975; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981), left open the possibility that terrestrial contamination at the fall site, or during the curation history of the meteorites, as well as analytical artefacts during the extraction, purification and derivatisation procedures, could have produced these compounds (van der Velden and Schwartz 1977).

Compound specific stable isotope compositions of hydrogen, carbon and nitrogen can be powerful discriminators of the origin of organic compounds in meteorites. For example, the ^{13}C isotope enrichment of amino acids and carboxylic acids in the Murchison meteorite has been critical to establish the extraterrestrial origin of these compounds (see e.g. Yuen *et al.* 1984; Engel *et al.* 1990; Pizzarello *et al.* 2004; Huang *et al.* 2005). Accordingly, to establish the origin (terrestrial vs. extraterrestrial) of the nucleobases in Murchison, the carbon isotopic ratio of these compounds must be determined. Compound specific isotope measurements of nucleobases in carbonaceous meteorites have not previously been reported. We subjected the Murchison meteorite (and appropriate controls) to a well-established extraction and isolation procedure (van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981) and supplemented it by analysing the extracts for the first time with modern compound-specific carbon isotope ratio instrumentation. For comparison, a soil sample collected in 1999 in the proximity of the meteorite's 1969 fall site was also subjected to the same extraction and isolation and analytical procedure. To our knowledge, soil from the Murchison meteorite fall site was not collected in 1969. Analysis of soil samples collected in the proximity of meteorite falls were previously shown to be critical in assessing the extent of terrestrial organic contamination in meteorites (e.g. Glavin *et al.* 1999). In this study, nucleobases were identified in Murchison meteorite and soil extracts using gas chromatography-quadrupole mass spectrometry (GC-QMS), and their carbon isotope ratios were

determined by gas chromatography-combustion-isotope ratio mass spectrometry (GC-C-IRMS).

3.2 Materials and Methods

3.2.1 Chemicals and reagents

Purine and pyrimidine standards comprised cytosine, uracil, hypoxanthine, guanine, xanthine, thymine (Sigma-Aldrich, $\geq 97\%$ purity) and adenine (Merck, $\geq 99\%$ purity). A stock solution of these nucleobases was prepared by dissolving each nucleobase in 0.1 N ammonium hydroxide. Ammonium formate, ammonium hydroxide, HPLC grade water, bis(trimethylsilyl)-trifluoroacetamide with 1% trimethylchlorosilane (BSTFA/TMCS), and anhydrous pyridine were purchased from Sigma-Aldrich. Formic acid p.a. (assay $\geq 99\%$) was purchased from Acros Organics, HPLC-S gradient grade acetonitrile from Biosolve Ltd., hydrochloric acid (37%) from Merck, and diethyl ether from Riedel-de-Haën. AG 50W-X8 cation exchange resin (100-200 mesh) was purchased from Bio-Rad and activated coconut charcoal (50-200 mesh) from Fisher. All glassware and ceramics used for sample processing were sterilized by annealing in aluminium foil at 500°C for 3 h. Biopur tips (Fisherbrand) and Eppendorf microcentrifuge tubes (Sigma-Aldrich) were supplied sterilised.

3.2.2 Extraction and cleaning procedure

A modification of previously published methods (van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981) was applied to our protocol for isolation, extraction and analysis of nucleobases. An interior piece of about 15 g of Murchison meteorite as well as 15 g of soil collected near the Murchison recovery location and a serpentine sample blank (heated to 500°C for 3 h) used as a procedural were separately crushed into powder using a ceramic mortar and pestle. Murchison meteorite powder and soil were placed separately inside Pyrex culture tubes (with Teflon lined screw caps), 1 g per tube. Samples were extracted by ultrasonication with formic acid (8 ml per tube, 3 times) for 1 h at 60°C. After centrifugation, the acid supernatants were transferred to 15 ml Pyrex tubes and dried under vacuum. Both meteorite and soil formic acid extracts were dissolved in 15 ml 1 M HCl, and added separately to columns of 0.6 x 5 cm activated charcoal. The extracts were then dried under vacuum and hydrolysed as described elsewhere (Stoks and Schwartz 1979, 1981). The hydrolysed extracts were diluted with 1 M HCl and extracted with ether, followed by charcoal cleaning of the aqueous fraction. The extracts were then dried under vacuum, dissolved in H₂O and subjected to ion-exchange separation with columns (0.4 x 6 cm) of 50W-X8 resin. Uracil and thymine eluted from these columns with H₂O, cytosine and all purines eluted with 5 M HCl. Both eluates were dried under vacuum.

3.2.3 Determination of the nucleobase recoveries

The efficiency of the cleaning process was tested by determining the yields of recovery for the different steps involved (charcoal filtration, hydrolysis and ion-exchange

separation) using solutions of standards of known concentrations. These results as well as the total nucleobase recoveries are displayed in Appendix (Table A 3.1). Results and technical implications are discussed in Appendix.

Charcoal filtration. The recoveries were determined by passing mixtures of nucleobase standards (5 µg of guanine and 4 µg each of uracil, thymine, xanthine, hypoxanthine and adenine, and a second mixture of nucleobase standards ten times more diluted) in 5 ml of 1 M HCl through columns of 0.6 x 5 cm activated charcoal beds. These columns were washed with 5 ml 1 M HCl and 5 ml H₂O each. The charcoal columns were first eluted with 5 ml of formic acid, and an additional elution of 5 ml of formic acid was collected. All collected fractions were dried under vacuum, brought up in 0.1 N NH₄OH and analysed by HPLC-UVS.

Acid hydrolysis. To test the hydrolysis step, individual nucleobase standards solutions were placed separately in different test tubes and then dried under vacuum. The amounts of nucleobases in the test tubes were 4 µg each of guanine and xanthine, and 5 µg of cytosine, uracil, thymine, hypoxanthine and adenine. The samples were hydrolysed with 8 ml 3 M HCl, at 110°C for 18 h. Subsequently, samples were dried, brought up in 0.1 N NH₄OH and analysed by HPLC-UVS.

Ion-exchange separation. The recoveries were determined by loading separate 50W-X8 resin columns (0.4 x 6 cm) each with 5 ml H₂O solutions containing 4 µg of guanine and xanthine, and 5 µg each of cytosine, uracil, thymine, hypoxanthine and adenine. The columns were first eluted two times with 5 ml of H₂O, then two times with 5 ml of 5 M HCl. All eluates were dried, re-dissolved in 0.1 N NH₄OH and analysed for purines and pyrimidines by HPLC-UVS.

HPLC-UVS analysis. HPLC separation was achieved using a dC18 reverse phase AtlantisTM column (Waters, 4.6 x 150 mm, 5 µm), at room temperature and a flow rate of 1 ml min⁻¹. Buffer A was 10 mM ammonium formate buffer and buffer B was acetonitrile. The gradient used was 0 to 0.01 min., 0% buffer B; 0.01 to 15 min., 0 to 3% buffer B; 15 to 15.01 min., 3 to 0% buffer B; 15.01 to 25 min., 0 % buffer B. UV absorption was monitored with diode array detection on a Shimadzu SPD-M10A_{VP}, (spectral acquisition range from 190 nm to 370 nm). Peaks were integrated using the maximum absorbance wavelength for each nucleobase: 257 nm for uracil, 248 nm for hypoxanthine, 246 nm for guanine, 265 nm for xanthine, 263 nm for thymine and 259 nm for adenine. Nucleobases were identified by comparison of the retention times of each peak at the respective wavelength with those of a standard.

3.2.4 GC-QMS analysis

The meteorite and soil extracts were dissolved in 500 µl of 0.1 N NH₄OH and 30 µl aliquots were dried under vacuum. 10 µl of anhydrous pyridine and 30 µl of BSTFA/TMCS were added to the dried extract residues. Derivatisation was carried out at 100°C for 90 min. 2 µl of the resulting solutions were each injected onto a GC-QMS (Thermo Finnigan Trace GC coupled to a Thermo Finnigan Trace DSQ QMS). Splitless

injection with He as carrier gas at a constant pressure of 13 PSI was used. Separation was performed on a HP Ultra 2 (25 m x 0.32 mm ID x 0.17 μ m film thickness) column. The GC oven temperature was held for 1 min at 75°C and ramped to 300°C at a rate of 5°C min⁻¹ and then held for 5 min. The presence of nucleobases was confirmed by retention time comparison to standards and by their unique mass fragmentation pattern.

3.2.5 GC-C-IRMS analysis

300 μ l aliquots of the meteorite and soil extracts (out of the 500 μ l) were carried through the same procedure as described for GC-QMS analysis. Carbon isotope analyses were performed using a Thermo Finnigan MAT Delta Plus XL GC-C-IRMS. Temperature program, carrier gas and pressure were the same as the GC-QMS analysis. The GC column in the GC-QMS was removed and then installed in the GC-C-IRMS. Compounds separated by the GC column were converted to CO₂ through an oxidation oven kept at 980°C. CO₂ reference gas with a known $\delta^{13}\text{C}$ value (-41.10‰ PDB) was injected via the interface to the IRMS, for the computation of $\delta^{13}\text{C}$ values of sample and standard compounds. Standards for the analyses included pyrene, with a $\delta^{13}\text{C}$ value of -24.03‰ (± 0.16 ‰) when measured by GC-C-IRMS. Additionally, individual nucleobases standards were subjected to the entire derivatisation procedure described above and run on the GC-C-IRMS, with typical standard deviation of ± 0.44 ‰. Corrections for carbon added from the BSFTA were calculated by mass balance: $\delta^{13}\text{C}$ nucleobase in sample derivatised = (% of carbon nucleobase) ($\delta^{13}\text{C}$ nucleobase sample) + (% of carbon BSTFA) ($\delta^{13}\text{C}$ BSTFA). The average $\delta^{13}\text{C}$ values used for BSTFA were -48.99‰ $\pm$ 0.1‰ (from uracil standards), -44.91‰ $\pm$ 0.38‰ (from thymine standards), and -40.47‰ $\pm$ 0.34‰ (from xanthine standards), and were obtained by mass balance: $\delta^{13}\text{C}$ nucleobase standard derivatised = (% of carbon nucleobase) (EA nucleobase standard) + (% of carbon BSTFA) ($\delta^{13}\text{C}$ BSTFA), where the EA nucleobase standard value corresponds to the $\delta^{13}\text{C}$ value of the nucleobase standard established by a Carlo Erba elemental analyser (EA)-IRMS with He as the carrier gas. The uncertainties in the $\delta^{13}\text{C}$ values (δx) are based on the standard deviation of the average value of between three and four separate measurements (N) with a standard error $\delta x = \sigma_x \cdot N^{-1/2}$.

3.3 Results and Discussion

Following the literature protocol (van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981), an interior fragment of the Murchison meteorite was extracted and purified for nucleobase isotopic analyses. This procedure substantially limited the presence of interfering compounds and was optimised for the detection of uracil and xanthine (see section 3.2 for details). A detailed study of the yields of recovery for each cleaning step during the purification process is described in section 3.2 (Materials and Methods) and Appendix. A relatively large quantity of Murchison meteorite (15 g) was necessary to perform carbon isotope measurements of nucleobases (see section 3.2). The limit of detection of the GC-C-IRMS, combined with the limited availability of meteorite samples, prevented us from performing stable nitrogen or hydrogen measurements of the nucleobases.




Fig. 3.1 - GC-C-IRMS analysis of the BSTFA-derivatised formic acid extract, water eluate of the Murchison meteorite. (a) The m/z 44 ($^{12}\text{CO}_2$) trace (bottom) and the ratio between the m/z 45 and m/z 44 ($^{13}\text{CO}_2/^{12}\text{CO}_2$) trace (top) for the GC-C-IRMS analysis are displayed. The insets show the uracil region of each chromatogram. The following peaks were tentatively identified by GC-QMS: 1. 2-hydroxyhexanoic acid; 2. butanedioic acid; 3. 2-methylbutanedioic acid; 4. unidentified; 5. 2,3-dimethylbutanedioic acid; 6. pentanedioic acid; 7. 2-methylpentanedioic acid; 8. 3-methylpentanedioic acid; 9. 3-ethylpentanedioic acid; 10. ethylpentanedioic acid; 11. hexanedioic acid; 12. heptanedioic acid; 13. 1,2-benzenedicarboxylic acid; 14. unidentified; 15. unidentified. (b) The GC-QMS mass spectrum for the peak assigned to BSTFA-derivatised uracil and its structure. The inset shows the mass spectrum of a BSTFA-derivatised uracil standard.




Fig. 3.2 - GC-C-IRMS analysis of the BSTFA-derivatised formic acid extract, hydrochloric acid eluate of the Murchison meteorite. (a) The m/z 44 ($^{12}\text{CO}_2$) trace (bottom) and the ratio between the m/z 45 and m/z 44 ($^{13}\text{CO}_2/^{12}\text{CO}_2$) trace (top) for the GC-C-IRMS analysis are displayed. The insets show the xanthine region of each chromatogram. The following peaks were tentatively identified by GC-QMS: 1. butanedioic acid; 2. 2-methylbutanedioic acid; 3. 2,3-dimethylbutanedioic acid; 4. pentanedioic acid; 5. 2-methylpentanedioic acid; 6. 3-methylpentanedioic acid; 7. unidentified; 8. 1,2-benzenedicarboxylic acid; 9. unidentified; 10. unidentified. (b) The GC-QMS mass spectrum for the peak assigned to BSTFA-derivatised xanthine and its structure. The inset shows the mass spectrum of a BSTFA-derivatised xanthine standard.

Table 3.1 - $\delta^{13}\text{C}$ values (‰) of nucleobases in the Murchison meteorite and soil samples.

$\delta^{13}\text{C}$	Uracil	Xanthine	Thymine
Murchison meteorite	$+44.5 \pm 2.3$	$+37.7 \pm 1.6$	n.d.
Soil	-10.6 ± 1.8	n.d.	-15.9 ± 1.1

n.d. - not determined due to low concentrations.

Two chromatographic traces obtained from the GC-C-IRMS analysis of the water eluate from the ion-exchange separation of the formic acid extract of the Murchison meteorite are shown in Fig. 3.1a: the m/z 44 ($^{12}\text{CO}_2$) trace (bottom) and the mass ratio 45/44 ($^{13}\text{CO}_2/^{12}\text{CO}_2$) (top). These traces include the peak corresponding to BSTFA-derivatised uracil, assigned by retention time comparison with BSTFA-derivatised authentic uracil standard analysed on the same instrument. Confirmation of this assignment was achieved by comparison of the mass fragmentation patterns of the corresponding peak in the meteorite extracts (Fig. 3.1b) with the mass spectra of a BSTFA-derivatised authentic uracil standard (Fig. 3.1b inset), analysed by GC-QMS using the same GC column and analytical conditions that were used for the GC-C-IRMS measurements. The same analysis was performed for the hydrochloric acid eluate from the ion-exchange separation. The two traces m/z 44 (bottom) and mass ratio 45/44 (top) that include the peak corresponding to BSTFA-derivatised xanthine are shown in Fig. 3.2a. The corresponding GC-QMS mass fragmentation patterns of the peaks in the meteorite extract and the authentic xanthine standard are shown in Fig. 3.2b. The unexpected peak shapes in the 45/44 traces (top in Fig. 3.1a and 3.2a) can be explained by the unavailability of GC-columns optimised for nucleobase compounds, and in particular for isotope measurements, as the 45/44 traces of BSTFA-derivatised authentic nucleobase standards analysed under the same conditions showed the same behaviour.

Analysis of the data from the GC-C-IRMS measurements yielded $\delta^{13}\text{C}$ values of $+44.5\text{‰}$ ($\pm 2.3\text{‰}$) for uracil and $+37.7\text{‰}$ ($\pm 1.6\text{‰}$) for xanthine in the Murchison meteorite (Table 3.1). These values fall within the range of those measured for extraterrestrial amino acids and carboxylic acids in carbonaceous chondrites (Yuen *et al.* 1984; Engel *et al.* 1990; Pizzarello *et al.* 2004; Huang *et al.* 2005). In order to constrain the contributions of terrestrial nucleobases to the carbon isotope values for uracil and xanthine measured in Murchison, GC-C-IRMS analyses were also carried out for the nucleobases present in the Murchison soil extract (Table 3.1). Soil uracil has a $\delta^{13}\text{C}$ value of -10.6‰ ($\pm 1.8\text{‰}$) and xanthine was below the detection limit of the GC-C-IRMS (~ 1 ppb). Thus, there should be no terrestrial contribution from the landing site soil to the value for xanthine measured in the meteorite. For uracil, any terrestrial contamination from the soil would decrease the measured $\delta^{13}\text{C}$ value in the Murchison meteorite extract. While our analytical methods were not optimised for the detection of other nucleobases, we were able to detect thymine in the soil ($\delta^{13}\text{C} = -15.9\text{‰} \pm 1.1\text{‰}$). The negative $\delta^{13}\text{C}$ values measured for uracil and thymine in the soil are in the range expected for terrestrial organic compounds of biological origin (for review see e.g. Sephton and Botta 2005) and are clearly distinct from the positive $\delta^{13}\text{C}$ values of uracil and xanthine we have measured in the Murchison meteorite.

Table 3.2 - $\delta^{13}\text{C}$ values (‰) of dicarboxylic acids in the Murchison meteorite.

Dicarboxylic acids	This study	Literature*
Butanedioic acid	+30.1	+28.1
2-Methylbutanedioic acid	+28.0	+26.5
Pentanedioic acid	+44.0	+26.8
2-Methylpentanedioic acid	+34.2	+27.9
3-Methylpentanedioic acid	+28.0	+19.1
Hexanedioic acid	+28.4	+21.4

*Pizzarello and Huang (2002)

Extraterrestrial dicarboxylic acids are the most abundant class of compounds detected in the Murchison meteorite extracts. Their measured $\delta^{13}\text{C}$ values were in the range of +28‰ to +44‰ (see Table 3.2 and section 3.3.1), consistent with previous results (Pizzarello and Huang 2002). These compounds were chromatographically separated from the nucleobases and therefore did not interfere with our carbon isotope measurements. Comparison of the mass fragmentation patterns of the meteorite extracts to standards indicates the possible presence of co-eluting compounds with the BSTFA-derivatised xanthine peaks. Of these, the most conspicuous interfering peak is the m/z 313 present in the Murchison xanthine spectrum (Fig. 3.2b). This mass fragment corresponds to BSTFA-derivatised hexadecanoic acid (a monocarboxylic acid), which is also observed at the same retention time in the soil extract, and has m/z 313 (Fig. 3.3). Thus, it is very likely that this compound in the meteorite has a terrestrial origin and carries a light isotopic signature. In addition, the high carbon number of hexadecanoic acid (C_{16}) is inconsistent with known extraterrestrial meteoritic monocarboxylic acids that range from C_2 up to C_{12} (Naraoka *et al.* 1999; Huang *et al.* 2005 and references given).

Based on these arguments, the measured carbon isotope values for uracil (possible contribution of terrestrial uracil from the soil) and xanthine (possible co-elution of terrestrial hexadecanoic acid) in the Murchison meteorite should be considered to be lower limits. Given the positive $\delta^{13}\text{C}$ values for uracil and xanthine measured for Murchison meteorite extracts, these interferences do not compromise the conclusion that these two nucleobases are of extraterrestrial origin.

3.3.1 Carboxylic acids in the Murchison meteorite and soil samples

The identification (by retention time and mass fragmentation patterns) of several peaks in the GC-QMS total ion current (TIC) was essential to determine whether the nucleobase peak assignments (both in the Murchison meteorite and in the soil) were correct and if the nucleobase peaks were separate and distinct from other compounds. The detection and carbon isotope composition of most of the dicarboxylic acids present in the Murchison meteorite have already been published (Lawless *et al.* 1974; Pelzer *et al.* 1984; Cronin *et al.* 1993; Pizzarello and Huang 2002). Despite the extensive purification procedure applied to isolate nucleobases from other compounds in the meteorite and soil extracts, GC-QMS analyses show that dicarboxylic acids are still present in the purified formic acid extracts of both the Murchison meteorite and soil. It is




Fig. 3.3 - GC-QMS mass spectrum corresponding to hexadecanoic acid present in the Murchison soil. The structure of BSTFA-derivatised hexadecanoic acid is shown.

not clear if additional purification steps would have removed these interferences. Moreover, additional processing would have led to further sample loss preventing the stable carbon isotope measurements of nucleobases. The presence of dicarboxylic acids in the eluates did not interfere with the isotopic analysis of the nucleobases since they were chromatographically separated on the GC-C-IRMS column. Dicarboxylic acids present on the GC-C-IRMS traces for both the H₂O eluate (Fig. 3.1a) and the HCl eluate of the Murchison meteorite (Fig. 3.2a) have been identified by GC-QMS as butanedioic acid, 2-methylbutanedioic acid, 2,3-dimethylbutanedioic acid, pentanedioic acid, 2-methylpentanedioic acid, 3-methylpentanedioic acid, and 1,2-benzenedicarboxylic acid (peaks 2, 3, 5 to 8, and 13 in the H₂O eluate, Fig. 3.1a; peaks 1 to 6, and 8 in the HCl eluate, Fig. 3.2a). Stable carbon isotope values of dicarboxylic acids present in the Murchison meteorite obtained in a previous study range from +19.1‰ to +28.1‰ (Pizzarello and Huang 2002). The $\delta^{13}\text{C}$ values for the Murchison meteorite dicarboxylic acids measured in this study are in agreement with these literature values, or are slightly higher (Table 3.2). The only exception is pentanedioic acid, whose $\delta^{13}\text{C}$ value of +44.0‰ is significantly higher than the $\delta^{13}\text{C}$ value of +26.8‰ published previously for Murchison (Pizzarello and Huang 2002). The difference between the two measurements could be due to a higher degree of terrestrial contamination in the sample of Murchison from the previous measurement, since pentanedioic acid is a common terrestrial contaminant found in the biosphere (Pizzarello and Huang 2002), leading to a small decrease in the $\delta^{13}\text{C}$ value. We cannot exclude the possibility of a small amount of an isotopically heavy compound co-eluting with pentanedioic acid in our analysis, which would increase the carbon isotope value for pentanedioic acid.

It is generally accepted that extraterrestrial nucleobases could have been formed by abiotic reaction mechanisms in a variety of cosmic environments. However, there is no evidence for this compound class in the interstellar and circumstellar medium (Kuan *et al.* 2003). Synthetic processes on the meteorite parent body during aqueous alteration are

more likely to be responsible for the presence of meteoritic nucleobases. A number of abiotic synthetic routes have been investigated in laboratory simulations. These include the polymerization of hydrogen cyanide (HCN) (Voet and Schwartz 1983; Schwartz and Baker 1989; Levy *et al.* 1999; Miyakawa *et al.* 2002), synthesis by quenching a CO-N₂-H₂O high-temperature plasma (Miyakawa *et al.* 2000), the reaction of cyanoacetylene with cyanate (Ferris *et al.* 1968) in relative dilute solution at pH 8 and room temperature, and reaction of cyanoacetaldehyde with urea in eutectic solution (Nelson *et al.* 2001). Other pathways are obviously also possible and a number of them might have occurred on the Murchison meteorite parent body. Degradation of nucleobases in the hydrated parent body environment also has to be considered. For example, cytosine degrades to uracil with a half-life of 17,000 years and guanine decomposes to xanthine with a half-life of 1.3 Ma (Levy and Miller 1998) at 0°C and pH 7. Consequently, meteoritic nucleobase distributions are the result of both synthetic and subsequent degradation reactions.

3.4 Conclusion

By demonstrating that one purine and one pyrimidine in the Murchison meteorite are extraterrestrial in origin, all of the key component classes in terrestrial biochemistry, including amino acids, sugar related compounds (Cooper *et al.* 2001), carboxylic acids and nucleobases, have been identified as indigenous components in the Murchison meteorite (for a review see e.g. Botta and Bada 2002; Sephton 2002). Our data advance proposals that life's raw materials were delivered to the early Earth and other planetary bodies by exogenous sources, including carbonaceous meteorites. In contrast, the endogenous synthesis of prebiotic organic compounds may have been constrained by the conditions on the young Earth, perhaps most importantly by the oxidation state of the atmosphere. For example, only low yields of amino acids were produced under non-reducing conditions in the Miller-Urey-type experiment (Stribling and Miller 1987). Yet, whatever the inventory of endogenous organic compounds on the ancient Earth, it would have been augmented by extraterrestrial material. It is estimated that these sources delivered $\sim 10^9$ kg of carbon per year to the Earth during the heavy bombardment phase 4.5–3.9 billion years ago (Chyba and Sagan 1992).

In modern biology uracil is ubiquitous as a nucleobase in RNA while the role of xanthine is limited in modern biochemistry (Kulikowska *et al.* 2004); most notably as an intermediate in the biosynthesis of guanosine and uric acid. It is also interesting to note that both xanthine and uracil are capable of self-association in monolayers, which might have been of importance in prebiotic chemistry on mineral surfaces on the early Earth (Sowerby and Petersen 1999). A continuous influx of meteoritic uracil and xanthine and possibly other nucleobases would have enriched the prebiotic organic inventory necessary for life to assemble on the early Earth. Following the birth of the solar system carbonaceous meteorite infall would have been common on all terrestrial planets. Consequently, nucleobases delivered to these worlds with sugar-related species and amino acids could have been critical to the origin of life on Earth, Mars, or elsewhere.

Acknowledgements

This project was supported by Fundação para a Ciência e a Tecnologia (scholarship SFRH/BD/10518/2002), NASA Astrobiology Institute, the NASA Exobiology and Evolutionary Biology Program and through cooperative agreement NNA04CC09A, Goddard Center for Astrobiology/NASA Astrobiology, European Space Agency, NWO-VI 016023003 and PPARC. The authors would like to thank Dr. Linda Welzenbach (Smithsonian National Museum of Natural History, Washington DC) for providing us with a Murchison sample and Zan Peeters for graphic support.

Appendix

Technical implications

Preliminary analysis of formic acid extracts of Murchison by HPLC-UVS showed the presence of underlying UV absorbing material, namely abundant aliphatic and aromatic carboxylic acids (identified by GC-MS), which interfered with the detection of the nucleobases. For this reason an isolation process subsequent to the formic acid extraction was performed based on the method used by van der Velden and Schwartz (1977) and Stoks and Schwartz (1979, 1981).

Minimal amounts of solvents were used in the isolation steps in order to minimize sample loss and contamination. Results show that it is necessary to use 10 ml of formic acid to completely elute 5 µg of guanine and 4 µg of uracil, thymine, xanthine, hypoxanthine and adenine from the charcoal columns. The overall average recovery (Table A 3.1) of the two solutions, one containing 5 µg of guanine and 4 µg of all the other nucleobases, and a solution diluted by a factor of ten, was 71% (values ranging from 47% for guanine to 97% for hypoxanthine). Our value is identical to that of Stoks and Schwartz (1981), who found an average nucleobases recovery for the formic acid extract after charcoal cleaning of 71%. Stoks and Schwartz (1979, 1981) used much larger volumes of solvent, which led to a more time consuming process with no apparent gain in the nucleobases recovery.

The average nucleobase recovery for acid hydrolysis was 88% (Table A 3.1), with values for each nucleobase of 81% for thymine, 82% for xanthine, 83% for adenine (hydrolysed to hypoxanthine), 85% for cytosine (hydrolysed to uracil), 95% for guanine (hydrolysed to xanthine) and hypoxanthine, and 96% for uracil. Our average recovery of purines and pyrimidines after 3 M HCl hydrolysis is substantially better than the 60% obtained by Stoks and Schwartz (1981).

Our analysis revealed that 5 ml of H₂O was necessary to elute 5 µg of uracil and 5 µg of thymine from the 50W-X8 resin, and that for all other nucleobases (4 µg of guanine and xanthine, and 5 µg each of cytosine, hypoxanthine and adenine) 10 ml of 5 M HCl was required. The overall average recovery was 72% (Table A 3.1) with values ranging from

Table A 3.1 - Summary of the percentage of nucleobase standard material recovered after each separate cleaning step (charcoal filtration, hydrolysis and ion-exchange separation) during the purification process of the Murchison meteorite extract and soil. Values are not cumulative.

Cleaning Step Nucleobases	Charcoal cleaning	Hydrolysis	Ion-exchange separation
Cytosine	n.d.	85	96
Hypoxanthine	97	95	69
Guanine	47	95	57
Xanthine	72	82	48
Adenine	53	83	78
Uracil	88	96	86
Thymine	69	81	68
NR	71	88	72

n.d. - not determined

NR (nucleobase recovery) - This value corresponds to the percentage of the yields of nucleobases after each specific purification step.

The total nucleobase recovery (TNR) was calculated by considering that we applied the charcoal filtration step twice, the hydrolysis step once and the ion-exchange step once. $TNR \text{ (in percentage)} = 71\% \text{ (charcoal filtration)} \times 88\% \text{ (hydrolysis)} \times 71\% \text{ (charcoal cleaning)} \times 72\% \text{ (ion-exchange)} = 32\%$

48% for xanthine to 96% for cytosine.

The overall recovery of nucleobases estimated from these individual steps was 32% (see Table A 3.1).

References

- Botta O. and Bada J. L. 2002. *Surv. Geophys.* 23, 411-467.
- Chyba C. and Sagan C. 1992. *Nature* 355, 125-132.
- Cooper G., Kimmich N., Belisle W., Sarinana J., Brabham K. and Garrel L. 2001. *Nature* 414, 879-883.
- Cronin J. R., Pizzarello S., Epstein S. and Krishnamurthy R. V. 1993. *Geochim. Cosmochim. Acta* 57, 4745-4752.
- Dworkin J. P., Lazcano A. and Miller S. L. 2003. *J. Theor. Evol.* 222, 127-134.
- Engel M. H., Macko S. A. and Silfer J. A. 1990. *Nature* 348, 47-49.
- Ferris J. P., Sanchez R. A. and Orgel L. E. 1968. *J. Mol. Biol.* 33, 693-704.
- Folsome C. E., Lawless J., Romiez M. and Ponnampuruma C. 1971. *Nature* 232, 108-109.
- Folsome C. E., Lawless J., Romiez M. and Ponnampuruma C. 1973. *Geochim. Cosmochim. Acta* 37, 455-465.
- Glavin D. P., Bada J. L., Brinton K. L. F. and McDonald G. D. 1999. *Proc. Natl. Acad. Sci. USA* 96, 8835-8838.
- Hayatsu R. 1964. *Science* 146, 1291-1293.
- Hayatsu R., Studier M. H., Oda, A., Fuse K. and Anders E. 1968. *Geochim. Cosmochim. Acta* 32, 175-190.
- Hayatsu R., Studier M. H., Moore L. P., and Anders E. 1975. *Geochim. Cosmochim. Acta* 39, 471-488.
- Huang Y., Wang Y., Alexandre M. R., Lee T., Rose-Petruck C., Fuller M. and Pizzarello S. 2005. *Geochim. Cosmochim. Acta* 69, 1073-1084.
- Kuan Y.-J., Yan C.-H., Charnley S. B., Kisiel Z., Ehrenfreund P., Huang H.-C. 2003. *Mon. Not. R. Astron. Soc.* 345, 650-656.
- Kulikowska E., Kierdaszuk B. and Shugar D. 2004. *Acta Biochim. Polon.* 51, 493-531.
- Lawless J. G., Folsome C. E. and Kvenvolden K. 1972. *Sci. Amer.* 226, 38-46.
- Lawless J. G., Zeitman B., Pereira W. E., Summons R. E. and Duffield A. M. 1974. *Nature* 251, 40-42.
- Levy M. and Miller S. L. 1998. *Proc. Natl. Acad. Sci. USA* 95, 7933-7938.
- Levy M., Miller S. L. and Oró J. 1999. *J. Mol. Evol.* 49, 165-168.
- Miyakawa S., Murasawa K., Kobayashi K. and Sawaoka A. B. 2000. *Orig. Life Evol. Biosph.* 30, 557-566.
- Miyakawa S., Cleaves H. J. and Miller S. L. 2002. *Orig. Life Evol. Biosph.* 32, 209-218.
- Naraoka H., Shimoyama A. and Harada K. 1999. *Orig. Life Evol. Biosph.* 29, 187-201.
- Nelson K. E., Robertson M. P., Levy M. and Miller S. L. 2001. *Orig. Life Evol. Biosph.* 31, 221-229.
- Peltzer E. T., Bada J. L., Schlesinger G. and Miller S. L. 1984. *Adv. Space Res.* 4, 69-74.
- Pizzarello S. and Huang Y. 2002. *Meteorit. Plan. Science* 37, 687-696.
- Pizzarello S., Huang Y. and Fuller M. 2004. *Geochim. Cosmochim. Acta* 68, 4963-4969.
- Schwartz A. W. and Bakker C. G. 1989. *Science* 245, 1102-1104.
- Sephton M. A. 2002. *Nat. Prod. Rep.* 19, 292-311.
- Sephton M. A. and Botta O. 2005. *Int. J. Astrobio.* 4, 269-276.
- Shimoyama A., Hagishita S. and Harada K. 1990. *Geochem. J.* 24, 343-349.
- Sowerby S. J. and Petersen G. B. 1999. *Orig. Life Evol. Biosph.* 29, 597-614.

Chapter 3

- Stoks P. G. and Schwartz A. W. 1979. *Nature* 282, 709-710.
- Stoks P. G. and Schwartz A. W. 1981. *Geochim. Cosmochim. Acta* 45, 563-569.
- Stribling R. and Miller S. 1987. *Orig. Life* 17, 261-273.
- Van Der Velden W. and Schwartz A. W. 1977. *Geochim. Cosmochim. Acta* 41, 961-968.
- Voet A. B. and Schwartz A. W. 1983. *Bioorg. Chem.* 12, 8-17.
- Watson J. D. and Crick F. H. C. 1953. *Nature* 171, 737-738.
- Yuen G., Blair N., Des Marais D. J. and Chang S. 1984. *Nature* 307, 252-254.