



Universiteit
Leiden
The Netherlands

Statistical methods for analysing complex genetic traits

El Galta, Rachid

Citation

El Galta, R. (2006, September 27). *Statistical methods for analysing complex genetic traits*. Retrieved from <https://hdl.handle.net/1887/4574>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/4574>

Note: To cite this publication please use the final published version (if applicable).

Bibliography

- Abecasis, G. R., S. S. Cherny, W. O. Cookson, and L. R. Cardon (2002). Merlin-rapid analysis of dense genetic maps using sparse gene flow trees. *Nature Genetics* 30, 97–101.
- Achenbach, T., S. McConaughy, and C. Howell. Child/adolescent behavioral and emotional problems: implications of cross-informant correlations for situational specificity. *Psychol Bull* 101, 213–132.
- Amos, C. I. (1994). Robust variance-components approach for assessing genetic-linkage in pedigrees. *American Journal of Human Genetics* 54, 535–543.
- Amos, C. I. and M. de Andrade (2001). Genetic linkage methods for quantitative traits. *Statistical Methods in Medical Research* 10, 3–25.
- Amos, C. I. and R. C. Elston (1989). Robust methods for the detection of genetic-linkage for quantitative data from pedigrees. *Genetic Epidemiology* 6, 349–360.
- Anderson, T. (1984). *An introduction to multivariate statistical analysis*. Second edition, Jhon Wiley & Sons, Inc.
- Aulchenko, Y. S., N. Vaessen, P. Heutink, J. Pullen, P. J. L. M. Snijders, A. Hofman, L. A. Sandkuijl, J. J. Honwing-Duistermaat, M. Edwards, S. Bennett, B. A. Oostra, and C. M. van Duijn (2003). A genome-wide search for genes involved in type 2 diabetes in a recently genetically isolated population from the netherlands. *Diabetes* 52, 3001–3004.
- Bacanu, S. (2005). Multipoint linkage analysis for a very dense set of markers. *Genetic Epidemiology* 29, 195–203.
- Bafna, V., D. Gusfield, G. Lancia, and S. Yooseph (2003). Haplotyping as perfect phylogeny: A direct approach. *Journal of Computational Biology* 10, 323–340.
- Bailey-Wilson, J. E. Almas, L., M. de Andrade, J. Bailey, H. Bickeböllner, H. J. Cordell, E. W. Daw, L. Goldin, E. L. Goode, C. Gray-McGuire, W. Hening, G. Jarvik, B. S. Mather, N. Mendell, A. D. Paterson, J. Rice, G. Satten, B. Suarez, V. Vieland, M. Wilcox, H. Zhang, A. Ziegler, and J. W. MacCluer (2005). Genetic analysis workshop 14: Microsatellite and snp marker loci for genome-wide scans. *BMC Genetics* 6, S1.
- Barkley, R. (1996). *Attention deficit/hyperactivity disorder. In: Child Psychopathology*. New York: Guilford Press.
- Barkley, R. A. (2003). Issues in the diagnosis of attention-deficit/hyperactivity disorder in children. *Brain and Development* 25(2), 77–83.

Bibliography

- Beaty, T. H. and K. Y. Liang (1987). Robust inference for variance-components models in families ascertained through probands .1. conditioning on probands phenotype. *Genetic Epidemiology* 4, 203–210.
- Becker, T., S. Cichon, and M. Jonson, E. Knapp (2005). Multiple testing in the context of haplotype analysis revisited: application to case-control data. *Annals of Human Genetics* 69, 747–56.
- Biernacka, J. (2004). *Statistical methods for studying two linked disease genes*. Ph. D. thesis, University of Toronto.
- Biernacka, J. M., L. Sun, and S. B. Bull (2005). Simultaneous localization of two linked disease susceptibility genes. *Genetic Epidemiology* 28, 33–47.
- Bonifati, V., P. Rizzu, F. Squitieri, E. Krieger, N. Vanacore, J. C. van Swieten, A. Brice, C. M. van Duijn, B. Oostra, G. Meco, and P. Heutink (2003). Dj-1 (park7), a novel gene for autosomal recessive, early onset parkinsonism. *Neurological Sciences* 24(3), 159–160.
- Bonney, G. E. (1986). Regressive logistic-models for familial disease and other binary traits. *Biometrics* 42, 611–625.
- Cannings, C. and E. A. Thompson (1977). Ascertainment in sequential sampling of pedigrees. *Clinical Genetics* 12, 208–212.
- Cantwell, D. P. (1996). Attention deficit disorder: A review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry* 35(8), 978–987.
- Cardon, L. R. and D. W. Fulker (1994). The power of interval mapping of quantitative trait loci, using selected sib pairs. *American Journal of Human Genetics* 55, 825–833.
- Carey, G. and J. Williamson (1991). Linkage analysis of quantitative traits increased power by using selected samples. *American Journal of Human Genetics* 49, 786–796.
- Carlson, C. S., S. F. Aldred, P. K. Lee, R. P. Tracy, S. M. Schwartz, M. Rieder, K. A. Liu, O. D. Williams, C. Iribarren, E. C. Lewis, M. Fornage, E. Boerwinkle, M. Gross, C. Jaquish, D. A. Nickerson, R. M. Myers, D. S. Siscovick, and A. P. Reiner (2005). Polymorphisms within the c-reactive protein (crp) promoter region are associated with plasma crp levels. *American Journal of Human Genetics* 77, 64–77.
- Clark, A. G. (2004). The role of haplotypes in candidate gene studies. *Genetic Epidemiology* 27, 321–333.
- Clayton, D. (2000). Linkage disequilibrium mapping of disease susceptibility genes in human populations. *International Statistical Review* 68, 23–43.
- Clayton, D., J. Chapman, and J. Cooper (2004). Use of unphased multilocus genotype data in indirect association studies. *Genetic Epidemiology* 27, 415–428.
- Collins, F. S., M. S. Guyer, and A. Chakravarti (1997). Variations on a theme: Cataloging human dna sequence variation. *Science* 278, 1580–1581.

- Commenges, D., H. Jacqmin, L. Letenneur, and C. M. van Duijn (1995). Score test for familial aggregation in probands studies: application to alzheimers-disease. *Biometrics* 51, 542–551.
- Cordell, H. J. (2001). Sample size requirements to control for stochastic variation in magnitude and location of allele-sharing linkage statistics in affected sibling pairs. *Annals of Human Genetics* 65, 491–502.
- Cordell, H. J. and D. G. Clayton (2005). Genetic associaton studies. *Lancet* 366, 1121–31.
- Cox, D. and D. V. Hinkley (1974). *Theoretical statistics*. Florida, USA: Chapman & Hall.
- Crystal, D. S., R. Ostrander, R. S. Chen, and G. J. August (2001). Multimethod assessment of psychopathology among dsm-iv subtypes of children with attention-deficit/hyperactivity disorder: Self-, parent, and teacher reports. *Journal of Abnormal Child Psychology* 29(3), 189–205.
- Czika, W. and B. Weir (2004). Properties of the multiallelic trend test. *Biometrics* 52, 69–74.
- Daly, M. J., J. D. Rioux, S. E. Schaffner, T. J. Hudson, and E. S. Lander (2001). High-resolution haplotype structure in the human genome. *Nature Genetics* 29, 229–232.
- Day, N. E. and M. J. Simons (1976). Disease susceptibility genestheir identification by multiple case family studies. *Tissue Antigens* 8, 109–119.
- de Andrade, M. and C. I. Amos (2000). Ascertainment issues in variance components models. *Genetic Epidemiology* 19, 333–344.
- de Nijs, P. F. A., R. E. Ferdinand, E. I. de Bruin, M. C. J. Dekker, C. M. van Duijn, and F. C. Verhulst (2004). Attention-deficit/hyperactivity disorder (adhd): parents' judgment about school, teachers' judgment about home. *European Child and Adolescent Psychiatry* 13(5), 315–320.
- Devlin, B. and N. A. Risch (1995). A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics*. 29, 311–22.
- Diggle, P. J., L. K. Y. and S. L. Zegger (1994). *Analysis of longitudinal data*. Oxford: Clarendon Press, Oxford Science Publication.
- Dudoit, S. and T. P. Speed (2000). A score test for the linkage analysis of qualitative and quantitative traits based on identity by descent data from sib-pairs. *Biostatistics* 1, 1–26.
- El Galta, R., T. Stijnen, and J. Houwing-Duistermaat (2004). Score statistic for analysis of association between disease and a multi-allelic marker [abstract]. *Genetic Epidemiology* 27, 268.
- Elston, R., J. Olson, and L. Palmer (Eds.) (2002). *Biostatistical Genetics and Genetic epidemiology*. John Wiley & Sons, Ltd.

Bibliography

- Elston, R. C. (2000). Introduction and overview. *Statistical Methods in Medical Research* 9, 527–541.
- Elston, R. C. and E. Sobel (1979). Sampling considerations in the gathering and analysis of pedigree data. *American Journal of Human Genetics* 31, 62–69.
- Epstein, M. P. and G. A. Satten (2003). Inference on haplotype effects in case-control studies using unphased genotype data. *American Journal of Human Genetics* 73, 1316–1329.
- Ewens, W., R. Griffiths, S. Ethier, S. Wilcox, and J. Graves (1992). Statistical analysis of in situ hybridization data: derivation and use of the zmax test. *Genomics*. 12, 675–82.
- Ewens, W. J. and N. C. E. Shute (1986). A resolution of the ascertainment sampling problem. i. theory. *Theoretical Population Biology* 30, 388–412.
- Faraone, S. V., J. Biederman, E. Mick, A. E. Doyle, T. Wilens, T. Spencer, E. Frazier, and K. Mullen (2001). A family study of psychiatric comorbidity in girls and boys with attention-deficit/hyperactivity disorder. *Biological Psychiatry* 50(8), 586–592.
- Faraone, S. V., J. Biederman, and M. C. Monuteaux (2000). Toward guidelines for pedigree selection in genetic studies of attention deficit hyperactivity disorder. *Genetic Epidemiology* 18(1), 1–16.
- Feingold, E. (2001). Methods for linkage analysis of quantitative trait loci in humans. *Theoretical Population Biology* 60, 167–180.
- Feingold, E., P. O. Brown, and D. Siegmund (1993). Gaussian models for genetic-linkage analysis using complete high-resolution maps of identity by descent. *American Journal of Human Genetics* 53, 234–251.
- Ferdinand, R. and J. van der Ende (2000). Nimh disc-iv: Diagnostic interview schedule for children. *Authorized Dutch translation.: Erasmus MC-Sophia: Rotterdam, the Netherlands*.
- Ferdinand, R. F., K. N. Hoogerheide, J. van der Ende, J. H. Visser, H. M. Koot, M. C. Kasius, and F. C. Verhulst (2003). The role of the clinician: three-year predictive value of parents', teachers', and clinicians' judgment of childhood psychopathology. *Journal Of Child Psychology And Psychiatry And Allied Disciplines* 44(6), 867–876.
- FitzGerald, P. E. B. and M. W. Knuiman (2000). Use of conditional and marginal odds-ratios for analysing familial aggregation of binary data. *Genetic Epidemiology* 18, 193–202.
- Forabosco, P., M. Falchi, and M. Devoto (2005). Statistical tools for linkage analysis and genetic association studies. *Expert Review of Molecular Diagnostics* 5, 781–796.
- Fullerton, S. M., A. G. Clark, K. M. Weiss, D. A. Nickerson, S. L. Taylor, J. H. Stengard, V. Salomaa, E. Vartiainen, M. Perola, E. Boerwinkle, and C. F. Sing (2000). Apolipoprotein e variation at the sequence haplotype level: Implications for the ori-

- gin and maintenance of a major human polymorphism. *American Journal of Human Genetics* 67, 881–900.
- Gabriel, S. B., S. F. Schaffner, H. Nguyen, J. M. Moore, J. Roy, B. Blumenstiel, J. Higgins, M. DeFelice, A. Lochner, M. Faggart, S. N. Liu-Cordero, C. Rotimi, A. Adeyemo, R. Cooper, R. Ward, E. S. Lander, M. J. Daly, and D. Altshuler (2002). The structure of haplotype blocks in the human genome. *Science* 296, 2225–2229.
- Gaub, M. and C. L. Carlson (1997). Gender differences in adhd: A meta-analysis and critical review. *Journal of the American Academy of Child and Adolescent Psychiatry* 36(8), 1036–1045.
- Green, J. R. and J. C. Woodrow (1977). Sibling method for detecting hla-linked genes in disease. *Tissue Antigens* 9, 31–35.
- Greenberg, D. A., J. Zhang, D. Shmulewitz, L. J. Strug, R. Zimmerman, V. Singh, and S. Marathe (2005). Construction of the model for the genetic analysis workshop 14 simulated data: genotype-phenotype relationships, gene interaction, linkage, association, disequilibrium, and ascertainment effects for a complex phenotype. *BMC Genetics* 6 (Suppl 1), S1, in press.
- Gudbjartsson, D. F., K. Jonasson, M. L. Frigge, and A. Kong (2000). Allegro, a new computer program for multipoint linkage analysis. *Nature Genetics* 25, 12–13.
- Haldane, J. (1939). The mean and variance of χ^2 when used as a test of homogeneity, when expectations are small. *Biometrika* 31, 346–355.
- Haseman, J. K. and R. C. Elston (1972). Investigation of linkage between a quantitative trait and a marker locus. *Behavior Genetics* 2, 3–19.
- Hastbacka, J., A. Delachapelle, I. Kaitila, P. Sistonen, A. Weaver, and E. Lander (1992). Linkage disequilibrium mapping in isolated founder populations - diastrophic dysplasia in finland. *Nature Genetics* 2, 204–211.
- Hill, J. C. and E. P. Schoener (1996). Age-dependent decline of attention deficit hyperactivity disorder. *American Journal of Psychiatry* 153(9), 1143–1146.
- Hofman, A., D. E. Grobbee, P. T. V. M. De Jong, and F. A. van den Ouweland (1991). Determinants of disease and disability in the elderly the rotterdam elderly study. *European Journal of Epidemiology* 7, 403–422.
- Hopper, J. L. and J. D. Mathews (1982). Extensions to multivariate normal-models for pedigree analysis. *Annals of Human Genetics* 46, 373–383.
- Houwing-Duistermaat, J. J., C. Bijkerk, L. Hsu, T. Stijnen, E. P. Slagboom, and C. M. van Duijn (2003). A unified approach to modelling linkage to quantitative and qualitative traits. *Annals of Human Genetics* 67, 457–463.
- Houwing-Duistermaat, J. J., B. H. F. Derkx, F. R. Rosendaal, and H. C. van Houwelingen (1995). Testing familial aggregation. *Biometrics* 51, 1292–1301.

Bibliography

- Houwing-Duistermaat, J. J. and R. C. Elston (2001). Linkage disequilibrium mapping of complex genetic diseases using multiallelic markers. *Genetic Epidemiology* 21S, 527–81.
- Johnson, R. and D. Wichern (1998). *Applied multivariate statistical analysis*. New jersey, USA: Prentice-Hall.
- Kizawa, H., I. Kou, A. Iida, A. Sudo, Y. Miyamoto, A. Fukuda, A. Kotani, A. Kawakami, S. Yamamoto, K. Uchida, A. Nakamura, K. Notoya, Y. Nakamura, and S. Ikegawa (2005). An aspartic acid repeat polymorphism in asporin inhibits chondrogenesis and increases susceptibility to osteoarthritis. *Nature Genetics* 37, 138–44.
- Koster, T., F. R. Rosendaal, H. Deronde, E. Briet, J. P. van den Broucke, and R. M. Bertina (1993). Venous thrombosis due to poor anticoagulant response to activated protein-cleiden thrombophilia study. *Lancet* 342, 1503–1506.
- Kruglyak, L., M. J. Daly, M. P. ReeveDaly, and E. S. Lander (1996). Parametric and nonparametric linkage analysis: A unified multipoint approach. *American Journal of Human Genetics* 58, 1347–1363.
- Lander, E. and L. Kruglyak (1995). Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nature Genetics* 11, 241–247.
- Lander, E. S. and N. J. Schork (1994). Genetic dissection of complex traits. *Science* 265, 2037–2048.
- Lange, K. (2002). *Mathematical and statistical methods for genetic analysis*. Second edition, Springer-Verlag.
- Lazzeroni, L. C. (2001). A chronology of fine-scale gene mapping by linkage disequilibrium. *Statistical Methods in Medical Research* 10, 57–76.
- le Cessie, S. and H. C. van Houwelingen (1995). Testing the fit of a regression-model via score tests in random effects models. *Biometrics* 51, 600–614.
- Lebecq, J., H. Putter, and J. C. van Houwelingen (2005). Potential bias in generalized estimating equations linkage methods under incomplete information. *To appear in Genetic Epidemiology*.
- Levy, F. and Hay, D., M. McStephen, C. Wood, and I. Waldman (1997). Attention-deficit hyperactivity disorder: a category or a continuum? genetic analysis of a large-scale twin study. *J Am Acad Child Adolesc Psychiatry* 36, 737–44.
- Liang, K. Y. and T. H. Beaty (1991). Measuring familial aggregation by using odds-ratio regression-models. *Genetic Epidemiology* 8, 361–370.
- Liang, K. Y. and T. H. Beaty (2000). Statistical designs for familial aggregation. *Statistical Methods in Medical Research* 9, 543–562.
- Liang, K. Y., Y. F. Chiu, and T. H. Beaty (2001). A robust identity-by-descent procedure using affected sib pairs: Multipoint mapping for complex diseases. *Human*

- Heredity* 51, 64–78.
- Liang, K. Y. and S. L. Zeger (1986). Longitudinal data-analysis using generalized linear-models. *Biometrika* 73, 13–22.
- Lopera, F., L. Palacio, I. Jimenez, P. Villegas, I. Puerta, D. Pineda, M. Jimenez, and M. Arcos-Burgos (1999). [discrimination between genetic factors in attention deficit] discriminacion de factores geneticos en el deficit de atencion. *Rev Neurol* 28, 660–664.
- Martin, N., J. Scourfield, and P. McGuffin (2002). Observer effects and heritability of childhood attention-deficit hyperactivity disorder symptoms. *British Journal of Psychiatry* 180, 260–265.
- McCullagh, P. and J. Nelder (1989). *Generalized linear models*. Second edition, London: Chapman and Hall.
- Meulenbelt, I., C. Bijkerk, S. C. De Wildt, H. S. Miedema, F. C. Breedveld, H. A. Pols, A. Hofman, C. M. Van Duijn, and P. E. Slagboom (1999). Haplotype analysis of three polymorphisms of the col2a1 gene and associations with generalised radiological osteoarthritis. *Annals of Human Genetics* 63, 393–400.
- Mitsis, E. M., K. E. McKay, K. P. Schulz, J. H. Newcorn, and J. M. Halperin (2000). Parent-teacher concordance for dsm-iv attention-deficit/hyperactivity disorder in a clinic-referred sample. *Journal of the American Academy of Child and Adolescent Psychiatry* 39(3), 308–313.
- Morton, N. E. (1959). Genetic tests under incomplete ascertainment. *American Journal of Human Genetics* 11, 1–16.
- Neuman, R. J., A. Heath, W. Reich, K. K. Bucholz, P. A. F. Madden, L. W. Sun, R. D. Todd, and J. J. Hudziak (2001). Latent class analysis of adhd and comorbid symptoms in a population sample of adolescent female twins. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 42(7), 933–942.
- Niu, T. H. (2004). Algorithms for inferring haplotypes. *Genetic Epidemiology* 27, 334–347.
- Ott, J. (1999). *Analysis of human genetic linkage*. Third edition, Johns Hopkins University Press, Baltimore.
- Pfeiffer, R. M., M. H. Gail, and D. Pee (2001). Inference for covariates that accounts for ascertainment and random genetic effects in family studies. *Biometrika* 88, 933–948.
- Putter, H., L. A. Sandkuijl, and J. C. van Houwelingen (2002). Score test for detecting linkage to quantitative traits. *Genetic Epidemiology* 22, 345–355.
- Rao, D. C. and R. Wette (1987). Nonrandom sampling in genetic epidemiology maximum-likelihood methods for multifactorial analysis of quantitative data ascertained through truncation. *Genetic Epidemiology* 4, 357–376.
- Rao, D. C., R. Wette, and W. J. Ewens (1988). Multifactorial analysis of family data as-

Bibliography

- certained through truncation - a comparative-evaluation of 2 methods of statistical-inference. *American Journal of Human Genetics* 42, 506–515.
- Reich, D. E. and E. S. Lander (2001). On the allelic spectrum of human disease. *Trends Genet.* 17, 502–10.
- Risch, N. (1990a). Linkage strategies for genetically complex traits .I. multilocus models. *American Journal of Human Genetics* 46, 222–228.
- Risch, N. (1990b). Linkage strategies for genetically complex traits .II. the power of affected relative pairs. *American Journal of Human Genetics* 46, 229–241.
- Risch, N. (1990c). linkage strategies for genetically complex traits .III. the effect of marker polymorphism on analysis of affected relative pairs. *American Journal of Human Genetics* 46, 242–253.
- Risch, N. (2000). Searching for genes in complex diseases: lessons from systemic lupus erythematosus. *Journal of Clinical Investigation* 105, 1503–1506.
- Risch, N. and H. P. Zhang (1995). Extreme discordant sib pairs for mapping quantitative trait loci in humans. *Science* 268, 1584–1589.
- Roks, G. P. and, v. S. J., M. Cruts, E. Boeren, L. Sandkuijl, P. Sniijders, C. Van Broeckhoven, B. Oostra, and C. van Duijn (2001). *Alzheimer's disease in a Dutch recent genetically isolated population. The GRIP Study, in Alzheimer's disease. A genetic epidemiologic approach.* Ph. D. thesis, p. 83-90. Erasmus MC: Rotterdam.
- Sasieni, P. D. (1997). From genotypes to genes: doubling the sample size. *Biometrics* 53, 1253–61.
- Satten, G. A. and M. P. Epstein (2004). Comparison of prospective and retrospective methods for haplotype inference in case-control studies. *Genetic Epidemiology* 27, 192–201.
- Schaid, D. J. (2004). Evaluating associations of haplotypes with traits. *Genetic Epidemiology* 27, 348–364.
- Schaid, D. J., C. M. Rowland, D. E. Tines, R. M. Jacobson, and G. A. Poland (2002). Score tests for association between traits and haplotypes when linkage phase is ambiguous. *American Journal of Human Genetics* 70, 425–434.
- Schaid, D. J., J. P. Sinnwell, and S. N. Thibodeau (2005). Robust multipoint identical-by-descent mapping for affected relative pairs. *American Journal of Human Genetics* 76, 128–138.
- Sebastiani, P., R. Lazarus, S. T. Weiss, L. M. Kunkel, I. S. Kohane, and M. F. Ramoni (2003). Minimal haplotype tagging. *Proceedings of the National Academy of Sciences of the United States of America* 100, 9900–9905.
- Self, S. and K. Liang (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under non-standard conditions. *JASA* 82, 605–610.

- Shaffer, D., P. Fisher, C. P. Lucas, M. K. Dulcan, and M. E. Schwab-Stone (2000). NIMH diagnostic interview schedule for children version iv (NIMH DISC-IV): Description, differences from previous versions, and reliability of some common diagnoses. *Journal of the American Academy of Child and Adolescent Psychiatry* 39(1), 28–38.
- Sham, P. C. (1998). *Statistics in human genetics*. New York: Oxford University Press Inc.
- Sham, P. C. and D. Curtis (1995). Monte Carlo tests for associations between disease and alleles at highly polymorphic loci. *Annals of Human Genetics* 59, 97–105.
- Sham, P. C., D. Curtis, and C. J. MacLean (1996). Likelihood ratio tests for linkage and linkage disequilibrium: asymptotic distribution and power. *American Journal of Human Genetics* 58, 1093–5.
- Shih, M. C. and A. S. Whittemore (2001). Allele-sharing among affected relatives: non-parametric methods for identifying genes. *Statistical Methods in Medical Research* 10, 27–55.
- Siegmund, D. (2001). Is peak height sufficient? *Genetic Epidemiology* 20, 403–408.
- Slager, S. L. and D. J. Schaid (2001). Case-control studies of genetic markers: Power and sample size approximations for Armitage's test for trend. *Human Heredity* 52, 149–153.
- Stram, D. O. (2004). Tag SNP selection for association studies. *Genetic Epidemiology* 27, 365–374.
- Stram, D. O., H. Lee, and D. C. Thomas (1993). Use of generalized estimating equations in segregation analysis of continuous outcomes. *Genetic Epidemiology* 10, 575–579.
- Teng, J. and D. Siegmund (1998). Multipoint linkage analysis using affected relative pairs and partially informative markers. *Biometrics* 54, 1247–1265.
- Terwilliger, J. D. (1995). A powerful likelihood method for the analysis of linkage disequilibrium between trait loci and one or more polymorphic marker loci. *American Journal of Human Genetics* 56, 777–87.
- Thapar, A., R. Harrington, K. Ross, and P. McGuffin (2000). Does the definition of ADHD affect heritability? *J Am Acad Child Adolesc Psychiatry* 39(6), 1528–1536.
- Thapar, A., A. Hervas, and P. McGuffin (1995). Childhood hyperactivity scores are highly heritable and show sibling competition effects: Twin study evidence. *Behaviour Genetics* 5, 537–544.
- Thapar, A., J. Holmes, K. Poulton, and R. Harrington (1999). Genetic basis of attention deficit and hyperactivity. *British Journal of Psychiatry* 174, 105–111.
- Thompson, E. A. (1975). Estimation of pairwise relationships. *Annals of Human Genetics* 39, 173–188.
- Todd, R. D., E. R. Rasmussen, R. J. Neuman, W. Reich, J. J. Hudziak, K. K. Bucholz,

Bibliography

- P. A. F. Madden, and A. Heath (2001). Familiarity and heritability of subtypes of attention deficit hyperactivity disorder in a population sample of adolescent female twins. *American Journal of Psychiatry* 158(11), 1891–1898.
- Tregouet, D. A. and L. Tiret (2000). Applications of the estimating equations theory to genetic epidemiology: a review. *Annals of Human Genetics* 64, 1–14.
- Tritchler, D., Y. Liu, and S. Fallah (2003). A test of linkage for complex discrete and continuous traits in nuclear families. *Biometrics* 59, 382–92.
- Uh, H. W., J. J. Houwing-Duistermaat, H. Putter, and van Houwelingen H. C. (2005). How to quantify information loss due to phase ambiguity in haplotype case-control studies. *BMC Genet.* 6 Suppl 1, S108.
- Uitte de Willige, S., M. C. de Visser, J. J. Houwing-Duistermaat, F. R. Rosendaal, H. L. Vos, and R. M. Bertina (2005). Genetic variation in the fibrinogen gamma gene increases the risk of deep venous thrombosis by reducing plasma fibrinogen gamma' levels. *Blood* 106, 4176–83.
- Vaessen, N., P. Heutink, J. J. Houwing-Duistermaat, P. J. L. M. Snijders, T. Rademaker, L. Testers, M. R. Batstra, L. A. Sandkuijl, C. M. van Duijn, and B. A. Oostra (2002). A genome-wide search for linkage-disequilibrium with type 1 diabetes in a recent genetically isolated population from the netherlands. *Diabetes* 51, 856–859.
- van der Ende, J. (1999). *Multiple informants: Multiple views*. In: Child psychiatric epidemiology. Accomplishments and future directions., Koot HM CA, Ferdinand RF, ed. Assen (The Netherlands): Van Gorcum & Company, pp 39-52.
- van der Meer, F. J. M., T. Koster, J. P. Vandenbroucke, E. Briet, and F. R. Rosendaal (1997). The leiden thrombophilia study (lets). *Thrombosis and Haemostasis* 78, 631–635.
- van Duijn, C., M. Dekker, V. Bonifati, R. Galjaard, J. Houwing-Duistermaat, P. Snijders, L. Testers, G. Breedveld, M. Horstink, L. Sandkuijl, J. van Swieten, B. Oostra, and P. Heutink (2001). Park7, a novel locus for autosomal recessive early-onset parkinsonism, on chromosome 1p36. *American journal of Human Genetics* 69, 629–34.
- Verbeke, G. and G. Molenberghs (1997). *Linear mixed models in practice: a sas-oriented approach*. New York: Springer-Verlag.
- Verhulst, F. C., H. M. Koot, and J. van der Ende (1994). Differential predictive value of parents and teachers reports of childrens problem behaviorsa longitudinal-study. *Journal of Abnormal Child Psychology* 22(5), 531–546.
- Verhulst, F. C., J. van der Ende, R. F. Ferdinand, and M. C. Kasius (1997). The prevalence of dsm-iii-r diagnoses in a national sample of dutch adolescents. *Archives of General Psychiatry* 54(4), 329–336.
- Wang, W. Y., B. J. Barratt, D. G. Clayton, and J. A. Todd (2005). Genome-wide associa-

- tion studies: theoretical and practical concerns. *Nature Reviews of Genetics* 6, 109–118.
- Whittemore, A. S. (1995). Logistic-regression of family data from case-control studies. *Biometrika* 82, 57–67.
- Wright, F. A. (1997). The phenotypic difference discards sib-pair qtl linkage information. *American Journal of Human Genetics* 60, 740–742.
- Zhang, H. P. and N. Risch (1996). Mapping quantitative-trait loci in humans by use of extreme concordant sib pairs: Selected sampling by parental phenotypes. *American Journal of Human Genetics* 59, 951–957.
- Zhao, H. Y. (2000). Family-based association studies. *Statistical Methods in Medical Research* 9, 563–587.
- Zhao, L. P., L. Hsu, S. Holte, Y. Chen, F. Quiaoit, and R. L. Prentice (1998). Combined association and aggregation analysis of data from case-control family studies. *Biometrika* 85, 299–315.
- Ziegler, A., C. Kastner, and M. Blettner (1998). The generalised estimating equations: an annotated bibliography. *Biometrical Journal* 40, 115–139.
- Zondervan, K. T. and L. R. Cardon (2004). The complex interplay among factors that influence allelic association. *Nat. Rev. Genet.* 5, 89–101.
- Zwetselaar, M. (2003). Pedkin, software to calculate kinship and inbreeding coefficients.

Bibliography

Curriculum Vitae

Rachid el Galta was born on 15 August 1972 in Elkbab, Morocco. In 1991, he obtained his Baccalaureate degree from Lycée technique, Beni Mellal, Morocco. In 1998 he graduated in mathematics at the University of Rabat, Morocco. In the same year he moved to the Netherlands. After he followed a Dutch language course for about eight months he started his master study in statistics at the University of Amsterdam. He also obtained a certificate for Dutch as second language (NT2). In the end of 2001 he received the MSc. degree in Business- and Industrial Statistics. In January 2002 he started the project described in this thesis at the department of Epidemiology and Biostatistics, Erasmus Medical Center, Rotterdam. During this period he followed several courses in statistical genetics. In December 2003 he moved along with the project to the department of Medical Statistics, Leiden University Medical Center, Leiden, where he completed this thesis. The content of this thesis has been presented in different international conferences, e.g. International Genetic Epidemiology Society Conference, New Orleans (2002), USA; Conference of the Royal Statistical Society (RSS 2003) - Statistical Genetics and Bioinformatics, Diepenbeek, Belgium; International Genetic Epidemiology Society Conference (2004), Noordwijkerhout, the Netherlands; IBS Multi-Regional Conference (2005), Leicester, UK; 25th meeting for statisticians (2005), Oslo, Norway. He also received the ANed Biometry Award 2006. In February 2006 he moved to London to join the Institute of Cancer Research, as scientific researcher.